Muon and Tau Anomalies Updated

Stephan Narison^a

^a Laboratoire de Physique Mathématique, Université de Montpellier II Place Eugène Bataillon, 34095 -Montpellier Cedex 05, France

and

Physics Division, National Center for Theoretical Sciences, Hsinchu, Taiwan 300, Republic of China. E-mail: qcd@lpm.univ-montp2.fr

We present a new independent evaluation of the hadronic vacuum polarization contributions $a_l^{had}(l.o) \equiv \frac{1}{2}(g-2)_l^{had}(l.o)$ to the anomalous magnetic moment (anomaly) of the muon and tau leptons using τ -decays and e^+e^- data. The alone theoretical input used for describing the high-energy region not accessible experimentally is perturbative QCD plus (negligible) additional effects due to the QCD vacuum condensates. We obtain: $a_{\mu}^{had}(l.o) = 7020.6(75.6) \times 10^{-11}$ and $a_{\tau}^{had}(l.o) = 353.6(3.8) \times 10^{-8}$, which we compare with previous determinations. Our analysis leads to the Standard Model (SM) prediction: $a_{\mu}^{SM} = 116$ 591 846.9(78.9) $\times 10^{-11}$. Confronting a_{μ}^{SM} with the recent BNL measurement leads to $a_{\mu}^{new} \equiv a_{\mu}^{exp} - a_{\mu}^{SM} = 176(170) \times 10^{-11}$. Combined with the mean existing determinations of $a_{\mu}^{had}(l.o)$, this leads to the conservative range: $-42 \leq a_{\mu}^{new} \times 10^{11} \leq 413$ at 90% CL, from which we derive lower bounds on the scales of some new physics. We also update our old and first predictions of the SM contributions to a_{τ} . Including QED to sixth order, higher order hadronic and electroweak contributions, we obtain $a_{\tau}^{SM} = 117$ 755(7) $\times 10^{-8}$, waiting for a future precise measurement of a_{τ} .

March, 19th 2001 PM/01-13 NSC-NCTS-010314

1. INTRODUCTION

The recent E821 BNL result [1] for the measurement of the muon anomaly averaged with older determinations [2,3] gives [4]:

$$a_{\mu}^{exp} = 116\ 592\ 023(151) \times 10^{-11} \ . \tag{1}$$

The announcement of a 2.6σ deviation [1,4] of this experimental result from the standard model (SM) prediction has stimulated intensive publications on the possible signal or/and constraints on new physics, but has also raised some criticisms [5]. In this paper, we consider that the only rational answer to the the criticisms of [5] is an independent re-evaluation of the hadronic vacuum polarisation contribution to the muon anomaly as an update of our old estimate in [6], which is the main motivation of this work. In order to avoid some specific theoretical dependences of the result, we shall mainly use the available data from τ decay and $e^+e^- \rightarrow$ hadrons, and limit ourselves to the use of perturbative QCD plus (negiligible) additional effects due to the QCD vacuum condensates for describing the high-energy region (QCD continuum) not accessible experimentally. We shall compare our results with previous determinations and analyze its implications for some new physics (e.g. supersymmetry, radiative muon, composite, extended Zee and leptoquark models) beyond the SM. Finally, we update our old and first [7] SM predictions for a_{τ} .

2. THE HADRONIC VACUUM POLARISA-TION CONTRIBUTION TO THE LEP-TON ANOMALY

Several papers have been devoted to the analysis of the contribution to the muon anomaly. Works prior 76 have been reviewed in [6], while more recent works (after 85) are reviewed in [4,5]. A partial historical review of the different determinations since 61 can be found in Figure 2 of [8], which we complete and update in Table 1, where only works published in journals with referee-policies have been considered. Using a disper-

$\mathbf{a}_{\mu}^{\mathbf{had}}(\mathbf{l.o}) imes\mathbf{10^{11}}$	Authors	Ref.	Comments $\pi^+\pi^-$ only	
≈ 3200	Bouchiat-Michel (61)	[9]		
$5500 \sim 11000$	Durand (62)	[10]	$\pi^+\pi^-$ only	
≈ 7500	Kinoshita-Oakes (67)	[11]	v	
3400^{+1900}_{-000}	Bowcock (68)	[12]		
6500(500)	Gourdin-de Rafael (69)	[13]		
6800(900)	Bramon-Etim-Greco (72)	[14]		
7300(1000)	Bailey et al. (75)	[15]		
6630(850)	Barger-Long-Olsson (75)	[16]		
6990(880)	Calmet-Narison-Perrottet-de Rafael (76&77)	[6]		
7020(800)*	Narison (78)	[7]		
6840(110)	Barkov et al. (85)	[17]		
7068(174)*	Kinoshita-Nizić-Ókamoto (85)	[18]		
7100(116)*	Casa-Lopez-Yndurain (85)	[19]		
7050(78)	Dubnicka-Martinovic (90)	[20]		
7024(153)*	Eidelman-Jegerlehner (95)	[21]		
7113(103)	Adel-Yndurain (95)	22		
7026(160)*	Brown-Worstell (96)	[23]		
6950(150)*	Alemany-Davier-Höcker (97)	[24]		
7011(94)	Alemany-Davier-Höcker (97)	[24]	+ au-decay	
6951(75)*	Davier-Höcker (98)	25	$+\tau$ -decay	
6924(62)	Davier-Höcker (98)	[26]	+Theoretical input	
7021(76)*	Narison (01)	This work	Average from e^+e^- and τ -decay (see section 3)	

Time and precision evolutions of the determinations of $a^{had}_{\mu}(l.o)$ from e^+e^- data. Results marked with * are our selected results which take into account update and results with minimal theoretical inputs.

Informative weighted mean values

 $6992(28)_{stat}$ $7008(41)_{stat}$

Table 1

sion relation, the hadronic vacuum polarisation contribution to the lepton anomaly a_l can be expressed as [9]-[13]:

$$a_l^{had}(l.o) = \frac{1}{4\pi^3} \int_{4m_\pi^2}^{\infty} dt \ K_l(t) \ \sigma_H(t) \ . \tag{2}$$

$$-K_l(t \ge 0)$$
 is the QED kernel function [27]:

$$K_l(t) = \int_0^1 dx \frac{x^2(1-x)}{x^2 + (t/m_l^2)(1-x)} , \qquad (3)$$

with the analytic forms:

$$\begin{aligned} K_l(t \le 4m_l^2) &= \frac{1}{2} - 4y_l - \\ &\quad 4y_l(1 - 2y_l)\log(4y_l) \\ &\quad -2\left(1 - 8y_l + 8y_l^2\right) \times \\ &\quad \sqrt{y_l(1 - y_l)}\arccos\sqrt{y_l} , \\ K_l(t \ge 4m_l^2) &= z_l^2\left(1 - \frac{z_l^2}{2}\right) + (1 + z_l)^2 \times \end{aligned}$$

Since 69 Selected results marked with *

$$\left(1 + \frac{1}{z_l^2}\right) \left[\log(1 + z_l) - z_l + \frac{z_l^2}{2}\right] + \left(\frac{1 + z_l}{1 - z_l}\right) z_l^2 \log z_l, \quad (4)$$

with:

$$y_l = \frac{t}{4m_l^2}, \quad z_l = \frac{(1-v_l)}{(1+v_l)} \text{ and } v_l = \sqrt{1-\frac{4m_l^2}{t}}.$$
 (5)

 $K_l(t)$ is a monotonically decreasing function of t. For large t, it behaves as:

$$K_l(t > m_l^2) \simeq \frac{m_l^2}{3t} , \qquad (6)$$

which will be useful for the analysis in the large t regime. Such properties then emphasize the importance of the low-energy contribution to $a_l^{had}(l.o)$ $(l \equiv e, \mu)$, where the QCD analytic calculations cannot be applied.

 $-\sigma_H(t) \equiv \sigma(e^+e^- \rightarrow \text{hadrons})$ is the $e^+e^- \rightarrow \text{hadrons}$

total cross-section which can be related to the hadronic two-point spectral function $\text{Im}\Pi(t)_{em}$ through the optical theorem:

$$R_{e^+e^-} \equiv \frac{\sigma(e^+e^- \to \text{hadrons})}{\sigma(e^+e^- \to \mu^+\mu^-)} = 12\pi \text{Im}\Pi(t)_{em} , \quad (7)$$

where:

$$\sigma(e^+e^- \to \mu^+\mu^-) = \frac{4\pi\alpha^2}{3t}.$$
(8)

Here,

$$\Pi_{em}^{\mu\nu} \equiv i \int d^4x \ e^{iqx} \ \langle 0 | \mathcal{T} J_{em}^{\mu}(x) \left(J_{em}^{\nu}(x) \right)^{\dagger} | 0 \rangle$$

= $- \left(g^{\mu\nu} q^2 - q^{\mu} q^{\nu} \right) \Pi_{em}(q^2)$ (9)

is the correlator built from the local electromagnetic current:

$$J^{\mu}_{em}(x) = \frac{2}{3}\bar{u}\gamma^{\mu}u - \frac{1}{3}\bar{d}\gamma^{\mu}d - \frac{1}{3}\bar{s}\gamma^{\mu}s + \dots$$
(10)

- In the following, we shall discuss in details the different hadronic contributions to $a_{\mu}^{had}(l.o)$. Our results will be given in Table 2.

3. HADRONIC VACUUM POLARISA-TION CONTRIBUTION TO THE MUON ANOMALY

3.1. LIGHT ISOVECTOR MESONS Region below 0.8 GeV²

Due to the *t*-behaviour of the kernel function $K_{\mu}(t)$, this region gives so far the most important contribution to $a_{\mu}^{had}(l.o) \approx 68\%$ of the total contribution), and also the largest source of the errors (77% when added quadratically). In our numerical analysis, we divide this region into three subregions.

- The first one is the region near the two pions threshold $4m_{\pi}^2 \leq t \leq 0.4 \text{ GeV}^2$, where the pion form factor $|F_{\pi}|^2(t)$ is constrained by universality $|F_{\pi}|^2(0) = 1$ and by the t expansion predicted by chiral perturbation theory:

$$|F_{\pi}|^{2}(t) \simeq 1 + \frac{1}{6} \langle r^{2} \rangle_{\pi} t + c_{\pi} t^{2} + \mathcal{O}(t^{3}) , \qquad (11)$$

or its resummed expression [28]. $\langle r^2 \rangle_{\pi} \simeq (0.431 \pm 0.026) \text{ fm}^2$ is the mean pion charge radius squared and $c_{\pi} \simeq (3.2 \pm 1.0) \text{ GeV}^4$ [29]. One can inspect that the ALEPH/OPAL [30,31] and e^+e^- data compiled in [21,24,34] satisfy both constraints.

- The second region is the one from 0.4 $\text{GeV}^2 \le t \le$ 0.6 GeV^2 on the top of the ρ resonance.

- For these two regions, we shall use CVC hypothesis which relates the electromagnetic to the charged current through an isospin rotation [32]:

$$\sigma_H(t) = \frac{4\pi\alpha^2}{t} v_1 \ . \tag{12}$$

We follow the notation of ALEPH [30], where:

$$\mathrm{Im}\Pi^{(1)}_{\bar{u}d,V} \equiv \frac{1}{2\pi} v_1 \ , \tag{13}$$

is the charged vector two-point correlator:

$$\Pi^{\mu\nu}_{\bar{u}d,V} \equiv i \int d^4x \ e^{iqx} \langle 0|\mathcal{T}J^{\mu}_{\bar{u}d}(x) \left(J^{\nu}_{\bar{u}d}(0)\right)^{\dagger} |0\rangle$$

$$= - \left(g^{\mu\nu}q^2 - q^{\mu}q^{\nu}\right) \Pi^{(1)}_{\bar{u}d,V}(q^2)$$

$$+ q^{\mu}q^{\nu}\Pi^{(0)}_{\bar{u}d,V}(q^2) , \qquad (14)$$

built from the local charged current $J^{\mu}_{\bar{u}d,V}(x) = \bar{u}\gamma^{\mu}d(x)$. We use the accurate semi-inclusive ALEPH/ OPAL data [30,31]. For a comparison, we also show the results obtained from the use of e^+e^- data in the whole region below 0.6 GeV².

- The third region is the one from 0.6 GeV² to 0.8 GeV², which is peculiar due to the $\omega - \rho$ mixing. In this region, we use either the τ -decay data with corrected $\omega - \rho$ mixing effect, or the e^+e^- data alone compiled in [24,30]. One can check from the data that the effect of the 4π is negligible in such a region.

Region from 0.8 to 3 GeV^2

For our fitting procedure, we use CVC and the ALEPH/OPAL semi-inclusive τ -decay data [30,31]. The dominant error from this region comes from the one between 2.1 and 3 GeV² (88% when the errors are added quadratically), due to the incaccuracy of the data near M_{τ} . For a comparison, we also show the results when using the sum of exclusive modes from e^+e^- compiled in [30,24], where in this case the errors come mainly from the region below 2 GeV².

3.2. LIGHT ISOSCALAR MESONS

ω and ϕ mesons

We treat these mesons in a narrow width approximation (NWA), which is expected to be a good approximation. Using the relation between the $e^+e^- \rightarrow$ hadrons total cross section and the leptonic width Γ_{ee} of the meson with a mass M_R :

$$\sigma_H(t)_{NWA} \simeq 12\pi^2 \frac{\Gamma_{ee}}{M_R} \delta(t - M_R^2) , \qquad (15)$$

one can write:

$$a_{\mu}^{had}(l.o)_{NWA} \simeq \frac{3}{\pi} \frac{\Gamma_{ee}}{M_R} K_{\mu}(M_R^2) , \qquad (16)$$

where, we shall use the PDG values of the electronic widths [33].

Region below 1.39^2 GeV²

We estimate the effect of this region by using the sum of exclusive I = 0 modes for the ratio $R_{e^+e^-}^{I=0}$ compiled in [34].

Region between 1.39^2 and **3** GeV²

In order to account for the multi-odd pions, $\bar{K}K$, $\bar{K}K\pi$,... modes, we estimate the effect of this region by assuming that it is mainly given by the $\omega(1419)$, $\omega(1662)$ and $\phi(1680)$, with their parameters measured by the DM2 [35] and DM1 [36] collaborations. We shall estimate their leptonic widths which are multiplied by the hadronic branching ratios in [35,36], by assuming (see PDG [33] and the indication from DM2 and DM1 data), that the $\omega(1419)$

Region in GeV ²	$\mathbf{a}_{\mu}^{\mathbf{had}}(\mathbf{l.o}) imes \mathbf{10^{11}}$		$\mathbf{a}_{ au}^{\mathbf{had}}(\mathbf{l.o}) imes \mathbf{10^8}$	Data input
	$\tau + \mathbf{e}^+ \mathbf{e}^-$	$\mathbf{e}^+\mathbf{e}^-$	$\tau + \mathbf{e}^+ \mathbf{e}^-$	
Light Isovector			•	
$4m_{\pi}^2 \rightarrow 0.8$	4794.6 ± 60.7	4730.2 ± 99.9	165.8 ± 1.5	[24, 30, 31]
$0.8 \xrightarrow{n} 2.1$	494.9 ± 15.8	565.0 ± 54.0	28.7 ± 1.1	[30,31]
$2.1 \rightarrow 3.$	202.0 ± 29.7	175.9 ± 16.0	17.0 ± 2.6	[30, 31]
Total Light I=1	5491.5 ± 69.4	5471.1 ± 114.7	211.5 ± 3.2	
Light Isoscalar				
Below 1.93				
ω	387.5 ± 13	387.5 ± 13	15.3 ± 0.5	NWA [33]
ϕ	393.3 ± 9.9	393.3 ± 9.9	21.0 ± 0.5	NWA 33
$0.66 \rightarrow 1.93$	79.3 ± 18.7	79.3 ± 18.7	4.3 ± 1.1	\sum exclusive [34]
From 1.93 to 3				
$\omega(1.42), \ \omega(1.65)$	31.3 ± 6.8	31.3 ± 6.8	2.6 ± 0.7	BW [35,33]
$\phi(1.68)$	42.4 ± 18.2	42.4 ± 18.2	3.8 ± 1.3	BW [35,36,33]
Total Light I=0	933.8 ± 31.5	933.8 ± 31.5	47.0 ± 2.0	
Heavy Isoscalar				
$J/\psi(1S \to 4.415)$	87.0 ± 4.7	87.0 ± 4.7	13.08 ± 0.69	NWA [33]
$\Upsilon(1S \rightarrow 11.020)$	0.95 ± 0.04	0.95 ± 0.04	0.23 ± 0.01	NWA [33]
Total Heavy I=0	88.0 ± 4.7	88.0 ± 4.7	13.3 ± 0.7	
QCD continuum				
$3. \to (4.57)^2$	407.0 ± 2.3	407.0 ± 2.3	49.4 ± 0.3	(u, d, s)
$(4.57)^2 \to (11.27)^2$	95.3 ± 0.5	95.3 ± 0.5	27.3 ± 0.1	(u, d, s, c)
$(11.27)^2 \to 4M_t^2$	20.5 ± 0.1	20.5 ± 0.1	5.87 ± 0.01	(u, d, s, c, b)
$4M_t^2 \to \infty$	$\approx 0.$	$\approx 0.$	$\approx 0.$	(u, d, s, c, b, t)
Total QCD Cont.	522.8 ± 2.4	522.8 ± 2.4	82.6 ± 0.3	
	7036.1(76.4)	7015.7(119.1)	354.4(3.8)	

Table 2 Determinations of $a_l^{had}(l.o)$ using combined e^+e^- and inclusive τ decay data (2nd and 4th columns) and averaged e^+e^- data (3rd column).

decays predominantly into $\rho\pi$ (we assume it to be approximately 90%); the $\omega(1662)$ decays mainly into $\rho\pi$ (44%) and $\omega\pi$ (56%), while the $\phi(1680)$ decays dominantly into K^*K (93%) and to $K\bar{K}$ (7%). In this way, we deduce the leptonic widths:

$$\Gamma_{\omega(1419)\to e^+e^-} \simeq (90 \pm 34) \text{ eV} ,
\Gamma_{\omega(1650)\to e^+e^-} \simeq (0.30 \pm 0.03) \text{ keV} ,
\Gamma_{\phi(1680)\to e^+e^-} \simeq (0.43 \pm .15) \text{ keV} .$$
(17)

We use a Breit-Wigner (BW) form for evaluating their contributions:

$$\sigma_H(t)|_{BW} \simeq 12\pi \frac{\Gamma_{ee}\Gamma_{tot}}{(t - M_R^2)^2 + M_R^2\Gamma_{tot}^2} , \qquad (18)$$

where we take M_R from PDG [33] and:

$$\Gamma_{tot}^{\omega(1420)} \simeq (174 \pm 59) \text{ MeV } [35],
\Gamma_{tot}^{\omega(1650)} \simeq (280 \pm 24) \text{ MeV } [35],
\Gamma_{tot}^{\phi(1680)} \simeq (150 \pm 50) \text{ MeV } [33].$$
(19)

3.3. THE J/ψ AND Υ FAMILIES

We consider the six J/ψ mesons 1S, 2S, 3770, 4040, 4160 and 4415, and the five Υ mesons 1S, 2S, 3S, 4S and 11020. We treat these mesons using NWA and the data on their electronic widths from PDG [33].

3.4. QCD CONTINUUM

As advertized previously, we shall treat the region not accessible experimentally by using perturbative QCD where it is expected to work and where nonperturbative effects like e.g. the quark and gluon condensates are negligible corrections.

Input for light quarks

In the case of massless quarks, the unit operator contribution to $R_{e^+e^-}$ is accurately known and reads, in the \overline{MS} scheme:

$$R_{e^+e^-}^{pert} = 3\sum_f Q_f^2 \left[1 + a_s + F_3 a_s^2 + F_4 a_s^3 \right]$$
(20)

with:

$$F_3 = 1.986 - 0.115 n_f [37],$$

$$F_4 = -6.637 - 1.200n_f - 0.005n_f^2 -1.240 \frac{\left(\sum_f Q_f\right)^2}{3\left(\sum_f Q_f^2\right)} \quad [38] , \qquad (21)$$

for n_f quark flavours; Q_f is the quark charge in units of e and $a_s \equiv (\alpha_s/\pi)$ is the QCD running coupling which we use to order a_s^2 (see e.g. the expression in [39,40]). We also use, the value of the QCD scale for three flavours:

$$\Lambda_3 = (375 \pm 50) \text{ MeV},$$
 (22)

obtained using the central value $\alpha_s(M_Z) = 0.119$ [33, 41].

To this perturbative correction, we add the light quark mass and non-perturbative corrections. We shall also consider the effects of a tachyonic gluon mass evaluated in [42] in order to take into account the effects of the truncation of the QCD perturbative series.

The dimension D = 2 (m_f^2 and tachyonic gluon) corrections are [39,42,43]:

$$tR_{e^+e^-}^{(2)} = -3\sum_f Q_f^2 \left[\bar{m}_f^2 \left[6 + 28a_s + (294.8 - 12.3n_f)a_s^2 \right] + 1.05a_s \lambda^2 \right], \quad (23)$$

where [42] $a_s \lambda^2 \simeq -(0.06 \sim 0.07) \text{ GeV}^2$.

The dimension D = 4 corrections are [44,39]:

$$t^{2}R_{e^{+}e^{-}}^{(4)} = 3\sum_{f}Q_{f}^{2}\left[\frac{2\pi}{3}\left(1-\frac{11}{18}a_{s}\right)\langle\alpha_{s}G^{2}\rangle +8\pi^{2}\left(1-a_{s}\right)m_{f}\langle\bar{\psi}_{f}\psi_{f}\rangle +\frac{32\pi^{2}}{27}a_{s}\sum_{f}m_{f}\langle\bar{\psi}_{f}\psi_{f}\rangle +\bar{m}_{s}^{4}\frac{8}{7}\left[-\frac{6}{a_{s}}+\frac{23}{4}\right] +a_{s}\left(\frac{2063}{24}-10\zeta_{3}\right)\right], \qquad (24)$$

where $\zeta_3 = 1.2020569...$

We use $\langle \alpha_s G^2 \rangle = (0.07 \pm 0.01) \text{ GeV}^4$ [45], and $(m_u + m_d) \langle \bar{u}u + \bar{d}d \rangle = -2f_\pi^2 m_\pi^2 (f_\pi = 92.6 \text{ MeV}).$ The running quark mass in the \overline{MS} scheme is defined as (see e.g. [40]):

$$\overline{m}_{i}(\nu) = \hat{m}_{i} \left(-\beta_{1} a_{s}(\nu)\right)^{-\gamma_{1}/\beta_{1}} \left\{ 1 + \frac{\beta_{2}}{\beta_{1}} \left(\frac{\gamma_{1}}{\beta_{1}} - \frac{\gamma_{2}}{\beta_{2}}\right) a_{s}(\nu) + \frac{1}{2} \left[\frac{\beta_{2}^{2}}{\beta_{1}^{2}} \left(\frac{\gamma_{1}}{\beta_{1}} - \frac{\gamma_{2}}{\beta_{2}}\right)^{2} - \frac{\beta_{2}^{2}}{\beta_{1}^{2}} \left(\frac{\gamma_{1}}{\beta_{1}} - \frac{\gamma_{2}}{\beta_{2}}\right) + \frac{\beta_{3}}{\beta_{1}} \left(\frac{\gamma_{1}}{\beta_{1}} - \frac{\gamma_{3}}{\beta_{3}}\right) \right] a_{s}^{2}(\nu) \right\}, \qquad (25)$$

where for three flavours the quark mass anomalous dimensions are (see e.g. [39,40]): $\gamma_1 = 2$, $\gamma_2 = 91/12$, $\gamma_3 = 24.8404$, while the coefficients of the β function are: $\beta_1 = -9/2$, $\beta_2 = -8$, $\beta_3 = -20.1198$. We shall use the value of the invariant strange quark mass [46]:

$$\hat{m}_s = (133.3 \pm 18.8) \text{ MeV}$$
 . (26)

We take into account the SU(3) breaking of the quark condensates $\langle \bar{s}s \rangle / \langle \bar{d}d \rangle \simeq 0.68^{+0.15}_{-0.29}$ [47] and consistently use $f_K = 1.2 f_{\pi}$.

Input for heavy quarks

In the case of heavy quarks, the perturbative spectral function can be accurately approximated by the Schwinger interpolating formula [48]:

$$R_{e^+e^-} = 3\sum_f Q_f^2 v_f \frac{(3-v_f^2)}{2} \left[1 + \frac{4}{3}\alpha_s f(v_f)\right], \quad (27)$$

where:

$$f(v_f) = \frac{\pi}{2v_f} - \frac{(3+v_f)}{4} \left(\frac{\pi}{2} - \frac{3}{4\pi}\right) , \qquad (28)$$

and:

$$v_f = \sqrt{1 - \frac{4m_f^2}{t}} , \qquad (29)$$

is the quark velocity; m_f is the heavy quark (pole) mass of flavour f. We shall see that, in the region where we shall work (away from threshold and for large t), these two parametrizations provide a sufficiently accurate description of the spectral function. For a conservative estimate, we shall consider the range of values spanned by the running and pole quark masses for the heavy quarks [49]:

$$m_c = (1.2 \sim 1.46) \text{ GeV}, \ m_b = (4.2 \sim 4.7) \text{ GeV}.$$
 (30)

Within the present accuracy of the values of the heavy quark masses, we find that it is not necessary to include the known α_s^2 corrections [50].

The (u, d, s) flavour contributions

We complete the contribution of the light quark channels by adding the QCD continuum from $t_c = 3$ to $(4.57 \text{ GeV})^2$. We shall use the value of Λ_3 given in Eq. (22). Our value of t_c is higher than $t_c \simeq 1.6 \text{ GeV}^2$ from global duality constraint in the I = 1 channel [51]¹. We have compared $a_{\mu}^{I=1}(1.6 \leq t \leq 3) \text{ GeV}^2$ from a QCD continuum parametrization and from the uses of data. They only differ by 16×10^{-11} , showing that the QCD continuum provides a good smearing of the data. We shall include this value into the systematics. One can also realize that the mass and non-perturbative corrections tend to cancel each others, whilst, individually, they are already small.

The (u, d, s, c) flavour contributions

We add the contribution of the charm quark about 150 MeV above the $J/\psi(4415)$ (\approx empirical mass-splitting

 $^{^1\}mathrm{I}$ thank Eduardo de Rafael for this remark. A similar value of the QCD continuum is obtained from the V- A channel [52].

between the radial excitations). In this region, the previous expressions give a good description of the spectral function, because at this energy, the charm quark is already relativistic with a velocity larger than 0.75. Additional threshold effects are taking into account by transforming the value of Λ_3 into Λ_4 .

The (u, d, s, c, b) flavour contributions

The contribution of the *b*-quark is added about 250 MeV above the $\Upsilon(11020)$ resonance (\approx empirical masssplitting between the radial excitations). Again at this energy, the *b* quark is already relativistic as its velocity is larger than 0.55, such that our perturbative parametrization remains a good approximation. We integrate until the $2M_t$ threshold, where we take $M_t \simeq 174.3(5.1)$ GeV as a best value given by PDG [33].

3.5. FINAL RESULT FOR $a_{\mu}^{had}(l.o)$ From combined τ -decay and e^+e^- data

Collecting all different contributions from Table 2, we deduce the final result at the end of that table, which we compile in Table 1 for a comparison with other previous determinations. We obtain 2 :

$$a_{\mu}^{had}(l.o) = 7036.1(76.4) \times 10^{-11}$$
, (31)

using the inclusive τ -decay data until 3 GeV² from ALEPH/OPAL [30,31] for the isovector channel and using e^+e^- data below 3 GeV² [33–36] for the isoscalar channel. The errors have been added quadratically. The effects of the heavy quark mesons have been treated using a narrow width approximation plus QCD continuum away from the quark-anti-quark thresholds.

From the alone e^+e^- data

If we use the isovector e^+e^- data below 3 GeV², we obtain:

$$a_{\mu}^{had}(l.o) = 7015.7(119.1) \times 10^{-11}$$
, (32)

which is slightly lower than the one from inclusive τ decay and less accurate. The difference in each region can be seen explicitly in Table 2 and easy to understand from the data given in [30,24].

Different source of errors

– The main error (80% when added quadratically) in our previous determinations comes from the ρ -meson region below 0.8 GeV². Hopefully, improved measurements of this region are feasible in the near future.

- The second source of errors comes from the region around M_{τ} for the inclusive τ -decay and between 1 GeV to M_{τ} for the e^+e^- data. These errors are about half of the one from the region below 0.8 GeV² in most different determinations. They can be reduced by improved measurements of inclusive τ -decay near M_{τ} (I = 1) and by improving the measurements of the odd multi-pions and $\bar{K}K$, $\bar{K}K\pi$, ... channels in the I = 0 channels from e^+e^- data.

- The contributions of the whole region above 3 ${\rm GeV^2}$

induce much smaller errors (7% of the total). There is a quite good consensus between different determinations in this energy region.

Improvements from τ -decay data

Our results for the isovector contributions, show that the precise inclusive τ -decay data have significantly improved by almost a factor two the accuracy of the determination compared with the one from e^+e^- data. However, they are amended by the inaccuracy near M_{τ} . Indeed, one can gain a bit in precision by using e^+e^- data in the region between 2.1 and 3 GeV² instead of τ -decay data, but we are aware of the disagreement among different e^+e^- data in this region. In this case of figure, one would get from Table 2:

$$a_{\mu}^{had}(l.o) = 7010.0(72.2) \times 10^{-11}$$
 (33)

Final result and additional systematic errors

We consider as a final result the average of the three results given in Eqs. (31), (32) and (33). We consider the accuracy given in Eq. (33), which avoids the weakness of τ -decay data near M_{τ} . We add to this error the following systematics:

– We take into account, an eventual deviation from the CVC assumption ³ (CVC has been also tested in [53]), and some other systematics in manipulating the isovector data by adding a systematic error of 15.5×10^{-11} coming from the largest difference of the central value of Eqs. (31) and (32) from the average value.

- We also add another 16×10^{-11} systematics due to the difference of the continuum threshold 3 GeV² value for three flavours used here and the one from QCD global duality in [51] (see section 3.4).

Adding the different errors quadratically, we deduce *our final estimate*:

$$a_{\mu}^{had}(l.o) = 7020.6(75.6) \times 10^{-11}$$
, (34)

It is amusing to notice that the central value of our different results almost coïncide with our 25 years old value [6,7] and with the *mean central value* given in Table 1.

3.6. COMPARISONS WITH PREVIOUS DE-TERMINATIONS OF $a_{\mu}^{had}(l.o)$

QCD continuum

- Our treatment of the QCD continuum in the heavy quark channels gives similar results than more involved parametrization [22] including non-perturbative gluon condensate and non-relativistic contributions. This is due to the fact that in the region where we work, the heavy quark-antiquark pairs are already relativistic with a quark velocity larger than 0.55.

- Comparing existing estimates, one can see that there is a consensus on the size of the QCD continuum effect. The departures from different determinations come mainly from the treatment of the low-energy region below 2 GeV.

 $^{^2\}mathrm{Here}$ and in the following, the errors will be added quadratically as usually done.

 $[\]overline{{}^{3}I}$ thank Paco Yndurain for mentioning this point, for checking some of the entries in Table 2 and for several communications.

Isoscalars

For the isoscalar resonances, one can realize by comparing the results with existing estimates that the narrow width approximation (NWA) gives enough accurate results comparable in magnitude and accuracy with more sophisticated parametrization including width effects [21].

Isovector channel and some global comparisons

In the following comparison, we shall use numbers in units of 10^{-11} .

- Comparison with [19, 22]

It has been argued in [19] that the contribution from this region can be improved by using constraint imposed by $\pi\pi$ scattering data in the spacelike region on the pion form factor. The result $a_{\mu}^{had}(l.o)(t \leq 0.8$ $\text{GeV}^2) = 4848(31)$ quoted by [19] is based on Watson theorem (extended to 0.8 GeV^2) for the $\pi\pi$ phase shift $\delta_1^1(t)$. The central value differs with our value 4795(61) obtained from the data in the timelike region by 53 MeV, but agrees with the result 4794(14) given in their Eq. (3.23) obtained from a fit of $\delta_1^1(t)$ from the data. Another discrepancy of 39(104) comes from the region between 0.8 and 2 GeV^2 , but is not significant due to the large errors. It is argued in [22] that error from high-energy region can be reduced .

- Comparison with [21]

Our result agrees within the errors with the final result 7024(153) of [21] from their Table 3a 4 , with the updated result 6989(111) quoted in [8], and with some of the results given in Table 2.

- Comparison with [24–26]

For a proper comparison with these different determinations, we shall only consider the two regions separated by the scale $\sqrt{t} = M_{\tau}$, and compare our result $a_{\mu}^{had}(l.o)(\sqrt{t} \leq M_{\tau}) = 6426(76)$ obtained from τ decay data. The central value of our result is relatively higher than the value 6365(74) in [25] and 6343(60) in [26]. Within the errors, the 1st result is consistent with ours while the second shows one σ discrepancy. For comparing with the result of [24], we subtract from their total result, the heavy quarks and QCD continuum contributions. In this way, we deduce the contribution 6405(94), much more closer to our result than the two formers, where one should also notice that our final results from the alone e^+e^- data agree.

The exact origin of the discrepancy of the central values is not easy to detect due to the complexity of the analysis in this region, though it is reassuring that results based on maximal data inputs [24,25] are consistent with ours within the errors.

We do not worry by the discrepancy with [26], as it uses more theoretical inputs than data ("determination with minimal data input" as quoted from [54]) for minimizing the errors , through the introduction (and subtraction) of some arbitrary polynomials in t(not coming from QCD first principles) for approaching the t = 0 region ⁵. On the other hand, the use of "local duality" in a given interval is stronger than the standard applications of QCD spectral sum rules à la SVZ [44] (for a review see e.g. [40]), where systematic uncertainties can be more severe. We do not expect that the result obtained in this way is much better (though apparently more accurate) than the ones previously obtained from pure data analysis.

One should also notice that, though one has been able to formulate the τ -decay width using QCD and the Operator Product Expansion (OPE) including the QCD vacuum condensates [39,55] à la SVZ [44], which has provided an accurate measurement of the QCD coupling α_s [30,31], the extension of such a program to the analysis of the lepton anomaly cannot be done in a straightforward way due the existence of a pole at t =0, which needs a nonperturbative QCD control of the contribution of the small circle at the origin when using the Cauchy contour for evaluating the muon anomaly in the complex-t plane. Such a control is not feasible at present from QCD first principles.

Final errors and comments on the uses of new data

We can conclude from the previous determinations in Table 1 and the present work that, with present data and with less theoretical inputs, a realistic error for extracting $a_{\mu}^{had}(l.o)$ cannot be less than 70×10^{-11} . After the completion of this work, we have checked [56] by doing similar analyses (hadronic contributions to α and to the muonium hyperfine splitting) that, using the new Novosibirsk [57] and BES [58] data, one does not improve noticeably the errors compared to the one obtained from the correlated averaged data and QCD continuum parametrizations used in this paper.

4. THEORY OF THE MUON ANOMALY

4.1. SUM OF THE THEORETICAL CON-TRIBUTIONS

- QED and SM electroweak contributions Using obvious notations, they read [4,59,5]:

$$a_{\mu}^{QED} = 116\ 584\ 705.7(2.9) \times 10^{-11} ,$$

$$a_{\mu}^{EW} = 151(4) \times 10^{-11} .$$
(35)

with QED up to 8th order and EW including two-loop corrections [60].

- Higher order hadronic contributions

For $a_{\mu}^{had}(h.o)$, we have used the contribution of the high-order vacuum polarizations $(h.o)_{V.P}$ recently estimated in [61], where we have checked that the central value coïncide with the original estimate in [6] which has the same result for $a_{\mu}^{had}(l.o)$ though the error is larger in [6] due to older data. This feature is reassuring for a self-consistency check. As the data used in [6,61] give smaller value of $a_{\mu}^{had}(l.o)$ by 1.00125 than

⁴I thank William Marciano for mentioning this correct result and for several communications.

⁵This procedure together with the analytic continuation method has been already criticized in [40]. Analogous comments apply to the estimate of $\alpha(M_Z)$.

the present work, we have rescaled the $(h.o)_{V.P}$ contribution by this small factor, and we deduce:

$$a_{\mu}^{had}(h.o)_{V.P} = -101.2(6.1) . \tag{36}$$

For the light by light scattering hadronic contributions, we use the average of the two estimates in [62] which agree each others. We have corrected the sign of the pseudoscalar and axial-vector contributions according to the result in [63], which has been confirmed in [64] 6 . Then, we obtain:

$$a_{\mu}^{had}(h.o)_{LL} = 70.8(21.0) \times 10^{-11}$$
 (37)

We have not included into the sum the contribution due to the quark loop, suspecting that adding this effect together with the pion loop and meson pole contributions can lead to a double counting (quark-hadron duality arguments ⁷). Instead, we have considered the quark loop effect incorporating vector mesons in the photon legs, which is about 15×10^{-11} (average of the two determinations in [62]) as another source of uncerainties of the estimate. Then, we deduce:

$$a_{\mu}^{had}(h.o) \equiv a_{\mu}^{had}(h.o)_{V.P} + a_{\mu}^{had}(h.o)_{LL} = -30.4(21.9) \times 10^{-11} , \qquad (38)$$

showing that the higher order hadronic effects tend to cancel each others.

- Total theoretical predictions in the SM

Adding the results in Eqs. (35) and (38), one obtains the SM theoretical contributions:

$$a_{\mu}^{SM} = 116\ 584\ 826.3(22.5) \times 10^{-11} + a_{\mu}^{had}(l.o)$$

= 116\ 591\ 846.9(78.9) \times 10^{-11} , (39)

where $a_{\mu}^{had}(l.o)$ is the lowest order hadronic contributions evaluated in this paper [see Eq. (34)].

4.2. THEORY VERSUS EXPERIMENT

In confronting the theoretical estimate with the experimental value in Eq. (1), we shall use, for reasons explained previously, the result in Eq. (34) with a moderate accuracy. We deduce:

$$a_{\mu}^{new} \equiv a_{\mu}^{exp} - a_{\mu}^{SM} = 176(170) \times 10^{-11} .$$
 (40)

The mean central value from our selected determinations given in Table 1 would give:

$$\langle a_{\mu}^{new} \rangle \equiv a_{\mu}^{exp} - \langle a_{\mu}^{SM} \rangle = 189(175) \times 10^{-11} , \qquad (41)$$

after adding to the "statistical" error in Table 1, the one from our determination in Eq. (34). Both values indicate an *eventual window* of about only σ for *new physics* beyond the standard model. A more definite claim is waiting for a more precise a_{μ} measurement. We translate the results in Eqs. (40) and (41) into the conservative range:

$$-42 \le a_{\mu}^{new} \times 10^{11} \le 413 \quad (90\% \text{ CL}) . \tag{42}$$

We shall discuss some numerical implications of this result by giving conservative lower bounds on the scales of new physics discussed recently in the literature [4, 66–69].

4.3. LOWER BOUNDS ON NEW PHYSICS Supersymmetry

Using the expression given in [4] and for $\tan \beta \equiv \langle \phi_2 \rangle / \langle \phi_1 \rangle$ (ratio of Higgs expectation values) ≥ 4 :

$$a_{\mu}^{SUSY} \simeq \frac{\alpha}{8\pi \sin^2 \theta_W} \frac{m_{\mu}^2}{\tilde{m}^2} \tan \beta , \qquad (43)$$

one obtains a conservative lower bound on the degenerate sparticle mass:

$$\tilde{m} \ge 113 \text{ GeV} \ (90\% \text{ CL}),$$
 (44)

which is comparable with the present experimental lower bound of 100 GeV.

Radiative mass, composite and extended Zee models

In these models where the typical contribution to a_{μ} is about $(m_{\mu}^2/M^2)\mathcal{O}(1)$ [4,66,67], one can derive the lower bound on the scale of the models:

$$M \ge 1.7 \text{ TeV}$$
 . (45)

This constraint can also be translated into a lower bound on the mass of a singlet scalar in the Zee Model with a coupling $1/96\pi^2$ [68]. One can deduce:

$$M_{S_2} \ge 55 \text{ GeV} . \tag{46}$$

Leptoquarks scenarios

Leptoquark effects on a_{μ} have been discussed recently in [69]. For an electromagnetic coupling at the leptonleptoquark-quark vertex, one can deduce the approximate expression: as:

$$a_{\mu}^{lq} \simeq \left(\frac{\alpha}{\pi}\right) \frac{m_c m_{\mu}}{M_{lq}^2} \log \frac{M_{lq}^2}{m_c^2} , \qquad (47)$$

giving:

$$M_{lq} \ge 1.1 \text{ TeV}$$
 . (48)

This bound is much larger than the present lower bounds of about $(200 \sim 300)$ GeV from direct search experiments at HERA and Tevatron.

We expect that these different bounds will be improved in the near future both from accurate measurements of a_{μ} and of e^+e^- data necessary for reducing the theoretical errors in the determinations of the hadronic contributions, being the major source of the theoretical uncertainties.

5. THEORY OF THE TAU ANOMALY

Here, we update our old and first work [7] on the τ lepton anomaly. We shall use $M_{\tau} = 1.77703$ GeV from PDG [33],

⁶We are also aware of the fact that the two results are modeldependent and further tests on their validity are needed. However, one can notice that this contribution is of the same size as the error of $a_{\mu}^{had}(l.o)$.

⁷The agreement of the result in Eq. (37) with the one $62(3) \times 10^{-11}$ obtained in [18] (see also [65]), using the quark constituent model (and no hadrons at all!) seems to support such arguments.

Lowest order hadronic contribution

The estimate from $e^+e^- + \tau$ -decay is given in Table 2. Taking the similar average as in the case of the muon (section 3.5), one obtains:

$$a_{\tau}^{had}(l.o) = 353.6(4.0) \times 10^{-8}$$
 (49)

One can see that the estimate $280(20) \times 10^{-8}$ in [70], using present values of the QCD condensates [45,40], is much lower than this result. This appears to be a general feature of this method based on minimal data inputs (see e.g. [26] for $a_{\mu}^{had}(l.o)$ discussed in the previous section). A comparison with the result of [21] shows that in both cases of muon and tau, this result is systematically higher than ours by the same scaling factor 1.0004. Previous results 370(40) [7] and 360(30) [71] in units of 10^{-8} are comparable with ours but inaccurate.

Higher order hadronic contributions

The higher order contributions due to vacuum polarization $a_{\tau}^{had}(h.o)_{V.P}$, can be obtained from the result of [61]. We rescale it by the factor 1/1.0083 like we have done for the muon case as explained in previous section 9. For the light by light scattering contribution, we use the result of [62] for the muon and we rescale it by the mass squared ratio $(m_{\tau}/m_{\mu})^2$, which is expected to be a good approximation from the semianalytical expression given in [6]. Then, we obtain:

$$a_{\tau}^{had}(h.o)_{V.P} = 7.6(0.2) \times 10^{-8} ,$$

$$a_{\tau}^{had}(h.o)_{L.L} = \left(\frac{m_{\tau}}{m_{\mu}}\right)^2 a_{\mu}^{had}(h.o)_{L.L}$$

$$= 20.0(5.8) \times 10^{-8} .$$
(50)

Therefore, we deduce:

$$a_{\tau}^{had}(h.o) = 27.6(5.8) \times 10^{-8},$$
 (51)

and:

$$a_{\tau}^{had} \equiv a_{\tau}^{had}(l.o) + a_{\tau}^{had}(h.o) = 381.2(7.0) \times 10^{-8}.$$
 (52)

Electroweak contributions

The lowest order electroweak SM contribution is well-known:

$$a_{\tau}^{EW}(l.o) = \frac{5}{3} \frac{G_{\tau} m_{\tau}^2}{8\pi^2 \sqrt{2}} \Big[1 + \frac{1}{5} (1 - 4\sin^2 \theta_W)^2 \Big], \qquad (53)$$

to order $(m_{\tau}/M)^2$ (*M* being the W or Higgs mass); We assume $G_{\tau} = G_{\mu} = 1.166 \ 39(1) \times 10^{-5}$ is the Fermi coupling; $\sin^2 \theta_W = 0.224$, where θ_W is the weak mixing angle. We add the two-loop contribution (2loop/l.o $\simeq -65\alpha/\pi$) [60], which induces a 15% reduction of the one loop result. We obtain:

$$a_{\tau}^{EW} \simeq 46.9(1.2) \times 10^{-8}$$
 (54)

QED contributions

- Generalities

These contributions have been first evaluated in [7] to

order α^3 . Here, we revise and improve these evaluations. In so doing, we use the relation [72,73]:

$$a_{\tau} - a_{e} = \sum_{l,l' \equiv e,\mu} \left[a_{4}(l) \left(\frac{\alpha}{\pi}\right)^{2} + \left[a_{6}(l) + a_{6}(ll') \right] \left(\frac{\alpha}{\pi}\right)^{3} \right],$$
(55)

where l indicates the internal fermion loop appearing in the photon propagator. $a_4(l)$ and $a_6(l)$ can be deduced from the result of the muon anomaly while $a_6(ll')$ is a new contribution involving one electron and one muon loop insertion in one photon propagator. At the same order of truncation of PT series, one has [73,59]:

$$a_e = 0.5 \left(\frac{\alpha}{\pi}\right) - 0.328 \ 478 \ 965 \left(\frac{\alpha}{\pi}\right)^2 + 1.181 \ 241 \ 456 \left(\frac{\alpha}{\pi}\right)^3 + \dots = 1 \ 159 \ 65.2 \times 10^{-8} \ , \tag{56}$$

where we have used the measurement:

$$\alpha^{-1} = 137.036\ 003\ 7(33)\ ,\tag{57}$$

from the quantum Hall effect [74].

- QED at fourth order

For evaluating this contribution, we use the dispersive representation:

$$a_4(l) = \frac{1}{3} \int_{4m_l^2}^{\infty} \frac{dt}{t^2} (t + 2m_l^2) \sqrt{1 - \frac{4m_l^2}{t}} K_\tau(t) , \qquad (58)$$

where $K_{\tau}(t)$ is the kernel function defined in Eq. (3). Then, we obtain:

$$a_4(e) = 2.024\ 29$$
 and $a_4(\mu) = 0.361\ 66$. (59)

The former result is well approximated by the known analytic approximate relation to order $(m_e/m_{\tau})^2$ given e.g. in [6], while the second result needs the inclusion of the not yet available $(m_{\mu}/m_{\tau})^3$ term⁸. Adding these two contributions, we obtain:

$$(a_{\tau} - a_e)_4 = 2.385 \ 95 \left(\frac{\alpha}{\pi}\right)^2 = 1287.3 \times 10^{-8}$$
 (60)

- QED at sixth order

Because of the accurate determinations of the hadronic and weak contributions, the inclusion of the sixth order contributions becomes necessary as they contribute with the same strength. The contributions of diagrams with vacuum polarizations and ladders, can be obtained from the analytic result given for the muon in [73]:

$$a_{6}(l) = \frac{2}{9} \log^{2} \frac{m_{\tau}}{m_{l}} - 1.113 \ 90 \log \frac{m_{\tau}}{m_{l}} + 4.307 \ 66 + \dots , \qquad (61)$$

which leads to:

$$a_6(e) = 10.000 \ 2 \ , \qquad a_6(\mu) = 2.934 \ 0 \ .$$
 (62)

⁸These results agree with the last paper in [71] confirming that the results in their two former papers are wrong.

 $a_6(\mu)$ differs from the value -.122 given in [71] where the origin of the negative value is not understandable from Eq. (61) but should come from a wrong term used there. We add the new class of contributions specific for a_{τ} given in [7]:

$$a_{6}(e\mu) = 2\left\{\frac{2}{9}\log\frac{m_{\tau}}{m_{e}}\log\frac{m_{\tau}}{m_{\mu}} - \frac{25}{54}\log\frac{m_{\tau}^{2}}{m_{e}m_{\mu}} + \frac{\pi^{2}}{27} + \frac{317}{324}\right\} = 2.753 \ 16 \ . \tag{63}$$

We parametrize the light by light scattering contribution as:

$$a_6(l)_{L.L} \simeq \frac{2\pi^2}{3} \log \frac{m_\tau}{m_l} - 14.13 ,$$
 (64)

by combining the known coefficient of the log-term [75] and the numerical value of the total contribution for the muon [76]. This leads to:

$$a_6(e)_{L.L} = 39.521\ 7\ , \qquad a_6(\mu)_{L.L} = 4.441\ 2\ ,\ (65)$$

This contribution is so far the most important at sixth order. Adding the different contributions from Eqs (61) to (65), one obtains to sixth order:

$$(a_{\tau} - a_e)_6 = 59.650 \left(\frac{\alpha}{\pi}\right)^3 = 74.758 \times 10^{-8} .$$
 (66)

This effect is about the same strength as the weak interaction effect and bigger than the higher order hadronic contributions.

Total QED contribution up to sixth order
 Adding the previous QED contributions, we deduce

$$a_{\tau}^{QED} = 117\ 327.1(1.2) \times 10^{-8}$$
, (67)

where the error comes from the measurement of α .

Final result

Summing up the previous different theoretical contributions, we deduce in the standard model:

$$a_{\tau}^{SM} = a_{\tau}^{had} + a_{\tau}^{EW} + a_{\tau}^{QED} = 117\ 755.2(7.2) \times 10^{-8} .$$
(68)

We consider this result as an improvement of our old [7] and other existing results. In [66], only an average of different existing hadronic contributions have been added to the QED and electroweak contributions. In [7,71], the hadronic contributions are inaccurate, while the value of the sixth order QED contribution is incorrect. This value in Eq. (68) can be compared with the present (inaccurate) experimental value [77]:

$$a_{\tau}^{exp} = 0.004 \pm 0.027 \pm 0.023 , \qquad (69)$$

which, we wish, will be improved in the near future.

6. SUMMARY

- We have re-evaluated the lowest order hadronic contribution to the muon and tau anomalies using the precise τ -decay data below M_{τ} and averaged e^+e^- data. Ours results are given in Table 2 and section 7, and our best choice in Eq. (34). Though the approach is not conceptually new, the present situation of a_{μ} (theory versus experiment) justifies a new independent reevaluation of the anomaly using minimal theoretical inputs.

- We have extensively discussed the different sources of the errors in the analysis, which are dominated by the region below M_{τ} . However, the relative weight of this region decreases for a_{τ} (see also detailed discussions in [7]) where a more precise theoretical prediction can then be provided.

– We have compared our analysis summarized in Table 2 and our final result in Eq. (34) with some of the existing estimates given in Table 1 ⁹.

- We have shortly discussed implications of our result to some models beyond the standard model, namely supersymmetry with large $\tan \beta$, radiative muon mass, composite, some extended Zee and leptoquark models. The lower bounds on the mass scale of these models from the muon anomaly is comparable or in some cases stronger than existing experimental lower bounds.

– We have completed our work by updating our old estimate [7] of the different theoretical contributions (QED up to sixth order, higher order hadronic and electroweak) to the τ lepton anomaly. Our result is given in Eq. (68). A precise future measurement of the τ -lepton anomaly is welcome as it will permit to probe the QED series at shorter distance, and where, the relative weights of different interaction effects are very different from the case of the muon.

ACKNOWLEDGEMENTS

It is a pleasure to thank Hsiang-nan Li and the theory group of NCTS-Hsinchu for the warm hospitality.

REFERENCES

- 1. H.N. Brown et al., hep-ex/0102017.
- 2. J. Bailey et al., Nucl. Phys. B 150 (1979) 1.
- 3. H.N. Brown et al., hep-ex/0009029.
- 4. A. Czarnecki and W.J. Marciano, hep-ph/0102122.
- 5. F.J. Yndurain, hep-ph/0102312.
- J. Calmet, S. Narison, M. Perrottet and E. de Rafael, *Rev. Mod. Phys.* 49 (1977) 21; *Phys. Lett.* B 161 (1976) 283; S. Narison, *Thèse de 3ème* cycle, Marseille (1976).
- 7. S. Narison, J. Phys. (Nucl. Phys.) G 4 (1978) 1840.
- 8. F. Jegerlehner, hep-ph/0104304.
- C. Bouchiat and L. Michel, J. Phys. Radium 22 (1961) 121.
- L. Durand III, *Phys. Rev.* **128** (1962) 441; erratum **129** (1963) 2835.
- T. Kinoshita and R. Oakes, *Phys. Lett.* B 25 (1967) 143.
- 12. J.E. Bowcock, Z. Phys. **211** (1968) 400.

⁹During the submission of this paper to the journal, some other papers appeared in the literature [78].

- M. Gourdin and E. de Rafael, Nucl. Phys. B 10 (1969) 667.
- A. Bramon, E. Etim and M. Greco, *Phys. Lett.* B 39 (1972) 514.
- 15. J. Bailey et al., *Phys. Lett.* B 55 (1975) 420.
- V. Barger, W.F. Long and M.G. Olsson, *Phys. Lett.* B 60 (1975) 89.
- 17. L.M. Barkov et al., Nucl. Phys. B 256 (1985) 365.
- T. Kinoshita, B. Nizić and Y. Okamoto, *Phys. Rev.* D 31 (1985) 2108.
- J.A. Casas, C. López and F.J. Ynduráin, *Phys. Rev.* D 32 (1985) 736.
- S. Dubnicka and L. Martinovic, *Phys. Rev.* D 42 (1990) 884.
- S. Eidelman and F. Jegerlehner, Z. Phys. C 67 (1995) 585, and private communication.
- K. Adel and F.J. Yndurain, hep-ph/9509378 (1995), *Rev. Acad. Ciencias (Esp)*, **92** (1998).
- D.H. Brown and W.A. Worstell, *Phys. Rev.* D 54 (1996) 3237.
- 24. R. Alemany, M. Davier and A. Höcker, *Eur. Phys. J.* C 2 (1998) 123.
- M. Davier and A. Höcker, *Phys. Lett.* B 419 (1998) 419.
- M. Davier and A. Höcker, *Phys. Lett.* B 435 (1998) 427.
- B. Lautrup and E. de Rafael, *Phys. Rev.* 174 (1968) 1835.
- D. Gómez Dumm, A. Pich and J. Portolés, hepph/000332.
- G. Colangelo, M. Finkemeir and R. Urech, *Phys. Rev.* D 54 (1996) 4403.
- The ALEPH collaboration, R. Barate et al., *Eur. Phys. J.* C 76 (1997) 15; C 4 (1998) 409; A. Hocker, hep-ex/9703004 (I thank Michel Davier for bringing the first reference to my attention).
- The OPAL collaboration, K. Ackerstaff et al., *Eur. Phys. J.* C 7 (1999) 571.
- F.J. Gilman and S.H. Rhie, *Phys. Rev.* D 31 (1985) 1066.
- PDG 2000, D.E. Groom et al., Eur. Phys. J. C 15 (2000) 1.
- 34. S. Dolinsky et al., Phys. Rep. C 202 (1991) 99.
- The DM2 collaboration, A. Antonelli et al., Z. Phys. C 56 (1992) 15; D. Bisello et al., Z. Phys. C 39 (1988) 13.
- The DM1 collaboration, F. Mane et al., *Phys. Lett.* B 112 (1982) 178; A. Cordier et al., *Phys. Lett.* B 110 (1982) 335.
- K.G. Chetyrkin, A.L. Kataev and F.V. Tkachov, *Phys. Lett.* B 85 (1979) 277; M. Dine and J. Sapirstein, *Phys. Rev. Lett.* 43 (1979) 668; W. Celmaster and R. Gonsalves, *Phys. Rev. Lett.* 44 (1980) 560.
- 38. S.G. Gorishny, A.L. Kataev and S.A. Larin, *Phys. Lett.* B 259 (1991) 144; See also : L.R. Surguladze and M.A. Samuel, *Phys. Rev. Lett.* 66 (1991) 560 and erratum 2416.
- E. Braaten, S. Narison and A. Pich, *Nucl. Phys.* B 373 (1992) 581.

- 40. S. Narison, Phys. Rep. 84 (1982) 263; QCD spectral sum rules, Lecture Notes in Physics, Vol 26 (1989) 19, ed. World Scientific; The QCD theory of hadrons, Cambridge Monograph Series (to appear in 2001).
- S. Bethke, Nucl. Phys. (Proc. Suppl.) B, C 39 (1995); ibid, B, A 54 (1997); hep-ex/0004201; M. Schmelling, ICHEP96, Varsaw, 1996; I. Hinchliffe and A. Manohar hep-ph/0004186.
- K.G. Chetyrkin, S. Narison and V.I. Zakharov, *Nucl. Phys.* B 550 (1999) 353.
- K.G. Chetyrkin and J.H. Kühn, *Phys. Lett.* B 248 (1990) 359.
- M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, Nucl. Phys B 147 (1979) 385, 448.
- S. Narison, *Phys. Lett.* B 387 (1996) 162; ibid B 361 (1995) 121 and references therein.
- S. Narison, Nucl. Phys. (Proc. Suppl.) B 86 (2000) 242; Phys. Lett. B 216 (1989) 191; H. Leutwyler, hep-ph/001049.
- 47. S. Narison, Phys. Lett. B 358 (1995) 113.
- J. Schwinger, Particles, Sources and Fields, Vol 2, Addison-Wesley (1973).
- 49. S. Narison, Phys. Lett. B 341 (1994) 73.
- K.G. Chetyrkin, J.H. Kühn and M. Steinhauser, *Nucl. Phys.* B 482 (1996) 213.
- S. Peris, M. Perrottet and E. de Rafael, Journ. High Ener. Phys. 05 (1998) 01.
- S. Narison, Nucl. Phys. B 593 (2001) 3; Nucl. Phys. (Proc. Suppl.) B 96 (2001) 364.
- S. Narison and A. Pich, *Phys. Lett.* B 304 (1993) 359.
- 54. K. Schilcher, in Conf. on Non-perturbative methods, Montpellier (1985) World Scientific, ed. S. Narison and references therein; S. Groote et al., Phys. Lett. B 440 (1998) 375.
- 55. F. Le Diberder and A. Pich, *Phys. Lett.* B 286 (1992) 147; B 289 (1992) 165.
- 56. S. Narison, hep-ph/0108065.
- The CMD collaboration, R.R. Akhmetshin et al., *Nucl. Phys.* A 675 (2000) 424c; S.I. Serednyakov, *Nucl. Phys. (Proc. Suppl.* B 96 (2001) 197.
- The BES collaboration, J.Z. Bai et al., *Phys. Rev. Lett.* 84 (2000) 594; hep-ex/0102003.
- V.W. Hughes and T. Kinoshita, *Rev. Mod. Phys.* 71, 2 (1999) S133.
- A. Czarnecki, B. Krause and W.J. Marciano, *Phys. Rev. Lett.* **76** (1996) 3267.
- 61. B. Krause, *Phys. Lett.* B **390** (1997) 392.
- M. Hayakawa and T. Kinoshita and A. Sanda, *Phys. Rev.* D 54 (1996) 3137; M. Hayakawa and T. Kinoshita, *Phys. Rev.* D 57 (1998) 465; J. Bi- jnens, E. Pallante and J. Prades, *Nucl. Phys.* B 474 (1996) 379 and private communication from J. Prades; E. de Rafael, *Phys. Lett.* B 322 (1994) 239.
- M. Knecht and A. Nyffeler, hep-ph/0111058;
 M. Knecht, A. Nyffeler, M. Perrottet and E. de Rafael, hep-ph/0111059.
- 64. M. Hayakawa and T. Kinoshita, hep-ph/0112102;

I. Blokland, A. Czarnecki and K. Melnikov, hep-ph/0112117.

- 65. J.F. de Trocóniz and F.J. Yndurain, hepph/0106025.
- A. Czarnecki and W.J. Marciano, Nucl. Phys. (Proc. Suppl.) B 76 (1999) 245.
- E. Eichten, K. Lane and J. Preskill, *Phys. Rev. Lett.* 47 (1980) 225; K. Lane, hep-ph/0102131 and private communication; X. Calmet, H. Fritzsch and D. Holtmannspötter, hep-ph/0103012.
- 68. D.A. Dicus, H-J. He and J.N. Ng, hep-ph/0103126 and private communication from J.N. Ng.
- K. Cheung, hep-ph/0102238 and references therein, and private communication; D. Chakraverty, D. Choudhury and A. Datta, hep-ph/0102180.
- 70. F. Hamzeh and N.F. Nasrallah, *Phys. Lett.* B 373 (1996) 211.
- G. Li, R. Mendel and M.A. Samuel, *Phys. Rev. Lett.* 67 (1991) 668; erratum 69 (1992) 995; *Phys. Rev.* D 47 (1993) 1723.
- 72. B.E. Lautrup, A. Peterman and E. de Rafael, *Phys. Rep.* C 3 (1972) 193.
- 73. W.J. Marciano and T. Kinoshita, in Quantum Electrodynamics, Advanced Series on Directions in High Energy Physics, Vol 7, World Scientific., ed. Kinoshita.
- 74. E. Krüger, W. Nistler and W. Weirauch, *Metrologia* **32** (1995) 117.
- 75. B.E. Lautrup and M.A. Samuel, *Phys. Lett.* B 72 (1977) 114.
- 76. T. Kinoshita, Phys. Rev. Lett. 61 (1988) 2898.
- L. Taylor, Nucl. Phys. (Proc. Suppl.) B 76 (1999) 273.
- 78. W.J. Marciano and B. Lee Roberts, hepph/0105056; J.F. de Trocóniz and F.J. Yndurain, hep-ph/0106025; K. Melnikov, hep-ph/0105267; E. Bartoš et al., hep-ph/0106084.