

# Sfermion Mass Degeneracy, Superconformal Dynamics and Supersymmetric Grand Unified Theories

Tatsuo KOBAYASHI,<sup>\*</sup> Hiroaki NAKANO,<sup>†</sup> Tatsuya NOGUCHI,<sup>‡</sup> and Haruhiko TERAO<sup>§</sup>

<sup>\*‡</sup>*Department of Physics, Kyoto University, Kyoto 606-8502, Japan*

<sup>†</sup>*Department of Physics, Niigata University, Niigata 950-2181, Japan*

<sup>\*‡</sup>*Institute for Theoretical Physics, Kanazawa University, Kanazawa 920-1192, Japan*

## Abstract

We discuss issues in a scenario that hierarchical Yukawa couplings are generated through strong dynamics of superconformal field theories (SCFTs). Independently of mediation mechanism of supersymmetry breaking, infrared convergence property of SCFTs can provide an interesting solution to supersymmetric flavor problem; sfermion masses are suppressed around the decoupling scale of SCFTs and eventually become degenerate to some degree, thanks to family-independent radiative corrections governed by the SM gaugino masses. We discuss under what conditions the degeneracy of sfermion mass can be estimated in a simple manner. We also discuss the constraints from lepton flavor violations. We then study explicitly sfermion mass degeneracy within the framework of grand unified theories coupled to SCFTs. It is found that the degeneracy for right-handed sleptons becomes worse in the conventional  $SU(5)$  model than in the MSSM. On the other hand, in the flipped  $SU(5) \times U(1)$  model, each right-handed lepton is still an  $SU(5)$ -singlet, whereas the bino mass  $M_1$  is determined by two independent gaugino masses of  $SU(5) \times U(1)$ . These two properties enable us to have an improved degeneracy for the right-handed sleptons. We also speculate how further improvement can be obtained in the SCFT approach.

---

<sup>\*</sup> E-mail: kobayash@gauge.scphys.kyoto-u.ac.jp

<sup>†</sup> E-mail: nakano@muse.hep.sc.niigata-u.ac.jp

<sup>‡</sup> E-mail: noguchi@gauge.scphys.kyoto-u.ac.jp

<sup>§</sup> E-mail: terao@hep.s.kanazawa-u.ac.jp

# 1 Introduction

Understanding the origin of hierarchical fermion masses and mixing angles is one of most important issues in particle physics. The Froggatt-Nielsen mechanism is a famous mechanism to realize hierarchical Yukawa couplings [1, 2]. Recently new ideas related to extra dimensions have also been discussed.

In models with softly-broken supersymmetry (SUSY), a mechanism that generates the hierarchical structure of Yukawa couplings generally affects the sfermion sector; one would have a characteristic pattern of sfermion masses and SUSY-breaking trilinear couplings. For example, the Froggatt-Nielsen mechanism with an extra  $U(1)$  gauge symmetry leads to the so-called  $D$ -term contribution to soft scalar masses,\* which are proportional to the charges under the broken  $U(1)$  symmetry. Such a pattern could be tested if superpartners will be discovered and sfermion masses as well as trilinear scalar couplings will be measured in future experiments. Even at present, soft SUSY-breaking parameters are constrained rather severely from the experimental bounds on flavor changing neutral current (FCNC) processes as well as CP violation. This is the SUSY flavor problem. For instance, the flavor-dependent  $D$ -term contributions are generically dangerous. In general, the SUSY flavor problem can be solved if either of the following three is realized at least for the first two families; 1) diagonal and degenerate sfermion masses, 2) decoupling of heavy sfermions, and 3) the alignment between the fermion and sfermion bases. Much effort has been devoted to realize the first solution by seeking a flavor-blind mediation mechanism of SUSY breaking.

Nelson and Strassler [5] have recently proposed an interesting mechanism to realize hierarchical Yukawa couplings. The setup is the minimal supersymmetric standard model (MSSM), or its extension, coupled to superconformal (SC) sector. The SC sector is strongly coupled and assumed to have an infrared (IR) fixed point [6, 7]. The first and second families of quarks and leptons gain a large and positive anomalous dimensions through the SC dynamics. Then their Yukawa couplings to electroweak Higgs fields are suppressed hierarchically at the scale  $M_C$  where the SC sector is assumed to decouple from the MSSM sector.

The SC fixed point has more interesting consequences. When pure superconformal field theory (SCFT) is perturbed by soft SUSY-breaking terms, general argument shows that such perturbation is exponentially suppressed toward the SC fixed point [8, 9, 10, 11]. Specifically one expects that a sfermion mass is suppressed at the decoupling scale  $M_C$  and eventually

---

\* See Ref. [3] for  $D$ -term contributions through GUT breaking and Ref. [4] for anomalous  $U(1)$  breaking.

receives radiative corrections governed by the SM gaugino masses, which are flavor-blind. Hence, we would have degenerate mass spectrum of sfermions like the ‘no-scale’ model (at least for the first two families). This possibility has already been mentioned in Ref. [5]. In this scenario, soft scalar masses are to be controlled by the flavor mechanism that generates hierarchical Yukawa couplings, no matter how SUSY breaking is mediated and no matter what initial conditions of soft SUSY-breaking terms are. This approach, which we shall pursue in this paper, is quite opposite to the usual scenario in which degenerate soft scalar masses are supposed to be derived by a flavor-blind mediation of SUSY breaking.

When the pure SCFT is perturbed by the SM gauge interactions and the SM gaugino masses, each sfermion mass is not completely suppressed, but converges on a flavor-dependent value [10, 11]. The convergent value is one-loop suppressed since the SM gauge couplings are perturbatively small. It is then plausible that sfermion masses at the weak scale may be calculated solely in terms of the SM gauge coupling constants and gaugino masses. In fact, under some assumptions, we can estimate the degeneracy factor  $\Delta_{\tilde{f}}$  of sfermion masses up to a single model-dependent parameter  $\Gamma_i$  [10]. It was found that for  $M_C > 10^{10}$  GeV, the degeneracy factor is 0.005 – 0.01 for squarks and 0.05 – 0.1 for sleptons. In particular, the right-handed sleptons are not well degenerate in mass. Note that these degeneracy factors are evaluated in the sfermion basis.

In this paper, we first examine the assumptions that are implicit in the estimation of sfermion mass degeneracy in the present SCFT approach. We also discuss to what extent the degeneracy is required in the Nelson-Strassler scenario. To this aim, we take into account the fact that FCNC processes, when correctly evaluated in the fermion basis, have additional suppression since the Nelson-Strassler mechanism leads to hierarchical Yukawa matrices.

It is natural to extend the analysis to grand unified theories (GUTs) and to examine how much degeneracy of sfermion masses is achieved by coupling to SCFTs. That is our second purpose; we explicitly study sfermion mass degeneracy within the framework of GUTs coupled to SC sectors. We take the  $SU(5)$  and the flipped  $SU(5) \times U(1)$  as a prototype of GUT models. We shall show how much degeneracy of sfermion masses is expected in each case. It turns out that a simple extension of the MSSM to the  $SU(5)$  makes the degeneracy of the right-handed sleptons worse, because each right-handed lepton is embedded into a higher-dimensional representation. The situation is different in the flipped  $SU(5) \times U(1)$  case, since the right-handed leptons remain  $SU(5)$ -singlets [12], and the bino mass  $M_1$  is determined by a combination of two independent gaugino masses of  $SU(5) \times U(1)$ .

This paper is organized as follows. After a brief review on the Nelson-Strassler mechanism in section 2.1, we outline in subsection 2.2 the results of Ref. [10] on degeneracy of sfermion masses in the sfermion basis within the framework of the MSSM coupled to SC sectors. We also discuss under what conditions the degeneracy factor can be estimated in a simple manner. The FCNC constraints are examined in subsection 2.3 by taking into account Yukawa-diagonalizing matrices. In sections 3 and 4, we extend the model to GUTs and investigate generic features. Specifically we argue that the degeneracy of the right-handed sleptons becomes worse in  $SU(5)$  models than the MSSM. We then show how the degeneracy for the right-handed sleptons can be improved in the flipped  $SU(5) \times U(1)$  models. In section 5, we briefly discuss various sorts of threshold effects, which might affect the previous results. This in particular includes  $D$ -term contributions on sfermion masses in the flipped  $SU(5) \times U(1)$  models. Section 6 is devoted to conclusion and discussion.

We note that Luty and Sundrum [13] discuss a scenario of ‘conformal sequestering’, which is another interesting approach to the SUSY flavor problem based on four-dimensional SCFTs. Although closely related to the present SCFT approach, it is slightly different in that the SCFT couples to messenger fields of SUSY breaking, not to quarks and leptons.

## 2 MSSM coupled to SC sector

### 2.1 Nelson-Strassler mechanism for Yukawa hierarchy

Here we give a brief review on the Nelson-Strassler (NS) mechanism that generates the hierarchical structure of Yukawa couplings [5]. We assume two sectors: One is the SM sector and the other is the SC sector. The SM sector has the gauge group  $SU(3) \times SU(2) \times U(1)_Y$ , or its extension, and contains three families of quarks and leptons  $\psi_i$  ( $i = 1, 2, 3$ ) as well as Higgs fields  $H$ . The SC sector has a gauge group  $G_{SC}$  and matter fields  $\Phi_r$ . The fields  $\psi_i$  and  $H$  are taken to be  $G_{SC}$ -singlets. The following superpotential is assumed;

$$W = y_{ij} \psi_i \psi_j H + \lambda_{irs} \psi_i \Phi_r \Phi_s + \dots , \quad (2.1)$$

where the first term describes the ordinary Yukawa couplings in the SM sector and the ellipsis denotes terms including only  $\Phi_r$ . The second term represents the couplings of quarks and leptons  $\psi_i$  to the SC sector, which we refer to as *messenger couplings*. For the messenger coupling  $\lambda_{irs}$  to be allowed by gauge invariance, either of  $\Phi_r$  or  $\Phi_s$  should belong to a nontrivial representation under the SM gauge group.

With sufficiently many matter fields, the  $G_{\text{SC}}$  gauge theory resides in ‘conformal window’; the SC gauge coupling  $g'$  has an IR fixed point [6, 7]. Suppose that the messenger couplings  $\lambda_{irs}$  as well as  $g'$  approach IR fixed points. At this new fixed point, the field  $\psi_i$  gains a large anomalous dimension  $\gamma_i^* = \mathcal{O}(0.1-1)$  through superconformal dynamics. As a result, the Yukawa couplings in the SM sector obey the power law and behave roughly like

$$y_{ij}(M_C) \approx y_{ij}(M_0) \left( \frac{M_C}{M_0} \right)^{\gamma_i^* + \gamma_j^*} . \quad (2.2)$$

Here  $M_0$  is the cut-off scale, at which  $y_{ij}(M_0) = \mathcal{O}(1)$  is expected. Thus the hierarchical structure of Yukawa couplings can be generated by family-dependent anomalous dimensions  $\gamma_i^*$ . The resultant Yukawa matrices are similar to the ones obtained by the Froggatt-Nielsen (FN) mechanism; large anomalous dimension in the NS mechanism corresponds to  $U(1)$  charge in the FN mechanism. As stressed in Ref. [5], the unitarity of the SCFT guarantees that the anomalous dimensions  $\gamma_i^*$  at the SC fixed point are always non-negative, whereas the non-negativity of  $U(1)$  charges is just the assumption in the conventional FN mechanism.

Since the top Yukawa coupling should not be suppressed, the top quark as well as the up-type Higgs field must not couple to the SC sector. Although the bottom quark and tau lepton as well as the down-type Higgs could couple, we will mainly consider, in what follows, the models in which only the first two families couple to the SC sector.

## 2.2 Degeneracy of sfermion masses

Next we outline, following Ref. [10], how convergent values of soft scalar masses and sfermion mass degeneracy can be estimated. See also Ref. [14] for a review. We also discuss some subtleties about such estimation.

The suppression of sfermion masses in the SCFT approach follows from a general property of the renormalization group equations (RGEs) of soft SUSY-breaking parameters [15, 14]. Let us concentrate for a moment on the diagonal elements  $m_i^2$  of a sfermion mass-squared matrix (in a flavor basis in which fermion Yukawa matrix takes the form (2.2)). Near an IR attractive fixed point of pure SCFT, the RGE of  $m_i^2$  takes the form

$$\mu \frac{dm_i^2}{d\mu} = \mathcal{M}_{ij} m_j^2 , \quad (2.3)$$

where the coefficient matrix  $\mathcal{M}_{ij}$  encodes full effects of the SC dynamics, and can be calculated by use of the ‘Grassmanian expansion method’ if the anomalous dimension  $\gamma_i$  is

known as a function of coupling constants  $g'$  and  $\lambda_{irs}$ . This matrix is positive-definite (non-negative) since the fixed point is IR attractive. It follows that certain combinations of  $m_i^2$  are exponentially suppressed. Moreover, *each*  $m_i^2$  is suppressed if its anomalous dimension  $\gamma_i^*$  is uniquely determined [5, 10] by fixed point equations  $\beta_{g'} = \beta_\lambda = 0$ . We assume that in each model considered below, there exists a proper set of couplings that satisfies this condition.

When we switch on the gauge couplings  $\alpha_a = g_a^2/8\pi^2$  and gaugino masses  $M_a$  in the SM sector ( $a = 1, 2, 3$ ), the RGE (2.3) gets modified, at the leading order, into

$$\mu \frac{dm_i^2}{d\mu} = \mathcal{M}_{ij} m_j^2 - \sum_{a=1,2,3} C_{ia} \alpha_a M_a^2, \quad (2.4)$$

where  $C_{ia} = 4C_2(R_{ia})$  with the quadratic Casimir coefficient  $C_2(R_{ia})$  of the  $R_{ia}$  representation, and we have neglected for simplicity a possible contribution from the  $U(1)_Y$  Fayet-Iliopoulos term  $\mathcal{S} \equiv \text{Tr}(Y m_i^2)$ . Eq. (2.4) implies that each sfermion mass eventually converges on one-loop suppressed value of the order  $\alpha_a M_a^2$ . The convergent values generally depend on detailed structure of the SC sector, since the above RGEs are coupled equations for soft scalar masses in the SC sector as well as those of squarks and sleptons.

We are interested in the convergent value of squark and slepton masses at the scale  $M_C$  where the SC sector decouples. Let  $m_{\tilde{f}_i}$  denote soft scalar mass of the  $i$ -th family of squark or slepton  $\tilde{f}$ . Then the convergent value of  $m_{\tilde{f}_i}^2$  at  $M_C$  can be written in the form [10]

$$m_{\tilde{f}_i}^2 \longrightarrow \frac{1}{\Gamma_{\tilde{f}_i}} \sum_{a=1,2,3} C_{fa} \alpha_a(M_C) M_a^2(M_C). \quad (2.5)$$

In this expression,  $C_{fa}$  is the Casimir factor for  $f = Q, u, d, L, e$ , and the prefactor  $\Gamma_{\tilde{f}_i}^{-1}$  summarizes the structure of the SC sector. Typically, we find  $\Gamma_i \sim \gamma_i^* = \mathcal{O}(0.1-1)$  for squarks and sleptons.<sup>†</sup> If  $\gamma_i^*$  and  $\gamma_j^*$  are different from each other, the factors  $\Gamma_{\tilde{f}_i}$  and  $\Gamma_{\tilde{f}_j}$  are also different. Only the prefactor  $\Gamma_{\tilde{f}_i}^{-1}$  has flavor-dependency. Consequently, the difference between the first and second families is given by

$$m_{\tilde{f}_2}^2(M_C) - m_{\tilde{f}_1}^2(M_C) = \left( \frac{1}{\Gamma_{\tilde{f}_2}} - \frac{1}{\Gamma_{\tilde{f}_1}} \right) \sum_{a=1,2,3} C_{fa} \alpha_a(M_C) M_a^2(M_C), \quad (2.6)$$

which is also one-loop suppressed compared with  $M_a^2(M_C)$ .

We have considered only the diagonal elements of sfermion mass-squared matrix. However, as was shown in Ref. [11] for pure SCFT, off-diagonal elements are also exponentially

---

<sup>†</sup> This is not always true for sfermions in SC sector.

suppressed. Even when we switch on SM effects, the RGEs of off-diagonal elements, unlike Eq. (2.4), contain no contribution from the SM gaugino masses at the leading order. Thus the off-diagonal elements converge on sufficiently small values, which we can safely neglect.

In Ref. [10], sfermion mass degeneracy was examined by assuming the MSSM field content below  $M_C$ . Here let us recall some results for later convenience. The sfermion mass receives radiative correction  $\Delta m_{\tilde{f}i}^2$  between  $M_C$  and  $M_Z$ , which is evaluated to be

$$\Delta m_{\tilde{f}i}^2(M_C \rightarrow M_Z) = \sum_{a=1,2,3} a_{fa} I_a(M_Z, M_C) M_a^2(M_C), \quad (2.7)$$

$$I_a(M_Z, M_C) \equiv 1 - \frac{\alpha_a^2(M_Z)}{\alpha_a^2(M_C)}, \quad a_{fa} \equiv \frac{C_{fa}}{2b_a}. \quad (2.8)$$

Here  $b_a$  are the MSSM gauge beta-function coefficients, and the factors  $a_{fa}$  ( $a = 1, 2, 3$ ) are shown in the second, third and fourth columns of Table 1 for each matter field. These radiative corrections are much larger than the convergent value  $m_{\tilde{f}i}^2(M_C)$ . It follows that

$$\left. \frac{m_{\tilde{f}2}^2 - m_{\tilde{f}1}^2}{m_{\tilde{f}2}^2 + m_{\tilde{f}1}^2} \right|_{M_Z} = \frac{1}{2} \left( \frac{1}{\Gamma_{\tilde{f}2}} - \frac{1}{\Gamma_{\tilde{f}1}} \right) \Delta_{\tilde{f}}, \quad (2.9)$$

where we define the degeneracy factor  $\Delta_{\tilde{f}}$  by

$$\Delta_{\tilde{f}} = \frac{\sum_a C_{\tilde{f}a} \alpha_a(M_C) M_a^2(M_C)}{\sum_a a_{fa} I_a(M_Z, M_C) M_a^2(M_C)}. \quad (2.10)$$

The factor  $\Delta_{\tilde{f}}$  serves as an estimate of how much degeneracy of sfermion masses is achieved in the present framework. Moreover, it is a calculable quantity independently of detailed structure of the SC sector, especially when the SM gaugino masses satisfy the ‘GUT’ relation,  $M_a/\alpha_a = \text{constant}$  for  $a = 1, 2, 3$ . For example, taking  $M_C = 10^{16}$  GeV gives [10]

$$\begin{aligned} \Delta_{\tilde{Q}} &= 8 \times 10^{-3}, & \Delta_{\tilde{u}} &= \Delta_{\tilde{d}} = 6 \times 10^{-3}, \\ \Delta_{\tilde{L}} &= 5 \times 10^{-2}, & \Delta_{\tilde{e}} &= 1 \times 10^{-1}. \end{aligned} \quad (2.11)$$

Unfortunately, the degeneracy factor  $\Delta_{\tilde{e}}$  for the right-handed slepton is rather large. The primary reason is that the radiative correction to the right-handed slepton mass, which involves only  $M_1$ , is smaller than the others.<sup>‡</sup> However, this result does not necessarily imply that the present SCFT approach to Yukawa hierarchy leads to significant lepton flavor-violation, because the actual size of lepton flavor-violating processes depends on an explicit form of lepton Yukawa matrix, as we discuss below.

<sup>‡</sup> The degeneracy factor  $\Delta_{\tilde{e}}$  for the right-handed sleptons can be somewhat reduced [10] if there is a sizable contribution from the  $U(1)_Y$  Fayet-Iliopoulos term  $\mathcal{S}$  (with a suitable sign). Note that we generally expect  $\mathcal{S} \neq 0$  since soft scalar masses of the third family and Higgs fields are not constrained by SC dynamics.

$f$	$a_{f3}$	$a_{f2}$	$a_{f1}$	rep. under $SU(5)$	$C_{f5}$
$Q$	$-8/9$	$3/2$	$1/198$	$\mathbf{10}$	$72/5$
$u$	$-8/9$	$0$	$8/99$	$\mathbf{10}$	$72/5$
$d$	$-8/9$	$0$	$2/99$	$\bar{\mathbf{5}}$	$48/5$
$L$	$0$	$3/2$	$1/22$	$\bar{\mathbf{5}}$	$48/5$
$e$	$0$	$0$	$2/11$	$\bar{\mathbf{10}}$	$72/5$

Table 1: Group-theoretical factors for  $\Delta_{\bar{f}}$ . Our normalization convention for the  $U(1)$  gauge coupling is the  $SU(5)$ -motivated one,  $\alpha_1 \equiv (5/3)\alpha_Y$ .

Some remarks are to be added here. The convergent value (2.5) is determined by the SM one-loop term in the RGE (2.4). In general, *higher-loop terms like  $(\lambda_*^2/8\pi^2)^n \times \alpha_a M_a^2$  are potentially large because the fixed point value of the messenger coupling  $\lambda_{irs}$  in Eq. (2.1) is not necessarily small*. In particular, slepton mass  $m_{\tilde{e}}^2(M_C)$  will be affected by a correction of order  $\alpha_3 M_3^2$  if the right-handed lepton  $e_R^c$  couples to colored SC fields through the messenger interactions. Nevertheless, the presence or absence of such corrections depends on the structure of the SC sector; for instance, there is no two-loop term of  $\lambda_e^2 \alpha_3 M_3^2$  if *both* of  $\Phi_r$  and  $\Phi_s$  are  $SU(3)$  singlets in the messenger interaction  $e_R^c \Phi_r \Phi_s$ . In the following analysis, we will assume that such higher-loop terms are negligible or at most comparable to the one-loop term.<sup>§</sup> Another remark is that there might arise large threshold effects when the strongly-coupled SC sector decouples. In the expression (2.6), we have implicitly assumed that these are negligible or flavor independent. We shall comment on such effects in section 5.

In passing, we note the sparticle mass spectrum at the weak scale. As far as the first two families of sfermions are concerned, the convergent values at  $M_C$  are quite small,  $m_{\tilde{f}_i}^2 \approx 0$ , and the mass spectrum is similar to that in ‘no-scale’ scenario, provided that there is no large correction at the SC threshold. For  $M_C = 10^{16}$  GeV, we have

$$(m_{\tilde{Q}}, m_{\tilde{L}}, m_{\tilde{e}}) = (0.91, 0.25, 0.13) M_3 . \quad (2.12)$$

The other  $SU(2)$ -singlet squarks have masses similar to  $m_{\tilde{Q}}$ . Again we have taken  $\mathcal{S} = 0$  for simplicity although nonzero  $\mathcal{S}$  is helpful to avoid charged lightest superparticle (LSP) [10].

<sup>§</sup> In the MSSM case with GUT relation of gaugino masses, we do not expect that the presence of the  $\alpha_3 M_3^2$  term significantly changes the previous estimation of  $\Delta_{\tilde{L}}$  and  $\Delta_{\tilde{e}}$ , unless it is associated with a large group-theoretical factor.



On the other hand, the third family, in particular the top quark, as well as Higgs fields do not couple to the SC sector. In general, their soft scalar masses depend on initial values as well as the details of the RG running of gauge and Yukawa couplings. Hence we restrict ourselves to the first and second families of sfermion masses in the following analysis.

### 2.3 Mixing angles and FCNC constraints

The factor  $\Delta_{\tilde{f}}$  represents a simple estimate of sfermion mass difference in the sfermion basis. To confront the NS scenario with the experimental bounds on FCNC processes [16], we still have to evaluate sfermion mass matrices in a basis that diagonalizes fermion Yukawa matrices; specifically we are interested in the mixings  $(\delta_{12})_{\text{LL,RR}}$  between the first two families of left-handed, or right-handed sfermions. In this respect, the NS scenario has an advantage that both of fermion and sfermion mass matrices are known if the anomalous dimensions are specified. If the first two families have the same anomalous dimension  $\gamma_{f_1}^* = \gamma_{f_2}^*$  at the SC fixed point, Yukawa-diagonalizing angle will be of  $\mathcal{O}(1)$ , but at the same time,  $\Gamma_{\tilde{f}_1} = \Gamma_{\tilde{f}_2}$  guarantees the complete mass degeneracy of sfermions. On the other hand, if  $\gamma_{f_1}^* > \gamma_{f_2}^*$  and  $\Gamma_{\tilde{f}_1} > \Gamma_{\tilde{f}_2}$ , the sfermions have non-degeneracy estimated by  $(\Gamma_{\tilde{f}_2}^{-1} - \Gamma_{\tilde{f}_1}^{-1}) \Delta_{\tilde{f}}$ , but the diagonalizing angle is as small as  $(M_C/M_0)^{\gamma_1^* - \gamma_2^*}$ , which gives an additional suppression to FCNC processes. In this way, we expect that approximate alignment happens.

For squarks, the degeneracy factors are fairly small already in the sfermion basis. Moreover, if the diagonalizing angles are of the order of the Cabibbo angle, we have  $(\delta_{12}^d)_{\text{LL,RR}} \sim 0.22 \times \Delta_{\tilde{f}} \approx 1 \times 10^{-3}$ . Using the constraint from the  $K^0 - \bar{K}^0$  system, *i.e.*,

$$(\delta_{12}^d)_{\text{LL,RR}} < 1.2 \times 10^{-3} \times \left( \frac{m_{\tilde{d}}}{500 \text{ GeV}} \right), \quad (2.13)$$

it is required that  $m_{\tilde{d}} \gtrsim 500 \text{ GeV}$ , which corresponds to  $M_3 \gtrsim 500 \text{ GeV}$  as well as  $m_{\tilde{e}} \gtrsim 50 \text{ GeV}$ . Thus FCNC constraints will easily be satisfied in squark sector.

On the other hand, the degeneracy factors are not small enough in slepton sector. Then lepton flavor-violating processes such as  $\mu \rightarrow e\gamma$  decay constrain the slepton masses. Of course, such constraints depend on the texture of lepton Yukawa matrix generated by the NS mechanism. To see this explicitly, let us concentrate on the first two families and denote by  $\theta_L$  and  $\theta_R$  the mixing angles for left- and right-handed leptons, respectively. First consider the case in which left-handed leptons receive the same anomalous dimension  $\gamma_{L1} = \gamma_{L2}$  from the SC dynamics. Then the first  $2 \times 2$  lepton Yukawa matrix takes, up to  $\mathcal{O}(1)$  factors, the

‘lopsided’ form

$$y_{ij} \approx y_{22} \begin{pmatrix} \varepsilon & 1 \\ \varepsilon & 1 \end{pmatrix}, \quad \varepsilon = \left( \frac{M_C}{M_0} \right)^{\gamma_{e1}^* - \gamma_{e2}^*} \sim \frac{m_e}{m_\mu} = 5 \times 10^{-3}, \quad (2.14)$$

where the lepton mass hierarchy originates from the anomalous dimensions of the right-handed leptons. In this case, the left-handed leptons have  $\mathcal{O}(1)$  mixing and that will be favorable from the viewpoint of neutrino oscillation. Interestingly, the left-handed sleptons are completely degenerate since  $\gamma_{L1}^* = \gamma_{L2}^*$  implies  $\Gamma_{L1} = \Gamma_{L2}$ . Moreover, the right-handed leptons have the mixing angle of  $\theta_R = \mathcal{O}(\varepsilon)$ , which gives an additional suppression to  $(\delta_{12}^\ell)_{RR}$ . Thus we will have no strong constraint.

If mixing angles are larger than  $\mathcal{O}(10^{-3})$ , the constraint becomes significant. For example, if both of mixing angles for left- and right-handed sectors are of the order

$$\theta_L \sim \theta_R \sim \sqrt{\frac{m_e}{m_\mu}} = 7 \times 10^{-2}, \quad (2.15)$$

we have  $(\delta_{12}^\ell)_{LL} \sim 7 \times 10^{-2} \Delta_{\tilde{f}} \approx 3 \times 10^{-3}$  and  $(\delta_{12}^\ell)_{RR} \sim 7 \times 10^{-2} \Delta_{\tilde{f}} \approx 7 \times 10^{-3}$ , and it is required that  $m_{\tilde{l}} \gtrsim 100$  GeV and  $m_{\tilde{e}} \gtrsim 200$  GeV from the constraint

$$(\delta_{12}^\ell)_{LL, RR} < 2.0 \times 10^{-3} \times \left( \frac{m_{\tilde{l}}}{100 \text{ GeV}} \right)^2. \quad (2.16)$$

The constraint on  $(\delta_{12}^\ell)_{RR}$  requires that the gluino and squarks are heavier than 2 TeV, whereas that on  $(\delta_{12}^\ell)_{LL}$  corresponds to the gluino and squarks heavier than 400 GeV. In general, the larger mixing angle the right-handed lepton sector has, the heavier sleptons are required from  $\mu \rightarrow e\gamma$  decay. For example, it is required that  $m_{\tilde{e}} \gtrsim 300$  GeV if  $\theta_R \gtrsim 0.2$ .

We remark that there are additional constraints coming from  $(\delta_{12})_{LR}$ , which are related to SUSY-breaking trilinear couplings  $h_{ij} \tilde{\psi}_i \tilde{\psi}_j H$ . In the present SCFT framework, soft scalar masses are well controlled by the SC dynamics, but the so-called  $A$ -terms  $A_{ij} \equiv h_{ij}/y_{ij}$  are not; the SC dynamics does give a suppression factor to  $h_{ij}$ , which is the same as the suppression factor of the corresponding Yukawa coupling  $y_{ij}$ . Therefore, even if the constraints on  $(\delta_{12})_{LL, RR}$  are satisfied, we should still take care of additional constraints on  $(\delta_{12})_{LR}$  from FCNC processes [11]. We need another mechanism to suppress  $A$ -terms, or to realize the complete alignment\* of  $A_{ij}$ . At any rate, we expect that the constraints coming from  $(\delta_{12})_{LR}$

---

\* One way to achieve this possibility was discussed in Ref. [17], where the Yukawa couplings  $y_{ij}$  by themselves are supposed to have infrared fixed points. In models with extra dimensions, it was also argued [18] that thanks to power-law behavior due to Kaluza-Klein modes, hierarchical Yukawa couplings can be obtained at the IR fixed points. In this case, however, it seems difficult to realize finite mixing angles.

will be less severe than those from  $(\delta_{12})_{\text{LL,RR}}$ , and to examine the latter is the subject of the following sections.

### 3 $SU(5)$ GUT coupled to SC sector

For  $M_C > 10^{16}$  GeV, we are naturally led to consider the embedding of the MSSM into SUSY GUTs. In this section, we study the Georgi-Glashow  $SU(5)$  model, which breaks down to the MSSM at the scale  $M_X = 2 \times 10^{16}$  GeV. We assume  $M_C \geq M_X$ , *i.e.*, the SC sector decouples at a larger scale than the GUT breaking scale.

As in the MSSM case of subsection 2.2, we expect that each sfermion mass converges on

$$m_{\tilde{f}_i}^2(M_C) = \frac{1}{\Gamma_{\tilde{f}_i}} C_{f5} \alpha_5(M_C) M_5^2(M_C) , \quad (3.1)$$

where  $\alpha_5(M_C)$  and  $M_5(M_C)$  are the  $SU(5)$  gauge coupling and gaugino mass at the scale  $M_C$ . Representations of quarks and leptons under the  $SU(5)_{\text{GG}}$  are shown in the fifth column of Table 1. The sixth column shows  $C_{f5}$  for each matter field. On the other hand, the radiative correction in the ‘GUT regime’ between  $M_C$  and  $M_X$  is obtained as

$$\Delta m_{\tilde{f}_i}^2(M_C \rightarrow M_X) = \frac{C_{f5}}{2b_5} I_5(M_X, M_C) M_5^2(M_C) , \quad (3.2)$$

where  $b_5$  is the beta-function coefficient of  $\alpha_5$ , and  $I_5(M_X, M_C)$  is defined by

$$I_5(M_X, M_C) \equiv 1 - \frac{\alpha_5^2(M_X)}{\alpha_5^2(M_C)} . \quad (3.3)$$

Then the sfermion mass degeneracy may be estimated by the use of

$$\Delta_{\tilde{f}} = \frac{C_{f5} \alpha_5(M_C) M_5^2(M_C)}{\Delta m_{\tilde{f}_i}^2(M_C \rightarrow M_X) + \Delta m_{\tilde{f}_i}^2(M_X \rightarrow M_Z)} , \quad (3.4)$$

where  $\Delta m_{\tilde{f}_i}^2(M_X \rightarrow M_Z)$  denotes the radiative correction in the MSSM regime, Eq. (2.7) with  $M_C = M_X$ . The value of  $\Delta_{\tilde{f}}$  depends on the flow of the gauge couplings,  $\alpha_5$  and  $\alpha_{a=1,2,3}$ , but not the gaugino masses.

Fig. 1 shows  $\Delta_{\tilde{f}}$  for squarks and Fig. 2 for sleptons. For definiteness, we have taken  $b_5 = -9/2$  corresponding to the minimal  $SU(5)$ , while different  $b_5$  lead to similar results.

Compared with the MSSM case, Eqs. (2.11)–(2.12), the mass spectrum at the weak scale is almost the same, but the degeneracy factors are not. First of all, the degeneracy factors




Figure 1:  $\Delta_{\tilde{Q}}$ ,  $\Delta_{\tilde{u}}$  and  $\Delta_{\tilde{d}}$  against  $M_C$  in the supersymmetric  $SU(5)$  GUT.

for squarks and left-handed slepton become slightly large. Note also that  $\Delta_{\tilde{Q}}$  and  $\Delta_{\tilde{u}}$  are larger than  $\Delta_{\tilde{d}}$ , since  $Q$  and  $u$  belong to the  $\mathbf{10}$  representation under the  $SU(5)$  whereas  $d$  belongs to the  $\mathbf{\bar{5}}$ , and the former has a larger convergent value than the latter. There is a small splitting between  $\Delta_{\tilde{Q}}$  and  $\Delta_{\tilde{u}}$  because  $m_{\tilde{Q}}^2$  gets slightly larger radiative corrections in the MSSM regime from the wino mass  $M_2$ .




Figure 2:  $\Delta_{\tilde{e}}$  and  $\Delta_{\tilde{L}}$  against  $M_C$  in the supersymmetric  $SU(5)$  GUT.

More importantly, the degeneracy for the right-handed slepton becomes drastically worse than before. The reason is very simple; the right-handed lepton is embedded into the  $\mathbf{10}$ , which has a large Casimir coefficient, so that the convergent value (3.1) is enlarged. Conse-

quently the FCNC constraint for  $(\delta_{12}^l)_{\text{RR}}$  cannot easily be satisfied, requiring heavier sleptons and/or a specific form of lepton Yukawa matrix. It is rather surprising that the simple group-theoretical fact precludes the  $SU(5)$  extension of the MSSM. At any rate, the embedding into the  $SU(5)$  discussed here would remove one of most interesting properties of the present SCFT approach, that is, approximate mass degeneracy of sfermions without specifying a mediation mechanism of SUSY breaking.

## 4 Flipped $SU(5) \times U(1)$ coupled to SC sector

We have observed that within the framework of the ordinary  $SU(5)$  GUT, the degeneracy factor  $\Delta_{\tilde{e}}$  is not small enough for the right-handed slepton because of its large Casimir coefficient. From this viewpoint, it is interesting to consider the flipped  $SU(5) \times U(1)$  models [12], where the right-handed lepton is not embedded into a nontrivial  $SU(5)$  representation, but is an  $SU(5)$ -singlet  $S_i$ .

Let us first recall the basic feature of the minimal flipped  $SU(5)_F \times U(1)_F$  model to fix our notation. The quantum number of matter and Higgs fields is shown in Table 2, and the embedding of each family of matter fields is explicitly given by

$$T = \begin{pmatrix} 0 & d_{\text{R}3}^c & -d_{\text{R}2}^c & u_1 & d_1 \\ & 0 & d_{\text{R}1}^c & u_2 & d_2 \\ & & 0 & u_3 & d_3 \\ & & & 0 & \nu_{\text{R}}^c \\ & & & & 0 \end{pmatrix}, \quad \bar{F} = \begin{pmatrix} u_{\text{R}1}^c \\ u_{\text{R}2}^c \\ u_{\text{R}3}^c \\ -e \\ \nu \end{pmatrix}, \quad S = e_{\text{R}}^c, \quad (4.1)$$

where  $\nu_{\text{R}}^c$  is a right-handed neutrino. If the  $\nu_{\text{R}}^c$  components of the 10-dimensional Higgs fields  $H$  and  $\bar{H}$  develop the vacuum expectation values, the symmetry breaking to the MSSM can be achieved with a simple implementation of the missing partner mechanism for doublet-triplet splitting [19]. Through this symmetry breaking, the 24-th generator  $T_{\text{F}}^{24} \equiv \sqrt{3/5} \tilde{Y}$  of  $SU(5)_F$  and the  $U(1)_F$  generator  $F \equiv \sqrt{40} \tilde{X}$  are related to the hypercharge  $Y$  as well as its orthogonal broken generator  $X$ . One way to describe this relation is to embed the gauge group into  $SO(10)$  and to represent  $U(1)$  generators in the  $SO(10)$  basis. By picking its  $SU(4)_{\text{PS}} \times SU(2)_{\text{L}} \times SU(2)_{\text{R}}$  subgroup, we have

$$\tilde{X} = \sqrt{\frac{3}{5}} T_{\text{PS}}^{15} + \sqrt{\frac{2}{5}} T_{\text{R}}^3, \quad \sqrt{\frac{3}{5}} \tilde{Y} = \sqrt{\frac{2}{5}} T_{\text{PS}}^{15} - \sqrt{\frac{3}{5}} T_{\text{R}}^3. \quad (4.2)$$

Here  $T_{\text{R}}^3$  is the third generator of  $SU(2)_{\text{R}}$  and  $T_{\text{PS}}^{15} = \sqrt{3/8} (B - L)$  is the 15-th generator of the Pati-Salam  $SU(4)$ , which is proportional to the  $B - L$  charge. Eq. (4.2) is precisely the

	MSSM content	$SU(5)_F$ rep.	$U(1)_F$ charge
$T_i$	$d_i, Q_i, \nu_{Ri}^c$	<b>10</b>	+1
$\bar{F}_i$	$u_i, L_i$	<b><math>\bar{5}</math></b>	-3
$S_i$	$e_{Ri}^c$	<b>1</b>	+5
$h$	$h_d^c, h_d$	<b>5</b>	-2
$\bar{h}$	$h_u^c, h_u$	<b><math>\bar{5}</math></b>	+2
$H$	—	<b>10</b>	+1
$\bar{H}$	—	<b><math>\bar{10}</math></b>	-1

Table 2: The minimal flipped  $SU(5)_F \times U(1)_F$  model: The last column shows  $F \equiv \sqrt{40} \tilde{X}$ , where  $\tilde{X}$  is the  $SO(10)$ -normalized  $U(1)_F$  charge of Eq. (4.2).

flipped version of the usual  $SO(10)$  relation

$$X = \sqrt{\frac{3}{5}} T_{\text{PS}}^{15} - \sqrt{\frac{2}{5}} T_{\text{R}}^3, \quad \sqrt{\frac{3}{5}} Y = \sqrt{\frac{2}{5}} T_{\text{PS}}^{15} + \sqrt{\frac{3}{5}} T_{\text{R}}^3, \quad (4.3)$$

where  $X$  is nothing but the  $U(1)$  generator of  $SO(10)/SU(5)_{\text{GG}}$ , often called  $U(1)_X$ . Since these two sets of  $U(1)$  generators are orthogonally related to each other, we have

$$X = \frac{1}{5} \tilde{X} + \frac{\sqrt{24}}{5} T_{\text{F}}^{24}, \quad \sqrt{\frac{3}{5}} Y = \frac{\sqrt{24}}{5} \tilde{X} - \frac{1}{5} T_{\text{F}}^{24}. \quad (4.4)$$

For notational simplicity, let  $\alpha'_5$  and  $\alpha'_1$  denote the gauge couplings of  $SU(5)_F \times U(1)_F$ . With the normalization of  $U(1)$  generators as above, the embedding into the  $SO(10)$  GUT would lead to  $\alpha'_1 = \alpha'_5$ , although we do not assume such a further unification.

We assume that the SC sector decouples at the scale  $M_C$ , below which the model is exactly the same as the minimal flipped  $SU(5) \times U(1)$  model. At the scale  $M_X$  where  $SU(5) \times U(1)$  breaks to the MSSM gauge group, the hypercharge gauge coupling  $\alpha_Y = (3/5) \alpha_1$  is matched with  $SU(5) \times U(1)$  gauge couplings  $\alpha'_5$  and  $\alpha'_1$  according to

$$\frac{15}{\alpha_Y} \equiv \frac{25}{\alpha_1} = \frac{1}{\alpha'_5} + \frac{24}{\alpha'_1}. \quad (4.5)$$

The measured value of the SM gauge couplings requires  $\alpha'_1(M_X) \approx \alpha'_5(M_X) \approx (2\pi \times 24.5)^{-1}$ . The matching condition for the corresponding gaugino masses is given by

$$\frac{M_1}{\alpha_1} = \frac{1}{25} \left( \frac{M'_5}{\alpha'_5} + \frac{24M'_1}{\alpha'_1} \right) = \left( \frac{1+24r}{25} \right) \frac{M'_5}{\alpha'_5} = \left( \frac{1+24r}{25r} \right) \frac{M'_1}{\alpha'_1}. \quad (4.6)$$

Here we have introduced an RG-invariant quantity

$$r \equiv \frac{M'_1/\alpha'_1}{M'_5/\alpha'_5} \approx \frac{M'_1(M_X)}{M'_5(M_X)}, \quad (4.7)$$

which parametrizes the relative size of two independent gaugino masses of  $SU(5) \times U(1)$ .  $SO(10)$  gauge symmetry would require  $r = +1$ , but we treat  $r$  as a free parameter. The gluino mass  $M_3$  and the wino mass  $M_2$  are matched as usual and are independent of  $M'_1$ . However, the bino mass  $M_1$  does depend on  $r$ , and the ratio  $M_1/M_3$  is given by

$$\frac{M_1}{M_3} = \left( \frac{1+24r}{25} \right) \frac{\alpha_1}{\alpha_3} = \left( \frac{1+24r}{25} \right) \frac{5\alpha_Y}{3\alpha_3}. \quad (4.8)$$

We expect that the degeneracy factor for the right-handed sleptons also depend on  $r$ , since their masses will be dominantly governed by the  $U(1)$  gaugino masses.

It is straightforward to estimate sfermion masses and confirm our expectation. The RGE for soft scalar mass, which includes the leading term of  $SU(5) \times U(1)$  gauge loop, is

$$\mu \frac{dm_i^2}{d\mu} = \mathcal{M}_{ij} m_j^2 - \sum_{a=5,1} C'_{ia} \alpha'_a M_a'^2, \quad (4.9)$$

where  $C'_{ia}$  denote the Casimir factors under  $SU(5) \times U(1)$ . Under the assumption that higher order corrections to the second term is small, we may estimate the convergent value of each sfermion mass at  $M_C$  and the degeneracy factor at the weak scale by

$$m_{\tilde{f}i}^2(M_C) = \frac{1}{\Gamma_{\tilde{f}i}} \sum_{a=5,1} C'_{fa} \alpha'_a(M_C) M_a'^2(M_C), \quad (4.10)$$

$$\Delta_{\tilde{f}} = \frac{\sum_{a=5,1} C'_{fa} \alpha'_a(M_C) M_a'^2(M_C)}{\Delta m_{\tilde{f}i}^2(M_C \rightarrow M_X) + \Delta m_{\tilde{f}i}^2(M_X \rightarrow M_Z)}. \quad (4.11)$$

The group-theoretical factors are shown in Table 3, and the radiative corrections in  $SU(5) \times U(1)$  regime can be calculated in a manner similar to Eq. (2.7) with  $b'_5 = -5$  and  $b'_1 = 15/2$ . The results does not strongly depend on  $M_C$ , and we will take  $M_C = M_X$  for definiteness.

We are especially interested in the expected degree of the degeneracy  $\Delta_{\tilde{e}}$  for the right-handed sleptons. The point is that the convergent value  $m_{\tilde{e}}^2(M_C)$  is governed by the flipped  $U(1)$  gaugino mass  $M'_1$ , whereas the radiative corrections  $\Delta m_{\tilde{e}}^2$  are determined by the bino mass  $M_1$ , which contains as a component the  $SU(5)$  gaugino mass  $M'_5$  as well as  $M'_1$  according to the matching condition (4.6). Consequently, the degeneracy factor  $\Delta_{\tilde{e}}$  behaves like

$$\Delta_{\tilde{e}} \propto \left( \frac{M'_1}{M_1} \right)^2 \propto \left( \frac{25r}{1+24r} \right)^2. \quad (4.12)$$

$f$	$SU(5) \times U(1)$	$C'_{f5}$	$C'_{f1}$	$6\tilde{Y}$	$\sqrt{40}X$
$Q$	$(\mathbf{10}, +1)$	$72/5$	$1/10$	$+1$	$+1$
$d_{\text{R}}^c$	$(\mathbf{10}, +1)$	$72/5$	$1/10$	$-4$	$-3$
$\nu_{\text{R}}^c$	$(\mathbf{10}, +1)$	$72/5$	$1/10$	$+6$	$+5$
$L$	$(\bar{\mathbf{5}}, -3)$	$48/5$	$9/10$	$-3$	$-3$
$u_{\text{R}}^c$	$(\bar{\mathbf{5}}, -3)$	$48/5$	$9/10$	$+2$	$+1$
$e_{\text{R}}^c$	$(\mathbf{1}, +5)$	$0$	$5/2$	$0$	$+1$

Table 3: The group-theoretical factors in the flipped  $SU(5) \times U(1)$  model: the last two column shows the ‘flipped hypercharge’  $\tilde{Y}$  and the  $U(1)_X$  charge  $X$  in Eqs. (4.2)–(4.3).

Numerical values for  $\Delta_{\tilde{e}}$  are shown in Fig. 3 as a function of  $r$ , and we find explicitly

$$\Delta_{\tilde{e}} = \begin{cases} 6.6 \times 10^{-2} & \text{for } r = 1/8 \\ 1.1 \times 10^{-1} & \text{for } r = 1 \end{cases}. \quad (4.13)$$

Compared with the MSSM case (2.11), a similar value of  $\Delta_{\tilde{e}}$  is obtained in the  $r = 1$  case, as is expected from the fact that  $r \approx 1$  corresponds to the usual GUT relation of gaugino masses. As we decrease  $r$  with keeping  $M'_5$  fixed,  $M'_1$  decreases linearly in  $r$ , whereas  $M_1$  reaches a nonzero value and thus  $\Delta_{\tilde{e}}$  becomes small. In other words, *the mass difference of  $\tilde{e}_{\text{R}}^c$  is determined by a small gaugino mass of the flipped  $U(1)$  while the  $SU(5)$  component of the bino mass effectively enhances the averaged slepton mass.* In this way, we find that the degeneracy factor  $\Delta_{\tilde{e}}$  is improved for  $r \lesssim 0.5$  compared with the MSSM case. For  $|r| \gg 1$ , the bino mass  $M_1$  becomes proportional to  $M'_1$ , and  $\Delta_{\tilde{e}}$  approaches its maximal 0.12.

At first sight, the degeneracy factor  $\Delta_{\tilde{e}}$  could be arbitrarily small if  $M'_1 \ll M'_5$ . However, this is not true because our estimation based in the RGE (4.9) is no longer reliable for a hierarchically small value of  $|r| \lesssim 0.1$ . At two-loop level, for instance, the RGE contains potentially dangerous terms like  $\lambda_e^2 \alpha'_5 M_5'^2$  and  $\alpha'_1 \alpha'_5 M'_1 M'_5$ , where  $\lambda_e$  is the messenger coupling of the  $SU(5)$ -singlet  $e_{\text{R}}^c$ . As mentioned in subsection 2.2, the former term is absent if we assume that only  $SU(5)$ -singlet fields couple to  $e_{\text{R}}^c$  through the messenger interaction. On the other hand, the latter term will give a substantial correction to  $\Delta_{\tilde{e}}$  for such a small value of  $|r|$ . Even if the latter correction is properly taken into account, the ratio  $M_1/M_3$  would be unacceptably small from the viewpoint of the naturalness.






Figure 3: The slepton degeneracy factor  $\Delta_{\tilde{e}}$  against  $r$  in the flipped  $SU(5) \times U(1)$  model. The dotted line shows the value in the MSSM case.




Figure 4:  $m_{\tilde{e}}/M_3$  against  $r$  in the flipped  $SU(5) \times U(1)$  model.

The slepton mass at the weak scale is approximately given by

$$m_{\tilde{e}} = 0.93M_1, \quad (4.14)$$

as long as the radiative correction  $\Delta m_{\tilde{e}}^2$  dominates over the convergent value. Eq. (4.8) then implies that the ratio  $m_{\tilde{e}}/M_3$  decrease as we decrease  $r$ , as is shown in Fig. 4.

For the other sfermions, the degeneracy factors and mass spectrum are determined dominantly by  $\alpha_a$  and  $M_a$  ( $a = 2, 3$ ), and are almost independent of  $r$  for a moderate range of the parameter  $r \lesssim \mathcal{O}(2.5)$ . Explicitly, we obtain for  $r = 1/8$ ,

$$\begin{aligned} \Delta_{\tilde{Q}} &= 1.5 \times 10^{-2}, & \Delta_{\tilde{L}} &= 1.3 \times 10^{-1}, \\ \Delta_{\tilde{d}} &= 1.6 \times 10^{-2}, & \Delta_{\tilde{u}} &= 1.1 \times 10^{-2}, \end{aligned} \quad (4.15)$$

which are almost the same as in the  $SU(5)$  case, except that  $\tilde{u}$  and  $\tilde{d}$  are interchanged. The

mass spectrum for squarks and left-handed sletpons is almost the same as in Eq. (2.12).

Noted that  $\Delta_{\tilde{L}}$  is larger than  $\Delta_{\tilde{e}} \leq 0.12$  owing to the embedding into  $\bar{\mathbf{5}}$ . As discussed in subsection 2.3, the constraints from lepton flavor violations depend on the lepton mixing angles. For mixing angles (2.15), the constraints from  $(\delta_{12}^\ell)_{\text{LL}}$  require  $m_{\tilde{e}} \gtrsim 200$  GeV, which should be compared with  $m_{\tilde{e}} \gtrsim 200$  GeV from  $(\delta_{12}^\ell)_{\text{RR}}$  in the MSSM with the same mixing angles. The former corresponds to  $M_3 \gtrsim 1$  TeV, while the latter requires  $M_3 \gtrsim 2$  TeV. Thus the flipped  $SU(5) \times U(1)$  model, unlike the  $SU(5)$  models, is on the level of the MSSM.

## 5 On threshold effects

Up to now, we have smoothly connected RG flows of soft scalar masses at  $M_X$  as well as  $M_C$ . We reexamine this procedure and briefly discuss possible threshold effects of various sorts.

### 5.1 $D$ -term contributions in flipped models

When the rank of gauge group is reduced, the gauge symmetry breaking induces the  $D$ -term contribution to soft scalar mass-squared,  $\Delta_D m_f^2 = q_f \langle D \rangle$ , which is proportional to the charge of broken  $U(1)$  symmetry [3, 4]. Here we consider effects of such additional terms at  $M_X$  induced by  $SU(5) \times U(1)$  breaking. Of course, the charges under the broken symmetry are the same between different families, and those do not create non-degeneracy. However, such additional contributions change overall magnitude of sfermion masses at the GUT threshold, and also change the mass spectrum and degeneracy factors at the weak scale.

The  $D$ -term contributions induced through the breaking of  $SU(5) \times U(1)$  can be expressed in terms of the broken generators  $F \equiv \sqrt{40} \tilde{X}$  and  $\tilde{Y} \equiv \sqrt{5/3} T_F^{24}$  in Eq. (4.2) as

$$\Delta_D m_{\tilde{f}}^2 = \frac{1}{5} \left( F_f \alpha_1'^2 + 24 \tilde{Y}_f \alpha_5'^2 \right) m_D^2 . \quad (5.1)$$

The magnitude of  $m_D^2$  can be calculated when we explicitly fix the model and its breaking pattern. To be generic, however, let us take  $m_D^2$  to be a free parameter. With the help of  $\alpha_5'(M_X) \approx \alpha_1'(M_X)$ , the above expression can be rewritten into

$$\Delta_D m_{\tilde{f}}^2 \approx \frac{\sqrt{40}}{5} \left( \tilde{X}_f + \sqrt{24} T_F^{24} \right) \alpha_1'^2 m_D^2 = \sqrt{40} X_f \langle D_X \rangle , \quad (5.2)$$

where  $X$  is the  $U(1)_X$  generator (4.4) in the  $SO(10)$  normalization, and  $\langle D_X \rangle \equiv \alpha_5'^2 m_D^2$ . Using the value of  $U(1)_X$  charges shown in Table 3, we have

$$\Delta_D m_{\tilde{L}}^2 = -3 \langle D_X \rangle , \quad \Delta_D m_{\tilde{e}}^2 = \langle D_X \rangle , \quad (5.3)$$

which give the opposite effect on left- and right-handed slepton masses. Thus it is not possible to improve the degeneracy for both of them at the same time by the  $D$ -term contribution.\*

Eq. (4.14) indicates that the right-handed sleptons tend to be the LSP. Therefore a positive contribution  $\Delta_D m_{\tilde{e}}^2 > 0$  is favored to make them heavier, and this slightly improves the degeneracy for the right-handed sleptons. However, the degeneracy for left-handed sleptons becomes worse. For instance, we have  $\Delta_{\tilde{L}} = 2.1 \times 10^{-1}$  if we take  $\langle D_\chi \rangle = 0.1 M_5^2(M_X)$ .

## 5.2 Note on SC threshold effects

When strongly-coupled SC sector decouples around the scale  $M_C$ , there arises another type of threshold corrections, which we refer to as SC threshold effects. We have little to say about such effects; one could in principle evaluate SC threshold effects once the SC sector is specified, but it requires hard work of understanding strong dynamics. Basically we do not expect that Yukawa hierarchy generated by large anomalous dimensions is modified so much by SC threshold effects, provided that the SC sector decouples quickly [5]. We have assumed that the same is true in our estimation of sfermion masses and degeneracy.

There is a special case in which SC threshold effects are reliably calculated. Consider for definiteness the MSSM coupled to SC sector, whose decoupling is caused by invariant mass terms of the order  $M_C$ . Suppose also that all the soft SUSY-breaking parameters are given *purely* by anomaly mediation [21]. Even in this special case, one can evaluate as before the ‘convergent value’ of sfermion mass, which is actually the same as anomaly-mediated one calculated by using beta-functions of the MSSM coupled to the SC sector. On the other hand, as was argued in Ref. [22], the mass parameter  $M_C$  should be extended to a background (non-dynamical) superfield, and its  $F$ -term will affect soft terms. One then expects that there arise SC threshold effects such that after the decoupling, sfermion masses are precisely anomaly-mediated one calculated in the MSSM framework. Thus the SC threshold effects are calculable in this case, although they are of no interest because of tachyonic sleptons. Note that the latter sfermion masses and the expected convergent values are of the same order of magnitudes,  $m_{\tilde{f}}^2 = \mathcal{O}(\alpha_a M_a^2)$ .

There is an interesting puzzle here. Originally we are interested in the SCFT approach because it can provide degenerate sfermion spectrum no matter how SUSY is mediated; if

---

\*  $D$ -term contribution through the breaking  $E_6 \rightarrow SO(10)$  can be used [20] to increase the slepton masses, which are originally tachyonic in a scenario of anomaly-mediated SUSY breaking [21]. Such contributions might also be helpful here since the broken  $U(1)$  charge takes the same sign for all of quarks and leptons.

we consider RG flows of each sfermion mass corresponding to various initial conditions at the cut-off scale, all the flows converge on a certain value thanks to the SC dynamics. Such RG flows would contain a special flow corresponding to anomaly-mediated spectrum. Now, suppose that SUSY breaking is mediated *not* purely by anomaly mediation. Nevertheless, the above convergence property of RG flows would imply that the sfermion spectrum at the decoupling scale, and thus SC threshold effects, be almost the same as anomaly-mediated one. Specifically the MSSM coupled to SC sector would always lead to tachyonic sleptons no matter how SUSY is initially mediated.

The resolution of this apparent puzzle is that the convergent value of each sfermion mass  $m_f^2$  is independent of the initial conditions for  $m_f^2$ , but does depend on the SM gaugino masses. Therefore sfermion mass spectrum can be different from anomaly-mediated one, provided that the SM gaugino masses are different.

At any rate, a lesson is that SC threshold effects would do affect sfermion masses, but their size would be at most comparable to the convergent values used in the previous sections. Therefore we do not expect that our results would be modified substantially. It is interesting to confirm this expectation by explicit calculations.

## 6 Conclusion and discussion

We have studied sfermion mass degeneracy within the framework of GUTs coupled to SC sectors. In the  $SU(5)$  model, the degeneracy factor  $\Delta_\varepsilon$  for the right-handed sleptons becomes worse than in the MSSM. The reason is that the right-handed lepton  $e_R^c$  is embedded into 10-dimensional representation and the convergent mass value is enlarged. Models with larger gauge group like  $SO(10)$  and  $E_6$  will have the same feature, as long as the SC sector decouples at the scale larger than the GUT breaking scale. One way to keep  $e_R^c$  a non-Abelian singlet is the flipped  $SU(5)_F \times U(1)_F$  model. In this model, the degeneracy of the right-handed sleptons can be improved if the  $U(1)_F$  gaugino mass  $M'_1$  is smaller than the  $SU(5)_F$  one; a small  $M'_1$  reduces the convergent value and the mass difference, while the  $SU(5)_F$  component of the bino mass enhances the averaged slepton mass.

In this paper, we have estimated sfermion mass degeneracy without specifying the model for the SC sector. To this end, we have assumed that the convergent value of sfermion masses at the decoupling scale is dominately determined by the SM one-loop terms in the RGEs. This assumption is plausible for the  $SU(5)$  case, but is crucial especially for the right-handed

sleptons in the MSSM and the flipped model. A clever model building will be required when the SC sector is vector-like under the SM-sector gauge group. We have also assumed that our estimation is not substantially modified by SC threshold effects. We expect that the size of such effects will be at most comparable to the estimated convergent value of sfermion masses. It is worth while examining these points by explicit calculations.

Finally let us speculate about possible extensions of the present work. We have neglected effects from the third family of quarks and leptons and their Yukawa couplings. In quark sector, it will be in safe to neglect the third family, since they have only small mixings with the others in many realistic Yukawa matrices. However, the same is not true in lepton sector. In particular, the flipped  $SU(5) \times U(1)$  model requires that the tau neutrino Yukawa coupling is as large as the top Yukawa coupling. In general, the large mixings in lepton sector induce a significant flavor violation  $\delta_{12}^\ell$  through the radiative corrections above the mass scale  $M_\nu$  of the right-handed neutrinos [23, 24, 25, 26], and that constrains sfermion mass spectrum and/or requires a specific form of Yukawa matrices. From this viewpoint, an interesting possibility is that a proper coupling to SC sector and subsequent IR convergence property can eliminate such flavor violation as well if  $M_C$  is taken below  $M_\nu = 10^{13} - 10^{15}$  GeV. We need further study concerning neutrino Yukawa couplings.\*

In models with a product gauge group like  $SU(5) \times U(1)$ , the degeneracy can be improved if the gaugino masses do not satisfy the usual GUT relation so that the convergent value of sfermion mass is much suppressed compared with the radiative corrections in the MSSM regime. The  $SU(5) \times U(3)_H$  models [28] will have a similar property. We speculate that even with a simple gauge group  $SU(5)$ , such a suppression of the convergent value could be realized if one supposes that the  $SU(5)$  gauge multiplets live in an extra-dimensional space-times. In this setup, the power-law evolution of gauge coupling and gaugino mass [29, 30, 31] makes them drastically small above  $M_X$  (in such a way that  $M_5/\alpha_5$  is still RG invariant [30]) provided that the gauge coupling is asymptotically free. Consequently the degeneracy would be improved if  $M_C$  is taken as the energy scale where  $\alpha_5$  and  $M_5$  are very small. To keep our RG calculations reliable in the SC regime, quarks and leptons as well as SC-sector fields are supposed to live in four dimensions. Such a setup could also explain why our gauge coupling  $\alpha_5$  is smaller than that in the SC sector, thanks to extra-dimensional volume suppression.

The Nelson-Strassler scenario, which we have focused in this paper, assumes that anomalous dimensions of quarks and leptons take flavor-dependent values at the SC fixed point.

---

\* An application of the SC idea to the neutrino as well as Higgs sectors was discussed in Ref. [27].

The difference between  $\gamma_i^*$  and  $\gamma_j^*$  leads to hierarchical Yukawa matrices, but at the same time, the difference between  $\Gamma_i$  and  $\Gamma_j$  leads to a distinctive non-degeneracy of sfermion masses. In the ‘Yukawa hierarchy transfer’ scenario [32], the SC dynamics is supposed to be flavor-blind at the fixed point,  $\gamma_i^* = \gamma_j^*$  and  $\Gamma_i = \Gamma_j$ , ensuring complete degeneracy of sfermion masses. The hierarchy among Yukawa couplings  $y_{ij}$  is derived by assuming the inverse hierarchy in the *messenger* couplings  $\lambda_{irs}$  at the cut-off scale and by *transferring* the initial hierarchy through the SC dynamics. Moreover, the assumed hierarchy can be derived, *e.g.*, through the FN mechanism in the SC sector *without* spoiling the sfermion degeneracy by family-dependent  $D$ -term contributions. This new scenario provides an alternative way of realizing degenerate sfermion spectrum and hierarchical Yukawa matrices at the same time.

## Acknowledgments

The authors would like to thank J. Kubo, N. Maekawa, J. Sato, D. Suematsu, M. Tanimoto and T. Yanagida for useful discussions. We would like to express our thanks to the organizers of Summer Institute 2001, Yamanashi, where a part of the work was done.

## References

- [1] C.D. Froggatt and H.B. Nielsen, Nucl. Phys. **B147** (1979) 277.
- [2] L.E. Ibáñez and G.G. Ross, Phys. Lett. **B332** (1994) 100.
- [3] M. Drees, Phys. Lett. **B181** (1986) 279;  
 J. S. Hagelin and S. Kelley, Nucl. Phys. **B342** (1990) 95;  
 Y. Kawamura, H. Murayama and M. Yamaguchi, Phys. Lett. **B324** (1994) 94;  
 Phys. Rev. D51 (1995) 1337.
- [4] H. Nakano, hep-th/9404033; Prog. Theor. Phys. Suppl. **123** (1996) 387;  
 Y. Kawamura and T. Kobayashi, Phys. Lett. **B375** (1996) 141, Erratum-ibid. **B388**  
 (1996) 867; Phys. Rev. **D56** (1997) 3844;  
 E. Dudas, S. Pokorski and C.A. Savoy, Phys. Lett. **B369** (1996) 255;  
 E. Dudas, C. Grojean, S. Pokorski and C.A. Savoy, Nucl. Phys. **B481** (1996) 85.
- [5] A.E. Nelson and M.J. Strassler, JHEP **0009** (2000) 030.

- [6] T. Banks and A. Zaks, Nucl. Phys. **B196** (1982) 189.
- [7] N. Seiberg, Nucl. Phys. **B435** (1995) 129.
- [8] A. Karch, T. Kobayashi, J. Kubo and G. Zoupanos, Phys. Lett. **B 441** (1998) 235.
- [9] M.A. Luty and R. Rattazzi, JHEP **9911** (1999) 001.
- [10] T. Kobayashi and H. Terao, Phys. Rev. **D64** (2001) 075003.
- [11] A. E. Nelson and M. J. Strassler, hep-ph/0104051.
- [12] S.M. Barr, Phys. Lett. **112** (1982) 219;  
 J.P. Derendinger, J.E. Kim and D.V. Nanopoulos, Phys. Lett. **139** (1984) 170;  
 I. Antoniadis, J. Ellis, J.S. Hagelin and D.V. Nanopoulos, Phys. Lett. **194** (1987) 231.
- [13] M.A. Luty and R. Sundrum, hep-th/0105137; hep-th/0111231.
- [14] H. Terao, hep-ph/0112021.
- [15] Y. Yamada, Phys. Rev. **D 50** (1994) 3537;  
 J. Hisano and M. Shifman, Phys. Rev. **D 56** (1997) 5475;  
 I. Jack and D.R.T. Jones, Phys. Lett. **B 415** (1997) 383; I. Jack, D.R.T. Jones and  
 A. Pickering, Phys. Lett. **B 426** (1998) 73; Phys. Lett. **B 432** (1998) 114;  
 L.V. Avdeev, D.I. Kazakov and I.N. Kondrashuk, Nucl. Phys. **B 510** (1998) 289;  
 T. Kobayashi, J. Kubo and G. Zoupanos, Phys. Lett. **B 427** (1998) 291;  
 N. Arkani-Hamed, G.F. Giudice, M.A. Luty and R. Rattazzi, Phys. Rev. **D 58** (1998)  
 115005;  
 D.I. Kazakov and V.N. Velizhanin, Phys. Lett. **B485** (2000) 393.
- [16] F. Gabbiani, E. Gabrielli, A. Masiero and L. Silverstrini, Nucl. Phys. **B477** (1996) 321;  
 See also for a review, *e.g.*,  
 M. Misiak, S. Pokorski and J. Rosiek, hep-ph/9703442; J.L. Feng, hep-ph/010112,  
 and references therein.
- [17] T. Kobayashi and K. Yoshioka, Phys. Rev. **D 62** (2000) 115003.
- [18] M. Bando, T. Kobayashi, T. Noguchi and K. Yoshioka, Phys. Lett. **B480** (2000) 187;  
 Phys. Rev. **D63** (2001) 113017.

- [19] A. Masiero, D.V. Nanopoulos, K. Tamvakis and T. Yanagida, Phys. Lett. **B115** (1982) 380.
- [20] M. Carena, K. Huitu and T. Kobayashi, Nucl. Phys. **B592** (2000) 164.
- [21] L. Randall and R. Sundrum, Nucl. Phys. **B557** (1999) 79;  
G.F. Giudice, M.A. Luty, H. Murayama and R. Rattazzi, JHEP **9812** (1998) 27.
- [22] A. Pomarol and R. Rattazzi, JHEP **9905** (1999) 013.
- [23] F. Borzumati and A. Masiero, Phys. Rev. Lett. **57** (1986) 961.
- [24] J. Hisano, T. Moroi, K. Tobe, M. Yamaguchi and T. Yanagida, Phys. Lett. **B357** (1995) 579;  
J. Hisano, T. Moroi, K. Tobe and T. Yanagida, Phys. Rev. **D53** (1996) 2442.
- [25] J. Sato, K. Tobe and T. Yanagida, Phys. Lett. **B498** (2001) 189;  
J. Sato and K. Tobe, Phys. Rev. **D63** (2001) 116010.
- [26] A. Kageyama, S. Kaneko, N. Shimoyama and M. Tanimoto, hep-ph/0110283.
- [27] J. Kubo and D. Suematsu, Phys. Rev. **D64** (2001) 115014.
- [28] T. Yanagida, Phys. Lett. **B344** (1995) 211;  
T. Hotta, Izawa K.-I. and T. Yanagida, Phys. Rev. **D3913** (1996); Phys. Rev. **D54** (1996) 6970;  
Izawa K.-I. and T. Yanagida, Prog. Theor. Phys. **97** (1997) 913;  
Y. Imamura, T. Watari and T. Yanagida, Phys. Rev. **D64** (2001) 065023.
- [29] K. Dienes, E. Dudas and T. Gherghetta, Phys. Lett. **B436** (1998) 311; Nucl. Phys. **B537** (1999) 47.
- [30] T. Kobayashi, J. Kubo, M. Mondragon and G. Zoupanos, Nucl. Phys. **B550** (1999) 99.
- [31] J. Kubo, H. Terao and G. Zoupanos, Nucl. Phys. **B574** (2000) 495.
- [32] T. Kobayashi, H. Nakano and H. Terao, Phys. Rev. **D65** (2002) 015006;  
T. Kobayashi, H. Nakano and H. Terao, in preparation.