

Crystalline Color Superconductivity: Effective Lagrangian and Phonon Dispersion Law

R. Casalbuoni^{a,b}, R. Gatto^c and G. Nardulli^{d,e}

^a*Dipartimento di Fisica, Università di Firenze, I-50125 Firenze, Italia*

^b*I.N.F.N., Sezione di Firenze, I-50125 Firenze, Italia*

^c*Départ. de Physique Théorique, Université de Genève, CH-1211 Genève 4,
Suisse*

^d*Dipartimento di Fisica, Università di Bari, I-70124 Bari, Italia*

^e*I.N.F.N., Sezione di Bari, I-70124 Bari, Italia*

Abstract

The recent study by Bowers and Rajagopal on the possible structures for the QCD crystalline color superconducting phase favors a cubic structure and allows for a first discussion of the QCD dynamics in the crystalline phase. The cubic structure leads to complete breaking of traslation invariance and to three phonon modes. We discuss the expected form of the effective Lagrangian and the phonon dispersion laws assuming the proposed crystal structure. Besides the interest for QCD and the related possible relevance to compact stars these considerations may become useful to atomic systems exhibiting crystalline superfluidity.

1 Introduction

In a recent paper Bowers and Rajagopal [1] have presented a comprehensive study of the different possible crystal structures for the crystalline color superconducting phase of QCD and concluded in favor of a face centered cubic condensate pattern. This study now allows discussion of the dynamics of QCD in the crystalline phase and this note is a first contribution in that direction.

The subject of the color superconducting phases of QCD [2] [3] has only recently had its most extensive developments [4]. At asymptotically large values of the chemical potential for quark number μ , cold QCD with three degenerate flavors lives in the color-flavor locked (CFL) phase [5] corresponding to condensation of quark pairs of opposite momenta and lying near the Fermi surfaces (BCS pairing).

For finite strange quark mass the CFL phase is expected to persist within some range below asymptotic densities. Decreasing the value of μ , the expected possibility becomes pairing between quarks of non vanishing total momentum, each quark momentum lying near the respective (now differing) Fermi surfaces [6] [7] [8] [9] [10]. Closeness to the Fermi surface will count for a lower free energy. Quantitative studies so far are mostly for two flavors and a difference in their chemical potential $\delta\mu$ is usually introduced to induce the difference in the Fermi momenta.

Similar possibilities of fermion pairing of non vanishing total momentum were already encountered in condensed matter (ferromagnetic systems with paramagnetic impurities) [11], so that the name LOFF phase is often used, from the initials of the authors of ref. [11].

In a LOFF phase translational and rotational invariance are spontaneously broken. The most relevant question for crystalline superconductivity is that of the space dependence of the Cooper pair condensate. Preliminary studies have been based on a plane wave ansatz: a single plane wave with the finite momentum of the Cooper pairs, assuming that all the momenta spontaneously choose a unique direction. These studies (see ref [6] and [11]) have lead to a picture of the LOFF phase existing within a certain window of $\delta\mu$ which starts with a first order phase transition and ends at the upper end with a second order transition.

The authors of [1] develop a Ginzburg-Landau expansion near the second order transition. They introduce a multiple plane-wave ansatz and find that the favorite crystal structure is that of a face-centered cubic crystal.

For the simple single plane wave structure, so far adopted as a provisional working hypothesis, we had constructed [8] the effective low energy phonon lagrangian and derived the phonon dispersion law. In that case only one phonon was expected from the partial breaking of translation invariance. For the cubic structure now derived in [1] one will have three phonons, due to the complete breaking of translation invariance.

The work [1] uses a simplified model with pointlike interaction and two massless quarks and, as we have said, a Ginzburg-Landau approximation. The

lowest value of the Ginzburg-Landau free energy corresponds to a crystal structure built up from eight plane waves directed to the corners of a cube. In spite of the approximations used one has the qualitative feeling that their proposed crystal structure would remain the favorite one also in a more general QCD setting (although not perhaps at asymptotic densities).

In this note we shall discuss the expected form of the effective Lagrangian and phonon dispersion laws assuming the cubic crystal structure of [1].

We have already mentioned the theoretical interest of color superconductivity in the crystalline phase as one of the general components of the QCD phase diagram. Physically, crystalline color superconductivity might be realized within compact stellar systems. In fact the quark chemical potential is expected to increase in penetrating towards the center of a compact star and other parameters, such as the strange quark mass, are also expected to evolve in such a way that one may reach the crystalline phase inside a core region of the star. In a rotating compact star the ensuing vortices may give rise to observable glitches, as discussed in ref. [6].

The considerations developed in connection to crystalline superconductivity in QCD, including those presented in this note, may also be useful in the field of atomic systems, in particular, as mentioned in ref. [1], for gases of fermionic atoms below their Fermi temperature [12] [13]. These systems present over QCD the advantage of a better parametric flexibility [14].

2 The LOFF phase of QCD

According to [1] the favored crystalline structure in the LOFF phase of QCD should be a face-centered-cubic one. This is characterized by eight \vec{q}_j vectors with equal length $q = \pi/a$ and directions corresponding to the eight vertices of the cube in Fig. 1:

$$\Delta(\vec{x}) = \Delta \sum_{j=1}^8 \exp\{2i\vec{q}_j \cdot \vec{x}\} = \Delta \sum_{\epsilon_i=\pm} \exp\left\{\frac{2\pi i}{a}(\epsilon_1 x_1 + \epsilon_2 x_2 + \epsilon_3 x_3)\right\} . \quad (1)$$




Fig. 1 - The figure shows some of the symmetry axes of the cube, denoted as C_2 , C_3 and C_4 (see the text). We show also the parameterization used for the coordinates of the vertices.

In Fig. 1 some of the symmetry axes of the cube are shown: they are denoted as C_4 (the three 4-fold axes), C_3 (the four 3-fold axes) and C_2 (the six 2-fold axes). For more details on the symmetry group of the cube see the book by Hamermesh [15].

We observe that the coordinates at the vertices are nothing but the directions of the \vec{q}_j vectors. The condensate (1) can be also written as follows:

$$\begin{aligned} \Delta(\vec{x}) = 2\Delta & \left[\cos \frac{2\pi}{a}(x_1 + x_2 + x_3) + \cos \frac{2\pi}{a}(x_1 - x_2 + x_3) \right. \\ & \left. + \cos \frac{2\pi}{a}(x_1 + x_2 - x_3) + \cos \frac{2\pi}{a}(-x_1 + x_2 + x_3) \right]. \end{aligned} \quad (2)$$

This condensate breaks both translations and rotations. It is however invariant under the discrete group O_h , the symmetry group of the cube. This can be seen by noticing that the previous expression is invariant under the following coordinate transformations

$$\begin{aligned} R_1 : & \quad x_1 \rightarrow x_1, \quad x_2 \rightarrow x_3, \quad x_3 \rightarrow -x_2, \\ R_2 : & \quad x_1 \rightarrow -x_3, \quad x_2 \rightarrow x_2, \quad x_3 \rightarrow x_1, \\ R_3 : & \quad x_1 \rightarrow x_2, \quad x_2 \rightarrow -x_1, \quad x_3 \rightarrow x_3, \\ I : & \quad x_1 \rightarrow -x_1, \quad x_2 \rightarrow -x_2, \quad x_3 \rightarrow -x_3, \end{aligned} \quad (3)$$

that is rotations of $\pi/2$ around the coordinate axes, and the inversion with respect to the origin. Since one can easily check that the group O_h is generated by the previous 4 elements the invariance follows at once.

We can then introduce 3 scalar fields $\Phi^{(i)}$ such that

$$\langle \Phi^{(i)} \rangle_0 = \frac{2\pi}{a} x_i \quad (4)$$

These fields should be defined on the lattice sites, but for the purpose of writing the effective action for such fields, we are interested only in the low energy behavior $p \ll 1/a$. Therefore we will treat them as continuous fields. The coupling of the Fermi fields generated by the condensate will be written as

$$\Delta\psi^T C \psi \sum_{\epsilon_i=\pm} \exp \left\{ i(\epsilon_1 \Phi^{(1)} + \epsilon_2 \Phi^{(2)} + \epsilon_3 \Phi^{(3)}) \right\} \quad (5)$$

making the theory invariant under translations and rotations. These invariances are broken spontaneously in the vacuum defined by eq. (4).

The effective lagrangian for the fields $\Phi^{(i)}$ has to enjoy the following symmetries: rotational and translational invariance; O_h symmetry on the fields $\Phi^{(i)}$. The latter requirement follows from the invariance of the coupling (5) under the group O_h acting upon $\Phi^{(i)}$.

We then define

$$\Phi^{(i)}(x) = \frac{2\pi}{a} x^i + \phi^{(i)}(x)/f, \quad (6)$$

with

$$\langle \phi^{(i)}(x) \rangle_0 = 0. \quad (7)$$

and f the decay constant of the phonon with dimension of an energy. It follows that, after the breaking, the phonon fields $\phi^{(i)}(x)$ and the coordinates x^i must transform under the diagonal discrete group obtained from the direct product of the rotation group acting over the coordinates and the O_h group acting over $\Phi^{(i)}(x)$. This is indeed the symmetry left after the breaking of translational and rotational invariance.

In order to build up the effective lagrangian, we start noticing that if X_i are quantities transforming as the same representation of the $\Phi^{(i)}$'s under O_h (a 3-dimensional representation), then all the invariant expressions can be obtained by the following three basic invariant expressions

$$I_2(X_i) = X_1^2 + X_2^2 + X_3^2, \quad I_4(X_i) = X_1^2 X_2^2 + X_2^2 X_3^2 + X_3^2 X_1^2, \quad I_6(X_i) = X_1^2 X_2^2 X_3^2. \quad (8)$$

The invariance is easily checked by noticing that these expressions are invariant under the three elementary rotations R_i and the inversion I (see eq. (3)). Quantities transforming as $\Phi^{(i)}$ are

$$\dot{\Phi}^{(i)} \equiv \frac{d\Phi^{(i)}}{dt}, \quad \vec{\nabla}\Phi^{(i)}. \quad (9)$$

We proceed now as in ref. [8], by noticing that the most general low-energy effective lagrangian, invariant under rotations, translations and O_h is given by

$$\mathcal{L} = \frac{f^2}{2} \sum_{i=1,2,3} (\dot{\Phi}^{(i)})^2 + \mathcal{L}_s(I_2(\vec{\nabla}\Phi^{(i)}), I_4(\vec{\nabla}\Phi^{(i)}), I_6(\vec{\nabla}\Phi^{(i)})) . \quad (10)$$

The reason for allowing all the possible spatial gradients in this expansion is that the vacuum expectation value of $\vec{\nabla}\Phi^{(i)}$ is given by x_i and it is not in general small.

Define the auxiliary quantities

$$X_i(\epsilon) = \frac{2\pi}{a} \vec{e}^{(i)} + \epsilon \vec{\nabla}\phi^{(i)} \quad (11)$$

such that

$$X_i(1/f) = \vec{\nabla}\Phi^{(i)}, \quad (12)$$

where $\vec{e}^{(i)}$ is a unit vector along the axis "i". The ϵ parameter has been introduced to keep track of the spatial derivatives. Since we are interested at the second order in these derivatives we need the quantity (first order terms are total derivatives)

$$\frac{1}{2} \frac{\partial^2 \mathcal{L}(\epsilon)}{\partial \epsilon^2} \Big|_{\epsilon=0} = \frac{1}{2} \left[\sum_{i,j=2,4,6} \frac{\partial^2 \mathcal{L}}{\partial I_i \partial I_j} \frac{\partial I_i}{\partial \epsilon} \frac{\partial I_j}{\partial \epsilon} + \sum_{i=2,4,6} \frac{\partial \mathcal{L}}{\partial I_i} \frac{\partial^2 I_i}{\partial \epsilon^2} \right]_{\epsilon=0} \quad (13)$$

where $\mathcal{L}(\epsilon)$ has been obtained by the substitution $\vec{\nabla}\Phi^{(i)} \rightarrow X_i(\epsilon)$. Since the derivatives of \mathcal{L} with respect to the invariant terms I_i , evaluated at $\epsilon = 0$ are just constant, using

$$\frac{\partial I_2}{\partial \epsilon} \Big|_{\epsilon=0} = 2 \sum_{i=1,2,3} \partial_i \phi^{(i)}, \quad \frac{\partial I_4}{\partial \epsilon} \Big|_{\epsilon=0} = 4 \sum_{i=1,2,3} \partial_i \phi^{(i)}, \quad \frac{\partial I_6}{\partial \epsilon} \Big|_{\epsilon=0} = 2 \sum_{i=1,2,3} \partial_i \phi^{(i)} \quad (14)$$

and

$$\begin{aligned}
\frac{1}{2} \frac{\partial^2 I_2}{\partial \epsilon^2} \Big|_{\epsilon=0} &= \sum_{i=1,2,3} |\vec{\nabla} \phi^{(i)}|^2 \\
\frac{1}{2} \frac{\partial^2 I_4}{\partial \epsilon^2} \Big|_{\epsilon=0} &= 4 \sum_{i<j=1,2,3} \partial_i \phi^{(i)} \partial_j \phi^{(j)} + 2 \sum_{i=1,2,3} |\vec{\nabla} \phi^{(i)}|^2 \\
\frac{1}{2} \frac{\partial^2 I_6}{\partial \epsilon^2} \Big|_{\epsilon=0} &= 4 \sum_{i<j=1,2,3} \partial_i \phi^{(i)} \partial_j \phi^{(j)} + \sum_{i=1,2,3} |\vec{\nabla} \phi^{(i)}|^2,
\end{aligned} \tag{15}$$

we see that the expression (13) depends only on the quantities

$$\sum_{i=1,2,3} |\vec{\nabla} \phi^{(i)}|^2, \quad \sum_{i=1,2,3} (\partial_i \phi^{(i)})^2, \quad \sum_{i<j=1,2,3} \partial_i \phi^{(i)} \partial_j \phi^{(j)}. \tag{16}$$

In conclusion one gets

$$\mathcal{L} = \frac{1}{2} \sum_{i=1,2,3} (\dot{\phi}^{(i)})^2 - \frac{a}{2} \sum_{i=1,2,3} |\vec{\nabla} \phi^{(i)}|^2 - \frac{b}{2} \sum_{i=1,2,3} (\partial_i \phi^{(i)})^2 - c \sum_{i<j=1,2,3} \partial_i \phi^{(i)} \partial_j \phi^{(j)}. \tag{17}$$

This expression could have been guessed by noticing that after the breaking one still has an invariance under the group O_h acting at the same time upon the coordinates and the phonon fields $\phi^{(i)}$.

3 Dispersion laws

To analyze the phonon dispersion law one has to diagonalize the matrix

$$M = \begin{pmatrix} a |\vec{p}|^2 + b p_1^2 & c p_1 p_2 & c p_1 p_3 \\ c p_1 p_2 & a |\vec{p}|^2 + b p_2^2 & c p_2 p_3 \\ c p_1 p_3 & c p_2 p_3 & a |\vec{p}|^2 + b p_3^2 \end{pmatrix}. \tag{18}$$

The dispersion law can be written as follows ($\vec{p} = \vec{n} p$):

$$E = v_r(\vec{n}) p \tag{19}$$

where $v_r^2(\vec{n})$ are given by

$$v_r^2(\vec{n}) = a + \frac{b}{3} - u_r + \frac{J_4 - \left(\frac{b}{3}\right)^2}{u_r} \tag{20}$$

and u_r are the three cubic roots defined by

$$u_r(\vec{n}) = \sqrt[3]{A(\vec{n}) + \sqrt{B(\vec{n})}} . \quad (21)$$

Here

$$\begin{aligned} A(\vec{n}) &= \frac{bJ_4 - J_6}{2} - \left(\frac{b}{3}\right)^3 \\ B(\vec{n}) &= J_4^3 - \frac{b^2}{12}J_4^2 - \frac{b}{2}J_4J_6 + \left(\frac{b}{3}\right)^3 J_6 + \frac{J_6^2}{4} \end{aligned} \quad (22)$$

and

$$\begin{aligned} J_4 &= \frac{b^2 - c^2}{3}I_4(\vec{n}) \\ J_6 &= (b + 2c)(b - c)^2I_6(\vec{n}) . \end{aligned} \quad (23)$$

These solutions display the same symmetries of the effective lagrangian because they depend on the directions \vec{n} only through the elementary symmetric functions $I_2(\vec{n}) = 1$, $I_4(\vec{n})$, $I_6(\vec{n})$. The corresponding eigenvectors can be computed in a straightforward manner; for example if

$$n_1 \neq 0 , \quad a - v_r^2 + bn_3^2 \neq 0 , \quad a - v_r^2 + (b - c)n_2^2 \neq 0 \quad (24)$$

one gets the eigenvectors:

$$\xi_r = \text{const.} \times \begin{pmatrix} n_1[a - v_r^2 + n_2^2(b - c)] \\ n_2[a - v_r^2 + n_1^2(b - c)] \\ -n_3c \frac{(a - v_r^2)(n_1^2 + n_2^2) + 2(b - c)n_1^2n_2^2}{a - v_r^2 + bn_3^2} \end{pmatrix} . \quad (25)$$

The dispersion law for the phonons depends on three arbitrary parameters a , b , c . This follows from the symmetry of the cubic crystal and could be also inferred from the general theory of the elastic waves in crystals. As matter of fact, the phonon amplitudes ϕ^j can be identified with the local deformations of the crystal with harmonic components

$$\varphi_j = \varphi_{0j} e^{i(\vec{p}\cdot\vec{x} - Et)} . \quad (26)$$

The equation of motion for the deformation is

$$\rho \frac{\partial^2 \varphi_j}{\partial t^2} = \partial_k \sigma_{jk} \quad (27)$$

where σ_{jk} is the stress tensor and ρ is the crystal density. The tensor σ_{jk} is related to the strain tensor $u_{\ell m}$:

$$u_{\ell m} = \frac{1}{2} (\partial_\ell \varphi_m + \partial_m \varphi_\ell + \partial_\ell \varphi_i \partial_m \varphi_i) , \quad (28)$$

by the relation

$$\sigma_{ik} = \lambda_{ik\ell m} u_{\ell m} . \quad (29)$$

$\lambda_{ik\ell m}$ is the elastic modulus tensor and the equation for elastic waves can be written as

$$\rho \frac{\partial^2 \varphi_j}{\partial t^2} = \lambda_{jk\ell m} \partial_k \partial_\ell \varphi_m , \quad (30)$$

from which the dispersion law follows:

$$Det \left(\rho E^2 \delta_{im} - \lambda_{ik\ell m} u_{\ell m} p_k p_\ell \right) = 0 . \quad (31)$$

Owing to its symmetry properties, the elastic modulus tensor depends in general on 21 parameters; for the cube group O_h only three of them are independent [16], for example λ_{1111} , λ_{1122} , λ_{1212} . The identification with the parameters of the effective lagrangian is as follows:

$$a = \frac{\lambda_{1212}}{\rho}, \quad b = \frac{\lambda_{1111} - \lambda_{1212}}{\rho}, \quad c = \frac{\lambda_{1122} + \lambda_{1212}}{\rho}. \quad (32)$$

The group velocity

$$\vec{v} = \frac{\partial E}{\partial \vec{p}} \quad (33)$$

is a function of the direction of the wave propagation \vec{n} but does not depend of the energy. For each fixed \vec{n} there are three velocities that are in general all different. For comparison in an isotropic medium $|\vec{v}|$ does not depend on \vec{n} and two of the velocities (the transverse ones) coincide, while being different and smaller than the velocity of the longitudinal waves.

Since the matrix M in (18) is real and symmetric, its eigenvalues are real as well. One has also to impose that $v_r^2 > 0$; for this to happen all the

principal minors

$$\Delta_1 = a + bn_i^2, \quad (i = 1, 2, 3) \quad (34)$$

$$\Delta_2 = (a + bn_i^2)(a + bn_j^2) - c^2 n_i^2 n_j^2, \quad (i \neq j) \quad (35)$$

$$\Delta_3 = a^2(a + b) + a(b^2 - c^2)I_4(\vec{n}) + (b + 2c)(b - c)^2 I_6(\vec{n}) \quad (36)$$

must be positive:

$$\Delta_i > 0 \quad (i = 1, 2, 3). \quad (37)$$

The constraint $\Delta_1 > 0$ gives the conditions

$$a > 0, \quad a + b > 0. \quad (38)$$

The most stringent constraint $\Delta_2 > 0$ is obtained for $n_i^2 = n_j^2 = 1/2$, which gives

$$b + 2a > |c|. \quad (39)$$

Finally the most stringent constraint $\Delta_3 > 0$ is obtained for $I_4(\vec{n}) = 1/3$ and $I_6(\vec{n}) = 1/27$, i.e. when \vec{n} is parallel to one of the 3-fold axes; since Δ_3 is the product of the three eigenvalues, one gets

$$a + \frac{b + 2c}{3} > 0. \quad (40)$$

These constraints are summarized in Fig. 2.

Let us specialize these results for momenta along the symmetry axes of the crystal i.e. along the three 4-fold axes, the four 3-fold axes and the six 2-fold axes.

We easily get for \vec{n} along one of the three 4-fold axes, e.g. $\vec{n} = (1, 0, 0)$,

$$v_1^2 = a + b \quad (41)$$

$$v_2^2 = v_3^2 = a \quad (42)$$

and the corresponding eigenvectors are

$$\xi_1 = \phi^{(1)} \quad (43)$$

while ξ_2, ξ_3 are in the subspace spanned by $\phi^{(2)}$ and $\phi^{(3)}$. The eigenvalues for the other two 3-fold axes with $\vec{n} = (0, 1, 0)$ and $\vec{n} = (0, 0, 1)$ are identical to (41) and (42) due to the cubic symmetry, while the corresponding eigenvectors are obtained by cyclic permutations.




Fig. 2 - The figure shows the regions where the square of all the velocities of propagation along the symmetry axes of the crystal are positive, in the plane (b, a) at fixed c . For $c < 0$ only the area with horizontal lines is allowed. For $c > 0$ both regions are permitted.

For \vec{n} along one of the four 3-fold axes, e.g. $\vec{n} = \frac{1}{\sqrt{3}}(1, 1, 1)$ one has

$$\begin{aligned} v_1^2 &= a + \frac{b+2c}{3} \\ v_2^2 &= v_3^2 = a + \frac{b-c}{3} \end{aligned} \quad (44)$$

with eigenvectors

$$\xi_1 = \frac{1}{\sqrt{3}}(\phi^{(1)} + \phi^{(2)} + \phi^{(3)}) \quad (45)$$

while ξ_2, ξ_3 are in the orthogonal subspace, i.e. the subspace spanned by

$$-\phi^{(1)} + x\phi^{(2)} + (1-x)\phi^{(3)}, \quad (46)$$

with x a real parameter.

The other three 3-fold axes are defined by the directions $\vec{n} = \frac{1}{\sqrt{3}}(1, -1, 1)$, $\vec{n} = \frac{1}{\sqrt{3}}(-1, 1, 1)$ and $\vec{n} = \frac{1}{\sqrt{3}}(-1, -1, 1)$; they can be obtained by the unit vector $\vec{n} = \frac{1}{\sqrt{3}}(1, 1, 1)$ by applying transformations in O_h ; due to the cubic symmetry the eigenvalues are again given by (44) and the corresponding

eigenvectors are obtained by applying the same transformation to the eigenvectors (45), (46); for example for $\vec{n} = \frac{1}{\sqrt{3}}(1, -1, 1)$ one has

$$\xi_1 = \frac{1}{\sqrt{3}} (\phi^{(1)} - \phi^{(2)} + \phi^{(3)}) \quad (47)$$

while ξ_2, ξ_3 are in the orthogonal subspace spanned by

$$\phi^{(1)} + x\phi^{(2)} + (1-x)\phi^{(3)} . \quad (48)$$

Finally we consider the six 2-fold axes defined by the directions $\vec{n} = \frac{1}{\sqrt{2}}(1, \pm 1, 0)$, $\vec{n} = \frac{1}{\sqrt{2}}(1, 0, \pm 1)$ and $\vec{n} = \frac{1}{\sqrt{2}}(0, 0, \pm 1)$. By way of example we take

$$\vec{n} = \frac{1}{\sqrt{2}}(1, \pm 1, 0) . \quad (49)$$

The eigenvalues are

$$v_1^2 = a + \frac{b+c}{2} \quad (50)$$

$$v_2^2 = a + \frac{b-c}{2} \quad (51)$$

$$v_3^2 = a . \quad (52)$$

The eigenvector corresponding to the eigenvalue (50) can be obtained by (25):

$$\xi_1 = \phi^{(1)} \pm \phi^{(2)} . \quad (53)$$

For the other two eigenvectors the conditions (24) are not satisfied and one has to solve the eigenvalue equations by assuming \vec{n} given by (49) from the very beginning; one obtains

$$\begin{aligned} \xi_2 &= \phi^{(1)} \mp \phi^{(2)} \\ \xi_3 &= \phi^{(3)} . \end{aligned} \quad (54)$$

The eigenvalues for \vec{n} along the other 2-fold axes are identical, due to the cubic symmetry; exactly as in the preceding cases the corresponding eigenvectors are obtained by (53), (54) by applying the same transformations that change (49) to the unit vectors of the other axes. All these results are summarized in Table 1 where we have introduced the unit vectors along the

coordinate axes, $\vec{e}^{(i)}$. We observe again that in the case of propagation along axes of type C_2 there are three different propagation velocities, where for the other two cases the independent velocities are only two. The case of propagation along axes of type C_3 is of particular interest since these directions are the same as the directions of the vectors \vec{q} specifying the lattice. One of the two eigenvalues is doubly degenerate, and the non-degenerate eigenvectors is proportional to

$$\sum_{i=1,2,3} p_i \phi^{(i)}. \quad (55)$$

The simplest case is for a propagation along the coordinate axes (type C_4), since in this case the matrix M is diagonal. Also in this case we have a doubly degenerate eigenvalue with eigenvectors the phonons corresponding to the plane perpendicular to the direction of propagation, whereas the non-degenerate eigenvector corresponds to the direction of propagation. It is interesting to notice that in all cases one of the eigenvectors is proportional to $\sum_i p_i \phi^{(i)}$.

4 Conclusions

Crystalline superconductivity arises from Cooper pairing between quarks of non vanishing total momentum, each quark momentum lying near the respective differing Fermi surfaces. Such a possibility (LOFF pairing) is known from study of certain condensed matter systems and recent theoretical studies have made it as highly probable within the QCD phase diagram. LOFF pairing leads to spontaneous breaking of translational and rotational invariance. Depending on the particular crystal structure a number of phonons are the lowest energy manifestation of the breaking.

A comparative study of the different possible crystal structures for the crystalline color superconducting phase of QCD has recently been published [1], with the conclusion that a face centered cubic crystal is the favorite structure. Assuming this condensate pattern we have derived the expected form of the effective low energy Lagrangian and the dispersion law for the the set of the three phonons. The parameters a , b and c appearing in the effective lagrangian can be evaluated following the methods illustrated in ref. [17]. The results will be given elsewhere.

References

- [1] J.A. Bowers and K. Rajagopal, hep-ph/0204079.
- [2] B. Barrois, Nucl. Phys. **B129** (1977) 390. S. Frautschi, Proceedings of workshop on hadronic matter at extreme density, Erice 1978. D. Bailin and A. Love, Phys. Rept. **107** (1984) 325.
- [3] M. Alford, K. Rajagopal and F. Wilczek, Phys. Lett. **B422**, 247 (1998), R. Rapp, T. Schäfer, E. V. Shuryak and M. Velkovsky, Phys. Rev. Lett. **81**, 53 (1998).
- [4] For a review see K.Rajagopal and F.Wilczek,in Handbook of QCD, M.Shifman ed. (World Scientific 2001), hep-ph/0011333.
- [5] M. Alford, K. Rajagopal and F. Wilczek, Nucl. Phys. **B537**, 443 (1999).
- [6] M. Alford, J. Bowers and K. Rajagopal, Phys. Rev. **D63**, 074016 (2001).
- [7] J. A. Bowers, J. Kundu, K. Rajagopal and E. Shuster, Phys. Rev. **D64**, 014024 (2001).
- [8] R. Casalbuoni, R. Gatto, M. Mannarelli and G. Nardulli, Phys. Lett. B **511**, 218 (2001) hep-ph/0101326.
- [9] A. K. Leibovich, K. Rajagopal and E. Shuster, Phys. Rev. **D64**, 094005 (2001). J. Kundu and K. Rajagopal, hep-ph/0112206.
- [10] For reviews see for instance: K. Rajagopal, in the proceedings of QCD@Work: International Workshop on QCD: Theory and Experiment, Martina Franca, Bari, Italy (2001), editors P. Colangelo and G. Nardulli, AIP Conference Proceedings, V. 602 (2001), hep-ph/0109135. R. Casalbuoni, ibidem, hep-th/0108195. G. Nardulli, Invited talk at Compact Stars in the QCD Phase Diagram, Copenhagen, Aug. 2001, hep-ph/0111178. M. Mannarelli, talk given at the Giselda Meeting, Frascati (Roma) (2001), hep-ph/0205098.
- [11] A. I. Larkin and Yu. N. Ovchinnikov, Sov. Phys. JETP **20**, 762 (1965), P. Fulde and R. A. Ferrell, Phys. Rev. **135**, A550 (1964).

- [12] F. Schreck *et al.* Phys. Rev. Lett. **87**, 080403 (2001), and references therein.
- [13] M. Holland, S. J. J. M. F. Kokkelmans, M. L. Chiafalo, and R. Walser, Phys. Rev. Lett. **87**, 120406 (2001), and references therein.
- [14] R. Combescot, Europhys. Lett. **55**, 150 (2001).
- [15] See, for instance, M. Hamermesh, *Group Theory and its application to Physical Problems*, Addison Wesley, 1964.
- [16] L.D. Landau and E. M. Lifshitz, *Course of Theoretical Physics*, Vol. 7, *Elasticity Theory*, Pergamon, London, 1959.
- [17] R.Casalbuoni, R. Gatto, M. Mannarelli and G. Nardulli, hep-ph/0201059.

Momentum	Eigenvalues	Eigenvectors
$\boxed{C_4}$ $\vec{p} = p \vec{e}^{(i)}$	$(a + b) p^2$ $a p^2$	$\phi^{(i)}$ $\phi^{(j)}, \phi^{(k)} \ (i \neq j \neq k)$
$\boxed{C_3}$ $\vec{p} = p (\vec{e}^{(1)} + \vec{e}^{(2)} + \vec{e}^{(3)})$	$(3a + b - c) p^2/3$ $(3a + b + 2c) p^2/3$	$-\phi^{(1)} + x\phi^{(2)} + (1 - x)\phi^{(3)}$ $\phi^{(1)} + \phi^{(2)} + \phi^{(3)}$
$\vec{p} = p (-\vec{e}^{(1)} + \vec{e}^{(2)} + \vec{e}^{(3)})$	$(3a + b - c) p^2/3$ $(3a + b + 2c) p^2/3$	$\phi^{(1)} + x\phi^{(2)} + (1 - x)\phi^{(3)}$ $-\phi^{(1)} + \phi^{(2)} + \phi^{(3)}$
$\vec{p} = p (\vec{e}^{(1)} - \vec{e}^{(2)} + \vec{e}^{(3)})$	$(3a + b - c) p^2/3$ $(3a + b + 2c) p^2/3$	$\phi^{(1)} + x\phi^{(2)} - (1 - x)\phi^{(3)}$ $\phi^{(1)} - \phi^{(2)} + \phi^{(3)}$
$\vec{p} = p (\vec{e}^{(1)} + \vec{e}^{(2)} - \vec{e}^{(3)})$	$(3a + b - c) p^2/3$ $(3a + b + 2c) p^2/3$	$\phi^{(1)} + x\phi^{(2)} + (1 + x)\phi^{(3)}$ $\phi^{(1)} + \phi^{(2)} - \phi^{(3)}$
$\boxed{C_2}$ $\vec{p} = p (\vec{e}^{(i)} + \vec{e}^{(j)})$ $i < j$	$a p^2$ $(2a + b - c) p^2/2$ $(2a + b + c) p^2/2$	$\phi^{(k)}$ $-\phi^{(i)} + \phi^{(j)}$ $\phi^{(i)} + \phi^{(j)}$
$\vec{p} = p (\vec{e}^{(i)} - \vec{e}^{(j)})$ $i < j$	$a p^2$ $(2a + b - c) p^2/2$ $(2a + b + c) p^2/2$	$\phi^{(k)}$ $\phi^{(i)} + \phi^{(j)}$ $-\phi^{(i)} + \phi^{(j)}$

Table 1 - Eigenvalues and eigenvectors for \vec{p} along the symmetry axes.