No-go for detecting CP violation via neutrinoless double beta decay

V. Barger¹, S.L. Glashow², P. Langacker³ and D. Marfatia²

¹Department of Physics, University of Wisconsin, Madison, WI 53706, USA

²Department of Physics, Boston University, Boston, MA 02215, USA

³Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104, USA

Abstract

We present a necessary condition on the solar oscillation amplitude for CP violation to be detectable through neutrinoless double beta $(0\nu\beta\beta)$ decay. It depends only on the fractional uncertainty in the $\nu_e-\nu_e$ element of the neutrino mass matrix. We demonstrate that even under very optimistic assumptions about the sensitivity of future experiments to the absolute neutrino mass scale, and on the precision with which nuclear matrix elements that contribute to $0\nu\beta\beta$ decay are calculable, it will be impossible to detect neutrino CP violation arising from Majorana phases.

If neutrinos are Majorana, the potentiality of detecting CP violation using neutrinoless double beta decay exists[[1](#page-6-0)]. We consider this in a scenario wherein there are exactly three left-handed neutrino states with Majorana masses. We derive a necessary condition that involves the solar oscillation amplitude and the uncertainty in the $\nu_e-\nu_e$ element of the neutrino mass matrix that must be satisfied for CP violation to be detectable. Assuming that this condition is satisfied and allowing for experimental and theoretical uncertainties that are unrealistically small in some cases, we show that it will not be possible to detect neutrino CP violation through this process.

The charged-current eigenstates are related to the mass eigenstates by a unitary transformation

$$
\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = UV \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix} = \begin{pmatrix} c_2c_3 & c_2s_3 & s_2e^{-i\delta} \\ -c_1s_3 - s_2s_1c_3e^{i\delta} & c_1c_3 - s_2s_1s_3e^{i\delta} & c_2s_1 \\ s_1s_3 - s_2c_1c_3e^{i\delta} & -s_1c_3 - s_2c_1s_3e^{i\delta} & c_2c_1 \end{pmatrix} V \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix},
$$
\n(1)

where s_i and c_i are the sines and cosines of θ_i , and V is the diagonal matrix $(1, e^{i\frac{\phi_2}{2}}, e^{i(\frac{\phi_3}{2}+\delta)})$. In Eq. (1), ϕ_2 and ϕ_3 are the Majorana phases that are not measurable in neutrino oscillations and which are either 0 or π if CP is conserved.

The solar neutrino data allow two solutions at the 3σ C.L. [\[2\]](#page-6-0):

$$
0.56 \le \sin^2 2\theta_3 \le 0.95 \,, \quad 2.0 \times 10^{-5} \le \Delta_s \le 2.3 \times 10^{-4} \text{ eV}^2 \,, \quad \text{(LMA)} \tag{2}
$$

$$
0.80 \le \sin^2 2\theta_3 \le 0.94 \,, \quad 9.0 \times 10^{-8} \le \Delta_s \le 2.0 \times 10^{-7} \text{ eV}^2 \,, \quad \text{(LOW)} \tag{3}
$$

of which the LMA is the only solution at the 99% C.L. No solution with $\theta_3 \ge \pi/4$ (cos $2\theta_3 \le 0$) isallowed at the 5 σ C.L. [[2](#page-6-0)]. Atmospheric neutrino data imply $\sin^2 2\theta_1 \geq 0.85$ and $1.1 \times$ 10^{-3} eV² $\leq \Delta_a \leq 5 \times 10^{-3}$ eV² at the 99% C.L. [[3\]](#page-6-0). The CHOOZ reactor experiment imposes theconstraint $\sin^2 2\theta_2 \leq 0.1$ at the 95% C.L. [[4\]](#page-6-0). Δ_s and Δ_a are the mass-squared differences relevant to solar and atmospheric neutrino oscillations, respectively.

vWe choose the mass ordering $m_1 < m_2 < m_3$ with m_i non-negative. There are two possible neutrino mass spectra:

$$
\Delta_s = m_2^2 - m_1^2, \qquad \Delta_a = m_3^2 - m_2^2, \qquad \text{(normal hierarchy)}, \tag{4}
$$

$$
\Delta_s = m_3^2 - m_2^2, \qquad \Delta_a = m_2^2 - m_1^2, \qquad \text{(inverted hierarchy)}, \tag{5}
$$

where in either case $\Delta_a \gg \Delta_s$ in accord with experimental data. For the normal hierarchy

(Case I), mixing is given by Eq.([1\)](#page-1-0). In this case solar neutrinos oscillate primarily between the two lighter mass eigenstates. For the *inverted hierarchy* (Case II), solar neutrinos oscillate primarily between the two nearly degenerate heavier states. The oscillation parameters retain the same import as in Case I if the columns of UV are permuted so that the third column takes the place of the first column.

It is well known that there are four combinations of the intrinsic CP parities of the mass eigenstates for which CP invariance holds [\[5](#page-6-0)]. If $\theta_2 = 0$, the number of such combinations is reduced to two. To minimize the confusion between CP violation and CP conservation, we neglect θ_2 which is already constrained to be small. This is justified as long as the mass spectrum is not hierarchical $(m_1 \ll m_2 \ll m_3)$. Since experiments will not be sensitive to the M_{ee} (< 0.01 eV) expected for a hierarchical spectrum our assumption that θ_2 is small is not only sound but also optimistic from the viewpoint of detecting CP violation via $0\nu\beta\beta$ decay. Of course, if θ_2 is small, long baseline neutrino experiments will find it harder to detect CP violation resulting from δ . In this sense $0\nu\beta\beta$ decay and long baseline experiments are complementary for detecting CP violation; if θ_2 is close to the CHOOZ bound, long baseline experiments have a greater chance of detecting CP violation and if θ_2 is tiny, $0\nu\beta\beta$ decay experiments have the better chance.

Under the assumption that θ_2 is negligible, the $\nu_e-\nu_e$ element of the neutrino mass ma-trix $[6]$ $[6]$, is

$$
M_{ee} = |c^2 m_1 + s^2 m_2 e^{i\phi}|, \quad \text{(Case I)}, \tag{6}
$$

$$
= |c^2 m_2 + s^2 m_3 e^{i\phi}|, \quad \text{(Case II)}, \tag{7}
$$

where we have dropped the subscripts on the sines and cosines of θ_3 and on ϕ_2 , since only one mixing angle and one Majorana phase is involved. The masses m_i may be determined from the lightest mass m_1 and the mass–squared differences. Since the solar mass–squared difference is very small it can be ignored; then setting $m_1 = m$ and $\Delta_a = \Delta$,

$$
m_e = m_2 = m
$$
, $m_3 = \sqrt{m^2 + \Delta}$, (Case I), (8)

$$
m_e = m_2 = m_3 = \sqrt{m^2 + \Delta}
$$
, (Case II). (9)

Here, the mass of the electron neutrino m_e is what tritium beta decay experiments seek to measure. It is related to the sum of neutrino masses $(\Sigma = \Sigma m_i)$ obtainable from data on weak lensing of galaxies and the CMB via

$$
\Sigma = 2m_e + \sqrt{m_e^2 \pm \Delta},\tag{10}
$$

where the plus sign applies to the normal hierarchy and the minus sign to the inverted hierarchy.Using Eqs. $(6-9)$ $(6-9)$, the Majorana phase in either case is given by

$$
\cos \phi = 1 - \frac{2}{\sin^2 2\theta} \left(1 - \frac{M_{ee}^2}{m_e^2} \right).
$$
 (11)

The second term on the right hand side quantifies the deviation of ϕ from 0. If $M_{ee}/m_e = 1$, $\phi = 0$ and if $M_{ee}/m_e = \cos 2\theta$, $\phi = \pi$. Thus,

$$
\cos 2\theta \le \frac{M_{ee}}{m_e} \le 1\,,\tag{12}
$$

with the boundaries of the interval corresponding to CP conservation. A measurement of M_{ee}/m_e that excludes the boundaries constitutes a detection of CP violation. A larger solar amplitude (a wider interval) is therefore more conducive to such a measurement. Let us evaluate the minimum $\sin^2 2\theta$ for which CP violation can be detected assuming a measurement $M_{ee}(1\pm x)$, where x is obtained by summing the theoretical uncertainty in the $0\nu\beta\beta$ nuclear matrix elements and the experimental uncertainty in quadrature. Then, for CP violation to be detectable the necessary condition is

$$
\frac{\cos 2\theta}{1-x} < \frac{M_{ee}}{m_e} < \frac{1}{1+x} \,,\tag{13}
$$

or

$$
\sin^2 2\theta > 1 - \left(\frac{1-x}{1+x}\right)^2.
$$
 (14)

The current solar data require $\sin^2 2\theta$ to be smaller than 0.95 at the 3σ C.L. Thus, x must be smaller than 0.63. It is a difficult task to reduce the factor of 3 uncertainty on the nuclear matrix elements (corresponding to $M_{ee}(1^{+2.0}_{-0.7})$) to such a degree especially since a reliable method for estimating the uncertainty does not exist¹ [\[7](#page-6-0)]. Conversely, for a realistically achievable improvement in x , it is unlikely that the solar amplitude is sufficiently large so as to satisfy Eq. (14) .

In what follows, we consider the remote possibility that Eq. (14) is in fact satisfied, and show that it is not sufficient to detect CP violation.

¹The often quoted factor of 3 uncertainty represents the range of calculated values of the matrix elements available in the literature.

We work under the following assumptions about the experimental and theoretical developments that might occur by 2020:

- 1. Experiments like GENIUS [\[8](#page-6-0)] and EXO [\[9](#page-6-0)] are sensitive to M_{ee} above 0.01 eV with a 25% experimental uncertainty[[7\]](#page-6-0) and are therefore not sensitive to the hierarchical neutrino mass spectrum. Also, since $M_{ee} \leq m_e$, we can draw meaningful conclusions only for $m_e \gtrsim 0.01$ eV or equivalently $\Sigma \gtrsim 0.08$ eV.
- 2. A breakthrough in the evaluation of the nuclear matrix elements has allowed an estimate of their uncertainty. The factor of 3 uncertainty that has plagued the matrix elements is reduced so that the combined theoretical and experimental uncertainty on M_{ee} is $x = 0.5$. Then Eq. ([14\)](#page-3-0) is satisfied for $\sin^2 2\theta > 0.89$.
- 3. Tritium beta decay experiments like KATRIN are sensitive to m_e above 0.35 eV with an uncertainty of 0.08 eV^2 on m_e^2 [[10](#page-6-0)].
- 4. Weak lensing of galaxies by large scale structure in conjunction with CMB data can measure Σ to an uncertainty of 0.04 eV [\[11\]](#page-6-0).
- 5. The KamLAND (Borexino) experiment has determined the solar oscillation amplitude to be 0.95 ± 0.04 where the precision in the LMA (LOW) region was estimated in Ref.[[12\]](#page-6-0) ([[13\]](#page-6-0)).
- 6. The JHF-Kamioka neutrino project has measured Δ to be 3×10^{-3} eV² to within 3% and has constrained $\sin^2 \theta_2$ to be smaller than 2×10^{-3} [\[14\]](#page-6-0).
- 7. The neutrino mass hierarchy is determined either from a superbeam experiment or from supernova neutrinos.

While some of these assumptions are overly optimistic, they serve well to show once and for all that it is not possible to detect CP violation from $0\nu\beta\beta$ decay in the foreseeable future.

Our assumptions clearly suggest that cosmology will not only probe smaller neutrino masses, but also with greater precision than tritium beta decay experiments. However, to have the possibility of an independent confirmation from a table-top experiment (in the regime of common sensitivity) is invaluable.

In our quantitative analysis, we assume the precision on Σ expected from cosmology. We fix $\Delta = 3 \times 10^{-3} \text{ eV}^2$ and assign no uncertainty to its value. We suppose that precise measurements of Σ and M_{ee} will be made such that the central value of M_{ee}/m_e lies in the interval of Eq.([12](#page-3-0)). One expects the extent to which CP violation can be detected to be dependent upon how close the central value of M_{ee}/m_e is to the middle of the interval.

For hypothetical measurements of Σ and M_{ee} , the regions allowed by scans within the 1σ uncertainties assumed for each of the measurements are shown in Fig. [1](#page-7-0) for the normal hierarchy. We have chosen the central values of M_{ee}/m_e to be 0.4, 0.6 and 0.8 so that they are not too close to either cos $2\theta (= 0.22)$ or 1 for which $\cos \phi = -1$ and $\cos \phi = 1$, respectively, are unavoidable. For $\Sigma \gg \sqrt{\Delta}$, the regions for both hierarchies are almost identical. For the case in which $\Sigma = 0.24$ eV, the regions for the inverted hierarchy do not extend to as high values of M_{ee}/m_e as for the normal hierarchy, but qualitatively they are the same in that $\cos \phi = -1$ is allowed.

We emphasize that while CP violation is not detectable via $0\nu\beta\beta$ decay, if the solar amplitude is found to be larger than current solar data suggest and if the precision on the various measurements and the refinement of the calculation of the nuclear matrix elements assumed by us is achieved, it might be possible to determine if ϕ is closer to 0 or to π (see Fig. [1\)](#page-7-0).

Acknowledgements. We thank S. Dodelson for drawing our attention to Ref. [\[11\]](#page-6-0). This work was supported in part by the NSF under grant No. NSF-PHY-0099529, in part by the U.S. Department of Energy under grant Nos. DE-FG02-91ER40676, DE-FG02-95ER40896 and DOE-EY-76-02-3071, and in part by the Wisconsin Alumni Research Foundation.

References

- [1] For related work see e.g., Y. Farzan, O. L. Peres and A. Y. Smirnov, Nucl. Phys. B 612, 59 (2001)[[arXiv:hep-ph/0105105\]](http://arxiv.org/abs/hep-ph/0105105); W. Rodejohann, [arXiv:hep-ph/0203214;](http://arxiv.org/abs/hep-ph/0203214) S. Pascoli and S. T. Petcov, [arXiv:hep-ph/0205022](http://arxiv.org/abs/hep-ph/0205022) and references therein.
- [2] Q. R. Ahmad et al. [SNO Collaboration], [arXiv:nucl-ex/0204009](http://arxiv.org/abs/nucl-ex/0204009); V. Barger, D. Marfatia, K. Whisnant and B. P. Wood, [arXiv:hep-ph/0204253.](http://arxiv.org/abs/hep-ph/0204253)
- [3] T. Toshito [SuperKamiokande Collaboration], [arXiv:hep-ex/0105023](http://arxiv.org/abs/hep-ex/0105023).
- [4] M. Apollonio *et al.* [CHOOZ Collaboration], Phys. Lett. B $466, 415$ (1999) [\[arXiv:hep](http://arxiv.org/abs/hep-ex/9907037)[ex/9907037](http://arxiv.org/abs/hep-ex/9907037)].
- [5] L. Wolfenstein, Phys. Lett. B 107, 77 (1981).
- [6] M. Doi, T. Kotani, H. Nishiura, K. Okuda and E. Takasugi, Phys. Lett. B 102, 323 (1981).
- [7] S. R. Elliott and P. Vogel, [arXiv:hep-ph/0202264](http://arxiv.org/abs/hep-ph/0202264).
- [8] H. V. Klapdor-Kleingrothaus et al. [GENIUS Collaboration], [arXiv:hep-ph/9910205.](http://arxiv.org/abs/hep-ph/9910205)
- [9] M. Danilov *et al.*, Phys. Lett. B 480 , 12 (2000) [\[arXiv:hep-ex/0002003](http://arxiv.org/abs/hep-ex/0002003)];
- [10] A. Osipowicz et al. [KATRIN Collaboration], [arXiv:hep-ex/0109033.](http://arxiv.org/abs/hep-ex/0109033)
- [11] W. Hu and M. Tegmark, Astrophys. J. Lett. 514 65 (1999).
- [12] V. Barger, D. Marfatia and B. P. Wood, Phys. Lett. B 498, 53 (2001) [\[arXiv:hep](http://arxiv.org/abs/hep-ph/0011251)[ph/0011251](http://arxiv.org/abs/hep-ph/0011251)].
- [13] A. de Gouvea, A. Friedland and H. Murayama, JHEP 0103, 009 (2001) [\[arXiv:hep](http://arxiv.org/abs/hep-ph/9910286)[ph/9910286](http://arxiv.org/abs/hep-ph/9910286)].
- [14] Y. Itow *et al.*, $arXiv:hep-ex/0106019$.

Figure 1: " 1σ " bands in the M_{ee}/m_e -cos ϕ plane for three possible measurements assuming the normal hierarchy: (a) $\Sigma = 0.24 \pm 0.04$ eV, $M_{ee} = 0.03(1 \pm 0.5)$ eV with the central value of $M_{ee}/m_e = 0.4$, (b) $\Sigma = 0.51 \pm 0.04$ eV, $M_{ee} = 0.10(1 \pm 0.5)$ eV with the central value of $M_{ee}/m_e = 0.6$, (c) $\Sigma = 1.12 \pm 0.04$ eV, $M_{ee} = 0.30(1 \pm 0.5)$ eV with the central value of $M_{ee}/m_e = 0.8$. Results are shown for the best possible solar amplitude (allowed by present solar data at 3σ) for the detection of CP violation, $\sin^2 2\theta = 0.95 \pm 0.04$. Δ is fixed at $3 \times 10^{-3} \text{ eV}^2$.