

Quantum fields and quantum groups

Christian Brouder^{1*} and Robert Oeckl^{2†}

¹Laboratoire de Minéralogie Cristallographie, CNRS UMR 7590, Universités Paris 6 et 7, IPGP, case 115, 4 place Jussieu, 75252 Paris cedex 05, France

²Centre de Physique Théorique, CNRS Luminy, F-13288 Marseille Cedex 9, France

(Dated: November 5, 2018)

Quantum fields are shown to be an example of infinite-dimensional quantum groups. A dictionary is established between quantum field and quantum group concepts: the expectation value over the vacuum is the counit, operator and time-ordered products are twisted products, Wick's theorem is the definition of a twisted product. Renormalization becomes the replacement of a coquasitriangular structure by a 2-coboundary. Quantum group methods are used to discuss quantum field theory with a correlated initial state.

Introduction – Quantum groups are a powerful generalization of groups and Lie algebras [1]. Since their creation in the mid-eighties, quantum groups have attracted considerable attention and more than four thousand papers were published on the subject in the mathematical literature. In physics, groups and Lie algebras are everywhere and it can be expected that quantum groups are useful too. Indeed, quantum groups play a role in solid-state physics (magnetic systems [2, 3, 4], phonons [5]) which is directly linked to their origin in integrable systems [6]. Furthermore, they are the basic symmetry of the quantum Hall effect [7, 8]. In this paper we first show that quantum groups underlie also a most successful and pervading tool of physics: quantum field theory. Then we use quantum group methods to solve a problem of quantum field theory with a correlated initial state, to interpret renormalization and to explain the origin of quasifree states.

Definitions – A quantum group is a Hopf algebra [9] equipped with a quasitriangular structure [1, 10]. A Hopf algebra H is an algebra: a vector space H with an associative product denoted by \cdot and a unit 1. It is also a coalgebra: there is a linear map $\Delta : H \rightarrow H \otimes H$, called a coproduct, which “splits” an element a of H into the sum of all its left and right parts: $\Delta a = \sum a_{(1)} \otimes a_{(2)}$. Coassociativity means that applying the coproduct again on the left or the right hand component yields the same result. In other words, Δ is coassociative if and only if $\sum (\Delta a_{(1)}) \otimes a_{(2)} = \sum a_{(1)} \otimes (\Delta a_{(2)})$. In a coalgebra, the coproduct is assumed coassociative. A coalgebra is also equipped with a counit ε , which is a map from H to \mathbb{C} such that $\sum \varepsilon(a_{(1)})a_{(2)} = \sum a_{(1)}\varepsilon(a_{(2)}) = a$. In a Hopf algebra, the coalgebra structure is compatible with the algebra structure: $\Delta(a \cdot b) = \sum (a_{(1)} \cdot b_{(1)}) \otimes (a_{(2)} \cdot b_{(2)})$ and $\varepsilon(a \cdot b) = \varepsilon(a)\varepsilon(b)$. The final ingredient of a Hopf algebra is the antipode, which is a map $S : H \rightarrow H$ such that $\sum S(a_{(1)}) \cdot a_{(2)} = \sum a_{(1)} \cdot S(a_{(2)}) = \varepsilon(a)1$. A Hopf algebra is commutative if the product \cdot is commutative, and cocommutative if $\sum a_{(1)} \otimes a_{(2)} = \sum a_{(2)} \otimes a_{(1)}$. A Hopf algebra is in some sense a self-dual structure, because the dual of a product is a coproduct and the dual of a unit is a counit, and vice-versa.

To complete the definitions, a quantum group is a Hopf algebra together with an element R of $H \otimes H$ called a quasitriangular structure or a universal R -matrix [10]. Because of the self-dual nature of Hopf algebras, quantum group can also be described by the dual of a universal R -matrix, called a coquasitriangular structure [1]. This is the concept we need in quantum field theory. A coquasitriangular structure is an invertible bilinear map $\mathcal{R} : H \times H \rightarrow \mathbb{C}$ such that

$$\mathcal{R}(a \cdot b, c) = \sum \mathcal{R}(a, c_{(1)})\mathcal{R}(b, c_{(2)}), \quad (1)$$

$$\mathcal{R}(a, b \cdot c) = \sum \mathcal{R}(a_{(1)}, c)\mathcal{R}(a_{(2)}, b). \quad (2)$$

For a commutative and cocommutative Hopf algebra, no other condition is required for \mathcal{R} . We restrict ourselves to this case in the following.

We use \mathcal{R} to define a twisted product, denoted by \circ ,

$$a \circ b = \sum \mathcal{R}(a_{(1)}, b_{(1)}) a_{(2)} \cdot b_{(2)}. \quad (3)$$

The twisted product is a special case of Sweedler's crossed product [11]. It can be checked that, when Eqs.(1) and (2) are satisfied and the coproduct is cocommutative, then the twisted product is associative [11].

The quantum group stage is now set, and we show that quantum field theory plays naturally on it. We take the example of a real scalar particle. The field operator is

$$\phi(x) = \int \frac{d\mathbf{k}}{(2\pi)^3 \sqrt{2\omega_k}} \left(e^{-ip \cdot x} a(\mathbf{k}) + e^{ip \cdot x} a^\dagger(\mathbf{k}) \right),$$

where $\omega_k = \sqrt{m^2 + |\mathbf{k}|^2}$, $p = (\omega_k, \mathbf{k})$, $a^\dagger(\mathbf{k})$ and $a(\mathbf{k})$ are the creation and annihilation operators acting on F_s , the symmetric Fock space of scalar particles [12]. To avoid ill-defined integrals, we consider the vector space of smoothed fields $V = \{\phi(f), f \in \mathcal{D}(\mathbb{R}^4)\}$, where $\phi(f) = \int dx f(x)\phi(x)$ and $\mathcal{D}(\mathbb{R}^4)$ is the space of smooth functions on \mathbb{R}^4 with compact support [12].

The quantum group of quantum fields – Now we establish a connection between the quantum group and quantum field concepts. From the vector space V , we build the symmetric Hopf algebra $S(V) = \bigoplus_{n=0}^{\infty} S^n(V)$ (see ref. [13]), where $S^0(V) = \mathbb{C}1$, $S^1(V) = V$ and

$S^n(V)$ is generated by the symmetric product of n elements of V . An element of $S^n(V)$ is said to have degree n . The symmetric product in $S(V)$ is denoted by \vee and is induced by the normal product of operators $a(\mathbf{k})$ and $a^\dagger(\mathbf{k})$: in the usual quantum field notation, $\phi(f)\vee\phi(g) = :\phi(f)\phi(g):$. Since \vee is associative and commutative, $S(V)$ is a commutative algebra with unit 1, where 1 is the unit operator acting on F_s . The coproduct is defined on V by $\Delta(\phi(f)) = \phi(f) \otimes 1 + 1 \otimes \phi(f)$, and extended to $S(V)$ by compatibility with the product. For example, $\Delta(\phi(f)\vee\phi(g)) = (\phi(f)\vee\phi(g)) \otimes 1 + \phi(f) \otimes \phi(g) + \phi(g) \otimes \phi(f) + 1 \otimes (\phi(f)\vee\phi(g))$. Note that this coproduct is cocommutative. The counit is $\varepsilon(a) = 0$ if $a \in S^n(V)$ with $n > 0$ and $\varepsilon(1) = 1$. In the quantum field case, we can prove the additional identity $\varepsilon(a) = \langle 0|a|0\rangle$, where a is any element of $S(V)$ and $|0\rangle$ is the vacuum [14]. Finally, the antipode is $S(a) = (-1)^n a$ for $a \in S^n(V)$. This makes $S(V)$ a commutative and cocommutative Hopf algebra. It is even a Hopf \star -algebra, with the involution $\phi(f)^\star = \phi(f^*)$, where f^* the complex conjugate of f .

To make $S(V)$ a quantum group, we must add a coquasitriangular structure. This structure is entirely determined by its value on V . More precisely, if $a = u_1 \vee \dots \vee u_m \in S^m(V)$ and $b = v_1 \vee \dots \vee v_n \in S^n(V)$, where u_i and v_j are elements of V , then Eqs.(1) and (2) imply that $\mathcal{R}(a, b) = 0$ if $m \neq n$ and $\mathcal{R}(a, b) = \text{per}(u_i, v_j)$ if $m = n$ [15]. The permanent $\text{per}(u_i, v_j)$ is a kind of determinant of the matrix $\mathcal{R}(u_i, v_j)$ without minus signs:

$$\text{per}(u_i, v_j) = \sum_{\sigma} \mathcal{R}(u_1, v_{\sigma(1)}) \cdots \mathcal{R}(u_n, v_{\sigma(n)}), \quad (4)$$

where the sum is over all permutations σ of $\{1, \dots, n\}$. Therefore, the most general Poincaré-invariant coquasitriangular structure on $S(V)$ is determined by its value on V , which can be written

$$\mathcal{R}(\phi(f), \phi(g)) = \int dx dy f(x) G(x-y) g(y), \quad (5)$$

where $G(x)$ is a Lorentz invariant distribution. As explained above, \mathcal{R} gives rise to a twisted product on $S(V)$ defined by (3). By choosing the proper $G(x)$, this twisted product can become the operator product or the time-ordered product. Explicitly, if

$$G(x) = G_+(x) = \int \frac{d\mathbf{k}}{(2\pi)^3 2\omega_k} e^{-ip \cdot x},$$

the twisted product defined by (3) is the usual product of field operators in quantum field theory. If $G(x)$ is the Feynman propagator $G_F(x)$, with

$$G_F(x) = i \int \frac{d^4 p}{(2\pi)^4} \frac{e^{-ip \cdot x}}{p^2 - m^2 + i\epsilon},$$

then the twisted product defined by (3) is the time-ordered product. Let us show this for the operator product. The twisted product of two fields is $\phi(f) \circ \phi(g) =$

$\phi(f)\vee\phi(g) + \mathcal{R}(\phi(f), \phi(g))$ and, by Wick's theorem, the operator product of two fields is $\phi(f)\phi(g) = \phi(f)\vee\phi(g) + \langle 0|\phi(f)\phi(g)|0\rangle$. Thus, the two products are identical if we define $\mathcal{R}(\phi(f), \phi(g)) = \langle 0|\phi(f)\phi(g)|0\rangle$. This identification gives us $G(x) = G_+(x)$. Once the products are identified for two fields, they can be identified for any number of fields by noticing that Eq.(3) is Wick's theorem under a quantum group disguise [14]: Eq.(3) can be described in words by "the twisted product of two normal products a and b is obtained by summing all possible pairings between a and b ", which is Wick's theorem for the product of a and b [16], the pairing being defined by $\langle 0|\phi(f)\phi(g)|0\rangle = \mathcal{R}(\phi(f), \phi(g))$. Since Wick's theorem is valid for operator products and time-ordered products [17], the twisted product is equal to the time-ordered product if we identify $\mathcal{R}(\phi(f), \phi(g))$ with $\langle 0|T\phi(f)\phi(g)|0\rangle$. This leads to $G(x) = G_F(x)$. Note that this type of relation between Wick's theorem and Hopf algebras was already pointed out by Fauser [18].

In this context, the normal product becomes the basic product of quantum field theory, while the operator product and the time-ordered product become deformed products. The advantage of this point of view was already stressed in the fifties by Gupta [19] and is the basis of the star-product approach to quantum field theory [20, 21].

Applications – We have shown that many concepts and tools of quantum field theory, such as normal product, expectation value over the vacuum, Wick's theorem, operator and time-ordered products, are natural elements of the quantum group point of view. Now we show that quantum group methods can solve elegantly quantum field theory problems. As a simple example, consider a system with a correlated initial state $|\Omega\rangle$. In solid-state and atomic physics, the initial state is usually described by a Slater determinant [22]. However, in the case of degenerate initial states [23, 24, 25], or when discussing non-equilibrium Green functions [26, 27, 28] or contracted Schrödinger equations (see [29, 30, 31] and references therein), a correlated initial state $|\Omega\rangle$ must be used. The main difficulty of this problem comes from the fact that the expectation value of normal products over $|\Omega\rangle$ is not zero, so that standard quantum field methods cannot be used. To solve this difficulty, Kutzelnigg and Mukherjee [32] proposed to define a normal product adapted to $|\Omega\rangle$: a transformation $a \rightarrow \tilde{a}$ such that $\langle \Omega|\tilde{a}|\Omega\rangle = \varepsilon(a)$. We can generalize this question by taking a linear map ω from $S(V)$ to \mathbb{C} such that $\omega(1) = 1$ and looking for a transformation $a \rightarrow \tilde{a}$ such that $\omega(\tilde{a}) = \varepsilon(a)$. The original problem is recovered when $\omega(a) = \langle \Omega|a|\Omega\rangle$, but ω can also represent a mixed initial state. This problem was solved explicitly for elements a of small degrees in [32] but the general solution can now be obtained by quantum group methods. We call 1-cochain a linear map ρ from $S(V)$ to \mathbb{C} such that $\rho(1) = 1$. In particular, ω is a 1-cochain. In a founda-

tional paper on Hopf algebra cohomology [11], Sweedler showed that two 1-cochains ω and ρ can be multiplied by defining $(\omega \star \rho)(a) = \sum \omega(a_{(1)})\rho(a_{(2)})$. The set of 1-cochains forms an Abelian group with unit element ε . If ω^{-1} is the unique 1-cochain defined by $\omega \star \omega^{-1} = \varepsilon$, then the normal products adapted to ω are obtained by defining $\tilde{a} = \sum \omega^{-1}(a_{(1)})a_{(2)}$. This solves the problem because $\omega(\tilde{a}) = (\omega^{-1} \star \omega)(a) = \varepsilon(a)$. If we compare this general solution with the first degrees given in [32], we observe that the quantum group method tames the combinatorial complexity of the problem and provides a way to manipulate the general term of the solution.

We saw that quantum groups fare well for the free scalar field and the extension to fermions is done by replacing Hopf algebras by Hopf superalgebras [33]. But what happens with interacting fields? We must extend our vector space V to include normal products of fields at a point $V_{\text{dp}} = \{\phi^{(n)}(f), f \in \mathcal{D}(\mathbb{R}^4)\}$, where $\phi^{(n)}(f) = \int dx f(x)\phi^{(n)}(x)$ and $\phi^{(n)}(x)$ is the divided power of n fields at x : $\phi^{(n)}(x) = (1/n!)\phi(x) \vee \dots \vee \phi(x)$. We choose divided powers instead of usual powers of fields because their coproduct is simpler:

$$\Delta\phi^{(n)}(x) = \sum_{k=0}^n \phi^{(k)}(x) \otimes \phi^{(n-k)}(x). \quad (6)$$

This coproduct is now extended to $S(V_{\text{dp}})$ by using $\Delta(a \vee b) = (\Delta a) \vee (\Delta b)$. The coquasitriangular structure of two divided powers is calculated from Eqs.(1) and (2):

$$\mathcal{R}(\phi^{(m)}(x), \phi^{(n)}(y)) = \frac{\delta_{m,n}}{n!} G(x-y)^n. \quad (7)$$

In general, $G(x)$ is a distribution and its power is not well defined. Using microlocal arguments, Hörmander proved that distributions can be multiplied if their wavefront sets satisfy a compatibility condition [12]. It turns out that $G_+(x)$ satisfies this condition, so that the operator product of powers of fields can be defined. In fact, Brunetti and Fredenhagen [34] argued that, up to smooth functions, $G_+(x)$ is the only distribution that solves the Klein-Gordon equation, that defines a state, and whose powers $G_+^n(x)$ exist as distributions. On the other hand, the Feynman propagator $G_F(x)$ does not satisfy the wavefront set condition and $G_F(x)^n$ is not defined. This is the origin of the renormalization problem. In fact, no coquasitriangular structure can reproduce the renormalized time-ordered product, and the quantum group approach seems to fail.

However, quantum group theory is quite resourceful. To see how it can be used to deal with renormalization, we use the Epstein-Glaser approach, which is a renormalization method without infinities. Epstein and Glaser [35] define a linear map T from $S(V_{\text{dp}})$ to $S(V_{\text{dp}})$, that we call the T -map (they call it T -products but we prefer T -map to avoid confusion). This map satisfies

the equations $T(1) = 1$, $T(u) = u$ for $u \in V_{\text{dp}}$ and $T(a) = \sum \varepsilon(T(a_{(1)}))a_{(2)}$ for any $a \in V_{\text{dp}}$. In these equations, which are also valid in curved space-times [34], $T(a)$ is renormalized (i.e. gives rise to finite quantities). As a side remark, the efficiency of the quantum group language can be appreciated by noting that our equation for $T(a)$ corresponds to a two-line formula in the original papers (Eq.(42) in [35], Eq.(26) in [34]). The importance of the T -map stems from the fact that it enables us to calculate the renormalized S-matrix. For an interacting Lagrangian $\lambda\phi^{(n)}(f)$, the renormalized S-matrix is [36]

$$S(f) = T\left(1 + \sum_{k=1}^{\infty} \frac{(-i\lambda)^k}{k!} \phi^{(n)}(f) \vee \dots \vee \phi^{(n)}(f)\right).$$

Let us investigate what quantum groups can tell us about the T -map. The function $t(a) = \varepsilon(T(a))$ is a linear map from $S(V_{\text{dp}})$ to \mathbb{C} such that $t(1) = 1$. Thus, it is a 1-cochain. For any 1-cochain ρ , Sweedler defines the 2-cocycle $(\partial\rho)(a, b) = \sum \rho(a_{(1)})\rho(b_{(1)})\rho^{-1}(a_{(2)} \vee b_{(2)})$. In particular, we call $\chi = \partial t^{-1}$ the 2-cocycle obtained from the inverse of t . In ref.[11], Sweedler shows that any 2-cocycle χ can be used to define an associative product $a \circ b = \sum \chi(a_{(1)}, b_{(1)})a_{(2)} \vee b_{(2)}$. Now we define the map \tilde{T} from $S(V_{\text{dp}})$ to $S(V_{\text{dp}})$ by $\tilde{T}(1) = 1$, $\tilde{T}(u) = u$ if $u \in V_{\text{dp}}$ and $\tilde{T}(a \vee b) = \tilde{T}(a) \circ \tilde{T}(b)$, so that $\tilde{T}(u_1 \vee \dots \vee u_n) = u_1 \circ \dots \circ u_n$ for u_i in V_{dp} . Then it can be shown that $\tilde{T} = T$. This fact has several consequences. From the quantum field point of view, it shows that the T -map arises from a renormalized product \circ which is associative. This fact seems not to have been realized before. From the quantum group point of view, renormalization is the replacement of a coquasitriangular structure by a more general 2-cocycle. However, not all 2-cocycles play a role in renormalization. The set of “renormalizing” 2-cocycles should be deduced from the Connes-Kreimer Hopf algebra of renormalization [37, 38, 39].

Let us give two last examples of the contact between quantum fields and quantum groups. We saw that any 1-cochain gives rise to a 2-cocycle. Since coquasitriangular structures are 2-cocycles, we could wonder which 1-cochains ω give rise to a coquasitriangular structure \mathcal{R} . The answer is very simple: the 1-cocycles that give rise to \mathcal{R} can be defined recursively by $\rho(a \vee b) = \sum \rho(a_{(1)})\rho(b_{(1)})\mathcal{R}(a_{(2)}, b_{(2)})$. If $\mathcal{R}(u^*, v^*) = \mathcal{R}(v, u)^*$, $\mathcal{R}(u^*, u) \geq 0$ and $\rho(u) = 0$ for any u, v in V , then ρ is a state for $S(V)$ (i.e. $\rho(a^* \vee a) \geq 0$ for any $a \in S(V)$). But these states are very special. They are called quasifree states. These states are fundamental to define a quantum field theory in curved spacetimes [40, 41]. In fact, the singular structure of any physically reasonable state of a scalar field in curved spacetimes is determined by a quasifree state [42]. The concept and importance of quasifree states emerged slowly in quantum field theory. Their peculiar nature is immediately clear from the quantum group point of view: they are the only states that

derive from a coquasitriangular structure. Finally, we consider a last problem posed by Kutzelnigg and Mukherjee [32]. When the normal products \tilde{a} adapted to ω are defined, how can we calculate their operator products? From the quantum group point of view, the answer is straightforward. If \mathcal{R} is the coquasitriangular structure used to define the product $a \circ b$, then the result is

$$\tilde{a} \circ \tilde{b} = \sum \chi(a_{(1)}, b_{(1)}) \widetilde{a_{(2)} \vee b_{(2)}},$$

where the 2-cocycle χ is $\chi = (\partial\omega^{-1}) \star \mathcal{R}$. Without the quantum group structure in sight, the same result can be obtained by hand for the first degrees, but its combinatorial complexity is appalling.

Conclusions – We showed that quantum groups provide a fresh look at many aspects of quantum field theory and they introduce powerful new concepts, such as coquasitriangular structures and cochains. On the practical side, quantum groups replace combinatorial proofs by algebraic proofs. On the conceptual side, quantum groups provide a second quantization without commutators that enables us to go beyond standard quantum field theory. For example, it allows the extension of quantum field theory to systems with quantum group symmetries [43]. That work employs a path integral approach which is dual (in the sense of quantum groups) to the present approach. (For comparison note that our \mathcal{R} is encoded there in γ .) In particular, this can be used to describe systems with anyonic particles [44] or models with non-commutative space-time structures [45].

We thank A. Frabetti, J.-L. Loday and B. Fauser for useful discussions, and M. Calandra and F. Mauri for their comments. R. O. was supported by a Marie Curie fellowship grant from the European Union. This is IPGP contribution #0000.

* Electronic address: brouder@lmcp.jussieu.fr

† Electronic address: oeckl@cpt.univ-mrs.fr

- [1] S. Majid, *Foundations of Quantum Group Theory* (Cambridge University Press, Cambridge, 1995).
- [2] F. C. Alcaraz, S. R. Salinas, and W. F. Wreszinski, *Phys. Rev. Lett.* **75**, 930 (1995).
- [3] M. A. Martín-Delgado and G. Sierra, *Phys. Rev. Lett.* **76**, 1146 (1996).
- [4] P. B. Wiegmann and A. V. Zabrodin, *Phys. Rev. Lett.* **72**, 1890 (1994).
- [5] F. Bonechi, E. Celeghini, R. Giachetti, E. Sorace, and M. Tarlini, *Phys. Rev. Lett.* **68**, 3718 (1992).
- [6] V. E. Korepin, N. M. Bogoliubov, and A. G. Izergin, *Quantum inverse scattering method and correlation functions* (Cambridge University Press, Cambridge, 1993).
- [7] G. Grensing, *Phys. Rev. B* **61**, 5483 (2000).
- [8] I. I. Kogan, *Int. J. Mod. Phys. A* **9**, 3887 (1994).
- [9] M. E. Sweedler, *Hopf Algebras* (W. A. Benjamin, New York, 1969).
- [10] C. Kassel, *Quantum Groups* (Springer Verlag, New York, 1995).
- [11] M. E. Sweedler, *Trans. Am. Math. Soc.* **133**, 205 (1968).
- [12] M. Reed and B. Simon, *Methods of Modern Mathematical Physics. II Fourier Analysis, Self-adjointness* (Academic Press, New York, 1975).
- [13] S. Dăscălescu, C. Năstăsescu, and Ş. Raianu, *Hopf Algebras* (Marcel Dekker Inc., New York, 2001).
- [14] C. Brouder, arXiv:math-ph/0201033.
- [15] F. D. Grosshans, G.-C. Rota, and J. A. Stein, *Invariant Theory and Superalgebras* (Amer. Math. Soc., Providence, 1987).
- [16] G. C. Wick, *Phys. Rev.* **80**, 268 (1950).
- [17] E. K. U. Gross, E. Runge, and O. Heinonen, *Many-Particle Theory* (Adam Hilger, Bristol, 1991).
- [18] B. Fauser, *J. Phys. A: Math. Gen.* **34**, 105 (2001).
- [19] S. N. Gupta, *Phys. Rev.* **107**, 1722 (1957).
- [20] J. Dito, *J. Math. Phys.* **33**, 791 (1992).
- [21] M. Dütsch and K. Fredenhagen, *Commun. Math. Phys.* **219**, 5 (2001).
- [22] A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw-Hill, Boston, 1971).
- [23] D. M. Esterling and R. V. Lange, *Rev. Mod. Phys.* **40**, 796 (1968).
- [24] T. T. S. Kuo, S. Y. Lee, and K. F. Ratcliff, *Nucl. Phys. A* **176**, 65 (1971).
- [25] L. S. Cederbaum and W. Domcke, *Adv. Chem. Phys.* **36**, 205 (1977).
- [26] A. G. Hall, *J. Phys. A: Math. Gen.* **8**, 214 (1975).
- [27] P. A. Henning, *Nucl. Phys. B* **337**, 547 (1990).
- [28] R. Fauser and H. H. Wolter, *Nucl. Phys. A* **584**, 604 (1995).
- [29] D. Mukherjee and W. Kutzelnigg, *J. Chem. Phys.* **114**, 2047 (2001).
- [30] D. Mukherjee and W. Kutzelnigg, *J. Chem. Phys.* **114**, 8226 (2001).
- [31] D. A. Mazziotti and R. M. Erdahl, *Phys. Rev. A* **63**, 042113 (2001).
- [32] W. Kutzelnigg and D. Mukherjee, *J. Chem. Phys.* **107**, 432 (1997).
- [33] C. Brouder, B. Fauser, A. Frabetti, and R. Oeckl, (unpublished).
- [34] R. Brunetti and K. Fredenhagen, in *Operator Algebras and Quantum Field Theory*, edited by S. Doplicher, R. Longo, J. Roberts, and L. Zsido (International Press of Boston, Cambridge, 1997), pp. 546–63.
- [35] H. Epstein and V. Glaser, *Ann. Inst. Henri Poincaré* **19**, 211 (1973).
- [36] G. Scharf, *Finite Quantum Electrodynamics* (Springer, Berlin, 1995), 2nd ed.
- [37] A. Connes and D. Kreimer, *Commun. Math. Phys.* **210**, 249 (2000).
- [38] A. Connes and D. Kreimer, *Commun. Math. Phys.* **216**, 215 (2001).
- [39] G. Pinter, *Lett. Math. Phys.* **54**, 227 (2000).
- [40] B. S. Kay and R. M. Wald, *Phys. Repts.* **207**, 49 (1991).
- [41] B. S. Kay, *J. Math. Phys.* **34**, 4519 (1993).
- [42] S. Hollands and W. Ruan, arXiv:gr-qc/0108032.
- [43] R. Oeckl, *Commun. Math. Phys.* **217**, 451 (2001).
- [44] R. Oeckl, *J. Geom. Phys.* **39**, 233 (2001).
- [45] R. Oeckl, *Nucl. Phys. B* **581**, 559 (2000).