

(Quasi)elastic electron–muon large-angle scattering to a two-loop approximation: eikonal type contributions. *

V.V. Bytev, E.A. Kuraev, B.G. Shaikhatdenov[†]

Joint Institute for Nuclear Research, Dubna, 141980, Russia

May 21, 2019

Abstract

Set of contributes to $e\mu$ large-angle high-energy scattering cross section is considered in quasi-elastic experimental set-up. Among them we had considered inelastic process with emission of one and two soft real photons as well as soft pion pairs. Virtual photon contributions are considered with logarithmical accuracy. Box-type Feynman amplitudes with vacuum polarization insertion of leptons and hadrons in the intermediate state are considered explicitly.

1 Introduction

The aim of our investigation is to calculate the next-to-leading contributions to the large-angle high energy (LAHE) cross section in quasi-elastic experimental set-up.

$$\frac{2E - E'_1 - E'_2}{2E} = \frac{\Delta E}{E} = \Delta \ll 1, \quad \Delta E \gg m_\mu, m_\pi \quad (1)$$

where E, E'_1, E'_2 are the energies of initial and scattered leptons in the frame of center of mass. The cross section will have in leading logarithm approximation the form of Drell-Yan process cross section [1]:

$$d\sigma(s, t) = \int dx_1 dx_2 D(x_1, \rho_t) D(x_2, \rho_t) d\sigma_0(sx_1x_2, tx_1x_3) D(x_3, \rho_t) D(x_4, \rho_t) dx_3 dx_4 \left(1 + \frac{\alpha}{\pi} k\right) \quad (2)$$

where $D(x, L)$ is the non-singlet structure function, which obey renorm-group (RG) equations. It's expansion in LLA

$$\frac{\alpha}{\pi}, \quad \frac{\alpha}{\pi} \rho_t \approx 1, \quad \rho_t = \ln \frac{-t}{m_e m_\mu}, \quad \rho_t \approx \rho_s = \ln \frac{s}{m_e m_\mu} \quad (3)$$

*supported in part by RFBR 01-02-17437 and by INTAS 00366 grant.

[†]on leave of absence from the Institute of Physics and Technology, Almaty-82, Kazakstan.

has a form:

$$D(x, \rho_t) = \delta(x - 1) + \frac{\alpha(\rho_t - 1)}{2\pi} P_1^{(1)}(x) + \dots + \frac{1}{n!} \left(\frac{\alpha(\rho_t - 1)}{2\pi} \right)^n P^{(n)}(x) + \dots, \quad (4)$$

In quasi-elastic set-up we can use only δ -part of $P_\Delta^{(i)}(x)$ of kernel $P^{(i)}(x)$:

$$P^{(i)}(x) = \int_x^1 \frac{dy}{y} P^{(1)}(y) P^{(i-1)}\left(\frac{x}{y}\right), \quad P^{(1)}(x) = P_\Delta^{(1)}(x) + P_\theta^{(1)}(x), \quad (5)$$

$$P_\Delta^{(1)}(x) = \delta(1 - x) P_\Delta^{(1)}, \quad P_\Delta^{(1)}(x) = 2 \ln \Delta + \frac{3}{2}, \quad P_\theta^{(1)}(x) = \theta(1 - x - \Delta) \frac{1 + x^2}{1 - x}.$$

So the structure function in that case has a form:

$$D(x, \rho_t) = \delta(1 - x) \left[1 + \sum_{k=1}^{\infty} \left(\frac{\alpha}{2\pi} (\rho_t - 1) \right)^k P_\Delta^{(k)} \right]. \quad (6)$$

So our aim is to calculate the k-factor in one and two-loop approximations.

Contributions to differential cross section are of two types. First is the one arising from virtual photon emission corrections up to two loop level, which require the calculations of real two-loop Feynman amplitudes.

Virtual -type contribution suffer from infrared divergencies, which are regularized by introduction of fictitious photon mass λ . Another type of contributions arise from emission of soft real photons and charged particle pairs.

General form of cross section can be presented as sum of three types: vertex part, eikonal part and decorated box type. Each of them contains virtual and real soft photon contribution and is free from infrared divergencies. All this contributions do not violate the LLA terms predicted by renorm-group despite that contributions of separate FD contain up to fourth power of large logarithm ρ_t in two-loop approximation.

In our previous paper [2] we had showed that the so-called vertex contributions provide the RG result 4 cited above.

This paper is devoted to investigation of some diagrams of E-type ones. For elastic process it corresponds to FD with one, two (box diagram) and three (double box FD) virtual photon exchanged by electron and muon.

Besides that we must take into account the box-type FD with vacuum polarization insertion to Green function to one of virtual photons (see fig.1). The single soft photon emission approximation must be applied to one-loop corrected Feynman amplitude. As well the emission of two soft photons (or charged particle pairs) must be taken into account.

In Section 2 we consider the vacuum polarization effects in box-type Feynman amplitudes with lepton pairs ($\mu^- \mu^+$, $e^- e^+$) or hadrons ($\pi^- \pi^+$) (fig.1). The relevant contributions from soft lepton pair and soft charged pion pair emission are considered as well as one and two soft real photon pair production (see fig.2,4).

In Section 3 the results of single and double soft photon emission are given as well as virtual corrections arising from the square of box-type amplitude (see fig. 3,4).

In Appendix A we give some details of virtual and soft real pair calculation and the complete set of scalar, vector and tensor integrals for box type FD with vacuum polarization insertion.

In Appendix B we give useful set of integrals for box type integrals.




Figure 1: Box-type FD with vacuum polarization insertion.




Figure 2: FD with soft lepton and pion pair production.




Figure 3: Box-type FD.




Figure 4: Some sorts of double soft photon emission FD.

2 Box-type FD with vacuum polarization insertion.

Vacuum polarization (VP) effects in Box-type Feynman amplitudes can be taken into account by replacement of photon Green function denominator $G(k)$ (see [3]):

$$\frac{1}{k^2} = \frac{\alpha}{3\pi} \int_0^1 \frac{dv\phi(v)}{(1-v^2)(k^2 - M^2(v))}, \quad M^2(v) = \frac{4\mu^2}{1-v^2}, \quad \phi(v) = 2 - (1-v^2)(2-v^2) \quad (7)$$

for the case of VP due to lepton with mass μ and

$$\frac{1}{k^2} = \frac{\alpha}{3\pi} \int_{4m_\pi^2}^\infty \frac{dM^2}{M^2} \frac{R(M^2)}{k^2 - M^2}, \quad R(M^2) = \frac{\sigma_{e\bar{e} \rightarrow \text{hadr}}(M^2)}{\sigma_{\mu\bar{\mu} \rightarrow \text{hadr}}(M^2)}. \quad (8)$$

for pion VP insertion. Here M is hadron's jet mass.

For matrix element squared we obtain:

$$\delta|M|_{vp(\text{lept})}^2 = \frac{2^8\alpha^4}{3t} \int_0^\infty \frac{dv\phi(v)}{1-v^2} \left(S(s, t, M^2(v)) - S(u, t, M^2(v)) \right) \quad (9)$$

for lepton VP, and

$$\delta|M|_{vp(\text{hadr})}^2 = \frac{2^8\alpha^4}{3t} \int_{4m_\pi^2}^\infty \frac{dM^2}{M^2} R(M^2) \left(S(s, t, M^2) - S(u, t, M^2) \right) \quad (10)$$

for hadron VP.

For the quantity $S(s, t, M)$ we have:

$$S(s, t, M^2) = \int \frac{d^4k}{i\pi^2} \frac{\text{Tr}(e) \cdot \text{Tr}(\mu)}{(1)(2)(3)(4)}, \quad (11)$$

where

$$(1) = k^2 - 2kp_1, \quad (2) = k^2 + 2kp_2, \quad (3) = k^2 - 2kq + \tilde{t}, \quad (4) = k^2 - \lambda^2, \quad (12)$$

$$\text{Tr}(e) = \frac{1}{4} Sp p_1 \gamma_\mu p'_1 \gamma_\nu (p_1 - k) \gamma_\lambda, \quad \text{Tr}(\mu) = \frac{1}{4} Sp p_2 \gamma_\mu p'_2 \gamma_\nu (p_2 + k) \gamma_\lambda,$$

$$p_1^2 = m_e^2, \quad p_2^2 = m_\mu^2, \quad s = (p_1 + p_2)^2, \quad t = (p_1 - p_2)^2, \quad \tilde{t} = t - M^2, \quad u = (p_1 - p'_2)^2.$$

Using the set of scalar, vector and tensor integrals for box-type four-momentum integrals (see App. A) we obtain the next result for the quantity S :

$$S(s, t, M^2) = u \left[\ln \frac{s}{-\tilde{t}} + \frac{M^2}{t} \ln \frac{-\tilde{t}}{M^2} \right] - (s(s-u) + \frac{t\tilde{t}}{2}) \left[I_{134} + I_{234} \right] + s(s^2 + u^2) I_{1234} \quad (13)$$

$$+ s(u + \frac{\tilde{t}}{2}) \left[-I_{123} - I_{124} + \tilde{t} I_{1234} \right],$$

where

$$I_{ijl} = \int \frac{d^4k}{i\pi^2} \frac{1}{(i)(j)(l)}, \quad I_{1234} = \int \frac{d^4k}{i\pi^2} \frac{1}{(1)(2)(3)(4)}. \quad (14)$$

Making integration and neglecting the terms of order $m_\mu^2/(-t) \ll 1$ we have:

$$\begin{aligned}
S(s, t, M^2) - S(u, t, M^2) \Big|_{-t \gg M^2} &= \frac{s^2 + u^2}{t^2} L_{us} (\rho_m - 2\rho_t - \rho_\lambda) \\
&+ (u - s) \left(\pi^2 + \rho_m^2 - \frac{1}{2} L_{st}^2 - \frac{1}{2} L_{ut}^2 + \ln^2 \frac{m_e}{m_\mu} \right) + u L_{st} - s L_{ut}, \\
\rho_m &= \ln \frac{M^2}{m_e m_\mu}, \quad \rho_\lambda = \ln \frac{m_e m_\mu}{\lambda^2}, \\
L_{st} &= \ln \frac{s}{-t}, \quad L_{ut} = \ln \frac{u}{t}, \quad L_{su} = \ln \frac{s}{-u},
\end{aligned} \tag{15}$$

for the case of big t . For the case of small t we have:

$$\begin{aligned}
S(s, t, M^2) - S(u, t, M^2) \Big|_{M^2 \gg -t} &= \frac{1}{M^2} \left[\frac{s^2 + u^2}{2} L_{us} (\rho_s + \rho_u + 2\rho_\lambda) + \frac{3}{2} (u^2 \rho_s - s^2 \rho_u) \right. \\
&\left. + t^2 L_{us} + t(u - s) \left(\frac{3}{2} \rho_M + \frac{7}{4} \right) \right].
\end{aligned} \tag{16}$$

Further integration for the case of lepton VP with mass $M^2 = 4\mu^2/(1 - v^2)$ (both cases $\mu = m_e, m_\mu$ are taken into account) leads to (with logarithmical accuracy):

$$\begin{aligned}
\frac{d\sigma_{vp}^{box}}{d\sigma_0} &= \frac{2\alpha^2}{3\pi^2} \rho_t \left[2L_{su} \left(\frac{3}{2} \rho_t + \rho_\lambda - \frac{10}{3} \right) + \frac{t(s - u)}{u^2 + s^2} (L_{st}^2 + L_{ut}^2 - 2\pi^2) \right. \\
&\left. + \frac{2t}{s^2 + u^2} ((t - s)L_{st} - (t - u)L_{ut}) \right].
\end{aligned} \tag{17}$$

To obtain the final result without infrared divergencies we must take into account the interference of soft photon emission amplitudes in Born level and one with lepton VP insertion:

$$\frac{d\sigma_{vp}^\gamma}{d\sigma_0} = -\frac{4\alpha^2}{3\pi^2} \left(\rho_t - \frac{5}{3} \right) \left[(2 \ln \Delta + \rho_\lambda) L_{su} + \rho_t L_{su} - \text{Li}_2 \left(\frac{1 - c}{2} \right) \right], \tag{18}$$

where Δ was defined in(1), and also consider the contribution from soft lepton pair emission with energy not exceeding $\Delta\epsilon$, $2\mu \ll \Delta\epsilon \ll \epsilon$:

$$\frac{d\sigma_{sp}}{d\sigma_0} = -\frac{2\alpha^2}{3\pi^2} \rho_t \left[L_{su} (\rho_t + L_{st} + L_{ut} + 2(2 \ln \Delta - \frac{5}{3})) - 2\text{Li}_2 \left(\frac{1 - c}{2} \right) \right]. \tag{19}$$

The summed result for total correction of e, μ VP effects and soft $e\bar{e}, \mu\bar{\mu}$ pair emission (see (17,18,19)):

$$\begin{aligned}
\frac{d\sigma_{vp,sp}}{d\sigma_0} &= \frac{2\alpha^2}{3\pi^2} \rho_t \left[-L_{su} (L_{st} + L_{ut} + 8 \ln \Delta) + \frac{t(s - u)}{s^2 + u^2} (L_{ut}^2 + L_{st}^2 - 2\pi^2) \right. \\
&\left. + \frac{2t}{s^2 + u^2} (tL_{su} - sL_{st} + uL_{ut}) + 4\text{Li}_2 \left(\frac{1 - c}{2} \right) \right], \quad c = \cos(\vec{p}_e, \vec{p}_e').
\end{aligned} \tag{20}$$

We see that this quantity contains only next-to leading terms of order $\alpha^2 \rho_t$ and does not contain infrared divergencies.

Let now return to soft pion pair production with total energy not exceeded $\Delta\epsilon$ and invariant square mass M^2 :

$$4m_\pi^2 \ll M^2 < (\Delta\epsilon)^2 \ll \epsilon^2 = s/4. \quad (21)$$

The corresponding contribution to differential cross section arises from interference of emission from electron and muon blocks:

$$\left. \frac{d\sigma}{dM^2 d\sigma_0} \right|_{soft\ pions} = 2 \left(\frac{4\pi\alpha}{M^2} \right)^2 \frac{d^4q}{M^2} \int \frac{d^3q_+ d^3q_-}{2\epsilon_+ 2\epsilon_-} \delta^4(q_+ + q_- - q) \left(\frac{Qp'_1}{p'_1q} - \frac{Qp_1}{p_1q} \right) \left(\frac{Qp'_2}{p'_2q} - \frac{Qp_2}{p_2q} \right), \quad (22)$$

$$q^2 = M^2, \quad Q = q_+ - q_-.$$

Perform the first invariant pion pair space integration we have:

$$\int \frac{d^3q_+ d^3q_-}{2\epsilon_+ 2\epsilon_-} \delta^4(q_+ + q_- - q) Q_\mu Q_\nu = \frac{1}{3} \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) Q^2 \frac{\pi\beta}{2}, \quad \beta = \sqrt{1 - \frac{4m^2}{q^2}}. \quad (23)$$

Arranging the phase volume as

$$\int \frac{d^4q}{dq^2} = \frac{1}{2} \int_{\sqrt{q^2}}^{\Delta\epsilon} dq_0 \sqrt{q_0^2 - q^2} dO_q \quad (24)$$

we put the r.h.s. of equation to the form:

$$\frac{\alpha^2}{3\pi^2} \int_M^{\Delta\epsilon} dq_0 \sqrt{q_0^2 - M^2} \int \frac{dO_q}{4\pi} \left(\frac{p_1 p_2}{p_1 q \cdot p_2 q} - \frac{p_1 p'_2}{p_1 q \cdot p'_2 q} \right) = \frac{\alpha^2}{3\pi^2} \left(L_{su} \ln \frac{2\Delta\epsilon}{M} + \mathcal{O}(1) \right). \quad (25)$$

The result is:

$$\left. \frac{d\sigma}{dM^2 d\sigma_0} \right|_{soft\ pions} = \frac{\alpha^2}{6\pi^2 M^2} L_{su} (\rho_t - \rho_M). \quad (26)$$

Contribution from box type FD with hadron VP cannot be obtained in analytical form due to presence of $R(M^2)$.

3 Soft photon emission contribution. Squared box contribution.

Interference of amplitudes of soft photon emission by electron and muon blocks can be evaluated by using the form:

$$I_{p_A p_B} = \frac{1}{4\pi} \int \frac{d^3k}{\omega} \frac{p_A p_B}{p_A k \cdot p_B k} \Big|_{\omega < \Delta\epsilon} \quad (27)$$

$$= (\ln \Delta + \frac{1}{2} \rho_\lambda) L_{AB} + \frac{1}{4} (L_{AB}^2 - L^2 - \frac{\pi^2}{3}) - \frac{\pi^2}{12} + \frac{1}{2} \text{Li}_2 \left(\frac{1+c}{2} \right),$$

where

$$L_{AB} = \ln \frac{2p_A p_B}{m_e m_\mu}, \quad \rho_\lambda = \ln \frac{m_e m_\mu}{\lambda^2}, \quad L = \ln \frac{m_\mu}{m_e}, \quad (28)$$

$$p_A p_B = \epsilon^2(1-c), \quad p_A^2 = m_e^2, \quad P_B^2 = m_\mu^2, \quad E_A = E_B = \epsilon.$$

Using the known results for interference of Born and box type elastic amplitudes and soft photon approximation we obtain for single photon emission contribution:

$$\frac{d\sigma_{vert}^\gamma}{d\sigma_0} = \left(\frac{\alpha}{\pi}\right)^2 \left[2L_{su}(\rho_t + \rho_\lambda) + \frac{t^2}{s^2 + u^2} \left(\frac{u}{t} L_{st} - \frac{s}{t} L_{ut} + \frac{s-u}{2t} (\pi^2 + L_{ut}^2 + L_{st}^2) \right) \right] \quad (29)$$

$$\times \left[-L_{su}\rho_t + \frac{1}{2}(L_{ut}^2 - L_{st}^2) - 2L_{su}(\ln \Delta + \frac{1}{2}\rho_\lambda) - \frac{pi^2}{6} + \text{Li}_2\left(\frac{1-c}{2}\right) \right].$$

For emission of two photons with total energy not exceeding $\Delta\epsilon$ we have:

$$\frac{d\sigma^{\gamma\gamma}}{d\sigma_0} = \left(\frac{2\alpha}{\pi}\right)^2 \left[\left[\frac{1}{2}\rho_t L_{su} + \frac{1}{4}(L_{st}^2 - L_{ut}^2) + L_{su}(\ln \Delta + \frac{1}{2}\rho_\lambda) + \frac{\pi^2}{12} - \frac{1}{2}\text{Li}_2\left(\frac{1-c}{2}\right) \right]^2 - \frac{\pi^2}{6} L_{su}^2 \right]. \quad (30)$$

Using the results of App. B we construct the contribution of box type square amplitude, which is relevant in this order:

$$\frac{d\sigma_{BB}}{d\sigma_0} = \frac{\alpha^2}{4\pi^2} \frac{t^2}{s^2 + u^2} \rho_t \left[A\rho_t + B \right], \quad (31)$$

where

$$A = 8 \frac{s^2 + u^2}{t^2} (L_{us}^2 + \pi^2), \quad B = \frac{16(s^2 + u^2)}{t^2} L_{us}^2 \rho_\lambda + 8L_{us} \left(\frac{s}{t} L_{ut} - \frac{u}{t} L_{st} \right) \quad (32)$$

$$+ 4 \frac{s-u}{t} (\pi^2 (2L_{st} - L_{su}) - L_{us} (L_{ut}^2 + L_{st}^2)) + \frac{8u}{t} \pi^2.$$

4 Summary

Here we consider some types of second-order contributes in large-angle $e-\mu$ scattering. For the case of box FD with polarization insertion we put the formulae (9,10) in which we show that contributions from interference of straight and cross "legs" of the FD with vacuum polarization insertion and Born FD are equal when one makes exchange $s \rightarrow u$ and changes sign of contribution to opposite.

Also we put the explicit expression for matrix element squared in terms of logarithms up to LLA. There we show two cases: for small (15) and big (16) masses M of virtual lepton and pion. By integrating over the quantity v and adding contribution of soft lepton pairs we achieve the free of infra-red discontinuities contribution of FD with lepton vacuum polarization insertion (20). This contribution is in agreement with RG predictions and does not contain large logarithms of third and fourth power. Also we put here expression for cross-section of soft pion production (26). Here we can not explicitly show the cancellation of L and NL logarithms with soft pion contribution due to partially non-analytical form of expression.

Also we bring here the expression for squared box contribution with addition relevant one and two soft photon emission contribution to avoid infra-red divergencies (31). The explicit expressions for soft photon emission are also written down (29,30).

The remnant part of contribution which arising from Born and eikonal box amplitudes will be considered in a next paper.

5 Appendix A

Here we put details of calculating box type diagram with VP insertion. For this we need integrals with mass virtual photon. For evaluating vector and tensor integrals we can use the results of Appendix B (box type contribution) in which all scalar integrals with three (I_{ijk}) and four (I) denominators change to the next integrals with one mass virtual photon: Let's consider the first the case of big mass M , $M^2 \gg s \approx -t$

$$\begin{aligned}
I_{123} &= \frac{1}{M^2} \left[-\ln \frac{M^2}{s} - 1 + \frac{s}{M^2} \left[\frac{1}{2} \ln \frac{M^2}{s} + \frac{1}{4} \right] \right], \\
I_{134} &= -\frac{1}{M^2} \left[\ln \frac{M^2}{m_e^2} + 1 + \frac{t}{M^2} \left[\frac{1}{2} \ln \frac{M^2}{m_e^2} + \frac{1}{4} \right] \right], \\
I_{234} &= -\frac{1}{M^2} \left[\ln \frac{M^2}{m_\mu^2} + 1 + \frac{t}{M^2} \left[\frac{1}{2} \ln \frac{M^2}{m_\mu^2} + \frac{1}{4} \right] \right], \\
I &= -\frac{1}{2sM^2} \left[2 \ln \frac{s}{m_e m_\mu} \ln \frac{m_e m_\mu}{\lambda^2} + \ln^2 \frac{s}{m_e m_\mu} - \frac{4\pi^2}{3} \right], \\
I_3 = I\tilde{t} - I_{124} &= \frac{1}{M^2} \left[-\frac{2t}{s} \ln \frac{s}{m_e m_\mu} + 1 - \ln \frac{s}{M^2} + \frac{s}{M^2} \left[\frac{1}{2} \ln \frac{s}{M^2} - \frac{1}{4} + \frac{t^2}{s^2} \ln \frac{s}{m_e m_\mu} \right] \right].
\end{aligned} \tag{33}$$

For the case of big $-t$, $-t \gg M^2$ we should use the next set of integrals:

$$\begin{aligned}
I_{134} &= \frac{1}{t} \left[\ln \frac{M^2}{m_e^2} \ln \frac{-t}{M^2} + \frac{\pi^2}{6} + \frac{1}{2} \ln^2 \frac{-t}{M^2} \right], \\
I_{234} &= \frac{1}{t} \left[\ln \frac{M^2}{m_\mu^2} \ln \frac{-t}{M^2} + \frac{\pi^2}{6} + \frac{1}{2} \ln^2 \frac{-t}{M^2} \right], \\
I_{123} &= -\frac{1}{2s} \left[2 \ln \frac{s}{m_e m_\mu} \ln \frac{M^2}{m_e m_\mu} + \frac{4\pi^2}{3} - \ln^2 \frac{s}{m_e m_\mu} + \ln^2 \frac{m_\mu}{m_e} \right], \\
I_3 &= \frac{1}{s} \left[2 \ln \frac{s}{m_e m_\mu} \ln \frac{-t}{M^2} - \ln \frac{s}{m_e m_\mu} \ln \frac{m_e m_\mu}{M^2} - \frac{1}{2} \ln^2 \frac{s}{m_e m_\mu} + \frac{1}{2} \ln^2 \frac{m_e}{m_\mu} + \frac{2\pi^2}{3} \right], \\
I_{1234} &= -\frac{2}{st} \ln \frac{s}{m_e m_\mu} \ln \frac{\lambda M}{-t}.
\end{aligned} \tag{34}$$

6 Appendix B

Here we put the details of box-box contribution calculation. The first of all we should distinguish three different cases: two box square with straight and crossed legs and one case with interference of amplitudes with crossed and straight legs.

For calculating the contributions we need to evaluate tensor, vector and scalar integrals with four and three denominators.

Lets first consider integral for box with straight legs. The vector integral we can write down in the form:

$$\int \frac{d^4 k k^\mu}{i\pi^2(1)(2)(3)(4)} = Ap_{1\mu} + Bp_{2\mu} + Cq_\mu, \tag{35}$$

where quantities (1), (2), (3), (4) were defined in (12), $q = p_1 - p'_1 = p'_2 - p_2$, $q^2 = t$ and coefficients A, B, C determined as follows:

$$\begin{aligned}
A &= \frac{1}{2stu}[-t^2a - t(2s+t)b - stc], \\
B &= \frac{1}{2stu}[-t(2s+t)a - t^2b + stc], \\
C &= \frac{1}{2stu}[-sta + stb - s^2c], \\
a &= I_{123} - I_{234}, \quad b = I_{134} - I_{123}, \quad c = tI.
\end{aligned} \tag{36}$$

Scalar integrals I, I_{ijk} reads:

$$\begin{aligned}
I &= \int \frac{d^4k}{i\pi^2(1)(2)(3)(4)} = \frac{2}{st} \left[\ln \frac{s}{mM} - i\pi \right] \ln \frac{-t}{\lambda^2}, \\
I_{123} &= \int \frac{d^4k}{i\pi^2(1)(2)(3)} = -\frac{1}{2s} \left[2 \left[\ln \frac{s}{mM} - i\pi \right] \ln \frac{\lambda^2}{mM} + \frac{\pi^2}{3} - \left[\ln \frac{s}{mM} - i\pi \right]^2 + \ln^2 \frac{M}{m} \right], \\
I_{134} &= \int \frac{d^4k}{i\pi(1)(3)(4)} = \frac{1}{t} \left[\frac{1}{2} \ln^2 \frac{-t}{m^2} + \frac{2\pi^2}{3} \right], \\
I_{234} &= \int \frac{d^4k}{i\pi(2)(3)(4)} = \frac{1}{t} \left[\frac{1}{2} \ln^2 \frac{-t}{M^2} + \frac{2\pi^2}{3} \right].
\end{aligned} \tag{37}$$

Tensor integral we consider by algebraical method:

$$\begin{aligned}
\int \frac{d^4k k_\mu k_\nu}{i\pi^2(1)(2)(3)(4)} &= a_g g_{\mu\nu} + a_{11} p_{1\mu} p_{1\nu} + a_{22} p_{2\mu} p_{2\nu} + a_{12} (p_{1\mu} p_{2\nu} + p_{1\nu} p_{2\mu}) \\
&\quad + a_{1q} (p_{1\mu} q_\nu + p_{1\nu} q_\mu) + a_{2q} (p_{2\mu} q_\nu + p_{2\nu} q_\mu) + a_{qq} q_\mu q_\nu.
\end{aligned} \tag{38}$$

To time the equation above to four-vectors p_1, p_2, q we can manage the system of algebraic equations, therefore we write down the quantities a_{ij} through scalar integrals:

$$\begin{aligned}
a_{22} &= \frac{1}{s}(A_2 - ta_{2q}), \\
a_{11} &= \frac{1}{s}(A_4 + ta_{1q}), \\
a_{12} &= \frac{1}{s}(A_1 - 2a_g - ta_{1q}), \\
a_g &= \frac{1}{2}(A_9 - 2ta_{qq} - ta_{1q} + ta_{2q}), \\
a_{1q} &= \frac{1}{t}(A_1 - A_5 - ta_{2q}), \\
a_{2q} &= \frac{1}{s+t}(A_3 + A_{10} - A_5 - A_9), \\
a_{qq} &= \frac{1}{t(s+t)}(tA_3 + s(A_5 + A_9 - A_{10})).
\end{aligned} \tag{39}$$

Quantities A_j read:

$$\begin{aligned}
A_1 &= \frac{1}{s}(I_{13} - I_{12}), \\
A_2 &= I_{234} + \frac{1}{s}(I_{12} - I_{23}) + \frac{1}{t}(2I_{34} - I_{23} - I_{24}), \\
A_3 &= I_{123} + \frac{1}{s}(2I_{12} - I_{13} - I_{23}) + \frac{1}{t}(-I_{23} + I_{34}), \\
A_4 &= I_{134} + \frac{1}{s}(I_{12} - I_{13}) + \frac{1}{t}(2I_{34} - I_{13} - I_{14}), \\
A_5 &= \frac{1}{s}(I_{23} - I_{12}), \\
A_9 &= tC + I_{123} + \frac{1}{s}(2I_{12} - I_{13} - I_{23}), \\
A_{10} &= I_{123}.
\end{aligned} \tag{40}$$

Here I_{ij} means scalar integrals with two denominators, I_{ijk} and I are determined above.

$$\begin{aligned}
I_{12} &= \int \frac{d^4k}{i\pi^2(1)(2)} = \ln \frac{\Lambda^2}{M^2} - \ln \frac{s}{M^2} + i\pi + 1 \\
I_{13} &= \int \frac{d^4k}{i\pi^2(1)(3)} = I_{14} = \int \frac{d^4k}{i\pi^2(1)(4)} = \ln \frac{\Lambda^2}{M^2} + \ln \frac{M^2}{m^2} + 1 \\
I_{23} &= \int \frac{d^4k}{i\pi^2(2)(3)} = I_{24} = \int \frac{d^4k}{i\pi^2(2)(4)} = \ln \frac{\Lambda^2}{M^2} + 1 \\
I_{34} &= \int \frac{d^4k}{i\pi^2(3)(4)} = \ln \frac{\Lambda^2}{M^2} - \ln \frac{-t}{M^2} + 1
\end{aligned} \tag{41}$$

While evaluating box type diagram square we must also consider complex conjugated amplitude. The integrals for it can be made by exchange $i\pi \rightarrow -i\pi$ in equations above.

For crossed legs in box type diagram we should evaluate the next integrals:

$$\int \frac{d^4k k^\mu}{i\pi^2(1)(\tilde{2})(3)(4)} = Ap_{1\mu} - Bp'_{2\mu} + Cq_\mu. \tag{42}$$

where $(\tilde{2}) = k^2 + p'_2 k$ and

$$\begin{aligned}
A &= \frac{1}{2stu}[-t^2a - t(2u + t)b - utc], \\
B &= \frac{1}{2stu}[-t(2u + t)a - t^2b + utc], \\
C &= \frac{1}{2stu}[-uta + utb - u^2c], \\
a &= I_{1\tilde{2}3} - I_{\tilde{2}34}, \quad b = I_{134} - I_{1\tilde{2}3}, \quad c = t\tilde{I}.
\end{aligned} \tag{43}$$

Here integrals read:

$$\begin{aligned}
I &= \int \frac{d^4k}{i\pi^2(1)(\tilde{2})(3)(4)} = \frac{2}{ut} \ln \frac{-u}{mM} \ln \frac{-t}{\lambda^2}, \\
I_{1\tilde{2}3} &= \int \frac{d^4k}{i\pi^2(1)(\tilde{2})(3)} = -\frac{1}{2u} \left[2 \ln \frac{-u}{mM} \ln \frac{\lambda^2}{mM} + \frac{\pi^2}{3} - \ln^2 \frac{-u}{mM} + \ln^2 \frac{M}{m} \right], \\
I_{134} &= \int \frac{d^4k}{i\pi(1)(3)(4)} = \frac{1}{t} \left[\frac{1}{2} \ln^2 \frac{-t}{m^2} + \frac{2\pi^2}{3} \right], \\
I_{\tilde{2}34} &= \int \frac{d^4k}{i\pi(\tilde{2})(3)(4)} = \frac{1}{t} \left[\frac{1}{2} \ln^2 \frac{-t}{M^2} + \frac{2\pi^2}{3} \right].
\end{aligned} \tag{44}$$

For tensor integral we have:

$$\begin{aligned}
\int \frac{d^4k k_\mu k_\nu}{i\pi^2(1)(\tilde{2})(3)(4)} &= a_g g_{\mu\nu} + a_{11} p_{1\mu} p_{1\nu} + a_{22} p'_{2\mu} p'_{2\nu} - a_{12} (p_{1\mu} p'_{2\nu} + p_{1\nu} p'_{2\mu}) \\
&\quad + a_{1q} (p_{1\mu} q_\nu + p_{1\nu} q_\mu) - a_{2q} (p'_{2\mu} q_\nu + p'_{2\nu} q_\mu) + a_{qq} q_\mu q_\nu,
\end{aligned} \tag{45}$$

here we use:

$$\begin{aligned}
a_{22} &= \frac{1}{u} (A_2 - t a_{2q}), \\
a_{11} &= \frac{1}{u} (A_4 + t a_{1q}), \\
a_{12} &= \frac{1}{u} (A_1 - 2a_g - t a_{1q}), \\
a_g &= \frac{1}{2} (A_9 - 2t a_{qq} - t a_{1q} + t a_{2q}), \\
a_{1q} &= \frac{1}{t} (A_1 - A_5 - t a_{2q}), \\
a_{2q} &= \frac{1}{u+t} (A_3 + A_{10} - A_5 - A_9), \\
a_{qq} &= \frac{1}{t(u+t)} (t A_3 + u(A_5 + A_9 - A_{10})).
\end{aligned} \tag{46}$$

Quantities A_j read:

$$\begin{aligned}
A_1 &= \frac{1}{u} (I_{13} - I_{12}), \\
A_2 &= I_{\tilde{2}34} + \frac{1}{u} (I_{1\tilde{2}} - I_{\tilde{2}3}) + \frac{1}{t} (2I_{34} - I_{\tilde{2}3} - I_{\tilde{2}4}), \\
A_3 &= I_{1\tilde{2}3} + \frac{1}{u} (2I_{1\tilde{2}} - I_{13} - I_{\tilde{2}3}) + \frac{1}{t} (-I_{\tilde{2}3} + I_{34}), \\
A_4 &= I_{134} + \frac{1}{u} (I_{1\tilde{2}} - I_{13}) + \frac{1}{t} (2I_{34} - I_{13} - I_{14}), \\
A_5 &= \frac{1}{u} (I_{\tilde{2}3} - I_{1\tilde{2}}), \\
A_9 &= tc + I_{1\tilde{2}3} + \frac{1}{u} (2I_{1\tilde{2}} - I_{13} - I_{\tilde{2}3}), \\
A_{10} &= I_{1\tilde{2}3}.
\end{aligned} \tag{47}$$

Here I_{ij} means scalar integrals with two denominators, I_{ijk} and I are determined above.

$$\begin{aligned}
I_{12} &= \int \frac{d^4k}{i\pi^2(1)(2)} = \ln \frac{\Lambda^2}{M^2} - \ln \frac{-u}{M^2} + 1 \\
I_{13} &= \int \frac{d^4k}{i\pi^2(1)(3)} = I_{14} = \int \frac{d^4k}{i\pi^2(1)(4)} = \ln \frac{\Lambda^2}{M^2} + \ln \frac{M^2}{m^2} + 1 \\
I_{23} &= \int \frac{d^4k}{i\pi^2(2)(3)} = I_{24} = \int \frac{d^4k}{i\pi^2(2)(4)} = \ln \frac{\Lambda^2}{M^2} + 1 \\
I_{34} &= \int \frac{d^4k}{i\pi^2(3)(4)} = \ln \frac{\Lambda^2}{M^2} - \ln \frac{-t}{M^2} + 1
\end{aligned} \tag{48}$$

By using this set of integral we can obtain for interference of Born-box matrix element :

$$\begin{aligned}
2 \sum M_B^* M_{box} &= \sum |M_B|^2 * \frac{\alpha}{\pi} \left[2L_{su}(\rho_t + \rho_\lambda) + \frac{t^2}{s^2 + u^2} \left[\frac{u}{t} L_{st} - \frac{s}{t} L_{ut} \right] \right. \\
&\quad \left. + \frac{s-u}{2t} \left[\pi^2 + L_{ut}^2 + L_{st}^2 \right] \right].
\end{aligned} \tag{49}$$

Adding to this expression the contribution arising from interference of soft photon emission by electron and muon block we arrive to expression K given in equation (16) of paper [2].

References

- [1] E. Kuraev, V. Fadin Sov. J. Nucl. Phys. 1985 v. 41 p. 466
E. Kuraev, V. Fadin Sov. J. Nucl. Phys. 1988 v. 47 p. 1009
- [2] V. Bytev, E. Kuraev, B. Shaikhatdenov JETP, vol. 9 (2002)
- [3] J. Schwinger Phys. Rev. 76 (1949) p. 1790