

$\Delta \rightarrow N\gamma$ in Large- N_c QCD

Elizabeth Jenkins*

*University of California, San Diego
9500 Gilman Drive, La Jolla, CA 92093-0319*

Xiangdong Ji†

*Department of Physics, University of Maryland,
College Park, MD 20742*

Aneesh V. Manohar‡

*University of California, San Diego
9500 Gilman Drive, La Jolla, CA 92093-0319*

(Dated: July 2002)

The decay $\Delta^+ \rightarrow p\gamma$ is studied in the $1/N_c$ expansion of QCD. The ratio of the helicity amplitudes is determined to be $A_{3/2}/A_{1/2} = \sqrt{3} + \mathcal{O}(1/N_c^2)$. Equivalently, the ratio $E2/M1$ of the multipole amplitudes is predicted to be order $1/N_c^2$.

One of the mysteries in hadronic physics is the success of the non-relativistic quark model in describing the structure and spectrum of the low-lying baryons. In the non-relativistic quark model, the spin structure of baryons is determined entirely by the spin of the constituent quarks, in clear contradiction to the result from polarized deep-inelastic scattering that only 20-30% of the nucleon spin is carried by the spin of the quarks [1]. Nevertheless, the non-relativistic quark model successfully predicts that the ratio of the neutron to proton magnetic moments is $-2/3$ and that the Δ to N electromagnetic transition is dominated by the M1 multipole. Recent advances in studying baryons in the $1/N_c$ expansion of QCD help to reconcile the success of the quark model predictions with the spin structure of the nucleon [2, 3, 4, 5].

The large- N_c limit of QCD validates many of the quark model spin-flavor symmetry results without assuming a non-relativistic quark dynamics for the baryon wave functions. An $SU(6)$ spin-flavor symmetry of baryons was derived in the large- N_c limit, which yields the same spin-flavor results as the non-relativistic quark model and Skyrme model without making any of the dynamical assumptions of these models. In the $1/N_c$ expansion of QCD, the ratio of the neutron and proton magnetic moments is predicted to be $-2/3$ up to a correction of order $1/N_c^2$ [2, 3, 4, 5]. Likewise, it is shown in this paper that the $E2/M1$ electromagnetic transition ratio between Δ and N is of order $1/N_c^2$. This result requires no assumption about the underlying quark orbital angular momentum, or intrinsic deformation, of the baryon.

The radiative decay $\Delta^+ \rightarrow p\gamma$ is described by two independent amplitudes. The standard convention [6] is to use helicity amplitudes $A_{1/2}$ and $A_{3/2}$, which are the am-

plitudes for a Δ^+ state with helicity $1/2$ or $3/2$ along its direction of motion to decay to a right-handed photon moving along this same direction, and a proton recoiling in the opposite direction. The helicity amplitudes for Δ^+ decay to a left-handed photon are related to $A_{1/2}$ and $A_{3/2}$ by parity conservation of the strong and electromagnetic interactions. The total decay rate for $\Delta^+ \rightarrow p\gamma$ is given by

$$\Gamma = \frac{k_\gamma^2}{\pi} \frac{2M_p}{(2J+1)M_\Delta} \left[|A_{1/2}|^2 + |A_{3/2}|^2 \right], \quad (1)$$

where $k_\gamma = 259$ MeV is the momentum of the photon, M_p is the proton mass, M_Δ is the Δ^+ mass, and $J = 3/2$ is the spin of the Δ^+ . Eq. (1) adheres to the normalization convention of the helicity amplitudes used by the Particle Data Group [6]. The experimental values for the helicity amplitudes are

$$\begin{aligned} A_{1/2} &= -0.135 \pm 0.006 \text{ GeV}^{-1/2}, \\ A_{3/2} &= -0.255 \pm 0.008 \text{ GeV}^{-1/2}, \end{aligned} \quad (2)$$

which yield the ratio

$$\frac{A_{3/2}}{A_{1/2}} = 1.89 \pm 0.10. \quad (3)$$

In this paper, we show that the large- N_c prediction for this ratio is

$$\frac{A_{3/2}}{A_{1/2}} = \sqrt{3} + \mathcal{O}\left(\frac{1}{N_c^2}\right) = 1.73 + \mathcal{O}\left(\frac{1}{N_c^2}\right), \quad (4)$$

in good agreement with the experimental result.

The $\Delta^+ \rightarrow p\gamma$ also can be described using a multipole expansion. The only multipoles which contribute are the magnetic moment (M1) and the electric quadrupole (E2). These multipoles are related to the helicity amplitudes by [6]

$$\begin{aligned} A_{1/2} &= -\frac{1}{2}(M1 + 3E2), \\ A_{3/2} &= -\frac{\sqrt{3}}{2}(M1 - E2). \end{aligned} \quad (5)$$

*Electronic address: ejenkins@ucsd.edu

†Electronic address: xji@physics.umd.edu

‡Electronic address: amanohar@ucsd.edu

The total decay rate Γ in Eq. (1) depends on

$$\left[|A_{1/2}|^2 + |A_{3/2}|^2 \right] = \left[|M1|^2 + 3|E2|^2 \right]. \quad (6)$$

The experimental value for the ratio $E2/M1$ is

$$\frac{E2}{M1} = -0.025 \pm 0.005. \quad (7)$$

We show that the large- N_c prediction for $E2/M1$,

$$\frac{E2}{M1} = \mathcal{O}\left(\frac{1}{N_c^2}\right), \quad (8)$$

is equivalent to the prediction Eq. (4) for the ratio of helicity amplitudes. The experimental measurement Eq. (7) is consistent with the prediction that $E2/M1$ is order $1/N_c^2$.

The large- N_c expansion for baryons gives an expansion in powers of $1/N_c$ for operators with definite spin and flavor transformation properties [4, 5, 7, 8]. In the $N_c \rightarrow \infty$ limit, the baryons are infinitely heavy and can be treated as static [9], with recoil effects included as $1/N_c$ corrections [10]. The static kinematics of the large- N_c limit makes it natural to use large- N_c methods to evaluate the multipole amplitudes, rather than the helicity amplitudes.

The $\Delta^+ \rightarrow p\gamma$ transition is an $I = 3/2 \rightarrow I = 1/2$ transition mediated by the isovector part of the electromagnetic current. The isovector electromagnetic current operator can be expanded in powers of the photon momentum (the multipole expansion), and is proportional to

$$(J_{EM})^{ia} \propto \mu^{ia} + Q^{(ij)a} k_\gamma^j + \dots \quad (9)$$

where i and $j = 1, 2, 3$ are vector spin indices, and $a = 1, 2, 3$ is a vector isospin index. In Eq. (9), μ^{ia} is the magnetic moment operator, which is an $(J, I) = (1, 1)$ operator, and $Q^{(ij)a}$ is the quadrupole moment operator, which is an $(J, I) = (2, 1)$ operator. The quadrupole operator $Q^{(ij)a}$ is a symmetric, traceless tensor in spin indices i and j . The $M1$ and $E2$ amplitudes are related to the matrix elements of the magnetic moment and quadrupole moment operators through

$$\begin{aligned} M1 &= ek_\gamma^{1/2} \langle N \uparrow | \mu_z | \Delta \uparrow \rangle, \\ E2 &= \frac{1}{12} ek_\gamma^{3/2} \langle N \uparrow | Q_{20} | \Delta \uparrow \rangle, \end{aligned} \quad (10)$$

where $\mu_z = \mu^{ia}$ for $i = 3, a = 3$, and Q_{20} is the $\ell = 2, m_\ell = 0$ component of $Q^{(ij)a}$ for $a = 3$.

As shown in the above equation, the $M1$ amplitude for $\Delta^+ \rightarrow p\gamma$ is proportional to the matrix element of the $(J, I) = (1, 1)$ operator μ^{ia} between Δ^+ and p . In the $N_c \rightarrow \infty$ limit, the states Δ^+ and p are both contained in the lowest-lying baryon spin-flavor multiplet consisting of baryons with $J = I$. The isovector magnetic moment operator has a $1/N_c$ expansion in terms of

operator products of the baryon $SU(4)$ spin-flavor generators G^{ia} , I^a and J^i . Each independent operator polynomial arises at a definite order in the $1/N_c$ expansion which is determined by its explicit factor of $1/N_c$ and the implicit N_c -dependence of its operator matrix elements. The $1/N_c$ expansion of the magnetic moment operator is given by

$$\mu^{ia} = \mu_1 G^{ia} + \mu_2 \frac{J^i I^a}{N_c} + \mu_3 \frac{1}{N_c^2} \{J^2, G^{ia}\} + \dots \quad (11)$$

where the coefficients $\mu_1, \mu_2, \mu_3, \dots$ have an expansion in $1/N_c$ beginning at order unity, and the ellipsis represents higher-body operators. The complete set of higher-body operators are of the form $\mathcal{O}_{n+2} = \{J^2, \mathcal{O}_n\}$ where $\mathcal{O}_1 = G^{ia}$ and $\mathcal{O}_2 = J^i I^a$. At leading order in $1/N_c$, the $1/N_c$ expansion of the magnetic moment operator can be truncated to the single operator $\mu_1 G^{ia}$. This truncation of the $1/N_c$ expansion yields the spin-flavor symmetry relation between the $\Delta^+ \rightarrow p\gamma$ transition moment and the isovector combination of the p and n magnetic moments [11],

$$\mu_{\Delta+p} = \frac{1}{\sqrt{2}} (\mu_p - \mu_n). \quad (12)$$

The matrix elements of the spin-flavor generator G^{ia} are of order N_c , so the transition magnetic moment matrix element for $\Delta^+ \rightarrow p\gamma$ is order N_c [18]. This power counting is in accordance with the $I = J$ rule [12, 13]. Note that the $M1$ amplitude is only order $\sqrt{N_c}$ when ignoring an N_c -counting for the electromagnetic coupling e , since the kinematic factor $\sqrt{k_\gamma}$ in the definition Eq. (10) gives a suppression factor of $\sqrt{1/N_c}$. The leading $1/N_c$ corrections to Eq. (12) are produced by the $J^i I^a/N_c$ and $\{J^2, G^{ia}\}/N_c^2$ operators which both yield corrections of order $1/N_c^2$ relative to the leading order N_c contribution since the matrix elements of the baryon spin-flavor generators obey the N_c power counting $G \sim N_c$, $I \sim 1$ and $J \sim 1$.

Since the $1/N_c$ expansion of the isovector magnetic moment is given by a unique operator G^{ia} to relative order $1/N_c^2$ and, as we will show, the $E2$ amplitude is order $1/N_c^2$ relative to the $M1$ amplitude, the ratio of the helicity amplitudes $A_{3/2}/A_{1/2}$ is predicted at leading order in the $1/N_c$ expansion. The value of the prediction is given in Eq. (4). It is straightforward to show that all higher order terms in the $1/N_c$ expansion of the $M1$ amplitude Eq. (11) do not affect the ratio $A_{3/2}/A_{1/2}$, although they do violate the $SU(4)$ symmetry relation Eq. (12).

The $E2$ amplitude for $\Delta^+ \rightarrow p\gamma$ is proportional to the matrix element of the $(J, I) = (2, 1)$ operator $Q^{(ij)a} k_\gamma^j$ between Δ^+ and p . The $E2$ amplitude has one additional power of the photon momentum k_γ relative to the $M1$ amplitude. Since the mass difference between Δ^+ and p is order $1/N_c$, k_γ is order $1/N_c$. This kinematic suppression of $E2/M1$ by one power of $1/N_c$ is not sufficient to explain the smallness of the measured ratio,

Eq. (7). However, there is an additional $1/N_c$ suppression beyond the kinematic suppression due to k_γ . The electric quadrupole amplitude is proportional to the matrix element of the $(J, I) = (2, 1)$ operator $Q^{(ij)a}$ between Δ^+ and p . The quadrupole moment operator has the $1/N_c$ expansion

$$Q^{(ij)a} = c_1 (q_u - q_d) \frac{\{G^{ia}, J^j\}}{N_c} + \dots \quad (13)$$

where the operator $\{G^{ia}, J^j\}/N_c$ is understood to have its trace subtracted, c_1 is an operator coefficient of order unity, and q_u and q_d are the u and d quark charges. The electric quadrupole operator $Q^{(ij)a}$ has matrix elements of order unity at leading order, since the matrix elements of the baryon spin-flavor generators obey the N_c power counting $G \sim N_c$, $I \sim 1$ and $J \sim 1$. This $1/N_c$ suppression is in accordance with the suppression of $I \neq J$ operators by $(1/N_c)^{|J-I|}$, as derived in [13], since $|J - I| = 1$ for the quadrupole operator. Thus $E2$ is order $(1/N_c)^{3/2}$ due to the factor of $(1/N_c)^{3/2}$ from $k_\gamma^{3/2}$ in Eq. (10), and $E2/M1$ is of order $1/N_c^2$ since $M1$ is order $\sqrt{N_c}$.

Note that the isoscalar quadrupole operator has the expansion,

$$Q^{(ij)} = c_0 (q_u + q_d) \frac{\{J^i, J^j\}}{N_c} + \dots, \quad (14)$$

where c_0 is of order unity. Since the matrix element of G^{ia} is of order N_c and of J^i is of order unity, the ratio of the isoscalar to isovector quadrupole moments is of order $1/N_c$ times $(q_u + q_d)/(q_u - q_d)$. If one assumes that $q_u = 2/3$ and $q_d = -1/3$, then $(q_u + q_d)/(q_u - q_d) = 1/3$ and the ratio of the the isoscalar to isovector quadrupole moments is order $1/N_c$. If, as in Ref. [14], one chooses $q_u = (N_c + 1)/(2N_c)$, and $q_d = (1 - N_c)/(2N_c)$, then $(q_u + q_d)/(q_u - q_d) = 1/N_c$ and the ratio of the isoscalar to isovector quadrupole moments is order $1/N_c^2$, in agreement with Ref. [14].

One concrete realization of the large- N_c spin-flavor symmetry of baryons in QCD is given by the Skyrme model. Thus, the Skyrme model can be used as an

explicit check of the general, model-independent large- N_c results. For instance, the Skyrme model predicts that the transition magnetic moment $\mu_{\Delta N} = \langle \mu \rangle_{I=1}/\sqrt{2}$, where the isovector magnetic moment is proportional to $f_\pi^{-1}e^{-3}$, and e is the coefficient of the Skyrme term. Since $f_\pi \sim \sqrt{N_c}$ and $e \sim 1/\sqrt{N_c}$, the transition magnetic moment scales like N_c [15]. Likewise, the Skyrme model predicts the transition quadrupole moment $Q_{\Delta N} = (\sqrt{2}/10)\langle r^2 \rangle_{I=1}$. Although the nucleon isovector charge radius $\langle r^2 \rangle_{I=1}$ diverges in the chiral limit, it is independent of N_c in the large- N_c limit. Therefore $Q_{\Delta N}$ is of order N_c^0 [16].

One can also apply the large- N_c counting rules to the charge radii. For the isovector charge radii, we have the expansion

$$\langle r^2 \rangle_{I=1}^a = a_1 (q_u - q_d) I^a + \dots, \quad (15)$$

where a_1 is a coefficient of order unity. Thus $\langle r^2 \rangle_{I=1}$ is of order N_c^0 , consistent with the Skyrme model prediction. For the isoscalar charge radii,

$$\langle r^2 \rangle_{I=0} = a_0 (q_u + q_d) N_c + \dots, \quad (16)$$

where a_0 is a coefficient of order unity. The ratio of the isoscalar to isovector charge radii is of order $N_c(q_u + q_d)/(q_u - q_d)$, in agreement with Ref. [17]. The above results also are consistent with the picture that the baryon radius is of order N_c^0 in the large- N_c limit. The charge radius $\langle r^2 \rangle$ for a quark is of order unity. The isoscalar charge radius is the sum of $\langle r^2 \rangle$ over N_c quarks, and is of order N_c , whereas the isovector charge radius is the sum of $\langle r^2 \rangle$ over the $(N_c + 1)/2$ u -quarks minus the sum of $\langle r^2 \rangle$ over the $(N_c - 1)/2$ d -quarks. The order N_c pieces cancels in the difference, so the isovector charge radius is order one.

Acknowledgments

E.J and A.M. were supported in part by the Department of Energy under grant DOE-FG03-97ER40546. X. J. thanks T. Cohen for useful discussions, and is supported in part by the DOE grant DOE-FG02-93ER40762.

-
- [1] For a recent review, see B. Filippone and X. Ji, *Adv. in Nucl. Phys.* **26**, 1 (2001).
 - [2] E. Jenkins, *Ann. Rev. Nucl. Part. Sci.* **48**, 81 (1998)
 - [3] A.V. Manohar, Large N_c QCD in Les Houches Session LXVIII, Probing the Standard Model of Particle Interactions, ed. F. David and R. Gupta, Amsterdam, Elsevier, 1998.
 - [4] R. Dashen and A.V. Manohar, *Phys. Lett. B* **315**, 425 (1993); **315**, 438 (1993); E. Jenkins, *Phys. Lett. B* **315**, 431 (1993).
 - [5] R. Dashen, E. Jenkins and A.V. Manohar, *Phys. Rev. D* **49**, 4713 (1994); **51**, 3697 (1995).
 - [6] Particle Data Group, D.E. Groom *et al*, *European Physical Journal C* **15**, 1 (2000).
 - [7] M.A. Luty and J. March-Russell, *Nucl. Phys. B* **426**, 71 (1994).
 - [8] C. Carone, H. Georgi and S. Osofsky, *Phys. Lett. B* **322**, 227 (1994).
 - [9] E. Jenkins and A.V. Manohar, *Phys. Lett. B* **255**, 558 (1991); *Phys. Lett. B* **259**, 353 (1991).
 - [10] S-L. Zhu, S. Puglia and M.J. Ramsey-Musolf, *Phys. Rev. D* **63**:034002 (2001).
 - [11] E. Jenkins and A.V. Manohar, *Phys. Lett. B* **335**, 452 (1994).

- [12] M.P. Mattis and M. Mukerjee, Phys. Rev. Lett. **61**, 1344 (1988); M.P. Mattis, Phys. Rev. D **39**, 994 (1989); M.P. Mattis and E. Braaten, *ibid.* **39**, 2737 (1989).
- [13] R.F. Dashen, E. Jenkins and A.V. Manohar, Phys. Rev. D **51**, 3697 (1995).
- [14] A.J. Buchmann, J.A. Hester and R.F. Lebed, hep-ph/0205108
- [15] G.S. Adkins, C.R. Nappi, and E. Witten, Nucl. Phys. B **228**, 552 (1983).
- [16] G.S. Adkins and C.R. Nappi, Nucl. Phys. B **249**, 507 (1985); T. D. Cohen and W. Broniowski, Phys. Rev. D **34**, 3472 (1986).
- [17] A.J. Buchmann and R.F. Lebed, Phys. Rev. D **62**, 096005 (2000).
- [18] The gyromagnetic ratio, the magnetic moment in units of $1/(2M_N)$, is of order N_c^2 since the nucleon mass M_N is of order N_c .