

# Polarized Parton Distributions in the Valon Model

S. Atashbar Tehrani<sup>\*(a)</sup> , Ali N. Khorramian<sup>†(b,c)</sup>

<sup>(a)</sup>Physics Department, Persian Gulf University 75168, Boushehr, Iran

<sup>(b)</sup> Institute for Studies in Theoretical Physics and Mathematics (IPM)

P.O.Box 19395-5531, Tehran, Iran

<sup>(c)</sup> Physics Department, Semnan University, Semnan, Iran

November 4, 2018

## Abstract

We calculated the polarized parton distributions in leading order approximation in QCD from valon model. Polarized structure function of the nucleon are analyzed in the polarized valon model in which a nucleon is assumed to be a bound state of three polarized valence quark cluster(valons). It is shown that the results of the model calculation agree rather well with the SMC and available experimental data . Our results for  $\Gamma_1$  of hadron and deuteron are in good agreement with experimental data for some value of  $Q^2$ .

## 1 INTRODUCTION

The idea of quark cluster is not new. In valon model, the hadron is envisaged as a bound state of valence quark cluster called valon [1, 2]. For example, the bound state of

---

\*atashbar@cic.aut.ac.ir

†khorramiana@theory.ipm.ac.ir

proton consists of two “up” and one “down” valon. These valons thus bear the quantum numbers of respective valence quarks. Hwa [1] found evidence in the deep inelastic neutrino scattering data that suggested their existence and applied it to a variety of phenomena. Hwa [2] had also successfully formulated a treatment of the low- $p_T$  reactions based on the structural analysis of the valons. Arash [3] has recently applied the valon model to study the contributions of the constituents of the proton to its spin in deep inelastic scattering (DIS). Results of the model calculation agree well with the European Muon Collaboration (EMC) results [4]. Arash adopt the valon point of view and calculate various contributions to the spin of the proton.

The valon description of the nucleon was developed by Hwa [5, 6] some years ago and was successfully used [3] to describe the contribution of constituents of the proton to its spin in order to understand the so-called *spin crisis*. In this paper, we investigate the calculation of polarized parton density of the nucleon using an improved model and apply the model to the study of the spin of nucleon. In Sec. 2, polarized valon distributions are derived. In this section we calculate polarized valon distributions as a function of  $y$  from unpolarized valon distributions. The moment of polarized parton distributions and using inverse Mellin transformation are discussed in Sec.3. In Sec. 4, we examine the polarized structure function,  $xg_1$ , for proton, neutron and deuteron and compare with the experimental data. Also our prediction for the first moment of these structure functions presents in this section too.

## 2 UNPOLARIZED AND POLARIZED VALON DISTRIBUTIONS

The polarized valon is defined to be a dressed polarized valence quark in QCD with the cloud of polarized gluons and sea quarks which can be resolved by high  $Q^2$  probes. In a scattering process the virtual emission and absorption of gluons in a valon become

bremsstrahlung and pair creation. The polarized structure function of a valon is determined by gluon bremsstrahlung and pair creation in QCD. At sufficiently low value of  $Q^2$  the internal polarized structure of a valon cannot be resolved and hence it behaves as a polarized valence quark.

To facilitate the phenomenological analysis Hwa [5] assumed a simple form for the exclusive valon distribution in unpolarized proton which is

$$G_{UUD/p}(y_1, y_2, y_3) = g_p (y_1 y_2)^\alpha y_3^\beta \delta(y_1 + y_2 + y_3 - 1) \quad (1)$$

$y_i$  is the momentum fraction of the  $i$ th valon, and  $g_p$  is

$$g_p = [B(\alpha + 1, \beta + 1)B(\alpha + 1, \alpha + \beta + 2)]^{-1}$$

where  $B(m, n)$  is the beta function. The valon distributions are obtained by integration

$$\begin{aligned} G_{U/p}(y) &= \int dy_2 \int dy_3 G_{UUD/p}(y, y_2, y_3) \\ &= g_p B(\alpha + 1, \beta + 1) y^\alpha (1 - y)^{\alpha + \beta + 1} \end{aligned} \quad (2)$$

$$\begin{aligned} G_{D/p}(y) &= \int dy_1 \int dy_2 G_{UUD/p}(y_1, y_2, y) \\ &= g_p B(\alpha + 1, \alpha + 1) y^\beta (1 - y)^{2\alpha + 1} \end{aligned} \quad (3)$$

Recently, R.C. Hwa [7] have recalculated the unpolarized valon distribution in proton with a new set of parameters. The new values of  $\alpha$ ,  $\beta$  are found to be  $\alpha = 1.76$  and  $\beta = 1.05$ . An alternative analysis of polarized structure function has been performed by Gehrmann and Stirling[8]. This was done in the same spirit as the unpolarized analysis [9], namely the polarized valon distribution chose to be of the general form

$$\delta G_{U/p}(y) = \delta F_U(y) \times G_{U/p}(y)$$

$$\delta G_{D/p}(y) = \delta F_D(y) \times G_{D/p}(y) \quad (4)$$

where  $\delta F_j(y)$  is define:

$$\delta F_j(y) = N_j y^{\alpha_j} (1-y)^{\beta_j} (1 + \gamma_j y + \delta_j y^{0.5}) \quad (5)$$

where the subscript  $j$  refer to  $U, D$ , and  $G_{\frac{v}{p}}(y)$  is defined in Eqs. (2,3). In order to write down the polarized valon distribution in a proton we assume that at some  $Q = Q_0$  a valon behaves as a polarized valence quark. The polarized quark distribution in a proton at a starting scale of  $Q_0^2 = 0.26 GeV^2$  is parameterized by GRV group [10]. We take that parameterization and extract all of the parameters as defined in Eq.(5). The corresponding parameters are given in Table I. In Figure (1) we present unpolarized and polarized valon distributions as a function of  $y$ .

### 3 POLARIZED PARTON DISTRIBUTIONS

We now go to the  $n$ -moment space and define the moment of polarized valon distribution  $\Delta M_{v/p}(n)$  as:

$$\Delta M_{v/p}(n) \equiv \int_0^1 y^{n-1} \delta G_{v/p}(y) dy \quad (6)$$

the subscription  $v$  stands for kind of  $U, D$  polarized valon. For the moments of polarized singlet and non singlet distributions we shall use, the leading order solutions of the renormalization group equation in QCD. They can be expressed entirely in terms of the evolution parameter  $s$

$$s = \ln \frac{\ln Q^2 / \Lambda^2}{\ln Q_0^2 / \Lambda^2} \quad (7)$$

where  $Q_0, \Lambda$  are scale parameters to be determined by experiments. From the theoretical standpoint both  $Q_0$  and  $\Lambda$ , depend on the order of the moments  $n$ . In this work we have assumed that they are independent of  $n$  and we fixed it by  $Q_0^2 = 0.26 GeV^2$  and  $\Lambda = 0.232 GeV$ .

The moments of singlet and non singlet polarized are

$$\Delta M^{NS}(n, Q^2) = \exp(-\delta d_{NS}^{(0)n} s) \quad (8)$$

$$\Delta M^S(n, Q^2) = \frac{1}{2}(1 + \delta\rho) \exp(-\delta d_+ s) + \frac{1}{2}(1 - \delta\rho) \exp(-\delta d_- s) \quad (9)$$

where  $\delta\rho$  and other associated parameters are as follows:

$$\begin{aligned} \delta\rho &= \frac{\delta d_{NS}^{(0)n} - \delta d_{gg}^{(0)n} + 4f\delta d_{gq}^{(0)n}}{\Delta} \\ \Delta &= \delta d_+ - \delta d_- = \sqrt{(\delta d_{NS}^{(0)n} - \delta d_{gg}^{(0)n})^2 + 8f\delta d_{gq}^{(0)n}\delta d_{gq}^{(0)n}} \\ \delta d_{\pm} &= \frac{1}{2}(\delta d_{NS}^{(0)n} + \delta d_{gg}^{(0)n} \pm \Delta) \\ b &= \frac{33 - 2f}{12\pi} \end{aligned} \quad (10)$$

and also the anomalous dimensions  $\delta d_{ij}^{(0)n}$  are simply the n-th moment of polarized LO splitting function are given by [11] and  $f$  is the number of active flavors.

$$\begin{aligned} \delta d_{NS}^{(0)n} &= \frac{1}{3\pi b} \left[ -3 - \frac{2}{n(n+1)} + 4S_1(n) \right] \\ \delta d_{gg}^{(0)n} &= \frac{-1}{4\pi b} \frac{n-1}{n(n+1)} \\ \delta d_{gq}^{(0)n} &= \frac{-2}{\pi b} \frac{1}{3} \frac{n+2}{n(n+1)} \end{aligned}$$

$$\delta d_{gq}^{(0)n} = -\frac{1}{2\pi b} \left\{ 3 \left[ \frac{11}{6} + \frac{4}{n(n+1)} - 2S_1(n) \right] - \frac{2f}{3} \right\}. \quad (11)$$

Here  $S_1(n) = \sum_{j=1}^n \frac{1}{j} = \psi(n+1) + \gamma_E$ ,  $\psi(n) = \frac{\Gamma'(n)}{\Gamma(n)}$  and  $\gamma_E = 0.577216$ .

The moment of polarized  $u$  and  $d$ -valence quark in a proton are:

$$\begin{aligned} \Delta M_{u_v}(n, s) &= 2\Delta M^{NS}(n, s)\Delta M_{U/p} \\ \Delta M_{d_v}(n, s) &= \Delta M^{NS}(n, s)\Delta M_{D/p} \end{aligned} \quad (12)$$

and the moment of polarized  $\Sigma$  and gluon distributions are as follows:

$$\Delta M_{\Sigma}(n, s) = \Delta M^s(n, s)(2\Delta M_{U/p} + \Delta M_{D/p})$$

$$\Delta M_g(n, s) = \Delta M_{gq}(n, s)(2\Delta M_{U/p} + \Delta M_{D/p}) \quad (13)$$

where  $\Delta M_{gq}(n, s)$  is the quark-to-gluon evolution function give by:

$$\begin{aligned} \Delta M_{gq}(n, s) = & \frac{(\delta d_+ - \delta d_{NS}^{(0)n})}{\Delta} \left(1 - \frac{\delta d_- - \delta d_{NS}^{(0)n}}{2f\delta d_{qg}^{(0)n}}\right) \exp -d_+ s \\ & - \frac{(\delta d_- - \delta d_{NS}^{(0)n})}{\Delta} \left(1 - \frac{\delta d_+ - \delta d_{NS}^{(0)n}}{2f\delta d_{qg}^{(0)n}}\right) \exp -d_- s \end{aligned} \quad (14)$$

In Figure (2) we present these moments as a function of  $n$  for some values of  $Q^2$ . To obtain the  $x$ -dependence of structure functions and parton distributions, usually required for practical purposes, from the above  $n$ -dependent exact analytical solutions in Mellin-moment space, one has to perform a numerical integral in order to invert the Mellin-transformation in (8) according to

$$\delta f(x, Q^2) = \frac{1}{\pi} \int_0^\infty dz \text{Im}[e^{i\phi} x^{-c-ze^{i\phi}} \Delta f^{n=c+ze^{i\phi}}(Q^2)] \quad (15)$$

where the contour of integration, and thus the value of  $c$ , has to lie to the right of all singularities of  $\Delta f^n(Q^2)$  in the complex  $n$ -plane, i.e.,  $c \geq 0$  since according to Eq. (13) the dominant pole of all  $\delta d_{ij}^n$  is located at  $n = 0$ . For all practical purpose one may choose  $c \simeq 1, \phi = 135^\circ$  and an upper limit of integration, for any  $Q^2$ , of about  $5 + 10/\ln x^{-1}$ , instead of  $\infty$ , which suffices guarantee and stable numerical results [12, 13]. In figure (3) our predictions for  $x$ -dependence spin densities for quarks and gluons at  $Q^2 = 2, 3, 5, 10 \text{ GeV}^2$  in a proton are shown.

Furthermore, in the polarized parton distribution we assume an unbroken  $SU(3)_f$  symmetric sea,

$$\delta \bar{q}(x, Q^2) \equiv \delta \bar{u} = \delta u_{sea} = \delta \bar{d} = \delta d_{sea} = \delta \bar{s} = \delta s_{sea} \quad (16)$$

Note that the proton include partons with positive and negative helicity which carry of fraction  $x$  of the proton momentum and to whom one can associate quark densities  $q_+(x, Q^2)$  and  $q_-(x, Q^2)$ . The difference

$$\delta q(x, Q^2) = q_+(x, Q^2) - q_-(x, Q^2) \quad (17)$$

measures how much the parton of flavor  $q$  “remembers” of its parent parton polarization. Similarly, one may define

$$\delta \bar{q}(x, Q^2) = \bar{q}_+(x, Q^2) - \bar{q}_-(x, Q^2) \quad (18)$$

for antiquarks.

## 4 POLARIZED STRUCTURE FUNCTION, $g_1$

We used polarized parton distribution from valon model for extracting structure function  $g_1$ . Within this model according Sec. (3), we calculated all of the polarized parton distributions in a proton. Now it is very easy to used these results for calculating  $g_1^p$ . In the quark model, to leading order (LO) in QCD,  $g_1^p$  can be written as a linear combination of  $\delta q$  and  $\delta \bar{q}$  [11, 14],

$$g_1^p(x, Q^2) = \frac{1}{2} \sum_q e_q^2 [\delta q(x, Q^2) + \delta \bar{q}(x, Q^2)] \quad (19)$$

where  $e_q$  are the electric charges of the (light) quark-flavors  $q = u, d, s$ .

Furthermore, Eq. (19) can be decomposed into a flavor nonsinglet (NS) and singlet (S) component

$$g_1^p(x, Q^2) = g_{1,NS}(x, Q^2) + g_{1,S}(x, Q^2) \quad (20)$$

where

$$g_{1,NS}(x, Q^2) = \frac{1}{2} \sum_q (e_q^2 - \langle e^2 \rangle) (\delta q + \delta \bar{q}) \quad (21)$$

with  $\langle e^2 \rangle = \frac{1}{f} \sum_q e_q^2$  (e.g. for  $f=3$  light  $u, d, s$  flavors  $\langle e^2 \rangle = \frac{2}{9}$ ) and

$$g_{1,S}(x, Q^2) = \frac{1}{2} \langle e^2 \rangle \sum_q (\delta q + \delta \bar{q}) \equiv \frac{1}{2} \langle e^2 \rangle \delta \Sigma. \quad (22)$$

where the sum usually runs over the light quark-flavors  $q = u, d, s$ , since the heavy quark contributions (c, b,...) have preferably to be calculated perturbatively from the intrinsic light quark ( $u, d, s$ ) and gluon ( $g$ ) partonic-constituent of the nucleon.

We are now in a position to present the results for the proton polarized structure function,  $g_1^p$ . In Eq. (19) we presented polarized proton structure function in a proton. In figure (4) the results are shown at some values of  $Q^2$ . The data points are from [15]. For the neutron structure function one only needs to interchange  $u_v$  and  $d_v$  in the Eq. (19). For deuteron it is given by:

$$g_1^d(x, Q^2) = \frac{1}{2} [g_1^p(x, Q^2) + g_1^n(x, Q^2)] (1 - \frac{3}{2} \omega_D) \quad (23)$$

where  $\omega_D \approx 0.058$  accounts for the D-state admixture in the deuteron wave function. Quantities of particular significance are the first moment ( $n = 1$ ) of polarized parton distributions  $\delta f(x, Q^2)$ ,

$$\Delta f(Q^2) \equiv \int_0^1 dx \delta f(x, Q^2) \quad , \quad f = q, \bar{q}, g \quad (24)$$

We calculated first moments (total polarizations)  $\Delta f(Q^2)$  of the polarized parton distributions. These results shown in Table II.

According to Eqs. (17) and (18),  $\Delta(q + \bar{q})$  is the net number of right-handed quarks of flavor  $q$  inside a right-handed proton and thus  $\frac{1}{2} \Delta \Sigma$ , is a measure of how much all quark flavors contribute to the spin of the proton. Similarly,  $\Delta g$  represents the total gluonic contribution to the spin of the nucleon.

Here we have defined the first moment of  $g_1$  (Ellis-Jaffe sum rule) by



$$\Gamma_1^{p,n,d}(Q^2) \equiv \int_0^1 dx g_1^{p,n,d}(x, Q^2)$$

Our values for  $\Gamma_1^{p,n,d}$  represents in Table III, which are in excellent agreement with the E143, E142, E154, SMC results from Ref. [16-19].

Figure (5) shows our results for  $xg_1^n$  at  $Q^2 = 2, 3, 5 \text{ GeV}^2$  which are in agreement with the results of experimental data from [15, 17, 18].

In figure (6) we have shown  $xg_1^d$  for some values of  $Q^2$ . The experimental data are from [15].

Finally by this model it is possible to attribute numbers  $\Delta\Sigma$  and  $\Delta g$  to the quark and gluon spin content of the nucleons which describe their total (integrated) contribution to the nucleon spin in the following sense

$$\frac{1}{2} = \frac{1}{2}\Delta\Sigma(Q^2) + \Delta g(Q^2) + L_z(Q^2) \quad (25)$$

where on the left hand side we have the spin ( $+\frac{1}{2}$ ) of a polarized nucleon state and on the right hand side a decomposition in terms of  $\Delta\Sigma(Q^2) = \Delta u + \Delta d + \Delta s + \Delta\bar{u} + \dots$ ,  $\Delta g(Q^2)$  and the relative orbital angular momentum  $L_z(Q^2)$  among all the quarks and gluons. Unfortunately, the decomposition in Eq. (25) cannot directly be measured in experiments. Instead various other combinations of  $\Delta\Sigma$  and  $\Delta g$  appear in experimental observables. Our calculation for study of nucleon spin presented in Fig. (7).

## 5 CONCLUSION

We have presented a model in which the polarized parton density is calculated. In this paper polarized valon distribution is derived from unpolarized valon distribution. After define the moment of polarized valon distribution and moments of all partons in proton, polarized parton density in a hadron are available. The results are used to evaluate the spin components of nucleon. It turns out that are results about polarized structure function

agree well with all available experimental data on  $g_1$  of proton ,neutron and deuteron.

Also our prediction for  $\Gamma_1$  is good agreement with experimental data.

## References

- [1] R. C. Hwa, Phys. Rev. D **22**, 759 (1980).
- [2] R. C. Hwa, Phys. Rev. D **22**, 1593 (1980).
- [3] Firooz Arash, Phys. Rev. D **50**, 1946 (1994).
- [4] J. Ashman *et al.*, Phys. Lett. B **206**, 364 (1988); Nucl. Phys. B**328**, 1 (1989).
- [5] R. C. Hwa and M. S. Zahir, Phys. Rev. D **23**, 2539 (1981).
- [6] R. C. Hwa and C. B. Yang, hep-ph/0204289.
- [7] R. C. Hwa and C. B. Yang, hep-ph/0202140 (submitted to Phys. Rev. C).
- [8] T. Gehrmann and W. J. Stirling, Phys. Rev. D**53**, 6100 (1996).
- [9] A. D. Martin, R. G. Roberts and W. J. Stirling, Phys. Lett. B**354**, 155 (1995).
- [10] M. Glück, E. Reya, M. Stratmann and W. Vogelsang Phys. Rev. D**63**, 094005 (2001).
- [11] G. Altarelli and Parisi, Nucl. Phys. B**126**, 298 (1977); B. Lampe and E. Reya, Phys.Rept. **332**,1 (2000).
- [12] M. Glück, E. Reya and A. Vogt, Z. Phys. C**48**, 471 (1991).
- [13] M. Glück, E. Reya and A. Vogt, Phys. Rev. D**45**, 3986 (1992).
- [14] G. Altarelli, Phys. Rep. **81**, 1 (1982)
- [15] E143 Collaboration, K. Abe *et al.*, Phys. Rev. D**58**, 112003 (1998); Spin Muon Collaboration, B. Adeva *et al.*, Phys.Rev. D**58**, 112001 (1998).
- [16] E143 Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **74**, 346 (1995)
- [17] E142 Collaboration, P. L. Anthony *et al.*, Phys. Rev. D**54**, 6620 (1996)

[18] E154 Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **79**, 26 (1997)

[19] Spin Muon Collaboration, D. Adams *et al.*, Phys. Rev. **D56**, 5330 (1997).

**Table Caption**

$v$	$N$	$\alpha$	$\beta$	$\gamma$	$\eta$
$U$	0.0475	-1.003	-0.998	0.653	-0.067
$D$	-0.015	-1.247	-0.951	8.351	-2.063

**Table I.** The parameters defined in Eqs. (4,5) for  $U$  and  $D$  polarized valon distribution.

$Q^2(GeV^2)$	$\Delta u_v$	$\Delta d_v$	$\Delta\Sigma$	$\Delta\bar{q}$	$\Delta g$
2	0.609	-0.306	0.329	0.004	0.876
3	0.609	-0.306	0.334	0.005	0.992
5	0.609	-0.306	0.341	0.006	1.139
10	0.609	-0.306	0.350	0.008	1.334

**Table II.** The first moments of polarized parton distributions,  $\Delta u_v$ ,  $\Delta d_v$ ,  $\Delta\Sigma$ ,  $\Delta\bar{q}$  and  $\Delta g$  for some value of  $Q^2$ .

$Q^2(GeV^2)$	$\Gamma_1^p$	$\Gamma_1^n$	$\Gamma_1^d$
2	0.1212	-0.0313	0.0412
3	0.1219	-0.0307	0.0417
5	0.1225	-0.0299	0.0424
10	0.1236	-0.0289	0.0433

**Table III.** The first moments of proton, neutron, and deuteron polarized structure function for  $Q^2 = 2, 3, 5, 10 GeV^2$ .




Figure 1: The unpolarized and polarized valon distributions as a function of  $y$ .




Figure 2: Moments of polarized parton distributions as a function of  $n$  for some values of  $Q^2$ .




Figure 3: Polarized parton distributions in proton at  $Q^2 = 2, 3, 5, 10\text{GeV}^2$  as a function of  $x$ .




Figure 4: The compares of our predictions for  $xg_1^p(x, Q^2)$  versus  $x$  from valon model with the measured structure functions in experiments on proton target. Data points are from [15].




Figure 5: Plots of neutron polarized structure function against  $x$  at  $Q^2 = 2, 3, 5 \text{ GeV}^2$ . Data points are from [15, 17, 18].




Figure 6: Plots of deuteron polarized structure function against  $x$  at  $Q^2 = 2, 3, 5$  and  $10 \text{ GeV}^2$ . Data points are from [15].






Figure 7: The variation of  $\frac{1}{2}\Delta\Sigma$ ,  $\Delta g$  and  $L_z$  as a function of  $Q^2$ .