

Decays $B \rightarrow \eta' K$ in R -parity violating supersymmetry

B. Dutta^{a*}, C. S. Kim^{b†} and Sechul Oh^{b‡}

^aCenter For Theoretical Physics, Department of Physics, Texas A&M University, College Station, TX 77843-4242, USA

^bDepartment of Physics and IPAP, Yonsei University, Seoul 120-479, Korea

In light of the recent experimental data from B factories, We try to explain the large branching ratio (compared to the Standard Model prediction) of the decay $B^\pm \rightarrow \eta' K^\pm$ in the context of R -parity violating (\mathcal{R}_p) supersymmetry. We investigate other observed $\eta^{(\prime)}$ modes and find that only two pairs of \mathcal{R}_p coupling can satisfy the requirements without affecting the other $B \rightarrow PP$ and $B \rightarrow VP$ decay modes except the mode $B \rightarrow \phi K$. We also calculate the CP asymmetry for the observed decay modes affected by the new couplings.

1. Introduction

Among the $B \rightarrow PP$ (P denotes a pseudoscalar meson) decay modes, the branching ratio for the decay $B^\pm \rightarrow \eta' K^\pm$ is observed to be still larger than that expected within the Standard Model (SM). The SM contribution is about 3σ smaller than the experimental world average (Fig.1). Among the $B \rightarrow VP$ (V denotes a vector meson) decay modes, the decay $B^\pm \rightarrow \phi K^\pm$ has been observed recently and the experimentally observed BR for the decay $B^0 \rightarrow \eta K^{*0}$ has been found to be 2σ larger than the SM. The BR for the newly observed decay $B^\pm \rightarrow \eta K^{*\pm}$ is also available.

In this work [1], we address these large BR problems of $B^{\pm(0)} \rightarrow \eta^{(\prime)} K^{\pm(*0)}$ systems using R -parity violating (\mathcal{R}_p) supersymmetric (SUSY) theories. The effects of \mathcal{R}_p couplings on B decays have been investigated previously in the literature[2]. Attempts were made to fit just the large BR for $B^\pm \rightarrow \eta' K^\pm$ [2]. At present, we have more results. Some of these results are con-

cerned with decay modes involving $\eta^{(\prime)}$ and these modes are influenced by the same \mathcal{R}_p coupling that affects $B^\pm \rightarrow \eta' K^\pm$. For example, the decay modes $B^\pm \rightarrow \eta K^{*\pm}$, $B^0 \rightarrow \eta K^{*0}$, $B^0 \rightarrow \eta' K^0$ are affected by the new couplings which cure the large BR problem of $B^\pm \rightarrow \eta' K^\pm$. Hence, it is natural to investigate these newly observed decay modes and try to see whether all the available data can be explained. We also need to be concerned about the other observed (not involving $\eta^{(\prime)}$) $B \rightarrow PP$ and $B \rightarrow VP$ decay modes, which could be influenced by these new couplings. Our effort is not to affect the other modes as much as possible, since except for $B \rightarrow \eta^{(\prime)} K^{(*)}$ decay modes, the other observed modes fit the available data well[3,4]. Further, using the preferred values of different parameters (e.g., new couplings etc.), we also make predictions for CP asymmetry for these observed modes which will be verified in the near future.

2. \mathcal{R}_p SUSY effects to decay amplitudes

The \mathcal{R}_p part of the amplitude of $B^\pm \rightarrow \eta' K^\pm$ decay is

$$\begin{aligned} \mathcal{M}_{\eta' K}^{\lambda'} &= (d_{121}^R - d_{112}^L) \xi A_{\eta'}^u & (1) \\ &+ (d_{222}^L - d_{222}^R) \left[\frac{\tilde{m}}{m_s} (A_{\eta'}^s - A_{\eta'}^u) - \xi A_{\eta'}^s \right] \end{aligned}$$

*The work of B.D. was supported in part by National Science Foundation grant No. PHY-0070964.

†The work of C.S.K. was supported in part by CHEP-SRC Program, Grant No. 20015-111-02-2 and Grant No. R03-2001-00010 of the KOSEF, in part by BK21 Program and Grant No. 2001-042-D00022 of the KRF, and in part by Yonsei Research Fund, Project No. 2001-1-0057.

‡The work of S.O. was supported by BK21 Program and Grant No. 2001-042-D00022 of the KRF.

$$\begin{aligned}
& + (d_{121}^L - d_{112}^R) \frac{\bar{m}}{m_d} A_{\eta'}^u \\
& + u_{112}^R \left[\xi A_{\eta'}^u - \frac{2m_K^2 A_K}{(m_s + m_u)(m_b - m_u)} \right],
\end{aligned}$$

where

$$\begin{aligned}
d_{jkn}^R &= \sum_{i=1}^3 \frac{\lambda'_{ijk} \lambda_{in3}^*}{8m_{\nu_{iL}}^2}, \\
d_{jkn}^L &= \sum_{i=1}^3 \frac{\lambda'_{i3k} \lambda_{inj}^*}{8m_{\nu_{iL}}^2}, \quad (j, k, n = 1, 2) \\
u_{jkn}^R &= \sum_{i=1}^3 \frac{\lambda'_{ijn} \lambda_{ik3}^*}{8m_{e_{iL}}^2}, \quad (j, k = 1, n = 2). \quad (2)
\end{aligned}$$

Here $\xi \equiv 1/N_c$ (N_c denotes the effective number of color), $\bar{m} \equiv m_{\eta'}^2/(m_b - m_s)$ and

$$A_{M_1} = \langle M_2 | J_b^\mu | B \rangle \langle M_1 | J_{l\mu} | 0 \rangle.$$

J stands for quark currents and the subscripts b and l indicate whether the current involves a b quark or only the light quarks. Analogous expressions hold for $B^\pm \rightarrow \eta K^\pm$ where we have to replace $A_{\eta'}^u$ by A_η^u , $A_{\eta'}^s$ by A_η^s and $m_{\eta'}$ by m_η . Replacing a pseudoscalar meson by a vector meson, we also get similar expressions for the amplitudes of $B^{\pm(0)} \rightarrow \eta' K^{*\pm(0)}$ modes. The \mathcal{R}_p part of the amplitude of $B \rightarrow \phi K$ decay mode involves only d_{222}^L and d_{222}^R .

$$\mathcal{M}_{\phi K}^{\lambda'} = (d_{222}^L + d_{222}^R) [\xi A_\phi], \quad (3)$$

where $A_\phi = \langle K | J_b^\mu | B \rangle \langle \phi | J_{l\mu} | 0 \rangle$.

3. Results

We use the effective Hamiltonian and the effective Wilson coefficients given in Ref. [5].

We first try to explain the large branching ratio of $B^\pm \rightarrow \eta' K^\pm$. The observed BR for this mode in three different experiments are [6–8]

$$\begin{aligned}
\mathcal{B}(B^\pm \rightarrow \eta' K^\pm) \times 10^6 &= 80_{-9}^{+10} \pm 7 \text{ (CLEO)}, \\
70 \pm 8 \pm 5 \text{ (BaBar)}, \quad 79_{-11}^{+12} \pm 9 \text{ (Belle)}. \quad (4)
\end{aligned}$$

The three results are close and we use the world average of them: (in 10^{-6}) 75 ± 7 . The maximum BR in SM that we find is 42×10^{-6} (Fig. 1). In

the \mathcal{R}_p SUSY framework, we find that the positive values of d_{222}^R and negative values of d_{222}^L can increase the BR keeping most of the other $B \rightarrow PP$ and $B \rightarrow VP$ modes unaffected. The other \mathcal{R}_p combinations are either not enough to increase in the BR or affect too many other modes. (An important role is played by the λ'_{32i} -type couplings, the constraints on which are relatively weak.) We divide our results into two cases,

Case 1: we use only d_{222}^R (positive values) and **Case 2:** we use a combination of d_{222}^R (positive values) and d_{222}^L (negative values).




Figure 1. The BR for the decay $B^\pm \rightarrow \eta' K^\pm$ vs ξ . The solid line is for the SM. The dashed, dotted and dot-dashed lines correspond to $|\lambda'_{323}| = |\lambda'_{322}| = 0.04, 0.06, 0.08$, respectively. The bold solid lines indicate the experimental world average bound.

Let us start with **Case 1**. We first discuss the case of $\gamma = 110^\circ$. In Fig. 1, we plot the BR for the decay $B^\pm \rightarrow \eta' K^\pm$ as a function of ξ . We have used $|\lambda'_{323}| = |\lambda'_{322}| = 0.04, 0.06, 0.08$ and $m_{\text{susy}} = 200$ GeV. We take d_{222}^R to be positive. The large branching ratio can be explained for $\lambda' \geq 0.05$.

The $\mathcal{B}(B \rightarrow X_s \nu \nu)$ can put bound on $\lambda'_{322} \lambda'_{323}^*$ in certain limits: $\lambda' \leq 0.07$.

In another scenario we can use smaller value of γ , e.g., $\gamma = 80^\circ$ to fit the $B \rightarrow \eta^{(\prime)} K^{(*)}$ data. In Table I, we present the BRs and the CP asymmetries for $B \rightarrow \eta^{(\prime)} K^{(*)}$ and $B \rightarrow \phi K$ for different values of δ and γ . Here the phase difference between λ'_{323} and λ'_{322} , δ , is defined by

$$\lambda'_{323} \lambda'_{322}^* = |\lambda'_{323} \lambda'_{322}^*| e^{i\delta}. \quad (5)$$

Table 1

Case 1: for $|\lambda'| = 0.06$ and $\xi = 0.25$.

mode	$\delta = 0,$ $\gamma = 110^0,$		$\delta = 15^0,$ $\gamma = 110^0$		$\delta = 0,$ $\gamma = 80^0,$		$\delta = 55^0,$ $\gamma = 80^0$	
	$\mathcal{B} \times 10^6$	\mathcal{A}_{CP}	$\mathcal{B} \times 10^6$	\mathcal{A}_{CP}	$\mathcal{B} \times 10^6$	\mathcal{A}_{CP}	$\mathcal{B} \times 10^6$	\mathcal{A}_{CP}
$B^+ \rightarrow \eta' K^+$	68.9	0.01	68.3	0.04	82.1	0.01	68.3	0.11
$B^+ \rightarrow \eta K^{*+}$	36.4	0.03	36.4	0.04	36.5	0.03	32.7	0.09
$B^0 \rightarrow \eta' K^0$	88.3	0.00	86.8	0.03	110.2	0.00	87.1	0.12
$B^0 \rightarrow \eta K^{*0}$	14.0	-0.39	14.6	-0.42	14.8	-0.28	20.4	-0.56
$B^+ \rightarrow \phi K^+$	7.11	0.00	6.97	0.04	7.10	0.00	5.76	0.14

Table 2

Case 2: for $|\lambda'| = 0.052$ and $\xi = 0.45$.

mode	$\delta = 0$		$\delta = 20^0$	
	$\mathcal{B} \times 10^6$	\mathcal{A}_{CP}	$\mathcal{B} \times 10^6$	\mathcal{A}_{CP}
$B^+ \rightarrow \eta' K^+$	69.3	0.01	68.0	0.05
$B^+ \rightarrow \eta K^{*+}$	27.9	0.04	27.8	0.05
$B^0 \rightarrow \eta' K^0$	107.4	0.00	104.5	0.05
$B^0 \rightarrow \eta K^{*0}$	20.5	-0.71	21.1	-0.72
$B^+ \rightarrow \phi K^+$	6.56	0.00	6.56	0.00

The maximum values of δ allowed by the BR of $B^\pm \rightarrow \eta' K^\pm$ are $\delta = 15^0$ for $\gamma = 110^0$ and $\delta = 55^0$ for $\gamma = 80^0$.

Case 2: We now use the combination of d_{222}^R and d_{222}^L with $\gamma = 110^0$. We assume $d_{222}^R = -d_{222}^L$. In this scenario, the \mathcal{R}_p coupling part of the amplitude in $B \rightarrow \phi K$ decay mode canceled exactly (Eq. 3). (In fact, our solution still works when the cancellation is incomplete by about 5%.) But we still have contributions to $B^\pm \rightarrow \eta' K^\pm$ (Eq. 2) and to increase the BR we choose d_{222}^R to be positive. There is no \mathcal{R}_p contribution to the other $B \rightarrow PP$ and $B \rightarrow VP$ modes in this case as well.

In Table II, we calculate the BRs and the CP asymmetries for $B \rightarrow \eta^{(\prime)} K^{(*)}$ and $B \rightarrow \phi K$ for different values of δ . The maximum value of δ allowed by the BR for $B^\pm \rightarrow \eta' K^\pm$ is $\delta = 20^0$ for $\gamma = 110^0$.

4. Conclusion

We have studied $B \rightarrow \eta^{(\prime)} K^{(*)}$ modes in the context of \mathcal{R}_p SUSY theories. We have found solutions for both large and small values of $\gamma = 110^0, 80^0$ and two different values of $\xi \simeq 0.25, 0.45$ for two different scenarios. For our solutions, we need $|\lambda'| \sim 0.05 - 0.06$ for

$m_{\text{susy}} = 200$ GeV.

REFERENCES

1. B. Dutta, C.S. Kim and Sechul Oh, Phys. Lett. **B535**, 249 (2002).
2. D. Choudhury, B. Dutta and A. Kundu, Phys. Lett. **B456**, 185(1999), and references therein.
3. H.-Y. Cheng and K.-C. Yang, Phys. Rev. **D62**, 054029 (2000).
4. B. Dutta and S. Oh, Phys. Rev. **D63**, 054016 (2001).
5. N.G. Deshpande, B. Dutta and S. Oh, Phys. Rev. **D57**, 5723 (1998); Phys. Lett. **B473**, 141 (2000).
6. S.J. Richichi *et al.* (CLEO collaboration), Phys. Rev. Lett **85**, 520 (2000).
7. K. Abe *et al.* (Belle collaboration), Phys. Lett. **B517**, 309 (2000).
8. B. Aubert *et al.* (BaBar collaboration), Phys. Rev. Lett. **87**, 221802 (2001).