

# Hydrodynamics with spontaneous symmetry breaking: application to relativistic heavy ion collisions

Y. Lallouet, D. Davesne and C. Pujol

*IPN Lyon, 43 Bd du 11 Novembre 1918, F-69622 Villeurbanne Cedex*

(Dated: October 30, 2018)

In this paper we apply hydrodynamics for systems with continuous broken symmetries to heavy ion collisions in the framework of (1+1) dimensional Bjorken model. The temperature profile with respect to proper time determined in that context exhibits no differences with the ideal fluid. On the contrary, it is shown that the profile obtained when Müller-Israel-Stewart second order theory of dissipation is included on top of standard hydrodynamics indicates a slower cooling of the system.

## I. INTRODUCTION

Relativistic hydrodynamics is a very useful tool in the field of heavy ion collisions when one wants to reproduce the space-time evolution of the hot and dense matter created in such events. Upon the hypothesis of local equilibrium, which can be questionable for such small and not long-lived systems, this approach directly gives in terms of usual quantities such as temperature, pressure..., an intuitive description of what happens during the whole evolution of the system until the freeze-out takes place. Moreover, it is usually assumed that the matter under consideration is almost perfect, so that standard hydrodynamics can be applied directly. However, the hot and dense matter produced is mainly constituted of pions and, with the exception of [1], no attention has been drawn to the fact that the breaking of chiral symmetry can affect the theory of hydrodynamics itself. One of the aims of this paper is thus to give in a simple approach a quantitative estimate of this modification. More precisely, we will determine the temperature profile with respect to time in the framework of the (1+1) dimensional boost-invariant Bjorken model with and without the effect of the spontaneous breaking of the symmetry. In a second step we also introduce the dissipation process and compare the previous results with those obtained when Müller-Israel-Stewart second order theory of dissipation is included on top of standard hydrodynamics.

## II. HYDRODYNAMICS EQUATIONS FOR CHIRAL FLUID

Following [1], the hydrodynamics degrees of freedom in a chiral fluid with chiral  $SU(2)$  symmetry spontaneously broken are the densities associated to conserved quantities, namely entropy density  $s$ , momentum density  $T^{0i}$ , baryonic number density  $n$ , left and right-handed charge densities which can be written as  $SU(2)$ -matrices  $\rho_L \equiv \rho_L^i \tau_i / 2$  and  $\rho_R \equiv \rho_R^i \tau_i / 2$  and finally the variables associated to the Goldstone modes introduced in the theory *via* a  $SU(2)$ -matrix  $\Sigma \equiv e^{i\vec{\tau} \cdot \vec{\pi} / f_\pi}$ . If we focus on the

central rapidity region of a heavy ion collision, we can neglect in a first step the baryonic density. Thus, the hydrodynamic equations taken from [1] become:

$$\partial_\mu T^{\mu\nu} = 0 \quad (1)$$

$$i\partial_\mu ((f_t^2 - f_s^2) u^\mu u^\nu \Sigma \partial_\nu \Sigma^\dagger + f_s^2 \Sigma \partial^\mu \Sigma^\dagger) = 0 \quad (2)$$

where the energy-momentum tensor  $T^{\mu\nu}$  is given by  $T^{\mu\nu} = (\epsilon + p) u^\mu u^\nu - p g^{\mu\nu} + f_s^2 / 4 \text{tr}(\partial^\mu \Sigma \partial^\nu \Sigma^\dagger + \partial^\nu \Sigma \partial^\mu \Sigma^\dagger)$ .

In the above equations,  $f_t$  ( $f_s$ ), which can be interpreted as the temporal (spatial) pion decay constant, are actually not constants but have a non-trivial temperature dependence. Nevertheless, in order to have an estimation of the effects induced by the presence of the spontaneous breaking of the chiral symmetry, we will adopt two simple approaches. The first one, for the sake of simplicity, is to consider that  $f_t = f_s = f_\pi = 93$  MeV. Of course, at finite temperature, this is not true and, for instance, for small temperatures, one knows that  $f_t^2$  and  $f_s^2$  differ from  $f_\pi^2$  by a factor proportional to  $T^2$  while the correction for  $f_t^2 - f_s^2$  is of order  $T^4$  so that, to first order,  $f_t^2 = f_s^2 = f_\pi^2 (1 - T^2 / (6f_\pi^2))$ . This second approach, more realistic, is based on this formula, extended to larger temperatures.

## III. THE BJORKEN PICTURE

Even if several important improvements concerning the geometric description of a heavy ion collision have been done since the pioneering paper of Bjorken [4], we will use here this simplified version. The Bjorken picture of a collision is based on the hypothesis that initial conditions are invariant under Lorentz transformations throughout the central rapidity plateau. This implies that, for a one-dimensional flow in the  $z$ -direction, all the quantities depend actually not on the time  $t$  and the position  $z$  but only on the proper time  $\tau = \sqrt{t^2 - z^2}$ . The space-time evolution of the system can be then represented by hyperbola  $\tau = cst$ . If the hydrodynamic velocity  $u^\mu$  is

taken as  $u^\mu = x^\mu/\tau = (t/\tau, 0, 0, z/\tau)$ , the hydrodynamic equations

$$\partial_\mu T^{\mu\nu} = 0 \quad (3)$$

with  $T^{\mu\nu} = (\epsilon+p)u^\mu u^\nu - pg^{\mu\nu}$  lead for an ultra relativistic gas to the following differential equations [5, 6]:

$$\frac{\partial s}{\partial \tau} + \frac{s}{\tau} = 0 \quad (4)$$

$$\frac{\partial T}{\partial \tau} + \frac{1}{3} \frac{T}{\tau} = 0 \quad (5)$$

$$\frac{\partial \epsilon}{\partial \tau} + \frac{4}{3} \frac{\epsilon}{\tau} = 0 \quad (6)$$

whose solutions give the following scaling behavior:

$$s(\tau) = s(\tau_0) \frac{\tau_0}{\tau} \quad (7)$$

$$T(\tau) = T(\tau_0) \left( \frac{\tau_0}{\tau} \right)^{1/3} \quad (8)$$

$$\epsilon(\tau) = \epsilon(\tau_0) \left( \frac{\tau_0}{\tau} \right)^{4/3}. \quad (9)$$

And, since  $p = \epsilon/3$ :

$$p(\tau) = p(\tau_0) \left( \frac{\tau_0}{\tau} \right)^{4/3}. \quad (10)$$

#### IV. RESULTS

We first take into account the effect of the spontaneous breaking of chiral symmetry where no renormalization of the pion decay constant occurs, i.e.  $f_t = f_s = f_\pi = 93$  MeV. We thus obtain from the hydrodynamic equations (1),(2) in the first non-vanishing order of the pion field :

$$\frac{\partial s}{\partial \tau} + \frac{s}{\tau} = 0 \quad (11)$$

$$\frac{\partial T}{\partial \tau} + \frac{1}{3} \frac{T}{\tau} = 0 \quad (12)$$

$$\frac{\partial \epsilon}{\partial \tau} + \frac{T s(\tau_0) \tau_0}{\tau^2} - \frac{1}{\tau} \sum_{i=1}^3 \left( \frac{\partial \pi^i}{\partial \tau} \right)^2 = 0 \quad (13)$$

$$\left( \frac{\partial}{\partial \tau} + \frac{1}{\tau} \right) \frac{\partial \pi^i}{\partial \tau} = 0. \quad (14)$$

and from the explicit expression of the pressure  $p$  given in [1]:

$$p = aT^4 + \frac{1}{2} \sum_{i=1}^3 \left( \frac{\partial \pi^i}{\partial \tau} \right)^2 \quad (15)$$

We see that there is no modification of the equation for the entropy density and for the temperature, that is the scaling behavior is exactly the same  $s(\tau) = s(\tau_0)\tau_0/\tau$  and  $T(\tau) = T(\tau_0) (\tau_0/\tau)^{1/3}$ . On the contrary, for the energy density and for the pressure, one can observe a modification due to a term proportional to  $\partial \pi^i/\partial \tau$  which can be determined from the last equation (14). The solution for the time derivative is  $\partial \pi^i/\partial \tau = A/\tau$  where  $A$  is a constant whose value is fixed by initial condition for  $\tau = \tau_0$ . If we now report in the above equations, we can solve analytically the resulting system to obtain:

$$\epsilon(\tau) = \epsilon(\tau_0) \left( \frac{\tau_0}{\tau} \right)^{4/3} - \frac{3}{2} \frac{A^2}{\tau_0^2} \left( \frac{\tau_0}{\tau} \right)^2 \quad (16)$$

$$p = p(\tau_0) \left( \frac{\tau_0}{\tau} \right)^{4/3} + \frac{3}{2} \frac{A^2}{\tau_0^2} \left( \frac{\tau_0}{\tau} \right)^2. \quad (17)$$

In order to fix the initial condition related to the value of  $A$ , let us now consider the density matrix  $\rho = \exp^{-\beta H}$ . The part of the Hamiltonian relevant for chiral hydrodynamics degrees of freedom can be taken from [1]. An explicit calculation with Bjorken modelisation of the collision gives:

$$\rho = \exp \left( -\beta \int d^3x T^{00} \right) \propto \exp \left( -\frac{\beta}{2} \int d^3x \sum_{i=1}^3 \left( \frac{\partial \pi^i}{\partial \tau} \right)^2 \right). \quad (18)$$

Assuming isospin symmetry, we then have:  $\rho \propto \exp \left( -\frac{3\beta}{2} \int d^3x \left( \frac{A}{\tau} \right)^2 \right)$ . For  $\tau = \tau_0$ , we thus immediately see that  $\rho$  is Gauss distributed. Actually, if we call  $V_0$  the initial volume in the rest-frame of the two colliding nuclei and  $\gamma_{cm}$  the Lorentz factor needed to go from that frame to the laboratory frame, we have:

$$\rho(\tau_0) \propto \exp \left( -\frac{3\beta}{2} \frac{V_0}{\gamma_{cm}} \left( \frac{A}{\tau_0} \right)^2 \right). \quad (19)$$

We can easily estimate  $V_0$  by geometric arguments:  $V_0 = \pi R^2 2\tau_0$  where  $R$  is the radius of the colliding nuclei. If we now put typical numbers on all these constants:  $R \simeq 5$  fm,  $\tau_0 = 1$  fm,  $\gamma_{cm} \simeq 10$  and  $T(\tau_0) = 200$  MeV, we can directly estimate the effect of the terms associated to the spontaneous symmetry breaking since  $\langle A^2 \rangle$  is simply related to the Gaussian distribution:  $\langle A^2 \rangle = T(\tau_0)\tau_0\gamma_{cm}/(6\pi R^2)$ . Numerically, using these inputs, we obtain a correction of order  $\simeq 1\%$  with respect to the perfect case.

We now introduce the explicit chiral perturbation  $T$ -dependence for  $f_t$  and  $f_s$ , i.e.  $f_t^2 = f_s^2 = f_\pi^2(1 - T^2/(6f_\pi^2))$ . The evolution of the temperature is then governed by the following equation :

$$4a\tau T^3 - \frac{\langle A'^2 \rangle}{\tau(1 - T^2/(6f_\pi^2))^2} \frac{T}{6f_\pi^2} = s(\tau_0)\tau_0 \quad (20)$$

where now  $\langle A'^2 \rangle = \langle A^2 \rangle (1 - T_0^2/(6f_\pi^2))$ . The numerical results are presented on figure 1. We can observe

that the correction, which induce a faster cooling of the system is of order 5%.

Even if the approach can be considered too simplistic, it is not reasonably expected that an improved treatment can modify noticeably the order of magnitude. One can thus conclude that the effect due the spontaneous breaking of chiral symmetry is quite negligible and without any experimental consequences.

## V. DISSIPATIVE EFFECTS

In the previous part we have seen that it is possible to neglect the effect of the breaking of the symmetry on the hydrodynamics theory itself, that is we can use "normal" hydrodynamics. Nevertheless an important question which has already been addressed in [2] is the effect of dissipation on top of this model of perfect fluid.

There are two non equivalent possibilities to take into account dissipation. First, one can use the usual formalism and just say that when an inhomogeneity appears in a system, the latter reacts and a flow proportional to the gradient develops itself in order to restore equilibrium. This is first order theory. However it is well-known that this approach presents instabilities and that it is necessary to consider a relaxation time for the flow. This goal is achieved through Müller-Israel-Stewart second order theory. In the following we will explore both methods.

In [2], the author showed that only the dissipative effects related to the shear viscosity  $\eta$  play a role for the temperature profile in the (1+1) dimensional Bjorken model. The main physical reason is that bulk viscosity vanishes for an ultra relativistic gas. He took the shear viscosity from [7] where the temperature dependence for  $\eta$  is  $\eta = \pi f_\pi^4 / (8T)$ . Then, the differential equations to be solved for the energy density  $\epsilon = 3aT^4$  are:

$$\frac{\partial \epsilon}{\partial \tau} = -\frac{4}{3} \frac{\epsilon}{\tau} + \frac{\Phi}{\tau} \quad (21)$$

where  $\Phi$  is related to the stress tensor (see [2] for further details) and  $\Phi = 0$  for the ideal (perfect) fluid,  $\Phi = \frac{4}{3} \frac{\eta}{\tau}$  for the first order and  $\tau_\pi \frac{\partial \Phi}{\partial \tau} = -\Phi + \frac{4}{3} \frac{\eta}{\tau}$  for the second order theory discussed before. In the last equation  $\tau_\pi$  is the relaxation time associated to  $\Phi$ .

We now use (21), using the same inputs as [2], namely massless pions and current algebra cross section. We can see on figure 2 that the correction due to viscosity compared to the ideal case is totally negligible not only for the (non-physical but usual) first order but also for the second order. Our results are different from those presented in the original paper [2]. The difference lies in fact in a small computational mistake in [2]. The main point is that, contrary to what is claimed in [2], there are, in this simple approach, no appreciable effects. If we would stop here, we could say, as in the first part dedicated to the breaking of chiral symmetry that a realistic descrip-

tion of a pion gas can be achieved with the perfect fluid model.

However, we stress that this conclusion only holds because of the hypothesis used in [2]. The temperature dependences of the viscosity  $\eta$  and of the relaxation time  $\tau_\pi$  are only valid for massless pions and when the cross-section for  $\pi-\pi$  interaction is given by current algebra. If we take instead the realistic case of massive pions and the experimental cross-section, these dependences are completely different. For instance,  $\eta$  increases with temperature [7, 8] instead of decreasing. With this realistic  $T$ -dependence for the viscosity and the relaxation time for the pions taken from [8], we obtain the curves plotted on figure 2. We can now observe that the first and the second order are still close but they are now around 20% above the ideal fluid case. This means that dissipative effects imply a slower cooling of the system and should be incorporated in the simulations of heavy ion collisions.

## VI. CONCLUSION

In the first part of this paper we have examined in a simple model an application of hydrodynamic equations with spontaneous breaking symmetry. In order to get an estimate of the magnitude of the effects, we have used the one dimensional Bjorken description of a heavy ion collision. We have shown that the model of the perfect fluid is almost correct : the effect of the breaking of the symmetry is to give a correction of about 1% when no  $T$ -dependence of the pion decay constant is taken into account and a few percents when we consider a crude  $T$ -dependence. In a second part we have considered an other deviation from the perfect fluid model, namely the dissipation. The dissipative effects, when taken into account in a simple description [2] are not so important as claimed in this reference but represent actually a few percents. On the contrary, we have obtained sizable effects with a correct  $T$ -dependence for the shear viscosity and the shear flow relaxation time. For example, for an initial temperature of 200 MeV, the temperature goes from about 110 MeV in the ideal case to 140 MeV with our dissipative effects for  $\tau/\tau_0 \simeq 6$ . The whole conclusion of this paper is that the temperature profile generated from the perfect fluid approach and used in many simulations is almost correct if one takes into account only the effects of symmetry breaking. On the contrary, if we consider dissipative effects the profile can be substantially modified, that is we have a slower cooling of the pion gas.

Acknowledgments : We thank M. Ericson, Emeritus Professor at University Claude Bernard and G. Chanfray for constant help and enlightening discussions. We also thank J. Serreau for many discussions.

- 
- [1] D.T.Son, Phys. Rev. Lett. **84** (2000) 3771.
- [2] A. Muronga, *Proceedings of 17th Winter Workshop on Nuclear Dynamics*, Park City, Utah, 10-17 March 2001.
- [3] We thank A. Muronga for discussions and clarifications on that point.
- [4] J.D. Bjorken Phys. Rev. **D27** (1983) 140-151.
- [5] L.P.Csernai, *Introduction to Relativistic Heavy Ions Collisions* (J. Wiley and Sons, New York, 1972).
- [6] Cheuk-Yin Wong *Introduction to High-Energy Heavy-Ion Collisions* (World Scientific, Singapore, 1994).
- [7] M. Prakash, M. Prakash, R. Venugopalan and G.W. Welke, Phys. Rep. **227** (1993) 321.
- [8] D. Davesne, *Phys. Rev.* **C53** (1996) 3069.




FIG. 1: Temperature profile with respect to proper time for two cases: without (solid line) and with (dashed line) effect of symmetry breaking.




FIG. 2: Temperature profile with respect to proper time for the perfect fluid (solid line), first order and second order theories.