

# Spacetime Structure of Massive Gravitinos

M. Kirchbach<sup>a</sup> and D. V. Ahluwalia<sup>a,b</sup>

*a.* Theoretical Physics Group

Facultad de Física, Universidad Autónoma de Zacatecas,

A. P. C-600, ZAC-98062, México

*b.* Inter-University Center for Astronomy and Astrophysics (IUCAA)

Post Bag 4, Ganeshkhind, Pune, India 411 007

**Abstract.** The talk presents an *ab initio* construct of the spacetime structure underlying massive gravitinos. We argue that single spin interpretation of massive gravitino is untenable, and that a spin measurement in the rest frame for an unpolarized ensemble of massive gravitinos, would yield the results  $3/2^-$  with probability one half, and  $1/2$  with probability one half. The latter is distributed uniformly, i.e. as  $1/4$ , among the two spin- $1/2^+$  and spin- $1/2^-$  states of opposite parities. From that we draw the conclusion that the massive gravitino should be interpreted as a particle of multiple spin. We expect that a natural extension of this work to finite-range gravity shall endow the graviton with spins 0, 1, and 2 components.

## 1. Introduction

This talk is adapted from our recent works [1, 2]. The goal is to review new insights into the spacetime properties of massive gravitinos. We examine the nature of the spinor-vector  $\psi^\mu$  that appears in supersymmetric theories as a fermionic gauge field, the so called gravitino, and show that its single-spin interpretation is unjustified. Rather, this particle represents itself as a multispin object. The presentation is organized as follows.

In the next Section we briefly outline the construction of the Dirac  $(1/2, 0) \oplus (0, 1/2)$  representation space. Section 3 is devoted to the massive gravitino. There we show that the so called auxiliary conditions to the spinor-vector  $\psi^\mu$  are nothing but defining conditions for the subspace of  $\psi^\mu$  that carries the maximal spin of  $3/2^-$  in the rest frame. In particular, we derive an equation for  $\psi^\mu \gamma_\mu$  showing that the latter auxiliary condition is not arbitrary at all and can not be set equal to zero unconditionally. We suggest sets of such defining-“auxiliary”- conditions for each one of the spin- $1/2^-$  and spin- $1/2^+$  rest-frame sectors of  $\psi^\mu$  and show that none of them is more or less physical than the other two. From that we derive the conclusion about the multi-spin character of the massive gravitino. The talk ends with a brief outlook.

## 2. Dirac’s $(1/2, 0) \oplus (0, 1/2)$ representation space

In this section we outline a general new procedure for obtaining the Dirac equation that relies on nothing more but the boost operators *and* a phase relationship between right and left handed Weyl spinors at rest. We begin with the rest-frame spinors and boost them using the following boosts

$$\kappa^{(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})} = \kappa^{(\frac{1}{2}, 0)} \oplus \kappa^{(0, \frac{1}{2})}, \quad (1)$$

with

$$\kappa^{(\frac{1}{2},0)} = \exp\left(+\frac{\boldsymbol{\sigma}}{2} \cdot \boldsymbol{\varphi}\right) = \sqrt{\frac{E+m}{2m}} \left(1_2 + \frac{\boldsymbol{\sigma} \cdot \mathbf{p}}{E+m}\right), \quad (2)$$

$$\kappa^{(0,\frac{1}{2})} = \exp\left(-\frac{\boldsymbol{\sigma}}{2} \cdot \boldsymbol{\varphi}\right) = \sqrt{\frac{E+m}{2m}} \left(1_2 - \frac{\boldsymbol{\sigma} \cdot \mathbf{p}}{E+m}\right). \quad (3)$$

In Eqs. (2) and (3) the boost parameter is defined as:

$$\cosh(\varphi) = \frac{E}{m}, \quad \sinh(\varphi) = \frac{|\mathbf{p}|}{m}, \quad \hat{\boldsymbol{\varphi}} = \frac{\mathbf{p}}{|\mathbf{p}|}. \quad (4)$$

The boosts take a particle at rest to a particle moving with momentum  $\mathbf{p}$  in the ‘‘boosted frame.’’ We use the notation in which  $1_n$  and  $0_n$  represent  $n \times n$  identity and null matrices, respectively. The remaining symbols carry their usual contextual meaning. We define the spin-1/2 helicity operator:  $\Sigma = (\boldsymbol{\sigma}/2) \cdot \hat{\mathbf{p}}$ , where  $\hat{\mathbf{p}} = \mathbf{p}/|\mathbf{p}|$ , and  $\mathbf{p} = |\mathbf{p}|(\sin(\theta) \cos(\phi), \sin(\theta) \sin(\phi), \cos(\theta))$ . Its positive and negative helicity states are:

$$\begin{aligned} h^+ &= N \begin{pmatrix} \cos(\theta/2) \exp(-i\phi/2) \\ \sin(\theta/2) \exp(i\phi/2) \end{pmatrix}, \\ h^- &= N \begin{pmatrix} \sin(\theta/2) \exp(-i\phi/2) \\ -\cos(\theta/2) \exp(i\phi/2) \end{pmatrix}. \end{aligned} \quad (5)$$

The rest-frame  $(1/2, 0) \oplus (0, 1/2)$  spinors are then chosen to be:

$$\begin{aligned} u_{+1/2}(\mathbf{0}) &= \begin{pmatrix} h^+ \\ h^+ \end{pmatrix}, \quad u_{-1/2}(\mathbf{0}) = \begin{pmatrix} h^- \\ h^- \end{pmatrix}, \\ v_{+1/2}(\mathbf{0}) &= \begin{pmatrix} h^+ \\ -h^+ \end{pmatrix}, \quad v_{-1/2}(\mathbf{0}) = \begin{pmatrix} h^- \\ -h^- \end{pmatrix}. \end{aligned} \quad (6)$$

The choice of the phases made in writing down these spinors has been determined by the demand of parity covariance [3]. The boosted spinors,  $u_{\pm 1/2}(\mathbf{p})$  and  $v_{\pm 1/2}(\mathbf{p})$  are obtained by applying the boost operator  $\kappa^{(\frac{1}{2},0) \oplus (0,\frac{1}{2})}$  to the above spinors, yielding:

$$\begin{aligned} u_{+1/2}(\mathbf{p}) &= \frac{N}{\sqrt{2m(m+E)}} \begin{pmatrix} \exp(-i\phi/2)(m+|\mathbf{p}|+E) \cos(\theta/2) \\ \exp(i\phi/2)(m+|\mathbf{p}|+E) \sin(\theta/2) \\ \exp(-i\phi/2)(m-|\mathbf{p}|+E) \cos(\theta/2) \\ \exp(i\phi/2)(m-|\mathbf{p}|+E) \sin(\theta/2) \end{pmatrix}, \\ u_{-1/2}(\mathbf{p}) &= \frac{N}{\sqrt{2m(m+E)}} \begin{pmatrix} \exp(-i\phi/2)(m-|\mathbf{p}|+E) \sin(\theta/2) \\ -\exp(i\phi/2)(m-|\mathbf{p}|+E) \cos(\theta/2) \\ \exp(-i\phi/2)(m+|\mathbf{p}|+E) \sin(\theta/2) \\ -\exp(i\phi/2)(m+|\mathbf{p}|+E) \cos(\theta/2) \end{pmatrix}, \\ v_{+1/2}(\mathbf{p}) &= \frac{N}{\sqrt{2m(m+E)}} \begin{pmatrix} \exp(-i\phi/2)(m+|\mathbf{p}|+E) \cos(\theta/2) \\ \exp(i\phi/2)(m+|\mathbf{p}|+E) \sin(\theta/2) \\ -\exp(-i\phi/2)(m-|\mathbf{p}|+E) \cos(\theta/2) \\ -\exp(i\phi/2)(m-|\mathbf{p}|+E) \sin(\theta/2) \end{pmatrix}, \\ v_{-1/2}(\mathbf{p}) &= \frac{N}{\sqrt{2m(m+E)}} \begin{pmatrix} \exp(-i\phi/2)(m-|\mathbf{p}|+E) \sin(\theta/2) \\ -\exp(i\phi/2)(m-|\mathbf{p}|+E) \cos(\theta/2) \\ -\exp(-i\phi/2)(m+|\mathbf{p}|+E) \sin(\theta/2) \\ \exp(i\phi/2)(m+|\mathbf{p}|+E) \cos(\theta/2) \end{pmatrix}. \end{aligned} \quad (7)$$

These satisfy the orthonormality and completeness relations (in standard notation):

$$\bar{u}_h(\mathbf{p}) u_{h'}(\mathbf{p}) = +2N^2 \delta_{hh'}, \quad \bar{v}_h(\mathbf{p}) v_{h'}(\mathbf{p}) = -2N^2 \delta_{hh'}, \quad (8)$$

$$\frac{1}{2N^2} \left[ \sum_{h=\pm 1/2} u_h(\mathbf{p}) \bar{u}_h(\mathbf{p}) - \sum_{h=\pm 1/2} v_h(\mathbf{p}) \bar{v}_h(\mathbf{p}) \right] = 1_4. \quad (9)$$

In order to obtain the wave equation satisfied by the  $u_h(\mathbf{p})$  and  $v_h(\mathbf{p})$  spinors we first note that ‡

$$\frac{1}{2N^2} \left[ \sum_{h=\pm 1/2} u_h(\mathbf{p}) \bar{u}_h(\mathbf{p}) + \sum_{h=\pm 1/2} v_h(\mathbf{p}) \bar{v}_h(\mathbf{p}) \right] \equiv \frac{\gamma_\mu p^\mu}{m}, \quad (10)$$

where we defined,

$$\gamma_0 = \begin{pmatrix} 0_2 & I_2 \\ I_2 & 0_2 \end{pmatrix}, \quad \gamma_i = \begin{pmatrix} 0_2 & \sigma_i \\ -\sigma_i & 0_2 \end{pmatrix}. \quad (11)$$

Adding/subtracting Eqs. (9) and (10), yields:

$$\frac{1}{2N^2} \sum_{h=\pm 1/2} u_h(\mathbf{p}) \bar{u}_h(\mathbf{p}) = \frac{\gamma_\mu p^\mu + m 1_4}{2m}, \quad (12)$$

$$\frac{1}{2N^2} \sum_{h=\pm 1/2} v_h(\mathbf{p}) \bar{v}_h(\mathbf{p}) = \frac{\gamma_\mu p^\mu - m 1_4}{2m}, \quad (13)$$

respectively. Multiplying Eq. (12) from the right by  $u_{h'}(\mathbf{p})$ , and Eq. (13) by  $v_{h'}(\mathbf{p})$ , and using Eqs. (8), immediately yields the well known momentum-space Dirac equation for the  $(1/2, 0) \oplus (0, 1/2)$  representation space,

$$(\gamma_\mu p^\mu \pm m 1_4) \psi_h(\mathbf{p}) = 0. \quad (14)$$

In Eq. (14), the minus sign is to be taken for,  $\psi_h(\mathbf{p}) = u_h(\mathbf{p})$ , and the plus sign for,  $\psi_h(\mathbf{p}) = v_h(\mathbf{p})$ . The essential element to note in regard to the derivation of Eq. (14) is that it follows directly from the explicit expressions for the  $u_h(\mathbf{p})$  and  $v_h(\mathbf{p})$ .

### 3. Spacetime properties of massive gravitino

A massive gravitino is described by  $\psi^\mu$ . As far as its spacetime properties are concerned, it transforms as a finite dimensional non-unitary representation of the Lorentz group,


$$\psi^\mu : \underbrace{\left[ \left( \frac{1}{2}, 0 \right) \oplus \left( 0, \frac{1}{2} \right) \right]}_{\text{SPINOR SECTOR}} \otimes \underbrace{\left( \frac{1}{2}, \frac{1}{2} \right)}_{\text{VECTOR SECTOR}}. \quad (15)$$

The unitarily transforming physical states are built upon this structure [4].

We enumerate two circumstances that motivate us to take an *ab initio* look at this representation space.

- (i) For the vector sector, it has recently been called to attention that the Proca description of the  $(1/2, 1/2)$  representation space is incomplete [1]. An *ab initio* construction of this sector reveals that the Stückelberg contribution to the propagator, so important for the renormalization of the gauge theories with massive bosons [5], is found to naturally reside in the  $(1/2, 1/2)$  representation space.

‡ This part of the calculation is best done if one starts with the momenta in the Cartesian co-ordinates, and takes the “axis of spin quantization” to be the z-axis.



**Figure 1.** Summary of the data on  $N$  and  $\Delta$  resonances. The breaking of the mass degeneracy for each of the clusters at about 5% may in fact be an artifact of the data analysis, as has been suggested by Höhler [12]. The filled circles represent known resonances, while the sole empty circle corresponds to a prediction.

- (ii) At the same time, the properties of the  $(1/2, 1/2)$ , along with that of the  $(1/2, 0) \oplus (0, 1/2)$ , representation space determine the structure of  $\psi^\mu$ . In order to impose a single-spin, i.e., spin  $3/2$ , interpretation on the latter, the lower spin- $1/2^+$  and spin- $1/2^-$  components of  $\psi^\mu$  are considered as redundant, unphysical, states that are claimed to be excluded from consideration by means of the two supplementary conditions:  $\gamma_\mu \psi^\mu(x) = 0$ , and  $\partial_\mu \psi^\mu(x) = 0$ , respectively. However, this time-honored framework was questioned by a recent empirical observation regarding the  $N$  and  $\Delta$  resonances [6]. The available data on high-spin resonances reveal an unexpected and systematic clustering in terms of the  $(j/2, j/2) \otimes [(1/2, 0) \oplus (0, 1/2)]$  representations with  $j = 1, 3$  and  $5$  *without* imposition of the supplementary conditions. For the  $N$  and  $\Delta$  resonances these results are summarized in Figure 1. For example, in the standard theoretical framework  $N(1440)$ ,  $N(1535)$ ,  $\Delta(1620)$  and  $\Delta(1750)$  should have been absent. Experimental data shows them to be present at statistically significant level.  $\Sigma$

In regard to the latter of the two enumerated circumstances, we take the position that any solution of the QCD Lagrangian for particle resonances must carry well-defined

$\Sigma$ The  $N(1440)$ ,  $N(1535)$ ,  $\Delta(1620)$  carry four star status, while at present  $\Delta(1750)$  simply has a one star significance [7].

transformation properties when looked upon from different inertial frames. This forces these resonances to belong to one, or the other, of various representation spaces of the Lorentz group. For this reason the data on particle resonances may furnish hints on physical interpretation of various Lorentz group representations that one needs in gauge theories, or theories of supergravity.

For exploring the spacetime structure of massive gravitinos the charge conjugation properties play an important role. Under the operation of charge conjugation, one may choose the spinor sector to behave as a Dirac object, and implement the Majorana nature of the massive gravitino at the level of the Fock space. This is standard, see, e.g., Ref. [8]. Or, from the very beginning choose the spinor sector to behave as a Majorana object. Since we wish to stress certain non-trivial aspects of massive gravitino that do not – at least qualitatively – depend on this choice, we shall here treat the spinor sector to be of Dirac type.

Very nature of our *ab initio* look at the representation space defined in Eq. (15), obliges us to present sufficient pedagogic details so that by the end of the lecture much that is needed to form an opinion on the arrived results is readily available. At the same time, length constraints of this manuscript would prevent us from delving into subtle details which are, for present, of secondary importance (but have been studied and are planned to be presented elsewhere).

We shall work in the momentum space. The notation will be essentially that introduced in Ref. [1].

### 3.1. $(1/2, 1/2)$ Representation space – An *ab initio* construct

We have constructed in Section 2 the spinorial sector entering  $\psi^\mu$  in Eq. (15). Therefore, our next task is to construct the  $(1/2, 1/2)$  representation space. As such, now we introduce the rest-frame vectors for the  $(1/2, 1/2)$  representation space,

$$\xi_1(\mathbf{0}) = h^+ \otimes h^+, \quad (16)$$

$$\xi_2(\mathbf{0}) = \frac{1}{\sqrt{2}} (h^+ \otimes h^- + h^- \otimes h^+) , \quad (17)$$

$$\xi_3(\mathbf{0}) = h^- \otimes h^- , \quad (18)$$

$$\xi_4(\mathbf{0}) = \frac{1}{\sqrt{2}} (h^+ \otimes h^- - h^- \otimes h^+) . \quad (19)$$

The boosted vectors are thus:  $\xi_\zeta(\mathbf{p}) = \kappa^{(\frac{1}{2}, \frac{1}{2})} \xi_\zeta(\mathbf{0})$ ,  $\zeta = 1, 2, 3, 4$ ,

$$\xi_1(\mathbf{p}) = \frac{N^2}{2} \begin{pmatrix} 2 \exp(-i\phi) \cos^2(\theta/2) \\ \sin(\theta) \\ \sin(\theta) \\ 2 \exp(i\phi) \sin^2(\theta/2) \end{pmatrix} ,$$

$$\xi_2(\mathbf{p}) = \frac{N^2}{\sqrt{2}m} \begin{pmatrix} \exp(-i\phi) E \sin(\theta) \\ -(|\mathbf{p}| + E \cos(\theta)) \\ |\mathbf{p}| - E \cos(\theta) \\ -\exp(i\phi) E \sin(\theta) \end{pmatrix} ,$$

$$\xi_3(\mathbf{p}) = \frac{N^2}{2} \begin{pmatrix} 2 \exp(-i\phi) \sin^2(\theta/2) \\ -\sin(\theta) \\ -\sin(\theta) \\ 2 \exp(i\phi) \cos^2(\theta/2) \end{pmatrix} ,$$

$$\xi_4(\mathbf{p}) = \frac{N^2}{\sqrt{2}m} \begin{pmatrix} \exp(-i\phi)|\mathbf{p}| \sin(\theta) \\ -(E + |\mathbf{p}| \cos(\theta)) \\ E - |\mathbf{p}| \cos(\theta) \\ -\exp(i\phi)|\mathbf{p}| \sin(\theta) \end{pmatrix}. \quad (20)$$

Here  $\kappa^{(1/2,1/2)} = \kappa^{(1/2,0)} \otimes \kappa^{(0,1/2)}$ .

In the notation of Ref. [1],<sup>||</sup> these satisfy the orthonormality and completeness relations, along with a new wave equation. The orthonormality and completeness relations are:

$$\begin{aligned} \bar{\xi}_\zeta(\mathbf{p})\xi_{\zeta'}(\mathbf{p}) &= -N^4 \delta_{\zeta\zeta'}, \quad \zeta = 1, 2, 3, \\ \bar{\xi}_\zeta(\mathbf{p})\xi_{\zeta'}(\mathbf{p}) &= +N^4 \delta_{\zeta\zeta'}, \quad \zeta = 4, \\ \frac{1}{N^4} \left[ \xi_4(\mathbf{p})\bar{\xi}_4(\mathbf{p}) - \sum_{\zeta=1,2,3} \xi_\zeta(\mathbf{p})\bar{\xi}_\zeta(\mathbf{p}) \right] &= 1_4, \end{aligned} \quad (21)$$

where

$$\bar{\xi}_\zeta(\mathbf{p}) \equiv \xi_\zeta(\mathbf{p})^\dagger \lambda_{00}, \quad (22)$$

with

$$\lambda_{00} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \quad (23)$$

The parity operator for the  $(1/2, 1/2)$  representation space is:

$$\begin{aligned} \mathcal{P} &= \lambda_{00} \exp[i\alpha] \mathcal{R}, \quad \mathcal{R} : \{\theta \rightarrow \pi - \theta, \phi \rightarrow \pi + \phi\} \\ &\alpha = \text{a real number}, \end{aligned} \quad (24)$$

while the helicity operator for this space is,  $\mathbf{J} \cdot \hat{\mathbf{p}}$ , with  $\mathbf{J}$  given by:

$$\begin{aligned} J_x &= \frac{1}{2} \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}, \quad J_y = \frac{1}{2} \begin{pmatrix} 0 & -i & -i & 0 \\ i & 0 & 0 & -i \\ i & 0 & 0 & -i \\ 0 & i & i & 0 \end{pmatrix}, \\ J_z &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \end{aligned} \quad (25)$$

We now must take a small definitional detour towards the notion of the dragged Casimirs for spacetime symmetries. It arises in the following fashion. The second Casimir operator,  $C_2$ , of the Poincaré group is defined as the square of the Pauli-Lubanski pseudovector:

$$\mathcal{W}^\mu = \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} M_{\nu\rho} P_\sigma, \quad (26)$$

where  $\epsilon^{\mu\nu\rho\sigma}$  is the standard Levi-Civita symbol in four dimensions, while  $M_{\mu\nu}$  denote generators of the Lorentz group,

$$M_{0i} = K_i, \quad M_{ij} = \epsilon_{ijk} J^k, \quad (27)$$

<sup>||</sup> In the cited work we presented the  $(1/2, 1/2)$  representation space in its parity realization. Here, the presentation is in terms of helicity realization. The two descriptions have mathematically similar but physically distinct structures, which, e.g., show up in their different behavior under the operation of Parity.

where each of the  $i, j, k$  runs over 1, 2, 3. The  $P_\mu$  are generators of the spacetime translations. In general, these have non-vanishing commutators with  $M_{\mu\nu}$ ,

$$[P_\mu, M_{\rho\sigma}] = i(\eta_{\mu\rho}P_\sigma - \eta_{\mu\sigma}P_\rho). \quad (28)$$

On using Eq. (28), we rewrite  $C_2$  as

$$C_2 = \frac{1}{4}\epsilon^{\mu\nu\rho\sigma}\epsilon_{\mu\lambda\kappa\zeta}M_{\nu\rho}M^{\lambda\kappa}P_\sigma P^\zeta + \left[ \frac{i}{4}\epsilon^{\mu\nu\rho\sigma}\epsilon_{\mu\lambda\kappa\zeta}M_{\nu\rho}\eta_\sigma^\lambda P^\kappa P^\zeta + \frac{i}{4}\epsilon^{\mu\nu\rho\sigma}\epsilon_{\mu\kappa\lambda\zeta}M_{\nu\rho}\eta_\sigma^\kappa P^\lambda P^\zeta \right]. \quad (29)$$

The squared bracket vanishes due to antisymmetry of the Levi-Civita symbol. As such, the space-time translation operators entering the definition of  $C_2$  can be moved to the very right. This observation allows for introducing the dragged Casimir  $\tilde{C}_2$  as an operator with the same form as  $C_2$  – the difference being that the commutator in Eq. (28) is now set to zero [as is appropriate for finite dimensional  $SU_R(2) \otimes SU_L(2)$  representations]. Consequently, while  $C_2$  and  $\tilde{C}_2$  carry same invariant eigenvalues when acting upon momentum eigenstates, their commutators with the Lorentz group generators are no longer identical. For the  $(1/2, 0) \oplus (0, 1/2)$  representation space,  $[\tilde{C}_2, \mathbf{J}^2]$  vanishes. For the  $(1/2, 1/2)$  representation space,  $[\tilde{C}_2, \mathbf{J}^2]$  does not vanish (except when acting upon rest states), and equals  $-4i E \mathbf{P} \cdot \mathbf{K}$ . This leads to the fact that while the former representation space is endowed with a well-defined spin, the latter is not:

As an immediate application,  $\tilde{C}_2$  for the  $(1/2, 1/2)$  representation space bifurcates this space into two sectors. The three states  $\xi_\zeta(\mathbf{p})$  with  $\zeta = 1, 2, 3$  are associated with the  $\tilde{C}_2$  eigenvalue,  $-2m^2$ ; while the,  $\zeta = 4$ , corresponds to eigenvalue zero. Thus, all the  $\xi_\zeta(\mathbf{p})$ , except for the rest frame, cease to be eigenstates of the  $(1/2, 1/2)$ 's  $\mathbf{J}^2$  and do not carry definite spins. This contrasts with the situation for the  $(1/2, 0) \oplus (0, 1/2)$  representation space, where the  $\psi_h(\mathbf{p})$  are eigenstates of the corresponding  $\mathbf{J}^2$ .

Now in order that the  $\xi_\zeta(\mathbf{p})$  carry the standard contravariant Lorentz index, we introduce a rotation in the  $(1/2, 1/2)$  representation space via [1]:

$$S = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & i & -i & 0 \\ -i & 0 & 0 & i \\ 1 & 0 & 0 & 1 \\ 0 & i & i & 0 \end{pmatrix}. \quad (30)$$

Then, the  $(1/2, 1/2)$  representation space is spanned by four Lorentz vectors:

$$\mathcal{A}_\zeta^\mu(\mathbf{p}) = S^{\mu\alpha} [\xi_\zeta(\mathbf{p})]_\alpha, \quad \zeta = 1, 2, 3, 4, \quad (31)$$

and the superscript  $\mu$  is the standard Lorentz index. Note that the  $\mathcal{W}^\mu$ 's by themselves are  $4 \times 4$  matrices in Lorentz index space, i.e.  $\mathcal{W}_{\nu\eta}^\mu$ . Following the procedure established in Sec. II, they can be shown to satisfy a new wave equation [1],

$$\left( \Lambda_{\mu\nu} p^\mu p^\nu \pm m^2 I_4 \right) \mathcal{A}_\zeta(\mathbf{p}) = 0, \quad (32)$$

**P** To avoid confusion, note that  $\tilde{C}_2$  is defined in the  $SU_R(2) \otimes SU_L(2)$ ; while  $C_2$  is defined in the Poincaré group.

where the plus sign is to be taken for,  $\zeta = 1, 2, 3$ , while the minus sign belongs to,  $\zeta = 4$ . The  $\Lambda_{\mu\nu}$  matrices are:  $\Lambda_{00} = \text{diag}(1, -1, -1, -1)$ ,  $\Lambda_{11} = \text{diag}(1, -1, 1, 1)$ ,  $\Lambda_{22} = \text{diag}(1, 1, -1, 1)$ ,  $\Lambda_{33} = \text{diag}(1, 1, 1, -1)$ , and

$$\begin{aligned} \Lambda_{01} &= \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \Lambda_{02} = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \Lambda_{03} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \\ \Lambda_{12} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \Lambda_{13} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, \\ \Lambda_{23} &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{pmatrix}. \end{aligned} \quad (33)$$

The remaining  $\Lambda_{\mu\nu}$  are obtained from the above expressions by noting:  $\Lambda_{\mu\nu} = \Lambda_{\nu\mu}$ . Parenthetically, we note that the S-transformed  $\lambda_{00}$  equals  $\Lambda_{00}$  and is nothing but the standard spacetime metric (for flat spacetime).

The massive (1/2,1/2) propagators that follow from the completeness relation within the (1/2,1/2) representation space in Eq. (21) read (in the notations of Eq. (31))

$$[\mathcal{A}_4(\vec{p})\bar{\mathcal{A}}_4(\vec{p})\Lambda_{00}]_{\mu\nu} = \frac{p_\mu p_\nu}{m^2}, \quad (34)$$

$$-\sum_{\zeta=1}^3 [\mathcal{A}_\zeta(\vec{p})\bar{\mathcal{A}}_\zeta(\vec{p})\Lambda_{00}]_{\mu\nu} = g_{\mu\nu} - \frac{p_\mu p_\nu}{m^2}. \quad (35)$$

One immediately realizes that the massive (1/2,1/2) propagator contains both the Stückelberg (34) and the Proca (35) terms [5].

This feature of the completeness relation within the (1/2,1/2) representation space appears quite appealing to us as it leads to a well behaved propagator of a massive gauge boson as arising in a spontaneously broken local gauge theory. Within the context of the scenario presented above, the Proca sector is characterized by vanishing of  $p^\mu \mathcal{A}_\mu = 0$ , while the Stückelberg sector is characterized by vanishing of  $p^\mu \widetilde{\mathcal{W}}_\mu = 0$  (see Table I).

It can also be seen that  $\xi_\zeta(\mathbf{p})$ , for  $\zeta = 1, 2, 3$ , coincide with the solutions of Proca framework (and are divergence-less); whereas  $\xi_4(\mathbf{p})$ , that gives the Stückelberg contribution to the propagator, lies outside the Proca framework:<sup>+</sup>

$\zeta$	$p_\mu \mathcal{A}^\mu(\mathbf{p})$	$\widetilde{\mathcal{W}}_\mu^{(1/2,1/2)} \mathcal{A}^\mu(\mathbf{p})$	$\lambda_\tau$	Remarks
1, 2, 3	$= 0$	$\neq 0$	2	Proca Sector
4	$\neq 0$	$= 0$	0	Stückelberg Sector

In the above table we have introduced the  $\lambda_\tau$  via the equation:

$$\widetilde{C}_2^{(1/2,1/2)} \mathcal{A}(\mathbf{p}) = -\lambda_\tau m^2 \mathcal{A}(\mathbf{p}). \quad (36)$$

<sup>+</sup> We define the dragged Pauli-Lubanski pseudovector,  $\widetilde{\mathcal{W}}_\mu$ , in a manner parallel to the introduction of the dragged second Casimir operator.



### 3.2. Construction of spinor vector: Massive gravitino

We now wish to present the basis vectors for the representation space defined by Eq. (15) in a language which is widely used [8]. This would allow the present analysis to be more readily available, and also bring out the relevant similarities and differences with the framework of Rarita and Schwinger [9].

In writing down the basis spinor-vectors, we will use the fact that in the  $(1/2, 1/2)$  representation space the charge conjugation is implemented by

$$\mathcal{C}^{(1/2,1/2)} : \mathcal{A}(\mathbf{p}) \rightarrow [\mathcal{A}(\mathbf{p})]^* . \quad (37)$$

In the  $(1/2, 0) \oplus (0, 1/2)$  representation space the charge conjugation operator is  $\mathcal{C}^{(1/2,0) \oplus (0,1/2)} : i\gamma^2 K$ , where  $K$  complex conjugates the spinor to its right. Then, we obtain:

$$\mathcal{C}^{(1/2,0) \oplus (0,1/2)} : \begin{cases} u_{+1/2}(\mathbf{p}) \rightarrow -v_{-1/2}(\mathbf{p}), u_{-1/2}(\mathbf{p}) \rightarrow v_{+1/2}(\mathbf{p}), \\ v_{+1/2}(\mathbf{p}) \rightarrow u_{-1/2}(\mathbf{p}), v_{-1/2}(\mathbf{p}) \rightarrow -u_{+1/2}(\mathbf{p}). \end{cases} \quad (38)$$

In the spirit outlined, the massive gravitino lives in a space spanned by sixteen spinor-vectors defined in items **A**, **B**, **C** below:

- A.** Of these, eight spinor-vectors have  $\tilde{C}_2$  – but not  $\mathbf{J}^2$  – eigenvalues,  $-\frac{15}{4}m^2$ . These can be further subdivided into particle,

$$\psi_a^\mu(\mathbf{p}) : \begin{cases} \psi_1^\mu(\mathbf{p}) = u_{+1/2}(\mathbf{p}) \otimes \mathcal{A}_1^\mu(\mathbf{p}), \\ \psi_2^\mu(\mathbf{p}) = \sqrt{\frac{2}{3}} u_{+1/2}(\mathbf{p}) \otimes \mathcal{A}_2^\mu(\mathbf{p}) + \sqrt{\frac{1}{3}} u_{-1/2}(\mathbf{p}) \otimes \mathcal{A}_1^\mu(\mathbf{p}), \\ \psi_3^\mu(\mathbf{p}) = \sqrt{\frac{1}{3}} u_{+1/2}(\mathbf{p}) \otimes \mathcal{A}_3^\mu(\mathbf{p}) + \sqrt{\frac{2}{3}} u_{-1/2}(\mathbf{p}) \otimes \mathcal{A}_2^\mu(\mathbf{p}), \\ \psi_4^\mu(\mathbf{p}) = u_{-1/2}(\mathbf{p}) \otimes \mathcal{A}_3^\mu(\mathbf{p}), \end{cases}$$

and antiparticle sectors:

$$[\psi_a^\mu(\mathbf{p})]^c : \begin{cases} \psi_5^\mu(\mathbf{p}) = -v_{-1/2}(\mathbf{p}) \otimes [\mathcal{A}_1^\mu(\mathbf{p})]^*, \\ \psi_6^\mu(\mathbf{p}) = -\sqrt{\frac{2}{3}} v_{-1/2}(\mathbf{p}) \otimes [\mathcal{A}_2^\mu(\mathbf{p})]^* + \sqrt{\frac{1}{3}} v_{+1/2}(\mathbf{p}) \otimes [\mathcal{A}_1^\mu(\mathbf{p})]^*, \\ \psi_7^\mu(\mathbf{p}) = -\sqrt{\frac{1}{3}} v_{-1/2}(\mathbf{p}) \otimes [\mathcal{A}_3^\mu(\mathbf{p})]^* + \sqrt{\frac{2}{3}} v_{+1/2}(\mathbf{p}) \otimes [\mathcal{A}_2^\mu(\mathbf{p})]^*, \\ \psi_8^\mu(\mathbf{p}) = v_{+1/2}(\mathbf{p}) \otimes [\mathcal{A}_3^\mu(\mathbf{p})]^*. \end{cases}$$

Here,  $[\psi_\tau^\mu(\mathbf{p})]^c = \mathcal{C}^{(1/2,0) \oplus (0,1/2)} \otimes \mathcal{C}^{(1/2,1/2)} \psi_\tau^\mu(\mathbf{p})$ ,  $\tau = a, b, c$ .

- B.** Four spinor-vectors have  $\tilde{C}_2$  – but not  $\mathbf{J}^2$  – eigenvalues,  $-\frac{3}{4}m^2$ :

$$\psi_b^\mu(\mathbf{p}) : \begin{cases} \psi_9^\mu(\mathbf{p}) = \sqrt{\frac{2}{3}} u_{-1/2}(\mathbf{p}) \otimes \mathcal{A}_1^\mu(\mathbf{p}) - \sqrt{\frac{1}{3}} u_{+1/2}(\mathbf{p}) \otimes \mathcal{A}_2^\mu(\mathbf{p}), \\ \psi_{10}^\mu(\mathbf{p}) = \sqrt{\frac{1}{3}} u_{-1/2}(\mathbf{p}) \otimes \mathcal{A}_2^\mu(\mathbf{p}) - \sqrt{\frac{2}{3}} u_{+1/2}(\mathbf{p}) \otimes \mathcal{A}_3^\mu(\mathbf{p}), \end{cases}$$

$$[\psi_b^\mu(\mathbf{p})]^c : \begin{cases} \psi_{11}^\mu(\mathbf{p}) = \sqrt{\frac{2}{3}} v_{+1/2}(\mathbf{p}) \otimes [\mathcal{A}_1^\mu(\mathbf{p})]^* + \sqrt{\frac{1}{3}} v_{-1/2}(\mathbf{p}) \otimes [\mathcal{A}_2^\mu(\mathbf{p})]^*, \\ \psi_{12}^\mu(\mathbf{p}) = \sqrt{\frac{1}{3}} v_{+1/2}(\mathbf{p}) \otimes [\mathcal{A}_2^\mu(\mathbf{p})]^* + \sqrt{\frac{2}{3}} v_{-1/2}(\mathbf{p}) \otimes [\mathcal{A}_3^\mu(\mathbf{p})]^*. \end{cases}$$

- C.** Another set of four spinor-vectors with  $\tilde{C}_2$  – but not  $\mathbf{J}^2$  – eigenvalues,  $-\frac{3}{4}m^2$ :

$$\psi_c^\mu(\mathbf{p}) : \begin{cases} \psi_{13}^\mu(\mathbf{p}) = u_{+1/2}(\mathbf{p}) \otimes \mathcal{A}_4^\mu(\mathbf{p}), \\ \psi_{14}^\mu(\mathbf{p}) = u_{-1/2}(\mathbf{p}) \otimes \mathcal{A}_4^\mu(\mathbf{p}), \end{cases}$$

$$[\psi_c^\mu(\mathbf{p})]^c : \begin{cases} \psi_{15}^\mu(\mathbf{p}) = -v_{-1/2}(\mathbf{p}) \otimes [\mathcal{A}_4^\mu(\mathbf{p})]^*, \\ \psi_{16}^\mu(\mathbf{p}) = v_{+1/2}(\mathbf{p}) \otimes [\mathcal{A}_4^\mu(\mathbf{p})]^*. \end{cases}$$

We have evaluated  $\gamma_\mu \psi^\mu(\mathbf{p})$ ,  $p_\mu \psi^\mu(\mathbf{p})$ , and  $\widetilde{\mathcal{W}}_\mu^{(1/2,1/2)} \psi^\mu(\mathbf{p})$ , for all of the above sixteen spinor vectors. The  $p_\mu \psi^\mu(\mathbf{p})$ , when transformed to the configuration space, tests the divergence of  $\psi^\mu(x)$ .

For  $\eta = 1, 4, 5, 8$ ,  $\gamma_\mu \psi^\mu(\mathbf{p})$  identically vanishes. Requiring it to vanish for  $\eta = 2, 3, 6, 7$  results in  $E^2 = |\mathbf{p}|^2 + m^2$ .

The  $\tau = b, c$  sectors, if (wrongly) imposed with the vanishing of,  $\gamma_\mu \psi^\mu(\mathbf{p})$  and  $p_\mu \psi^\mu(\mathbf{p})$ , results in kinematically acausal dispersion relation (i.e., in  $E^2 \neq |\mathbf{p}|^2 + m^2$ ). This could be the source of the well-known problems of the Rarita-Schwinger framework as noted in works of Johnson and Sudarshan [10], and those of Velo and Zwanziger [11]. In this context one may wish to recall that interactions can induce transitions between different  $\tau$  sectors.

The analysis for all the  $\tau$  sectors of the  $\psi^\mu(\mathbf{p})$  can be summarized in the following table:

$\tau$	$p_\mu \psi^\mu(\mathbf{p})$	$\gamma_\mu \psi^\mu(\mathbf{p})$	$\widetilde{\mathcal{W}}_\mu^{(1/2,1/2)} \psi^\mu(\mathbf{p})$	$\lambda_\tau$	$\beta_\tau$	$\alpha_\tau$	Remarks
$a$	$= 0$	$= 0$	$\neq 0$	$15/4$	$2$	$1$	Rarita-Schwinger Sector
$b$	$= 0$	$\neq 0$	$\neq 0$	$3/4$	$2$	$-2$	
$c$	$\neq 0$	$\neq 0$	$= 0$	$3/4$	$0$	$0$	

The table clearly illustrates that there is no particular reason – except (the unjustified) insistence that each particle of nature be associated with a definite spin – to favor one  $\tau$  sector over the other. Each of the  $\tau$  sectors is endowed with specific properties. The Rarita-Schwinger sector has no more, or no less, physical significance than the other two sectors. While, for instance, the Rarita-Schwinger sector can be characterized by vanishing of the  $p_\mu \psi^\mu(\mathbf{p})$  and  $\gamma_\mu \psi^\mu(\mathbf{p})$ ; the  $\tau = c$  sector is uniquely characterized by vanishing of  $\widetilde{\mathcal{W}}_\mu^{(1/2,1/2)} \psi^\mu(\mathbf{p})$ . The  $\tau = b$  sector allows for vanishing of  $p_\mu \psi^\mu(\mathbf{p})$  only.

Except for the rest frame, the  $\psi_\tau^\mu(\mathbf{p})$ , in general, are not eigenstates of the  $\mathbf{J}^2$  for representation space (15). Instead, the three  $\tau$  sectors of the representation space under consideration correspond to the following inertial-frame independent values of the associated dragged second Casimir invariant:

$$\widetilde{C}_2^{[(1/2,0) \oplus (0,1/2)] \otimes (1/2,1/2)} \psi_\tau(\mathbf{p}) = -m^2 \lambda_\tau \psi_\tau(\mathbf{p}). \quad (39)$$

Note that the latter equation is a matrix equation in the Lorentz index space. For this reason the index  $\mu$  of the spinor-vector does not show up. Yet,  $\psi_\tau(\mathbf{p})$  is still the vector spinor and should not be confused with an ordinary Dirac spinor. Also there is no summation over the index  $W^2$ -sector index  $\tau$ . In fact, the defining Eq. (39) of the various  $\widetilde{C}_2^{[(1/2,0) \oplus (0,1/2)] \otimes (1/2,1/2)}$  sectors of  $\psi^\mu(\mathbf{p})$  translates into an equation for  $\gamma_\mu \psi_\tau^\mu(\mathbf{p})$ . To establish it, we note that  $\widetilde{C}_2^{[(1/2,0) \oplus (0,1/2)] \otimes (1/2,1/2)}$  is given as the squared sum of the (dragged) Pauli-Lubanski pseudovectors  $\widetilde{W}^{(1/2,0) \oplus (0,1/2)}$  and  $\widetilde{W}^{(1/2,1/2)}$ . In other words, one has

$$\widetilde{C}_2^{[(1/2,0) \oplus (0,1/2)] \otimes (1/2,1/2)} \psi_\tau(\mathbf{p}) = \left( \widetilde{W}^{(1/2,0) \oplus (0,1/2)} + \widetilde{W}^{(1/2,1/2)} \right)^2 \psi_\tau(\mathbf{p}). \quad (40)$$

The action of  $\widetilde{C}_2^{[(1/2,0) \oplus (0,1/2)] \otimes (1/2,1/2)}$  upon  $\psi_\tau(\mathbf{p})$  (with  $\tau = a, b, c$ ) takes the form

$$\begin{aligned} \widetilde{C}_2^{[(1/2,0) \oplus (0,1/2)] \otimes (1/2,1/2)} \psi_\tau(\mathbf{p}) &= \left( \widetilde{C}_2^{(1/2,0) \oplus (0,1/2)} + \widetilde{C}_2^{(1/2,1/2)} \right. \\ &\quad \left. + 2\widetilde{W}_\beta^{(1/2,0) \oplus (0,1/2)} \widetilde{W}^{\beta(1/2,1/2)} \right) \psi_\tau(\mathbf{p}) \\ &= \left( -m^2 \frac{1}{2} \left( \frac{1}{2} + 1 \right) - m^2 \beta_\tau - m^2 \alpha_\tau \right) \psi_\tau(\mathbf{p}) \\ &= -m^2 \lambda_\tau \psi_\tau(\mathbf{p}), \quad \beta_\tau = 0, 2. \end{aligned} \quad (41)$$

The latter equation shows that the values of  $\alpha_\tau$  and  $\beta_\tau$ , which in turn determine the eigenvalues of  $2\widetilde{W}^{\beta(1/2,1/2)}\widetilde{W}_\beta^{(1/2,0)\oplus(0,1/2)}$  and  $\widetilde{C}^{(1/2,1/2)}$  with respect to  $\psi_\tau(\mathbf{p})$  are well suited to label the various  $\widetilde{C}_2^{[(1/2,0)\oplus(0,1/2)]\otimes(1/2,1/2)}$  sectors. Indeed, one can replace Eq. (39) by

$$2\widetilde{W}^{\beta(1/2,1/2)}\widetilde{W}_\beta^{(1/2,0)\oplus(0,1/2)}\psi_\tau(\mathbf{p}) = -m^2\alpha_\tau\psi_\tau(\mathbf{p}),$$

$$\alpha_\tau = \lambda_\tau - \beta_\tau - \frac{3}{4}. \quad (42)$$

Insertion of the explicit expression for  $\widetilde{W}^{(1/2,0)\oplus(0,1/2)}$  into (42) and usage of the Dirac equation leads to

$$2\widetilde{W}_\nu^{(1/2,1/2)}\left(-\frac{1}{4}\gamma_5[\not{p}, \gamma^\nu]_-\right)\psi_\tau(\mathbf{p}) = -m^2\alpha_\tau\psi_\tau(\mathbf{p}),$$

$$\widetilde{W}_\nu^{(1/2,1/2)}\gamma_5(\not{p}\gamma^\nu - \gamma^\nu\not{p})\psi_\tau(\mathbf{p}) = 2m^2\alpha_\tau\psi_\tau(\mathbf{p})$$

$$\widetilde{W}_\nu^{(1/2,1/2)}\gamma_5(2g^{\alpha\nu}p_\lambda - 2\gamma^\nu\not{p})\psi_\tau(\mathbf{p}) = 2m^2\alpha_\tau\psi_\tau(\mathbf{p})$$

$$\widetilde{W}^{(1/2,1/2)}\cdot p\gamma_5\psi_\tau(\mathbf{p}) + \widetilde{W}^{(1/2,1/2)}\cdot\gamma\gamma_5m\psi_\tau(\mathbf{p}) = m^2\alpha_\tau\psi_\tau(\mathbf{p})$$

$$\left(\widetilde{W}^{(1/2,1/2)}\cdot p + \widetilde{W}^{(1/2,1/2)}\cdot\gamma\right)\gamma_5m\psi_\tau(\mathbf{p}) = m^2\alpha_\tau\psi_\tau(\mathbf{p}) \quad (43)$$

In taking Lorentz contraction of both sides of the last equation with  $\gamma^\epsilon$  and in accounting for  $\widetilde{W}^{(1/2,1/2)}\cdot p = 0$ , one arrives at the following equation for the  $\gamma\cdot\psi_\tau(\mathbf{p})$  spinor

$$\frac{1}{m\alpha_\tau}\gamma^\epsilon\left(\widetilde{W}_{\epsilon\eta}^{(1/2,1/2)}\cdot\gamma\gamma_5\right)\psi_\tau^\eta(\mathbf{p}) = \gamma\cdot\psi_\tau(\mathbf{p}). \quad (44)$$

The non-relativistic counterpart of Eq. (44) reads

$$\left(1 + \frac{2}{\alpha_\tau}\right)\boldsymbol{\sigma}\cdot\psi_\tau(\mathbf{0}) = 0. \quad (45)$$

For  $\alpha_a = 1$  one finds  $\boldsymbol{\sigma}\cdot\psi_a(\mathbf{0}) = 0$  (corresponding to  $\gamma\cdot\psi_a = 0$ ) while for  $\alpha_b = -2$ , where the numerical factor in (45) vanishes, one encounters  $\boldsymbol{\sigma}\cdot\psi_b(\mathbf{0}) \neq 0$  (corresponding to  $\gamma\cdot\psi_b \neq 0$ ). In our opinion, the troubles with the supplementary conditions in the Rarita-Schwinger framework is that after gauging, the full equation (44) does not reduce any longer to  $\gamma\cdot\psi(\mathbf{p})$ .

For each one of the  $\tau$  sectors, the  $\lambda_\tau$ ,  $\beta_\tau$ , and  $\alpha_\tau$  are given in the table above. Stated differently, the boosted  $\tau = b, c$  sectors do not carry spin one half. Similarly, the,  $\tau = a$ , sector is not a spin three half sector. The consequence is that the boosted  $\tau = b, c$  sector, in particular, should not be treated as a Dirac representation space. The correct wave equation for  $\psi^\mu(\mathbf{p})$  is:

$$\left[(\gamma_\mu p^\mu \pm mI_4)\right] \otimes \left(\Lambda_{\mu\nu}p^\mu p^\nu \pm m^2I_4\right)_{\eta\epsilon} \psi^\epsilon(\mathbf{p}) = 0. \quad (46)$$

In the standard Rarita-Schwinger framework  $\partial_\mu\psi^\mu(x)$  and  $\gamma_5\gamma_\mu\psi^\mu(x)$  do indeed behave as Dirac spinors, and do indeed satisfy the Dirac equation. However, they are not identical to the  $\tau = b, c$  sectors (which do not carry a characterization in terms of spin one half). If one (mistakenly) makes this identification, and sets  $\partial_\mu\psi^\mu(x)$  and  $\gamma_5\gamma_\mu\psi^\mu(x)$  to zero, one introduces an element of kinematic acausality. The *covariant* quantum numbers that are appropriate for labeling the basis vectors of the spinor-vector (and consequently for carrying out the quantization procedure) are

$$|\lambda_\tau \beta_\tau h\rangle, \quad (47)$$

where  $h$  is the eigenvalue of the helicity ( $\mathbf{J}\cdot\hat{\mathbf{p}}$ ) operator in the  $\tau$  sector under consideration.

### 3.3. Interpretation of the massive gravitino as a particle of multiple spin

If one is to respect the mathematical completeness of the spinor-vector representation space associated with  $\psi^\mu(x)$ , the Rarita-Schwinger framework cannot be considered to describe the full physical content of the representation space associated with a massive gravitino. This circumstance is akin to Dirac's observation that a part of a representation space [which would have violated the completeness of the  $(1/2, 0) \oplus (0, 1/2)$ ] cannot be "projected out" without introducing certain mathematical inconsistencies, and losing its physical content (i.e. antiparticles, or particles). Further, the same qualitative remarks apply to the  $(1/2, 1/2)$  representation space when in the Proca framework one only confines to the divergence-less vectors. The "projecting out" of the divergence-full vector, throws away the Stückelberg contribution to the propagator, and in addition leaves the  $(1/2, 1/2)$  representation space mathematically incomplete. Now, we suggest that for the representation space defined by Eq. (15), one needs to consider all three  $\tau$  sectors of  $\psi^\mu(x)$  as physical, and necessary for its mathematical consistency. The suggested framework already carries consistency with the known data on the  $N$  and  $\Delta$  resonances, and asks that massive gravitino be considered as an object that is better described by the eigenvalues of the dragged second Casimir operator. In its rest frame it is endowed with a spin three half, and two spin half, components. A spin measurement for unpolarized ensemble of massive gravitinos at rest would yield the results  $3/2$  with probability one half, and  $1/2$  with probability one half. The latter probability is distributed uniformly, i.e. as one quarter, over each of the,  $\tau = b$ , and,  $\tau = c$ , sectors.

## 4. Outlook

The systematic and self-contained description of the spacetime structure of fundamental particles presented in this talk calls for a more detailed analysis of the consequences of the multi-spin character of the massive gravitino for the renormalizability of supergravity and the phenomenology of the early universe. We also conjectured in [13] that gauge bosons of a quantum theory of gravity shall have a well pronounced multi-spin character. In particular, the above considerations suggest that a natural extension of this work to finite-range gravity shall endow the graviton with spins 0, 1, and 2 components, a possibility that has been partly entertained only very recently in [14, 15].

## Acknowledgments

Work supported by Consejo Nacional de Ciencia y Tecnología (CONACyT, Mexico) under grant number 32067-E. The organizers of Beyond02 deserve special thanks for having created that pervading friendly atmosphere which makes scientific events so stimulating.

## References

- [1] D. V. Ahluwalia and M. Kirchbach, *Mod. Phys. Lett.* **A16**, 1377 (2001).
- [2] M. Kirchbach and D. V. Ahluwalia, *Phys. Lett.* **B529**, 124 (2002).
- [3] D. V. Ahluwalia, *Found. Phys.* **29**, 527 (1998).
- [4] S. Weinberg, *The quantum theory of fields, Vol. I and III* (Cambridge University Press, Cambridge, 1995 and 2000).
- [5] M. Veltman, *Int. J. Mod. Phys.* **A15**, 4557 (2000).
- [6] M. Kirchbach, *Mod. Phys. Lett.* **A12**, 2373 (1997);  
M. Kirchbach, *Nucl. Phys.* **A689**, 157c (2001);  
M. Kirchbach, M. Moshinsky, and Yu. F. Smirnov, *Phys. Rev.* **D64**, 114005 (2001).
- [7] Particle Data Group, *Eur. Phys. J.* **C15**, 1 (2000).

- [8] T. Moroi, Ph. D. thesis (Tohoku University, Japan, 1995), hep-ph/9503210.
- [9] W. Rarita and J. Schwinger, Phys. Rev. **60**, 61 (1941).
- [10] K. Johnson and E. C. G. Sudarshan, Ann. Phys. (N. Y. )**13**, 126 (1961)
- [11] G. Velo and D. Zwanzinger, Phys. Rev. **186**, 1337 (1969).
- [12] G. Höhler, in *Pion-Nucleon Scattering* (Springer Publishers, Heidelberg, 1983), Landolt-Börnstein Vol. I/9b2, Ed. H. Schopper.
- [13] D. V. Ahluwalia, N. Dadhich, and M. Kirchbach, “On the spin of gravitational bosons,” gr-qc/0205072.
- [14] M. Visser, Gen. Rel. Grav. **30**, 1717 (1998).
- [15] S. V. Babak, L. P. Grishchuk, “Finite-Range Gravity and Its Role in Gravitational Waves, Black Holes and Cosmology,” gr-qc/0209006.