Running Neutrino Masses, Leptonic Mixing Angles and CP-Violating Phases: From M_Z to Λ_{GUT}

Jianwei Mei

CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, China and Institute of High Energy Physics, Chinese Academy of Sciences, P.O. Box 918 (4), Beijing 100049, China * (Electronic address: jwmei@mail.ihep.ac.cn)

Abstract

We derive renormalization group equations for neutrino masses, leptonic mixing angles and CP-violating phases running at energies above the seesaw threshold, both in the Standard Model and in the Minimal Supersymmetric Standard Model extended with three right-handed neutrinos. With these equations, we carry out a systematic study of the radiative correction that may arise to neutrino parameters, via their renormalization group evolution from the electroweak scale (M_Z) to the scale of Grand Unified theories (Λ_{GUT}). We study in detail three typically interesting neutrino mass patterns: normal hierarchy, near degeneracy and inverted hierarchy. Magnitudes of possible corrections in each case are carefully investigated. We also emphasize the significance of CP-violating phases in controlling the evolution behavior of all neutrino parameters.

PACS number(s): 14.60.Pq, 13.10.+q, 25.30.Pt

Typeset using $\text{REVT}_{\text{E}}X$

^{*}Mailing address

I. INTRODUCTION

Experimental information on neutrino masses and mixings has opened up a new playground for efforts [1] of a better understanding of the Yukawa couplings in the Standard Model (SM) and its extensions. In the lepton sector, it is conceptually natural to use the elegant seesaw mechanism [2–4] to give masses to the SM neutrinos. In the most simple version of this mechanism, the dimension 5 neutrino mass term is the low energy relic of some more fundamental theories with very heavy right-handed neutrinos. So in this framework, we often need to relate physics at vastly different energy scales.

The only way to compare the high energy theoretical prediction and the low energy experimental observation is to use renormalization group equations (RGEs). It has been found that corrections arising from renormalization group (RG) evolution can be very significant for leptonic mixing angles and neutrino mass splitting, especially in the case of nearly degenerate left-handed neutrinos. So in principle, the RG correction should not be neglected in the discussion of models suggested at high energy scales. With RGEs derived in Refs. [5–7], early discussions [8–16] of the RG running effect are mainly concerned with neutrino mass (or Yukawa coupling) matrices. The evolution of neutrino masses, leptonic mixing angles and possible CP-violating phases is studied by diagonalizing the relevant Yukawa matrices at different energy scales, and the behaviors are discussed in a numerical or semi-analytical way. However, authors of Refs. [17-20] have also emphasized the significance of RGEs for individual neutrino parameters. Such equations not only make it possible to predict the evolution behavior of each parameter [21–24], but also can help people appreciate interesting features such as the existence of (pseudo-) fixed points in the evolution of mixing angles and CP-violating phases [14,16,17,25,26]. The derivation of these equations below the seesaw threshold has been done in Refs. [17,18,27,23]. And based on such equations, a comprehensive study of the RG evolution of neutrino parameters from the electroweak scale to the seesaw threshold has been carried out in Ref. [27].

However, RG corrections above the seesaw threshold sometimes are as important as or even more significant than those below the threshold [8,15,28–30]. Since the physics responsible for neutrino mass generation is more likely to exist at the scale of Grand Unified theories, a systematic study of the RG correction above the seesaw threshold should be necessary. And this is one of our main concerns in this work. In much the same spirit as of Ref. [27], we derive RGEs for individual neutrino parameters running above the seesaw threshold, under the condition that eigenvalues of the Yukawa coupling matrix that connects leftand right-handed neutrinos are hierarchical. The contribution from the largest eigenvalue of this matrix is explicitly shown. By setting this contribution to zero, we can regain RGEs obtained earlier in Refs. [27,23], which are valid at energies below the seesaw threshold.

The second purpose of this work is to carry out a systematic study of the radiative correction that may arise via the RG evolution in the full energy range from the electroweak scale (M_Z) to the scale of Grand Unified theories (Λ_{GUT}) . To demonstrate main features of possible corrections, we study in detail three typically allowed neutrino mass patterns: normal hierarchy, near degeneracy and inverted hierarchy.

The paper is organized as follows. We write down *full* one-loop RGEs for individual neutrino masses, leptonic mixing angles and CP violating phases in Section II, with a brief discussion. Then in Section III, we carry out a systematic study of the correction that may

arise during the RG evolution from M_Z to Λ_{GUT} , in theories with the three neutrino mass patterns mentioned above. Section IV is devoted to a Summary.

II. ANALYTICAL FORMULAE FOR NEUTRINO PARAMETERS RUNNING ABOVE THE SEESAW THRESHOLD

Extended with three right-handed neutrinos, the Lagrangian giving mass to leptons in the SM is

$$-\mathcal{L}_{l} = \overline{E_{L}}HY_{l}l_{R} + \overline{E_{L}}H^{c}Y_{\nu}\nu_{R} + \frac{1}{2}\overline{\nu_{R}^{c}}M_{R}\nu_{R} + h.c.; \qquad (1)$$

and in the MSSM is

$$-\mathcal{L}_{l} = \overline{E_{L}}H_{1}Y_{l}l_{R} + \overline{E_{L}}H_{2}Y_{\nu}\nu_{R} + \frac{1}{2}\overline{\nu_{R}^{c}}M_{R}\nu_{R} + h.c., \qquad (2)$$

where, E_L , l_R and ν_R denote $SU(2)_L$ -doublets, right-handed charged leptons and righthanded neutrinos, respectively. Both in Eqs.(1) and (2), the scale of M_R is expected to be extremely high, since there is not a protective symmetry. Around this energy scale, mass eigenstates of M_R are successively integrated out at their respective masses ($M_1 < M_2 < M_3$), giving rise to a series of effective theories at different energy scales [28]. Then at energies below the lightest right-handed neutrino mass, we obtain the dimension 5 effective mass term for left-handed neutrinos

$$-\mathcal{L}_{\nu} = -\frac{1}{2}\overline{E_L}\Phi \cdot \kappa \cdot \Phi^T E_L^c + h.c., \qquad (3)$$

where κ is the effective Yukawa coupling matrix, and Φ is H^c in the SM but is H_2 in the MSSM. Since κ is calculated from Y_{ν} and M_R by decoupling right-handed neutrinos at successive energy scales, step by step, the relation between κ and Y_{ν} , M_R is complicated. Only in the most simplified procedure when all right-handed neutrinos are decoupled at a common scale, can we obtain (at that chosen scale) a simple equation because of the tree-level matching condition [28]:

$$\kappa = Y_{\nu} M_R^{-1} Y_{\nu}^T . \tag{4}$$

Then, when the Higgs field acquires a non-zero vacuum expectation value $\langle \Phi \rangle = (v \ 0)^T$ during the electroweak symmetry breaking, Eq.(3) yields an effective mass matrix $M_{\nu} = v^2 \kappa$ for left-handed neutrinos. In the SM, $v \simeq 174$ GeV; and in the MSSM, $v \simeq 174 \sin \beta$ GeV.

Since right-handed neutrinos are to be decoupled at their respective thresholds, it will be too complicated to derive RGEs for neutrino parameters between these thresholds. So we shall be less ambitious than solving the whole problem, but shall simplify it by (a) decoupling all right-handed neutrinos at a *common* scale, which we take to be M_3 , and (b) limiting our derivation only to the case when eigenvalues of Y_{ν} are *hierarchical*. Here, assumption (a) can be justified when the RG evolution through right-handed neutrino thresholds is not too dramatic (As has been demonstrated in Refs. [15,28], this is not always the case). And assumption (b) is also well motivated since in a large class of high energy models considered in literature (such as those based on U(1) symmetry or those base on SO(10) Grand Unified Theories), there is often a certain similarity or even identification between Y_{ν} and the upquark Yukawa coupling matrix. Such a similarity of matrices should lead to some likeness between their eigenvalues.

One-loop RGEs for κ running from $M_{\rm Z}$ to M_1 and those for Y_{ν} and M_R running through right-handed neutrino thresholds to $\Lambda_{\rm GUT}$ have been given in Ref. [28]. For readers' convenience, we have collected a part of them together with those for Y_l in Appendix A. To discuss the RG evolution of neutrino parameters in the full energy range from $M_{\rm Z}$ to $\Lambda_{\rm GUT}$, it is convenient to make use of κ also at energies above the seesaw threshold [24,30]. In this energy range, we find from Eqs.(4), (A.6) and (A.7)

$$16\pi^2 \frac{\mathrm{d}\kappa}{\mathrm{d}t} = \alpha_\kappa \kappa + N_\kappa \kappa + \kappa N_\kappa^T,\tag{5}$$

where $t = \ln \mu$ with μ being the energy scale, and details of α_{κ} and N_{κ} are given in Eq.(A.15) in the Appendix.

For neutrino masses, leptonic mixing angles and CP-violating phases, κ is diagonalized by a unitary matrix U_{κ} :

$$\kappa = U_{\kappa} \kappa' U_{\kappa}^{T}; \quad \kappa' = \operatorname{diag}\left\{k_{1}, k_{2}, k_{3}\right\}, \tag{6}$$

where k_i (for i = 1, 2, 3) at M_Z are proportional to left-handed neutrino masses : $k_i (M_Z) \equiv m_i (M_Z) / v^2$. At energy scales below M_1 , Y_l is always diagonal during the RG evolution, if it is diagonal at the beginning. In such a basis, the leptonic mixing matrix is $U_{\rm MNS} \equiv U_{\kappa}$. However, Y_l can not be kept diagonal above the seesaw threshold when Eq.(A.5) is used for its RG evolution. In this case, there is the contribution to $U_{\rm MNS}$ from diagonalizing $H_l = Y_l Y_l^{\dagger}$:

$$H_{l} = U_{l}H_{l}^{\prime}U_{l}^{\dagger}; \quad H_{l}^{\prime} = \text{diag}\left\{y_{e}^{2}, y_{\mu}^{2}, y_{\tau}^{2}\right\},$$
(7)

$$\implies U_{\rm MNS} = U_l^{\dagger} U_{\kappa}. \tag{8}$$

For the MNS matrix, a convenient parametrization can be found in Ref. [31]

$$U_{\rm MNS} = \begin{pmatrix} 1 & & \\ c_y & s_y \\ -s_y & c_y \end{pmatrix} \begin{pmatrix} c_z & s_z \\ e^{-i\delta} \\ -s_z & c_z \end{pmatrix} \begin{pmatrix} c_x & s_x \\ -s_x & c_x \\ & & 1 \end{pmatrix} \begin{pmatrix} e^{i\rho} & & \\ e^{i\sigma} & & \\ & & 1 \end{pmatrix} = \begin{pmatrix} c_x c_z & & c_z s_x & s_z \\ -c_y s_x e^{-i\delta} - c_x s_y s_z & c_x c_y e^{-i\delta} - s_x s_y s_z & c_z s_y \\ s_x s_y e^{-i\delta} - c_x c_y s_z & -c_x s_y e^{-i\delta} - c_y s_x s_z & c_y c_z \end{pmatrix} \begin{pmatrix} e^{i\rho} & & \\ e^{i\sigma} & & \\ & & 1 \end{pmatrix},$$
(9)

where $c_x \equiv \cos \theta_x$, $s_x \equiv \sin \theta_x$ and so on. ¹The merit of this parametrization is that the Dirac phase δ does not appear in the neutrinoless double beta decay, while Majorana phases

¹Comparing to another parametrization used in the literature [32], apart from the notation of mixing angles $\theta_{12} \Leftrightarrow \theta_x$, $\theta_{23} \Leftrightarrow \theta_y$ and $\theta_{13} \Leftrightarrow \theta_z$, the main difference resides only in the Majorana phases: $\alpha_1 \Leftrightarrow 2(\rho - \delta)$, $\alpha_2 \Leftrightarrow 2(\sigma - \delta)$, while the Dirac phase δ is the same.

 ρ and σ do not contribute to the leptonic CP violation in neutrino oscillations. Thus two different types of phases can be separately studied in different types of experiments.

At energies below the seesaw threshold, RGEs for the running of left-handed neutrino masses (m_1, m_2, m_3) , leptonic mixing angles $(\theta_x, \theta_y, \theta_z)$ and CP-violating phases (δ, ρ, σ) have been derived in Refs. [17,18,27,23]. In order to obtain the same kind of formulae at energies above the seesaw threshold, we also need a parametrization of Y_{ν} . We find that the derivation is most straightforward if Y_{ν} is parameterized in the diagonal basis of κ by $(H_{\nu} \equiv Y_{\nu} Y_{\nu}^{\dagger})$:

$$H_{\nu} = U_{\nu} H_{\nu}' U_{\nu}^{\dagger}; \quad H_{\nu}' = y_{\nu}^2 \cdot \text{diag}\left\{r_1^2, r_2^2, 1\right\} , \qquad (10)$$

where (with $c_1 = \cos \theta_1$, $s_1 = \sin \theta_1$ and so on)

$$U_{\nu} = \begin{pmatrix} e^{i\phi_1} & & \\ & e^{i\phi_2} & \\ & & 1 \end{pmatrix} \begin{pmatrix} 1 & & \\ & c_1 & s_1 \\ & -s_1 & c_1 \end{pmatrix} \begin{pmatrix} c_2 & & s_2 \\ & e^{-i\delta_{\nu}} & \\ -s_2 & & c_2 \end{pmatrix} \begin{pmatrix} c_3 & s_3 & \\ -s_3 & c_3 & \\ & & 1 \end{pmatrix}.$$
 (11)

As mentioned above, we shall concentrate on cases in which eigenvalues of Y_{ν} are hierarchical, i.e. $r_1^2 << r_2^2 << 1$. So the contribution of r_1^2 and r_2^2 to the running of neutrino parameters can always be neglected. This is equivalent to taking $r_1^2 = r_2^2 = 0$ when deriving RGEs. From Eqs.(10) and (11), it is obvious that only $\theta_1, \theta_2, \phi_1$ and ϕ_2 in U_{ν} contribute to the evolution of parameters $m_1, m_2, m_3, \theta_x, \theta_y, \theta_z, \delta, \rho$ and σ .

Before writing down all the analytical formulae, we remark that RG corrections to mixing angles and CP-violating phases come from three separable sources. To clarify this point, we need Eq.(A.15): $N_{\kappa} = C_{\kappa}^{l}H_{l} + C_{\kappa}^{\nu}H_{\nu}$. In Eq.(5), only N_{κ} contributes to the evolution of mixing angles and CP-violating phases. So each of the two terms in N_{κ} is a source of RG corrections. Also, there is a contribution from diagonalizing Y_{l} , and this is the third one.

- The contribution from $C_{\kappa}^{l}H_{l}$ in N_{κ} is proportional to $C_{\kappa}^{l}y_{\tau}^{2}$ (we have omitted the contributions from y_{e}^{2} and y_{μ}^{2} for obvious reasons). This contribution is *exactly the same as* that governs the evolution of neutrino parameters at energies below the seesaw threshold. Analytical formulae of this contribution have been derived and extensively discussed in Refs. [27,23].
- The contribution from $C_{\kappa}^{\nu}H_{\nu}$ in N_{κ} is proportional to $C_{\kappa}^{\nu}y_{\nu}^{2}$. While y_{τ} can be of $\mathcal{O}(1)$ only in the MSSM when $\tan \beta$ is large, it is quite natural for y_{ν} to be of the same magnitude as the top Yukawa coupling. Furthermore, just like the contribution from $C_{\kappa}^{l}H_{l}$, the contribution from $C_{\kappa}^{\nu}H_{\nu}$ can also be resonantly enhanced when eigenvalues of κ are nearly degenerate. This is because both contributions contain such enhancing factors as ζ_{ij}^{-1} (for i < j; i, j = 1, 2, 3). Note that

$$\zeta_{ij} \equiv \frac{k_i - k_j}{k_i + k_j} ; \quad i, j = 1, 2, 3 .$$
(12)

• The third contribution comes from diagonalizing Y_l and is proportional to $C_l^{\nu} y_{\nu}^2$. Different from that of $C_{\kappa}^{\nu} H_{\nu}$, there are no enhancing factors in this contribution other

than functions of mixing angles, such as $s_z^{-1} \equiv (\sin \theta_z)^{-1}$ or $s_x^{-1} \equiv (\sin \theta_x)^{-1}$ etc., which appear mostly in RGEs of CP-violating phases and are important only when s_z (or s_x etc.) << O(1).

Now following the same procedure as described in Refs. [18,27] but including contributions from Y_{ν} and Y_l (yet with the above explained simplifications), we obtain the following full one-loop RGEs for left-handed neutrino masses (m_1, m_2, m_3) , leptonic mixing angles $(\theta_x, \theta_y, \theta_z)$ and CP-violating phases (δ, ρ, σ) running at energies above the seesaw threshold.

²For the running of left-handed neutrino masses (at M_Z : $m_i \equiv v^2 k_i$; i = 1, 2, 3):

$$k_{i} = a_{i}k_{i} , \quad i = 1, 2, 3 ;$$

$$a_{1} \equiv \alpha_{\kappa} + 2C_{\kappa}^{l}y_{\tau}^{2} \left(s_{x}^{2}s_{y}^{2} - 2c_{\delta}c_{x}c_{y}s_{x}s_{y}s_{z} + c_{x}^{2}c_{y}^{2}s_{z}^{2}\right) + 2C_{\kappa}^{\nu}y_{\nu}^{2}s_{2}^{2} ;$$

$$a_{2} \equiv \alpha_{\kappa} + 2C_{\kappa}^{l}y_{\tau}^{2} \left(c_{x}^{2}s_{y}^{2} + 2c_{\delta}c_{x}c_{y}s_{x}s_{y}s_{z} + c_{y}^{2}s_{x}^{2}s_{z}^{2}\right) + 2C_{\kappa}^{\nu}y_{\nu}^{2}c_{2}^{2}s_{1}^{2} ;$$

$$a_{3} \equiv \alpha_{\kappa} + 2C_{\kappa}^{l}y_{\tau}^{2}c_{y}^{2}c_{z}^{2} + 2C_{\kappa}^{\nu}y_{\nu}^{2}c_{1}^{2}c_{2}^{2} . \qquad (13)$$

For the running of leptonic mixing angles and CP-violating phases $(c_{\delta} \equiv \cos \delta, s_{\delta} \equiv \sin \delta, c_{(\delta-\rho)} \equiv \cos (\delta-\rho), s_{(\delta-\rho)} \equiv \sin (\delta-\rho)$ and so on):

$$\begin{split} \dot{\theta}_{x} &= C_{\kappa}^{l} y_{\tau}^{2} \left\{ \frac{c_{(\rho-\sigma)}}{\zeta_{12}} \left[c_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(c_{(\delta+\rho-\sigma)} c_{x}^{2} - c_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \right. \\ &+ \zeta_{12} \cdot s_{(\rho-\sigma)} \left[s_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(s_{(\delta+\rho-\sigma)} c_{x}^{2} + s_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \\ &- \left[\frac{c_{\rho}}{\zeta_{13}} \left(c_{(\delta-\rho)} s_{x} s_{y} - c_{\rho} c_{x} c_{y} s_{z} \right) - \zeta_{13} \cdot s_{\rho} \left(s_{(\delta-\rho)} s_{x} s_{y} + s_{\rho} c_{x} c_{y} s_{z} \right) \right] c_{y} s_{x} s_{z} \\ &- \left[\frac{c_{\sigma}}{\zeta_{23}} \left(c_{(\delta-\sigma)} c_{x} s_{y} + c_{\sigma} c_{y} s_{x} s_{z} \right) - \zeta_{23} \cdot s_{\sigma} \left(s_{(\delta-\sigma)} c_{x} s_{y} - s_{\sigma} c_{y} s_{x} s_{z} \right) \right] c_{x} c_{y} s_{z} \right\} \\ &+ C_{\kappa}^{\nu} y_{\nu}^{2} \left\{ - \left(\frac{c_{(\rho-\sigma)} c_{(\phi_{1}-\phi_{2})}}{\zeta_{12}} - \zeta_{12} \cdot s_{(\rho-\sigma)} s_{(\phi_{1}-\phi_{2})} \right) c_{2} s_{1} s_{2} \right. \\ &- \left(\frac{c_{\rho} c_{\phi_{1}}}{\zeta_{13}} - \zeta_{13} \cdot s_{\rho} s_{\phi_{1}} \right) \frac{c_{1} c_{2} s_{2} s_{x} s_{z}}{c_{z}} + \left(\frac{c_{\sigma} c_{\phi_{2}}}{\zeta_{23}} - \zeta_{23} \cdot s_{\sigma} s_{\phi_{2}} \right) \frac{c_{1} c_{2}^{2} c_{x} s_{1} s_{z}}{c_{z}} \right\} \\ &+ C_{l}^{\nu} y_{\nu}^{2} \left\{ \left[c_{(\rho+\phi_{1})} s_{2} s_{x} - c_{(\sigma+\phi_{2})} c_{2} c_{x} s_{1} \right] \frac{c_{1} c_{2} s_{z}}{c_{z}} - c_{x} s_{x} \left(c_{2}^{2} s_{1}^{2} - s_{2}^{2} \right) \right\} ; \end{split}$$

$$\begin{split} \dot{\theta}_y &= C_{\kappa}^l y_{\tau}^2 \left\{ \left[\frac{c_{(\delta-\rho)}}{\zeta_{13}} \left(c_{(\delta-\rho)} s_x s_y - c_{\rho} c_x c_y s_z \right) + \zeta_{13} \cdot s_{(\delta-\rho)} \left(s_{(\delta-\rho)} s_x s_y + s_{\rho} c_x c_y s_z \right) \right] c_y s_x \\ &+ \left[\frac{c_{(\delta-\sigma)}}{\zeta_{23}} \left(c_{(\delta-\sigma)} c_x s_y + c_{\sigma} c_y s_x s_z \right) + \zeta_{23} \cdot s_{(\delta-\sigma)} \left(s_{(\delta-\sigma)} c_x s_y - s_{\sigma} c_y s_x s_z \right) \right] c_x c_y \right\} \end{split}$$

²Throughout this work, for what ever F: $\dot{F} \equiv 16\pi^2 \frac{dF}{dt}$; $t = \ln\mu$, with μ being the energy scale.

$$+C_{\kappa}^{\nu}y_{\nu}^{2}\left\{\left(\frac{c_{(\delta-\rho)}c_{\phi_{1}}}{\zeta_{13}}+\zeta_{13}\cdot s_{(\delta-\rho)}s_{\phi_{1}}\right)\frac{c_{1}c_{2}s_{2}s_{x}}{c_{z}}-\left(\frac{c_{(\delta-\sigma)}c_{\phi_{2}}}{\zeta_{23}}+\zeta_{23}\cdot s_{(\delta-\sigma)}s_{\phi_{2}}\right)\frac{c_{1}c_{2}^{2}c_{x}s_{1}}{c_{z}}\right\}$$

$$+C_{l}^{\nu}y_{\nu}^{2}\left\{c_{y}s_{y}\left(-c_{1}^{2}c_{2}^{2}c_{z}^{2}+s_{2}^{2}\left(s_{x}^{2}-c_{x}^{2}s_{z}^{2}\right)+c_{2}^{2}s_{1}^{2}\left(c_{x}^{2}-s_{x}^{2}s_{z}^{2}\right)\right)$$

$$+2c_{\delta}c_{x}c_{y}^{2}\left(c_{2}^{2}s_{1}^{2}-s_{2}^{2}\right)s_{x}s_{z}+2\left(c_{(\rho+\phi_{1})}c_{x}s_{2}+c_{(\sigma+\phi_{2})}c_{2}s_{1}s_{x}\right)c_{1}c_{2}c_{y}c_{z}s_{y}s_{z}$$

$$+\left(c_{(\delta-\rho-\phi_{1})}s_{2}s_{x}-c_{(\delta-\sigma-\phi_{2})}c_{2}c_{x}s_{1}\right)\frac{c_{1}c_{2}}{c_{z}}\left(c_{z}^{2}\left(c_{y}^{2}-s_{y}^{2}\right)-s_{z}^{2}\right)$$

$$+2\left[-c_{(\rho-\sigma+\phi_{1}-\phi_{2})}c_{x}s_{x}s_{y}\left(1+s_{z}^{2}\right)\right]$$

$$+\left(c_{(\delta+\rho-\sigma+\phi_{1}-\phi_{2})}c_{x}^{2}-c_{(\delta-\rho+\sigma-\phi_{1}+\phi_{2})}s_{x}^{2}\right)c_{y}s_{z}\right]c_{2}c_{y}s_{1}s_{2}\right\};$$
(15)

$$\begin{aligned} \dot{\theta}_{z} &= C_{\kappa}^{l} y_{\tau}^{2} \left\{ -\left[\frac{c_{\rho}}{\zeta_{13}} \left(c_{(\delta-\rho)} s_{x} s_{y} - c_{\rho} c_{x} c_{y} s_{z} \right) - \zeta_{13} \cdot s_{\rho} \left(s_{(\delta-\rho)} s_{x} s_{y} + s_{\rho} c_{x} c_{y} s_{z} \right) \right] c_{x} c_{y} c_{z} \\ &+ \left[\frac{c_{\sigma}}{\zeta_{23}} \left(c_{(\delta-\sigma)} c_{x} s_{y} + c_{\sigma} c_{y} s_{x} s_{z} \right) - \zeta_{23} \cdot s_{\sigma} \left(s_{(\delta-\sigma)} c_{x} s_{y} - s_{\sigma} c_{y} s_{x} s_{z} \right) \right] c_{y} c_{z} s_{x} \right\} \\ &+ C_{\kappa}^{\nu} y_{\nu}^{2} \left\{ - \left(\frac{c_{\rho} c_{\phi_{1}}}{\zeta_{13}} - \zeta_{13} \cdot s_{\rho} s_{\phi_{1}} \right) c_{1} c_{2} c_{x} s_{2} - \left(\frac{c_{\sigma} c_{\phi_{2}}}{\zeta_{23}} - \zeta_{23} \cdot s_{\sigma} s_{\phi_{2}} \right) c_{1} c_{2}^{2} s_{1} s_{x} \right\} \\ &+ C_{l}^{\nu} y_{\nu}^{2} \left\{ \left(c_{x}^{2} s_{2}^{2} - c_{2}^{2} \left(c_{1}^{2} - s_{1}^{2} s_{x}^{2} \right) \right) c_{z} s_{z} - \left(c_{(\rho+\phi_{1})} c_{x} s_{2} + c_{(\sigma+\phi_{2})} c_{2} s_{1} s_{x} \right) c_{1} c_{2} \left(c_{z}^{2} - s_{z}^{2} \right) \\ &+ 2 c_{(\rho-\sigma+\phi_{1}-\phi_{2})} c_{2} c_{x} c_{z} s_{1} s_{2} s_{x} s_{z} \right\} ; \end{aligned}$$

$$(16)$$

$$\begin{split} \dot{\delta} &= C_{\kappa}^{l} y_{\tau}^{2} \left\{ \frac{s_{(\rho-\sigma)}}{\zeta_{12}} \left[c_{(\rho-\sigma)} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(c_{(\delta+\rho-\sigma)} c_{x}^{2} - c_{(\delta-\rho+\sigma)} s_{x}^{2} \right) \frac{c_{y} s_{y} s_{z}}{c_{x} s_{x}} \right] \right. \\ &- \zeta_{12} \cdot c_{(\rho-\sigma)} \left[s_{(\rho-\sigma)} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(s_{(\delta+\rho-\sigma)} c_{x}^{2} + s_{(\delta-\rho+\sigma)} s_{x}^{2} \right) \frac{c_{y} s_{y} s_{z}}{c_{x} s_{x}} \right] \\ &+ \zeta_{13}^{-1} \cdot \left(c_{(\delta-\rho)} s_{x} s_{y} - c_{\rho} c_{x} c_{y} s_{z} \right) \left[\frac{s_{\rho} c_{y}}{c_{x} s_{z}} \left(c_{x}^{2} - s_{x}^{2} s_{z}^{2} \right) - s_{(\delta-\rho)} \frac{s_{x}}{s_{y}} \left(c_{y}^{2} - s_{y}^{2} \right) \right] \\ &+ \zeta_{13} \cdot \left(s_{(\delta-\rho)} s_{x} s_{y} + s_{\rho} c_{x} c_{y} s_{z} \right) \left[\frac{c_{\rho} c_{y}}{c_{x} s_{z}} \left(c_{x}^{2} - s_{x}^{2} s_{z}^{2} \right) + c_{(\delta-\rho)} \frac{s_{x}}{s_{y}} \left(c_{y}^{2} - s_{y}^{2} \right) \right] \\ &- \zeta_{23}^{-1} \cdot \left(c_{(\delta-\sigma)} c_{x} s_{y} + c_{\sigma} c_{y} s_{x} s_{z} \right) \left[\frac{s_{\sigma} c_{y}}{s_{x} s_{z}} \left(s_{x}^{2} - c_{x}^{2} s_{z}^{2} \right) - c_{(\delta-\sigma)} \frac{c_{x}}{s_{y}} \left(c_{y}^{2} - s_{y}^{2} \right) \right] \right\} \\ &- \zeta_{23} \cdot \left(s_{(\delta-\sigma)} c_{x} s_{y} - s_{\sigma} c_{y} s_{x} s_{z} \right) \left[\frac{c_{\sigma} c_{y}}{s_{x} s_{z}} \left(s_{x}^{2} - c_{x}^{2} s_{z}^{2} \right) - c_{(\delta-\sigma)} \frac{c_{x}}{s_{y}} \left(c_{y}^{2} - s_{y}^{2} \right) \right] \right\} \\ &+ C_{\kappa}^{\nu} y_{\nu}^{2} \left\{ - \left(\frac{c_{(\phi+-\phi_{2})} s_{(\rho-\sigma)}}{\zeta_{12}} + \zeta_{12} \cdot c_{(\rho-\sigma)} s_{(\phi+-\phi_{2})} \right) \frac{c_{2} s_{1} s_{2}}{c_{x} s_{x}} \\ &+ \left[\frac{c_{\phi_{1}}}{\zeta_{13}} \left(s_{\rho} c_{y} s_{y} \left(c_{x}^{2} - s_{x}^{2} s_{z}^{2} \right) - s_{(\delta-\rho)} \left(c_{y}^{2} - s_{y}^{2} \right) c_{x} s_{x} s_{z} \right) \\ &+ \zeta_{13} \cdot s_{\phi_{1}} \left(c_{\rho} c_{y} s_{y} \left(c_{x}^{2} - s_{x}^{2} s_{z}^{2} \right) + c_{(\delta-\rho)} \left(c_{y}^{2} - s_{y}^{2} \right) c_{x} s_{x} s_{z} \right) \\ &+ \left[\frac{c_{\phi_{2}}}}{\zeta_{23}} \left(s_{\sigma} c_{y} s_{y} \left(s_{x}^{2} - c_{x}^{2} s_{z}^{2} \right) + s_{(\delta-\sigma)} \left(c_{y}^{2} - s_{y}^{2} \right) c_{x} s_{x} s_{z} \right) \right]$$

$$+\zeta_{23} \cdot s_{\phi_2} \left(c_{\sigma} c_y s_y \left(s_x^2 - c_x^2 s_z^2 \right) - c_{(\delta-\sigma)} \left(c_y^2 - s_y^2 \right) c_x s_x s_z \right) \right] \frac{c_1 c_2^2 s_1}{c_y c_z s_x s_y s_z} \right\}$$

$$+C_l^{\nu} y_{\nu}^2 \left\{ -2s_{\delta} \left(c_2^2 s_1^2 - s_2^2 \right) \frac{c_x c_y s_x s_z}{s_y} + \left[s_{(\rho+\phi_1)} s_2 s_x \left(c_x^2 \left(c_z^2 - s_z^2 \right) + s_x^2 s_z^2 \right) \right. \right. \\ \left. + s_{(\sigma+\phi_2)} c_2 c_x s_1 \left(c_x^2 s_z^2 + s_x^2 \left(c_z^2 - s_z^2 \right) \right) \right] \frac{c_1 c_2}{c_x c_z s_x s_z} \right. \\ \left. - \left[s_{(\delta-\rho-\phi_1)} s_2 s_x \left(c_z^2 - \left(c_y^2 - s_y^2 \right) s_z^2 \right) - s_{(\delta-\sigma-\phi_2)} c_2 c_x s_1 \left(1 - 2 c_y^2 s_z^2 \right) \right] \frac{c_1 c_2}{c_y c_z s_y} \right. \\ \left. - s_{(\rho-\sigma+\phi_1-\phi_2)} \left(c_x^2 - s_x^2 \right) \frac{c_2 s_1 s_2}{c_x s_x} \right. \\ \left. - 2 \left[s_{(\delta+\rho-\sigma+\phi_1-\phi_2)} - 2 c_{(\rho-\sigma+\phi_1-\phi_2)} s_{\delta} s_x^2 \right] \frac{c_2 c_y s_1 s_2 s_z}{s_y} \right\} ; \qquad (17)$$

$$\begin{split} \dot{\rho} &= C_{\kappa}^{l} y_{\tau}^{2} \left\{ \frac{s_{(\rho-\sigma)}}{\zeta_{12}} \left[c_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(c_{(\delta+\rho-\sigma)} c_{x}^{2} - c_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \frac{s_{x}}{c_{x}} \\ &- \zeta_{12} \cdot c_{(\rho-\sigma)} \left[s_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(s_{(\delta+\rho-\sigma)} c_{x}^{2} + s_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \frac{s_{x}}{c_{x}} \\ &+ \left[\frac{s_{\rho}}{\zeta_{13}} \left(c_{(\delta-\rho)} s_{x} s_{y} - c_{\rho} c_{x} c_{y} s_{z} \right) + \zeta_{13} \cdot c_{\rho} \left(s_{(\delta-\rho)} s_{x} s_{y} + s_{\rho} c_{x} c_{y} s_{z} \right) \right] \frac{c_{y} \left(c_{x}^{2} c_{z}^{2} - s_{z}^{2} \right)}{c_{x} s_{z}} \\ &- \left[\frac{s_{\sigma}}{\zeta_{23}} \left(c_{(\delta-\sigma)} c_{x} s_{y} + c_{\sigma} c_{y} s_{x} s_{z} \right) + \zeta_{23} \cdot c_{\sigma} \left(s_{(\delta-\sigma)} c_{x} s_{y} - s_{\sigma} c_{y} s_{x} s_{z} \right) \right] \frac{c_{y} c_{z}^{2} s_{x}}{s_{z}} \right\} \\ &+ C_{\kappa}^{\nu} y_{\nu}^{2} \left\{ - \left[\frac{c_{(\phi-1-\phi_{2})} s_{(\rho-\sigma)}}{\zeta_{12}} + \zeta_{12} \cdot c_{(\rho-\sigma)} s_{(\phi_{1}-\phi_{2})} \right] \frac{c_{2} s_{1} s_{2} s_{x}}{c_{x}} \\ &+ \left(\frac{c_{\phi_{1}} s_{\rho}}{\zeta_{13}} + \zeta_{13} \cdot c_{\rho} s_{\phi_{1}} \right) \frac{c_{1} c_{2} s_{2} \left(c_{x}^{2} c_{z}^{2} - s_{z}^{2} \right)}{c_{x} c_{z} s_{z}} + \left(\frac{c_{\phi_{2}} s_{\sigma}}{\zeta_{23}} + \zeta_{23} \cdot c_{\sigma} s_{\phi_{2}} \right) \frac{c_{1} c_{2}^{2} c_{z} s_{1} s_{x}}{s_{z}} \right\} \\ &+ C_{l}^{\nu} y_{\nu}^{2} \left\{ s_{(\rho+\phi_{1})} \frac{c_{1} c_{2} s_{2} \left(1 - c_{z}^{2} s_{x}^{2} \right)}{c_{x} c_{z} s_{z}} + s_{(\sigma+\phi_{2})} \frac{c_{1} c_{2}^{2} c_{z} s_{1} s_{x}}{s_{z}} + s_{(\rho-\sigma+\phi_{1}-\phi_{2})} \frac{c_{2} s_{1} s_{2} s_{x}}}{c_{x}} \right\} ; \quad (18)$$

$$\begin{split} \dot{\sigma} &= C_{\kappa}^{l} y_{\tau}^{2} \left\{ \frac{s_{(\rho-\sigma)}}{\zeta_{12}} \left[c_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(c_{(\delta+\rho-\sigma)} c_{x}^{2} - c_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \frac{c_{x}}{s_{x}} \\ &- \zeta_{12} \cdot c_{(\rho-\sigma)} \left[s_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(s_{(\delta+\rho-\sigma)} c_{x}^{2} + s_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \frac{c_{x}}{s_{x}} \\ &+ \left[\frac{s_{\rho}}{\zeta_{13}} \left(c_{(\delta-\rho)} s_{x} s_{y} - c_{\rho} c_{x} c_{y} s_{z} \right) + \zeta_{13} \cdot c_{\rho} \left(s_{(\delta-\rho)} s_{x} s_{y} + s_{\rho} c_{x} c_{y} s_{z} \right) \right] \frac{c_{x} c_{y} c_{z}^{2}}{s_{z}} \\ &- \left[\frac{s_{\sigma}}{\zeta_{23}} \left(c_{(\delta-\sigma)} c_{x} s_{y} + c_{\sigma} c_{y} s_{x} s_{z} \right) + \zeta_{23} c_{\sigma} \left(s_{(\delta-\sigma)} c_{x} s_{y} - s_{\sigma} c_{y} s_{x} s_{z} \right) \right] \frac{c_{y} \left(c_{z}^{2} s_{x}^{2} - s_{z}^{2} \right)}{s_{x} s_{z}} \right\} \\ &+ C_{\kappa}^{\nu} y_{\nu}^{2} \left\{ - \left(\frac{c_{(\phi_{1}-\phi_{2})} s_{(\rho-\sigma)}}{\zeta_{12}} + \zeta_{12} \cdot c_{(\rho-\sigma)} s_{(\phi_{1}-\phi_{2})} \right) \frac{c_{2} c_{x} s_{1} s_{2}}{s_{x}} \\ &+ \left(\frac{c_{\phi_{1}} s_{\rho}}{\zeta_{13}} + \zeta_{13} \cdot c_{\rho} s_{\phi_{1}} \right) \frac{c_{1} c_{2} c_{x} c_{z} s_{2}}{s_{z}} + \left(\frac{c_{\phi_{2}} s_{\sigma}}{\zeta_{23}} + \zeta_{23} \cdot c_{\sigma} s_{\phi_{2}} \right) \frac{c_{1} c_{z}^{2} s_{1} \left(c_{z}^{2} s_{x}^{2} - s_{z}^{2} \right)}{c_{z} s_{x} s_{z}} \right\} \end{split}$$

$$+C_{l}^{\nu}y_{\nu}^{2}\left\{s_{(\rho+\phi_{1})}\frac{c_{1}c_{2}c_{x}c_{z}s_{2}}{s_{z}}+s_{(\sigma+\phi_{2})}\frac{c_{1}c_{2}^{2}s_{1}\left(c_{z}^{2}s_{x}^{2}+s_{z}^{2}\right)}{c_{z}s_{x}s_{z}}-s_{(\rho-\sigma+\phi_{1}-\phi_{2})}\frac{c_{2}c_{x}s_{1}s_{2}}{s_{x}}\right\} .$$
 (19)

An outline of the derivation is given in Appendix B. Concerning these equations, two remarks are in order:

- As mentioned above, we can obtain exactly the same formulae as in Ref. [23] (and also in Ref. [27] but with a slightly different phase convention) if we set $y_{\nu} = 0$. This serves as a check of our derivation, at least for the part below the seesaw threshold.
- ρ and σ are only determined up to nπ (n = 0, 1, 2, 3, ···) in Eqs.(6) and (9). Such an ambiguity in ρ and σ also leads to an ambiguity in φ₁ and φ₂ defined in Eqs.(10) and (11). However, both ambiguities cancel on the right hand side of Eqs.(14)-(19). So this ambiguity is harmless, as it should be.

In addition to equations given above, a knowledge of the RGE evolution behavior of $\theta_1, \theta_2, \phi_1$ and ϕ_2 will be helpful in our following discussions. So for completeness, we have also derived one-loop RGEs for parameters in U_{ν} under the condition that eigenvalues of Y_{ν} are hierarchical. We find that RG corrections to $\theta_1, \theta_2, \theta_3, \delta_{\nu}, \phi_1$ and ϕ_2 can be strongly enhanced by the factors ζ_{ij}^{-1} defined in Eq.(12), even if M_3 and Λ_{GUT} are only one or two orders apart in magnitude. The full analytical formulae are given in Appendix C.

III. RG CORRECTION TO NEUTRINO PARAMETERS WITH THREE TYPICAL MASS PATTERNS: FROM M_Z TO Λ_{GUT}

We have verified our analytical formulae in Eqs.(13)-(19) by comparing their numerical solution with those obtained in the more conventional way, which is to integrate RGEs for Y_l and κ numerically, and then to calculate left-handed neutrino masses, leptonic mixing angles and CP-violating phases by diagonalizing these two matrices at different energy scales.

In this section, we compare main features of the numerical result with those predicted in Eqs.(13)-(19). We try to clarify what corrections are possible for neutrino parameters, during the RG evolution from M_Z to Λ_{GUT} . Such a study is important in that, it helps us understand what values are *possible* or even *preferred* for neutrino parameters at the scale of Grand Unified theories. In the numerical calculation, we follow a bottom-up procedure. We start with the best fit values of neutrino parameters at the low energy, and numerically integrate Eqs.(13)-(19) to obtain left-handed neutrino masses, leptonic mixing angles and CP-violating phases at different high energy scales. Though such an approach may not seem well motivated from the perspective of a fundamental theory, it is quite advantageous for our present task. In this way, we no longer have to tune parameters at the high energy scale to meet low energy constraints.

At the present time, low energy experiments have measured leptonic mixing angles and neutrino mass squared differences to a reasonable degree of accuracy(best fit values and 3σ errors [33]):

$$\theta_x \approx 33.2^{\circ} {}^{+4.8^{\circ}}_{-4.6^{\circ}}; \quad \theta_y \approx 45.0^{\circ} {}^{+10.6^{\circ}}_{-9.3^{\circ}}; \quad \theta_z < 13.0^{\circ}; \\ \Delta m_{21}^2 = m_2^2 - m_1^2 \approx \left(7.9 {}^{+1.0}_{-0.8}\right) \times 10^{-5} \text{eV}^2; \\ \left|\Delta m_{31}^2\right| = \left|m_3^2 - m_1^2\right| \approx \left(2.2 {}^{+1.1}_{-0.8}\right) \times 10^{-3} \text{eV}^2.$$

$$(20)$$

However, we still lack a lot of information. We do not know about the smallest mixing angle θ_z and the absolute scale of neutrino masses, except for a few upper bounds. And we know nothing about leptonic CP-violating phases and the Yukawa coupling matrix Y_{ν} at all. We can only speculate that Y_{ν} might have some similarity to quark Yukawa coupling matrices. For such reasons, it is not practical to scan the whole parameter space for all possibilities.

On the other hand, earlier works [17,18,27] have emphasized that enhancing factors like ζ_{ij}^{-1} play a significant role in the RG evolution of mixing angles and CP-violating phases, if left-handed neutrinos are nearly degenerate. So to be relevant and illustrative, we discuss in detail the RG correction in theories with three typically interesting neutrino mass patterns:

(i) Normal Hierarchy: $m_1 \ll m_2 \ll m_3$.

With Eq.(20), we find

$$m_2 \simeq \sqrt{\Delta m_{21}^2} \simeq 0.009 \text{eV}; \quad m_3 \simeq \sqrt{|\Delta m_{31}^2|} \simeq 0.047 \text{eV}.$$
 (21)

Then $m_1 \lesssim 10^{-3} \text{eV}$ is small enough to make neutrino masses hierarchical:

$$\zeta_{12}^{-1} \approx -1.25, \quad \zeta_{13}^{-1} \approx -1.0, \quad \zeta_{23}^{-1} \approx -1.5 \quad : m_1 = 10^{-3} \text{eV}; \zeta_{12}^{-1} \approx -1.0, \quad \zeta_{13}^{-1} \approx -1.0, \quad \zeta_{23}^{-1} \approx -1.5 \quad : m_1 \le 10^{-4} \text{eV}.$$
(22)

Furthermore, from the determinant of Eq.(4)

$$M_{3} = \frac{v^{2}}{m_{a}} \cdot \sqrt[3]{\varsigma y_{\nu}^{2}}; \quad \varsigma \equiv (r_{1}r_{2})^{2} \left/ \left(\frac{m_{x}m_{y}}{m_{a}^{2}}\frac{M_{1}M_{2}}{M_{3}^{2}}\right) \right.$$
(23)

where $m_a \equiv \max[m_1, m_2, m_3]$, while m_x and m_y denoting the other two lighter lefthanded neutrino masses. When $\varsigma \sim 1$, we have in the most interesting case $y_{\nu} \sim \mathcal{O}(1)$,

$$M_3 \sim \frac{174^2}{m_3} \text{GeV} \simeq 6.4 \times 10^{14} \text{GeV}.$$
 (24)

(ii) Near Degeneracy: $m_1 \lesssim m_2 \lesssim m_3$.

Left-handed neutrinos are nearly degenerate if the absolute mass scale is much larger than values given in Eq.(21). β decay, $\beta\beta_{0\nu}$ decay, and cosmological and astrophysical observations have all set upper bounds on certain combinations of left-handed neutrino masses [34]. A rather stringent upper bound on nearly degenerate left-handed neutrino masses is [35]

$$m_{1,2,3} < 0.34 \text{eV}.$$
 (25)

If left-handed neutrino masses lie rightly beneath this bound, we find that

$$\zeta_{12}^{-1} \approx -5600, \quad \zeta_{13}^{-1} \approx -201, \quad \zeta_{23}^{-1} \approx -208.$$
 (26)

From Eq.(23), when $\varsigma \sim 1$ and $y_{\nu} \sim \mathcal{O}(1)$,

$$M_3 \sim \frac{174^2}{m_3} \text{GeV} \simeq 8.9 \times 10^{13} \text{GeV} \sim 10^{14} \text{GeV}.$$
 (27)

Note that the pattern $m_3 \lesssim m_1 \lesssim m_2$ is also possible, we shall consider this case in a different work.

(iii) Inverted Hierarchy: $m_3 \ll m_1 \lesssim m_2$.

In this case

$$m_1 \simeq \sqrt{|\Delta m_{31}^2|} \simeq 0.047 \text{eV}, \quad m_2 \approx \sqrt{m_1^2 + \Delta m_{21}^2} \simeq 0.048 \text{eV}.$$
 (28)

Then $m_3 \lesssim 5 \times 10^{-3}$ eV is small enough to make a hierarchy between itself and the other two masses:

$$\begin{aligned} \zeta_{12}^{-1} &\approx -112, \quad \zeta_{13}^{-1} \approx 1.2, \quad \zeta_{23}^{-1} \approx 1.2 \quad : m_3 = 5 \times 10^{-3} \text{eV}; \\ \zeta_{12}^{-1} &\approx -111, \quad \zeta_{13}^{-1} \approx 1.0, \quad \zeta_{23}^{-1} \approx 1.0 \quad : m_3 \le 10^{-3} \text{eV}. \end{aligned}$$
(29)

Furthermore, when $\varsigma \sim 1$ and $y_{\nu} \sim \mathcal{O}(1)$,

$$M_3 \sim \frac{174^2}{m_2} \text{GeV} \simeq 6.4 \times 10^{14} \text{GeV}.$$
 (30)

Note that in cases (i) and (iii), we shall always refer to the first lines of Eqs.(22) and (29) in our numerical calculation. And we shall also need the value of M_3 (mostly in the factor $\ln M_3$) in our discussions. Since varying $(\ln M_3)$ by one order of magnitude requires changing the magnitude of M_3 by a factor of $\mathcal{O}(10^4)$, the values of M_3 given in Eqs.(24), (27) and (30) are "precise" enough for an order of magnitude estimation. However, it should be stressed that although we always assume $y_{\nu}^2 \sim \mathcal{O}(1)$, there can be significant errors in the subsequent discussions if $\varsigma >> \mathcal{O}(10^4)$ or $\varsigma << \mathcal{O}(10^{-4})$ in Eq.(23). But such errors are easily corrected by taking into account the precise value of ς .

In the following subsections, we firstly discuss the evolution behavior of left-handed neutrino masses and of enhancing factors ζ_{ij}^{-1} defined in Eq.(12), then we study RG corrections to leptonic mixing angles and CP-violating phases in theories with each of the three neutrino mass patterns listed above.

A. RG Evolution of Neutrino Masses and Mass Ratios ζ_{ij}^{-1}

From Eq.(13), it is easy to obtain

$$\frac{m_i \left(\Lambda_{\rm GUT}\right)}{m_i \left(M_{\rm Z}\right)} = \exp\left[\frac{1}{16\pi^2} \int_{t_0 = \ln M_{\rm Z}}^{t' = \ln \Lambda_{\rm GUT}} a_i \left(t\right) dt\right], \quad i = 1, 2, 3.$$
(31)

Since α_{κ} dominates a_i (for i = 1, 2, 3) in general, the RG evolution of left-handed neutrino masses is mainly governed by a common scaling [25,27]

$$\frac{m_i \left(\Lambda_{\rm GUT}\right)}{m_i \left(M_{\rm Z}\right)} \simeq \exp\left[\frac{1}{16\pi^2} \int_{t_0=\ln M_{\rm Z}}^{t'=\ln \Lambda_{\rm GUT}} \alpha_\kappa\left(t\right) dt\right], \quad i=1,2,3.$$
(32)

There is a sudden change in the direction of the common scaling at the point where $y_{\nu} \sim \mathcal{O}(1)$ is turned on, since y_{ν} contributes to $\alpha_{\kappa}(t)$. This feature is quite obvious in Figure 1, where m_1 in cases (i) and (ii) and m_3 in case (iii) are plotted as functions of the energy scale, normalized by their values at M_Z .

Appreciable deviations from the common scaling may occur when y_{τ} or y_{ν} is large. Below the seesaw threshold, significant deviations are possible only in the MSSM when $\tan \beta$ is large, in which case $y_{\tau} \sim \mathcal{O}(1)$. Above the seesaw threshold, appreciable deviations are generally possible since it is natural to have $y_{\nu} \sim \mathcal{O}(1)$. In the following, we shall elaborate on this problem in a new but more meaningful way, i.e. by studying the RG evolution of the factors ζ_{ij}^{-1} (for i < j, i, j = 1, 2, 3) defined in Eq.(12).

1. Magnitudes of RG Corrections to ζ_{ii}^{-1}

As already mentioned, enhancing factors as ζ_{ij}^{-1} can be very important in the RG evolution of mixing angles and CP-violating phases. So it is necessary to study their own RG evolution behavior. From Eqs.(12) and (13), we find

$$\frac{\zeta_{ij}^{-1}}{\zeta_{ij}^{-1}} = -\frac{\dot{\zeta}_{ij}}{\zeta_{ij}} = -\left(\zeta_{ij}^{-1} - \zeta_{ij}\right)\frac{a_i - a_j}{2}.$$
(33)

Obviously, ζ_{ij} would be constants if all left-handed neutrino masses varied by an exactly common scaling, i.e. $a_i = a_j$ (for i, j = 1, 2, 3), which is of course not realistic. To estimate magnitudes of possible corrections to ζ_{ij} (or ζ_{ij}^{-1}) from M_Z to M_3 , we find that³

$$\frac{\Delta\zeta_{ij}^{-1}}{\zeta_{ij}^{-1}} \sim -\left(\zeta_{ij}^{-1} - \zeta_{ij}\right) \cdot \frac{a_i - a_j}{2} \cdot \frac{\Delta t}{16\pi^2} \sim -\left(\zeta_{ij}^{-1} - \zeta_{ij}\right) \frac{C_{\kappa}^l y_{\tau}^2 \ln\left(M_3/M_{\rm Z}\right)}{16\pi^2},\tag{34}$$

$$\sim \left(\zeta_{ij}^{-1} - \zeta_{ij}\right) \cdot 10^{-5};$$
 : in the SM (35)

$$\sim -\left(\zeta_{ij}^{-1} - \zeta_{ij}\right) \cdot \left(1 + \tan^2\beta\right) \cdot 10^{-5}, \qquad : \text{in the MSSM}$$
(36)

³Throughout this work, we shall omit the difference between $M_{\rm Z}$ and the SUSY breaking scale for simplicity.

where we have used Eq.(33) and have taken $y_{\tau} \sim m_{\tau} (M_Z) / (174 \text{GeV}) \approx 0.01$ in the SM, and $y_{\tau} \approx m_{\tau} (M_Z) / (\cos \beta \cdot 174 \text{GeV}) \approx 0.01 / \cos \beta$ in the MSSM.

With Eqs.(22), (26) and (29), we find that ζ_{ij}^{-1} vary little from M_Z to M_3 in the SM, even in case (ii). If Eq.(26) is used, the variation of ζ_{12}^{-1} is most significant: $|\Delta\zeta_{12}^{-1}/\zeta_{12}^{-1}| \sim 5.6 \times 10^{-2}$, while variations of ζ_{13}^{-1} and ζ_{23}^{-1} are much smaller: $|\Delta\zeta_{13}^{-1}/\zeta_{13}^{-1}| \sim |\Delta\zeta_{23}^{-1}/\zeta_{23}^{-1}| \sim 2 \times 10^{-3}$. In the MSSM, variations of ζ_{ij}^{-1} are amplified when $\tan \beta$ is large. However, $\tan^2 \beta$ alone is not large enough to generate appreciable excitation of $\zeta_{ij}^{-1} = 2.5 \times 10^{-2}$.

In the MSSM, variations of ζ_{ij}^{-1} are amplified when $\tan \beta$ is large. However, $\tan^2 \beta$ alone is not large enough to generate appreciable variations of ζ_{ij}^{-1} : $|\Delta \zeta_{ij}^{-1}/\zeta_{ij}^{-1}| \sim 2.5 \times 10^{-2}$ in the case when $\tan \beta \sim 50$ but $|\zeta_{ij}^{-1}| \sim \mathcal{O}(1)$. So corrections to ζ_{ij}^{-1} in case (i) are always negligible from M_Z to M_3 . Large variations of ζ_{ij}^{-1} are possible only when $|\zeta_{ij}^{-1}|$ are also large enough. In the limit $|\zeta_{ij}^{-1}| \to \infty$, we find from Eq.(33)

$$\dot{\zeta}_{ij} \simeq \frac{a_i - a_j}{2}, \quad \Longrightarrow \Delta \zeta_{ij} \sim \frac{a_i - a_j}{2} \cdot \frac{\Delta t}{16\pi^2} \sim \frac{C_\kappa^l y_\tau^2 \ln\left(M_3/M_Z\right)}{16\pi^2}, \tag{37}$$

$$\implies \frac{\zeta_{ij}^{-1}(M_{\rm Z})}{\zeta_{ij}^{-1}(M_{\rm 3})} = \frac{\zeta_{ij}(M_{\rm Z}) + \Delta\zeta_{ij}}{\zeta_{ij}(M_{\rm Z})} \sim \frac{(1 + \tan^2\beta) \cdot 10^{-5}}{\zeta_{ij}(M_{\rm Z})}.$$
 (38)

When $\tan \beta \sim 10$, only ζ_{12}^{-1} in case (ii) is modified: $|\zeta_{12}^{-1}(M_Z)/\zeta_{12}^{-1}(M_3)| \sim 5.6$. In contrast, significant variations of ζ_{ij}^{-1} are popular when $\tan \beta \sim 50$. Now there are roughly $|\zeta_{12}^{-1}(M_Z)/\zeta_{12}^{-1}(M_3)| \sim 140$, $|\zeta_{13}^{-1}(M_Z)/\zeta_{13}^{-1}(M_3)| \sim |\zeta_{23}^{-1}(M_Z)/\zeta_{23}^{-1}(M_3)| \sim 5$ in case (ii), and $|\zeta_{12}^{-1}(M_Z)/\zeta_{12}^{-1}(M_3)| \sim 2.5$ in case (iii). It is spectacular that $|\zeta_{12}^{-1}|$ is diminished to orders of magnitude smaller at M_3 than at M_Z in case (ii). Numerically, we find that the typical magnitude of ζ_{12}^{-1} at M_3 is of $\mathcal{O}(100)$, while $|\zeta_{13}^{-1}|$ and $|\zeta_{23}^{-1}|$ are of a few tens⁴.

Since the common scaling of k_i (for i = 1, 2, 3) from M_Z to Λ_{GUT} is only by a factor of $\mathcal{O}(1)$ and that $\Delta m_{21}^2 \propto k_2^2 - k_1^2 = -\zeta_{12} (k_1 + k_2)^2$, the strong reduction in $|\zeta_{12}^{-1}|$ means that Δm_{21}^2 is orders of magnitude larger at M_3 than at M_Z . This point applies also to $|\Delta m_{32}^2|$ (or $|\Delta m_{31}^2|$) when $|\zeta_{23}^{-1}|$ (or $|\zeta_{13}^{-1}|$) is strongly damped.

From M_3 to Λ_{GUT} , $y_{\nu} \sim \mathcal{O}(1)$ usually dominates corrections to ζ_{ij}^{-1} both in the SM and in the MSSM with $\tan \beta \sim 10$, while y_{τ} is comparable to y_{ν} in the MSSM when $\tan \beta \sim 50$. In whichever case, the dominant contribution is of the order $\mathcal{O}(y_{\nu})$, so we can make an estimation as in Eq.(38)

$$\frac{\zeta_{ij}^{-1}(M_3)}{\zeta_{ij}^{-1}(\Lambda_{\rm GUT})} \sim 1 + \frac{y_{\nu}^2 \ln\left(\Lambda_{\rm GUT}/M_3\right)}{16\pi^2 \cdot \zeta_{ij}(M_3)} \sim \frac{3 \times 10^{-2}}{\zeta_{ij}(M_3)} : \qquad M_3 \sim 10^{14} \text{GeV}.$$
(39)

Obviously, corrections to ζ_{ij}^{-1} in case (i) are again negligible from M_3 to $\Lambda_{\rm GUT}$. But it is remarkable that substantial corrections to ζ_{ij}^{-1} are now possible in the SM, both in cases (ii) and (iii). With Eqs.(26) and (29) and also considering the fact $|\zeta_{ij}^{-1}(M_3)| \approx |\zeta_{ij}^{-1}(M_Z)|$, we find $|\zeta_{12}^{-1}(M_3)/\zeta_{12}^{-1}(\Lambda_{\rm GUT})| \sim 170$, $|\zeta_{13}^{-1}(M_3)/\zeta_{13}^{-1}(\Lambda_{\rm GUT})| \sim |\zeta_{23}^{-1}(M_3)/\zeta_{23}^{-1}(\Lambda_{\rm GUT})| \sim 6$ in case (ii), and $|\zeta_{12}^{-1}(M_3)/\zeta_{12}^{-1}(\Lambda_{\rm GUT})| \sim 3$ in case (iii). In the MSSM when $\tan \beta \sim 10$, the

⁴An interesting exception to this simple estimation is that ζ_{23}^{-1} may vary little from M_Z to M_3 . Such a behavior occurs when θ_x is swiftly diminished to stay with θ_z at very near zero, while θ_y is kept at about 45° during the most part of the energy range from M_Z to M_3 . One can easily understand this point with Eqs.(37) and (44).

situation is quite similar. In this case only ζ_{12}^{-1} of case (ii) is moderately damped from M_Z to M_3 , but it still is a powerful enhancing factor in Eq.(39). For ζ_{13}^{-1} and ζ_{23}^{-1} of case (ii) and for ζ_{12}^{-1} of case (iii), the situation is the same as in the SM. In contrast, in the MSSM when $\tan \beta \sim 50$, $|\zeta_{ij}^{-1}|$ in general are strongly damped from M_Z to M_3 . As a result, only ζ_{12}^{-1} of case (ii) may vary significantly from M_3 to Λ_{GUT} , by the ratio $|\zeta_{12}^{-1}(M_3)/\zeta_{12}^{-1}(\Lambda_{GUT})| \sim 3$. However, there can also be appreciable modification in ζ_{13}^{-1} or ζ_{23}^{-1} of case (ii) or in ζ_{12}^{-1} of case (iii) if any of them is not so strongly damped from M_Z to M_3 .

2. Signs of RG Corrections to ζ_{ij}^{-1}

Now we turn to discuss the signs of corrections to ζ_{ij}^{-1} . By Eq.(37), we find

$$\zeta_{ij}\left(M_{3}\right) \approx \zeta_{ij}\left(M_{\rm Z}\right) + \frac{a_{i} - a_{j}}{2} \cdot \frac{\Delta t_{1}}{16\pi^{2}} ; \qquad (40)$$

$$\zeta_{ij} \left(\Lambda_{\rm GUT} \right) \approx \zeta_{ij} \left(M_3 \right) + \frac{a_i - a_j}{2} \cdot \frac{\Delta t_2}{16\pi^2} , \qquad (41)$$

where $\Delta t_1 = \ln(M_3/M_Z)$, $\Delta t_2 = \ln(\Lambda_{GUT}/M_3)$ and

$$\frac{a_1 - a_2}{2} = -C_{\kappa}^l y_{\tau}^2 \left[\left(c_x^2 - s_x^2 \right) \left(s_y^2 - c_y^2 s_z^2 \right) + 4c_{\delta} c_x c_y s_x s_y s_z \right] - C_{\kappa}^{\nu} y_{\nu}^2 \left(c_2^2 s_1^2 - s_2^2 \right) ; \qquad (42)$$

$$\frac{a_1 - a_3}{2} = -C_{\kappa}^l y_{\tau}^2 \left[c_y^2 \left(c_z^2 - c_x^2 s_z^2 \right) - s_x^2 s_y^2 + 2c_\delta c_x c_y s_x s_y s_z \right] - C_{\kappa}^{\nu} y_{\nu}^2 \left(c_1^2 c_2^2 - s_2^2 \right) ; \qquad (43)$$

$$\frac{a_2 - a_3}{2} = -C_{\kappa}^l y_{\tau}^2 \left[c_y^2 \left(c_z^2 - s_x^2 s_z^2 \right) - c_x^2 s_y^2 - 2c_\delta c_x c_y s_x s_y s_z \right] - C_{\kappa}^{\nu} y_{\nu}^2 \left(c_1^2 - s_1^2 \right) c_2^2 . \tag{44}$$

In these equations, one should let $y_{\nu} = 0$ at energies below the seesaw threshold (or M_3 equivalently, in this work). Left-handed neutrino masses will keep their order of sequence if $(a_i - a_j)/2$ are of the same signs as $\zeta_{ij} (M_Z)$ (for i < j; i, j = 1, 2, 3, respectively).

From M_Z to M_3 , signs of $(a_i - a_j)/2$ are determined by that of the $C_{\kappa}^l y_{\tau}^2$ term. As estimated above, significant variations of ζ_{ij}^{-1} are possible only in the MSSM, where $C_{\kappa}^l = 1$. It has been shown in Ref. [27] that θ_x tends to evolve toward zero whenever the correction is large; θ_y is always smaller than but not far away from about 45°; and θ_z is always smaller than about 15°. By this we can conclude that the $C_{\kappa}^l y_{\tau}^2$ term in Eqs.(42)-(44) are usually *negative*, and so that $|\zeta_{12}|$ of cases (ii) and (iii) is enlarged with increasing energy scales in general. A positive correction to ζ_{12} (which diminishes $|\zeta_{12}|$) is possible only when s_z is large and c_{δ} is negative in Eq.(42), just as observed in Ref. [27]. However, it is numerically more difficult to arrange a positive correction to ζ_{12} in the whole energy range from M_Z to M_3 than a negative one. For the same reason, ζ_{13} and ζ_{23} of case (ii) usually receive corrections of their own signs during the RG evolution.

From M_3 to Λ_{GUT} , the contribution from y_{ν} is important both in the SM and in the MSSM. In the SM, the y_{ν} term dominates $(a_i - a_j)/2$. Since θ_1 and θ_2 are totally arbitrary parameters, we can choose $(a_i - a_j)/2$ to be negative or positive at M_3 as we like. In the MSSM, the y_{τ} term is also important when $\tan \beta$ is large. But we still can change signs of $(a_i - a_j)/2$ by adjusting θ_1 and θ_2 . Furthermore, mixing angles and CP-violating phases often vary dramatically at energies above M_3 when $|\zeta_{ij}^{-1}|$ are large. So $(a_i - a_j)/2$ may also

change their signs during the evolution from M_3 to Λ_{GUT} . In general, both positive and negative $(a_i - a_j)/2$ are possible at energy scales above the seesaw threshold. However, as will be clear in the following, signs of ζ_{ij} are not likely to be changed despite this fact.

In Figure 2, we illustrate the typical evolution behavior of $\left(-\zeta_{ij}^{-1}\right) = |\zeta_{ij}^{-1}|$ in the SM and the MSSM. It is obvious that $\left(-\zeta_{ij}^{-1}\right)$ vary in a way just as discussed above.

3. RG Evolution of ζ_{ii}^{-1} in Nearly Singular Situations

As explained above, signs of $(a_i - a_j)/2$ can be either negative or positive from M_3 to Λ_{GUT} , depending mostly on θ_1 and θ_2 . Immediately, there comes the question of whether it is possible to change signs of ζ_{ij} at energies above M_3 . We find that the answer is likely to be negative.

When corrections to ζ_{ij} are positive, $|\zeta_{ij}|$ are diminished and $|\zeta_{ij}^{-1}|$ may be enhanced to extremely large values. Then the situation is nearly singular. However, we find that $|\zeta_{ij}^{-1}|$ are always dramatically diminished after they have reached some extremely large values. In such cases, extremely high but very narrow peaks appear in the plot of $|\zeta_{ij}^{-1}|$ against the energy scale. There seems to be a protecting mechanism that keeps $|\zeta_{ij}^{-1}|$ from going to infinity (or equivalently, keeps ζ_{ij} from going to zero).

One can understand this point most easily in the SM, in which y_{τ} is small and so that Eqs.(42)-(44) are dominated by the $C_{\kappa}^{\nu}y_{\nu}^{2}$ term. We also need Eqs.(C.1) and (C.2) in the limit $|\zeta_{ij}^{-1}| \to \infty$:

$$\dot{\theta}_1 = C_{\kappa}^{\nu} y_{\nu}^2 \left(-\frac{c_{(\phi_1 - \phi_2)}^2 s_2^2}{\zeta_{12}} + \frac{c_{\phi_1}^2 s_2^2}{\zeta_{13}} + \frac{c_{\phi_2}^2 c_2^2}{\zeta_{23}} \right) c_1 s_1 + \cdots;$$
(45)

$$\dot{\theta}_2 = C_{\kappa}^{\nu} y_{\nu}^2 \left(\frac{c_{(\phi_1 - \phi_2)}^2 s_1^2}{\zeta_{12}} + \frac{c_{\phi_1}^2 c_1^2}{\zeta_{13}} \right) c_2 s_2 + \cdots$$
(46)

It is crucial that, with ζ_{ij}^{-1} given in Eqs.(26) and (29), the dominant correction to θ_2 is always *negative*, while θ_1 can either be enlarged by the ζ_{12}^{-1} term or be reduced by terms of ζ_{13}^{-1} and ζ_{23}^{-1} . As a result, θ_2 is always driven toward 0° when $|\zeta_{12}^{-1}|$ or $|\zeta_{13}^{-1}|$ is large. In case (ii), since all three factors $|\zeta_{ij}^{-1}|$ are large, θ_1 can be driven toward either 0° or 90°, depending on the competition among different terms. In case (iii), since only $|\zeta_{12}^{-1}|$ is significant, θ_1 is always driven toward 90°.

Corrections to ζ_{ij} in case (i) are always negligible, so we only need to consider cases (ii) and (iii). We shall firstly discuss case (ii) in detail.

In Eqs.(42), (43) and (44), a nearly singular situation is most likely to occur when any of $\theta_1 \sim 0^\circ$ or 90° or $\theta_2 \sim 90^\circ$ is satisfied: if there is only $\theta_1 \sim 0^\circ$, a peak of $|\zeta_{12}^{-1}|$ may appear; if there is only $\theta_1 \sim 90^\circ$, peaks of $|\zeta_{13}^{-1}|$ and $|\zeta_{23}^{-1}|$ may appear; and if there is only $\theta_2 \sim 90^\circ$, peaks of $|\zeta_{12}^{-1}|$, $|\zeta_{13}^{-1}|$ and $|\zeta_{13}^{-1}|$ may appear; in case (ii)

$$\zeta_{13}^{-1} \approx \frac{1}{\zeta_{12} + \zeta_{23}} , \qquad (47)$$

a peak of $|\zeta_{13}^{-1}|$ should always coexist with peaks of $|\zeta_{12}^{-1}|$ and $|\zeta_{23}^{-1}|$. Such a situation is in fact quite rare.

Now we can discuss how signs of ζ_{ij} are protected.

Firstly, if ζ_{12} is driven to very near zero and $|\zeta_{12}^{-1}|$ develops a peak, terms led by ζ_{12}^{-1} dominate Eqs.(45) and (46). Then θ_2 is dramatically driven to near 0°, while θ_1 is swiftly enhanced toward 90°. The smaller $|\zeta_{12}|$ is, the more efficient this mechanism can take effect. As a result, $(a_1 - a_2)/2$ in Eq.(42) is quickly driven to be negative, leaving no chance for ζ_{12} to reach zero or even to get its sign changed.

Secondly, if ζ_{23} is driven to very near zero and $|\zeta_{23}^{-1}|$ develops a peak for some reasons, the ζ_{23}^{-1} term dominates Eq.(45). Then θ_1 is dramatically driven toward 0°. The smaller $|\zeta_{23}|$ is, the more efficient this mechanism can take effect. Since $\theta_2 \sim 90^\circ$ is not preferred in Eq.(46), $c_2^2 \neq 0$ in general. As a result, $(a_2 - a_3)/2$ in Eq.(43) is quickly driven to be negative, leaving no chance for ζ_{23} to reach zero or even become positive.

Finally, if $|\zeta_{13}^{-1}|$ develops a peak (so there are also peaks of $|\zeta_{12}^{-1}|$ and $|\zeta_{23}^{-1}|$), then ζ_{12}^{-1} and ζ_{13}^{-1} in Eq.(46) can dramatically drive θ_2 to 0°. Since terms led by ζ_{12}^{-1} and ζ_{13}^{-1} in Eq.(45) are suppressed by s_2^2 , the ζ_{23}^{-1} term then dominates Eq.(45) and drives θ_1 to 0°. Then $(a_i - a_j)/2$ in Eqs.(42), (43) and (44) all become negative quickly. As a result, none of ζ_{12} , ζ_{13} and ζ_{23} will vanish or even get its sign changed.

So it is not likely that signs of ζ_{ij} can be changed. However, we should stress that such a discussion is only a way to understand how signs of ζ_{ij} can always be preserved, while the latter point is in fact not proved.

In case (iii), only $|\zeta_{12}^{-1}|$ is important. A nearly singular situation is possible if $\theta_1 \sim 0^\circ$ or $\theta_2 \sim 90^\circ$. At the peak of $|\zeta_{12}^{-1}|$, θ_2 is dramatically driven to 0° and θ_1 is driven to 90° . Then there will be no more peaks.

In the MSSM when $\tan \beta \sim 50$, y_{τ} is comparable to y_{ν} and the contribution from y_{τ} is not negligible in general. But numerically, we find that the mechanism discussed above still works as long as y_{ν} is large.

Numerical examples illustrating these features are given in Figure 3.

B. The Normal Hierarchy Case

When left-handed neutrino masses are hierarchical, $|\zeta_{ij}^{-1}|$ are of $\mathcal{O}(1)$. So there are no enhancing factors in the evolution of mixing angles in Eqs.(14)-(16), while the evolution of CP-violating phases in Eqs.(17)-(19) can still be enhanced by the factor s_z^{-1} . In the limit $s_z \longrightarrow 0$:

$$\dot{\delta} \approx \dot{\rho} \approx \dot{\sigma} = C_{\kappa}^{l} y_{\tau}^{2} \left[\left(\frac{c_{(\delta-\rho)} s_{\rho}}{\zeta_{13}} + \zeta_{13} c_{\rho} s_{(\delta-\rho)} \right) - \left(\frac{c_{(\delta-\sigma)} s_{\sigma}}{\zeta_{23}} + \zeta_{23} c_{\sigma} s_{(\delta-\sigma)} \right) \right] \frac{c_{x} c_{y} s_{x} s_{y}}{s_{z}} \\ + C_{\kappa}^{\nu} y_{\nu}^{2} \left[\left(\frac{c_{\phi_{1}} s_{\rho}}{\zeta_{13}} + \zeta_{13} c_{\rho} s_{\phi_{1}} \right) c_{x} s_{2} + \left(\frac{c_{\phi_{2}} s_{\sigma}}{\zeta_{23}} + \zeta_{23} c_{\sigma} s_{\phi_{2}} \right) c_{2} s_{1} s_{x} \right] \frac{c_{1} c_{2}}{s_{z}} \\ + C_{l}^{\nu} y_{\nu}^{2} \left(s_{(\rho+\phi_{1})} c_{x} s_{2} + s_{(\sigma+\phi_{2})} c_{2} s_{1} s_{x} \right) \frac{c_{1} c_{2}}{s_{z}} + \cdots,$$
(48)

where we have neglected all terms that are not enhanced by s_z^{-1} . It is remarkable that dominant corrections to δ, ρ and σ are exactly the same. It is also interesting that the term proportional to $C_{\kappa}^l y_{\tau}^2$ vanishes in the limit $\zeta_{13} \approx \zeta_{23} \approx -1$, while those led by y_{ν}^2 become proportional to $(C_l^{\nu} - C_{\kappa}^{\nu})$. Since $C_{\kappa}^{\nu} = C_l^{\nu} = 1$ in the MSSM, the contribution from y_{ν} also vanishes in the limit $\zeta_{13} \approx \zeta_{23} \approx -1$ in this case.

1. RG Corrections in the SM

In the SM, $y_{\tau} \approx m_{\tau} (M_{\rm Z}) / (174 {\rm GeV}) \approx 0.01$, and the typical contribution is

$$\frac{y_{\tau}^2}{16\pi^2} \ln \frac{\Lambda_{\rm GUT}}{M_{\rm Z}} \,({\rm rad}) \sim 10^{-5} \,({\rm rad}) \sim \left(10^{-3}\right)^\circ. \tag{49}$$

Since we are mostly interested in cases with a large y_{ν} , we shall consider only $y_{\nu} \sim \mathcal{O}(1)$ in this work. When $y_{\nu} \sim \mathcal{O}(1)$, the contribution is

$$\frac{y_{\nu}^2}{16\pi^2} \ln \frac{\Lambda_{\rm GUT}}{M_3} \,({\rm rad}) \sim \begin{cases} 0.014 \,({\rm rad}) \sim 1.1^\circ : & M_3 \sim 10^{15} {\rm GeV} \\ 0.029 \,({\rm rad}) \sim 1.9^\circ : & M_3 \sim 10^{14} {\rm GeV} \end{cases} \,. \tag{50}$$

Apart from these, all other miscellaneous terms (besides ζ_{ij}^{-1}) on the right-hand side of Eqs.(14)-(19) usually damp the values given above strongly. Through out this work, we shall assume that the net effect of all these terms is equivalent to a factor of $\mathcal{O}(0.1)$. Though such an assumption is mainly based on our numerical experience and is far from precise, it serves as a crude estimation and can help us understand the most important part of RG corrections. With this assumption, we estimate that only corrections of $(10^{-4})^{\circ}$ are possible for mixing angles running from $M_{\rm Z}$ to M_3 , while corrections of $\mathcal{O}(0.1^{\circ})$ are possible from M_3 to $\Lambda_{\rm GUT}$, if $y_{\nu} \sim \mathcal{O}(1)$.

For CP-violating phases, significant corrections are possible in the energy range from M_3 to $\Lambda_{\rm GUT}$, since there is the enhancing factor $s_z^{-1} \gtrsim 5$.

Firstly, the RG correction to θ_z is negligible in the whole range from M_Z to Λ_{GUT} , when θ_z is of $\mathcal{O}(1^\circ \sim 10^\circ)$ at M_Z . In this case, $s_z^{-1} \sim \mathcal{O}(5 \sim 50)$. From Eq.(50) and the $\mathcal{O}(0.1)$ factor explained above, corrections to CP-violating phases are of $\mathcal{O}(0.5^\circ \sim 5^\circ)$.

Secondly, when θ_z is of $\mathcal{O}(0.1^{\circ} \sim 1^{\circ})$ at M_Z , the RG correction to it is not negligible from M_3 to Λ_{GUT} . However, the order of magnitude of s_z^{-1} is not changed by the correction. So in this case, corrections to CP-violating phases are of $\mathcal{O}(5^{\circ} \sim 50^{\circ})$, given that $y_{\nu} \sim \mathcal{O}(1)$.

Finally, though a value much smaller than possible radiative corrections may not seem natural for θ_z , we shall consider the case $\theta_z \ll 0.1^\circ$ for completeness. Since the radiative correction can enlarge θ_z to $\mathcal{O}(0.1^\circ \sim 1^\circ)$ at energies above M_3 , corrections to CP-violating phases in this energy range are roughly the same as in the second case. But if θ_z is not magnified above M_3 , the corrections can be extraordinarily large. At energies below M_3 , corrections from y_{τ} to CP-violating phases are still negligible when θ_z is of $(0.01^\circ \sim 0.1^\circ)$. But when $\theta_z < 0.01^\circ$, i.e. $s_z^{-1} > 0.5 \times 10^4$, the contribution from y_{τ} can be so strongly enhanced as to become appreciable or even important.

In Figure 4, we illustrate each of the possibilities discussed above. It is obvious that the phases δ , ρ and σ in the figure always vary by approximately the *same* size, just as predicted in Eq.(48).

2. RG Corrections in the MSSM when $\tan \beta \sim 10$

In the MSSM when $\tan \beta \sim 10$, $y_{\tau} \approx m_{\tau} (M_{\rm Z}) / (\cos \beta \cdot 174 \text{GeV}) \approx 0.1$, and the typical contribution from which is

$$\frac{y_{\tau}^2}{16\pi^2} \ln \frac{\Lambda_{\rm GUT}}{M_Z} \,(\text{rad}) \sim 0.1^\circ.$$
(51)

The contribution form y_{ν} is the same as given in Eq.(50). Then corrections to mixing angles are of $\mathcal{O}(0.01^{\circ})$ in the range from $M_{\rm Z}$ to M_3 , but are still of $\mathcal{O}(0.1^{\circ})$ from M_3 to $\Lambda_{\rm GUT}$. For CP-violating phases, RG corrections from $M_{\rm Z}$ to M_3 can already be appreciable when θ_z is of $\mathcal{O}(0.1^{\circ} \sim 1^{\circ})$. This is in vast contrast to the SM case, where to make the contribution from y_{τ} appreciable, θ_z should be about one hundred times smaller than what is given here.

Now we have a novel possibility that there can be large corrections to CP-violating phases both at energies *above* and *below* M_3 . When $\theta_z \sim \mathcal{O}(0.01^\circ)$ at M_Z , the RG correction does not change the order of magnitude of θ_z in the energy range from M_Z to M_3 , so $s_z^{-1} \sim \mathcal{O}(5 \times 10^3)$ can enhance corrections to CP-violating phases to be of $\mathcal{O}(50^\circ)$. At energies above M_3 , θ_z may acquire a correction of $\mathcal{O}(0.1^\circ)$ in general, but it is still of $\mathcal{O}(0.01)$ at M_3 . As a result, CP-violating phases can vary dramatically above M_3 , and corrections of $\mathcal{O}(50^\circ)$ are readily possible for them. In contrast, in the MSSM when $\tan \beta \sim 50$, θ_z usually is magnified to be well above 0.1° in the energy range from M_Z to M_3 . So corrections (from terms led by y_{ν}^2) to CP-violating phases are less enhanced and the phases vary little at energies above M_3 . To conclude, it is only in the MSSM when $\tan \beta \sim 10$ that large corrections to CPviolating phases are most probable, both at energies *below* and *above* M_3 . This observation is supported in Figure 5.

3. RG Corrections in the MSSM when $\tan \beta \sim 50$

In the MSSM when $\tan \beta \sim 50$, $y_{\tau} \approx m_{\tau} (M_{\rm Z}) / (\cos \beta \cdot 174 {\rm GeV}) \approx 0.5$, and the correction from which is

$$\frac{y_{\tau}^2}{16\pi^2} \ln \frac{\Lambda_{\rm GUT}}{M_{\rm Z}} \,(\rm rad) \sim 2.9^\circ.$$
(52)

The correction from y_{ν} is still the same as given in Eq.(50). As a result, corrections of $\mathcal{O}(0.3^{\circ})$ are possible for mixing angles running from $M_{\rm Z}$ to M_3 . From M_3 to $\Lambda_{\rm GUT}$, corrections to mixing angles are still of $\mathcal{O}(0.1^{\circ})$. But for CP-violating phases, RE corrections can be extraordinarily large now.

Firstly, the RG correction does not change the order of magnitude of s_z^{-1} when θ_z is of $\mathcal{O}(0.1^\circ \sim 10^\circ)$ at M_Z . Then corrections to CP-violating phases are of $\mathcal{O}(1.5^\circ)$, $\mathcal{O}(15^\circ)$ and $\mathcal{O}(150^\circ)$ when $\theta_z \sim \mathcal{O}(10^\circ)$, $\mathcal{O}(1^\circ)$ and $\mathcal{O}(0.1^\circ)$ respectively. When possible corrections are as large as $\mathcal{O}(150^\circ)$, CP-violating phases are often driven to near their (pseudo-) fixed points: the phases keep varying dramatically until the right hand side of Eq.(48) vanishes, and then they become stable against the energy scale. RG corrections to CP-violating phases can be largely damped because of this behavior.

Secondly, when $\theta_z \ll 0.1^\circ$ at M_Z , the RG correction can enlarge θ_z to be of $\mathcal{O}(0.1^\circ \sim 1^\circ)$ along the way from M_Z to M_3 . However, extraordinarily large corrections to CP-violating phases can arise in a very narrow energy range at near M_Z , if θ_z is extremely small at the beginning. In this case, (pseudo-) fixed points of CP-violating phase are often swiftly reached, and then the phases evolve little with the energy scale. Above M_3 , the phases start to vary again when the contribution from y_{ν} is turned on. However, the dominant contribution still comes from y_{τ} . This is because θ_z is generally enlarged to $\mathcal{O}(0.1^\circ \sim 1^\circ)$ below M_3 , and so that the contribution from y_{ν} is less enhanced.

Numerical examples illustrating these points are given in Figure 5.

Apart from the magnitudes, signs of RG corrections to CP-violating phases are also important. In Eq.(48), the signs are determined by those of C_{κ}^{l} , C_{κ}^{ν} , C_{l}^{ν} , ζ_{13} and ζ_{23} , and by values of CP-violating phases. It may not worth while discussing all possibilities, but one case is simple and interesting. In the SM, since $C_{\kappa}^{\nu} = 1/2$, $C_{l}^{\nu} = -3/2$ and since ζ_{13} and ζ_{23} are negative, terms led by y_{ν}^{2} in Eq.(48) are always negative when both $\rho + \phi_{1}$ and $\sigma + \phi_{2}$ are in the range $(0 \sim \pi)$, but the contribution from these terms is positive when the range is $(\pi \sim 2\pi)$. One can check this point with Figure 4.

C. The Near Degeneracy Case

When neutrino masses are nearly degenerate, corrections to mixing angles and CPviolating phases can be resonantly enhanced since enhancing factors $|\zeta_{ij}^{-1}|$ in Eq.(26) are large. Before the discussion of any specific models, we remark that

- In Eqs.(14)-(19), corrections to θ_x , δ , ρ and σ are enhanced by ζ_{12}^{-1} , ζ_{13}^{-1} and ζ_{23}^{-1} , while corrections to θ_y and θ_z are only enhanced by ζ_{13}^{-1} and ζ_{23}^{-1} .
- In Eqs.(14)-(19), the factors ζ_{ij}^{-1} appear in terms led by y_{ν} via three common combinations: $\zeta_{12}^{-1} \cdot c_2 s_1 s_2 c_{(\phi_1 \phi_2)}, \zeta_{13}^{-1} \cdot c_1 c_2 s_2 c_{\phi_1}$ and $\zeta_{23}^{-1} \cdot c_1 c_2^2 s_1 c_{\phi_2}$. So there will be no resonantly enhanced contribution from y_{ν} if $\theta_1 = \theta_2 = 0$ (or $\theta_1 = \theta_2 = 90^\circ$ etc).
- Just like c_{ϕ_1} , c_{ϕ_2} and $c_{(\phi_1-\phi_2)}$ in the contribution from y_{ν} , certain functions of δ, ρ and σ can also be factorized out together with the enhancing factors ζ_{ij}^{-1} , both in contributions from y_{τ} and from y_{ν} . The result is collected in Table 1. With this, it is easy to find out what CP-violating phases can best damp the resonantly enhanced correction to a specific mixing angle or CP-violating phase.
- In the contribution from y_{ν} in Eqs.(14)-(19), the association of ζ_{ij}^{-1} with CP-violating phases is rather simple. So it is easy to tell signs of corrections enhanced by different factors ζ_{ij}^{-1} . We collect the result in Table 2, where we have assumed that minus signs in Eqs.(14)-(19) belong to phase factors.

In the following we discuss RG corrections to mixing angles and CP-violating phases both in the SM and in the MSSM.

1. RG Corrections in the SM

From M_Z to M_3 , RG corrections are determined by the contribution from y_{τ} . With the help of Eqs.(26) and (49) and also considering the overall $\mathcal{O}(0.1)$ factor explained below Eq.(50), we estimate that the correction (enhanced by ζ_{12}^{-1}) to θ_x is of $\mathcal{O}(0.5^\circ)$, and corrections (enhanced by ζ_{13}^{-1} and ζ_{23}^{-1}) to θ_y and θ_z are of $\mathcal{O}(0.02^\circ)$.

From M_3 to Λ_{GUT} , the contribution from $y_{\nu} \sim \mathcal{O}(1)$ is dominant and the magnitude is given in Eq.(50). Since $\zeta_{ij}^{-1}(M_3) \approx \zeta_{ij}^{-1}(M_Z)$, the correction (enhanced by ζ_{12}^{-1}) to θ_x is of $\mathcal{O}(1100^\circ)$, and corrections (enhanced by ζ_{13}^{-1} and ζ_{23}^{-1}) to θ_y and θ_z are of $\mathcal{O}(40^\circ)$. The $\mathcal{O}(40^\circ)$ correction to θ_z is a quite generous gift: a correction of $\mathcal{O}(1^\circ)$ could be interesting when θ_z is really small at M_Z [21,27,30]. We shall discuss this problem in more detail in the MSSM when $\tan \beta \sim 50$ (within this subsection, i.e. also for the near degeneracy case). For θ_x , however, the $\mathcal{O}(1100^\circ)$ correction is overestimated. As we have discussed in the first subsection and also is vivid in Figure 2, in a large part of the energy range from M_3 to Λ_{GUT} , $|\zeta_{12}^{-1}|$ is more than an order of magnitude smaller than it is at M_3 . So the correction enhanced by ζ_{12}^{-1} should be about an order smaller in magnitude than estimated above.

In cases when there are nearly singular situations, i.e. when $|\zeta_{ij}^{-1}|$ develop peaks, the "protective mechanism" discussed in the first subsection becomes important again in damping extraordinarily enhanced corrections to mixing angles and CP-violating phases. As mentioned in the beginning of this subsection, corrections from y_{ν} that can be enhanced by $|\zeta_{ij}^{-1}|$ are always controlled by s_1 and s_2 : i.e. ζ_{12}^{-1} by s_1s_2 , ζ_{13}^{-1} by s_2 and ζ_{23}^{-1} by s_1 . θ_1 is dramatically driven to near zero whenever $|\zeta_{23}^{-1}|$ develops a peak, and θ_2 is dramatically driven to near zero whenever $|\zeta_{12}^{-1}|$ develops a peak. As a result, enhancing effect of all peaks of $|\zeta_{ij}^{-1}|$ are strongly damped. However, the contribution from y_{ν} can still be very large in such a case (numerically, we find that a value of $\mathcal{O}(100^\circ)$ is popular).

Signs of corrections to mixing angles are determined by CP-violating phases δ , ρ , σ , ϕ_1 and ϕ_2 , and the competition among different contributions led by ζ_{12}^{-1} , ζ_{13}^{-1} and ζ_{23}^{-1} . For an interesting example, we discuss in detail the case when all of the five phases are in the range $(0, \pi/2)$. As discussed above, only the contribution from y_{ν} at energies above M_3 is important.

- For θ_x : In Eq.(14), the contributions from y_{ν} led by ζ_{12}^{-1} and ζ_{13}^{-1} are positive, while that led by ζ_{23}^{-1} is negative. Since $|\zeta_{12}^{-1}|$ is generally larger than $|\zeta_{13}^{-1}|$ and $|\zeta_{23}^{-1}|$, the net correction to θ_x is positive in general, at near M_3 . However, since θ_2 usually is more swiftly driven to near zero than θ_1 , negative contributing terms led by ζ_{23}^{-1} in Eq.(14) often have a chance to catch up with the positive contributions led by ζ_{12}^{-1} and ζ_{13}^{-1} . As a result, sometimes θ_x may make a change of its evolution direction and evolve toward zero.
- For θ_y : In Eq.(15), the part of the correction from y_{ν} led by ζ_{13}^{-1} is negative but that led by ζ_{23}^{-1} is positive. Since $|\zeta_{13}^{-1}|$ and $|\zeta_{23}^{-1}|$ are comparable to each other in general, the two parts of the correction usually cancel each other to a large extent. So the net correction to θ_y is relatively small. Furthermore, since θ_2 is often more swiftly diminished than θ_1 , the part led by ζ_{23}^{-1} often is dominant and the correction to θ_y is positive.

• For θ_z : In Eq.(16), the correction from y_{ν} to θ_z is *positive*. So θ_z only *increases* with the energy scale in this case. This feature leads to a quite interesting possibility that θ_z is comparable to the other two angles at Λ_{GUT} , but it is diminished when the energy scale is decreased.

Numerical examples illustrating these features are given in Figure 6. In the figure, θ_y is decreased above M_3 . This is because $\delta > \pi/2$ in that energy range.

For CP-violating phases, from M_Z to M_3 , the factors $|\zeta_{ij}^{-1}|$ alone can enhance corrections from y_{τ} to be of $\mathcal{O}(0.5^{\circ})$. Large corrections to CP-violating phases are possible only when θ_z is small enough. To make this point clear, we find (similar to Eq.(48)) in the limit $s_z \to 0$:

$$\dot{\delta} \approx \dot{\rho} \approx \dot{\sigma} = C_{\kappa}^{l} y_{\tau}^{2} \left[\frac{c_{(\delta-\rho)} s_{\rho}}{\zeta_{13}} - \frac{c_{(\delta-\sigma)} s_{\sigma}}{\zeta_{23}} \right] \frac{c_{x} c_{y} s_{x} s_{y}}{s_{z}} + C_{\kappa}^{\nu} y_{\nu}^{2} \left[\frac{c_{\phi_{1}} s_{\rho} c_{x} s_{2}}{\zeta_{13}} + \frac{c_{\phi_{2}} s_{\sigma} c_{2} s_{1} s_{x}}{\zeta_{23}} \right] \frac{c_{1} c_{2}}{s_{z}} + C_{l}^{\nu} y_{\nu}^{2} \left(s_{(\rho+\phi_{1})} c_{x} s_{2} + s_{(\sigma+\phi_{2})} c_{2} s_{1} s_{x} \right) \frac{c_{1} c_{2}}{s_{z}} + \cdots , \qquad (53)$$

where we have omitted terms that are not enhanced by s_z^{-1} (including those led by ζ_{12}^{-1}). The contribution from y_{τ} can be enhanced to $\mathcal{O}(0.02^{\circ})$ by ζ_{13}^{-1} and ζ_{23}^{-1} , so $\theta_z \sim \mathcal{O}(0.1^{\circ})$ is small enough to help generating corrections of $\mathcal{O}(10^{\circ})$. However, since $|\zeta_{13}^{-1}|$ and $|\zeta_{23}^{-1}|$ are comparable to each other in general, corrections from y_{τ} are strongly damped if $\rho \approx \sigma$.

From M_3 to Λ_{GUT} , s_z^{-1} in general is not an enhancing factor since a correction of $\mathcal{O}(40^\circ)$ is possible to θ_z . With ζ_{ij}^{-1} in Eq.(26) being the only enhancing factors, the situation for CP-violating phases is much like that for mixing angles discussed above. So in this energy range, corrections of $\mathcal{O}(100^\circ)$ are possible for CP-violating phases. However, there is still a notable difference between the evolution of CP-violating phases and that of mixing angles. At energies right above M_3 , θ_z is still very small. So the enhancing effect of s_z^{-1} combined with that of ζ_{13}^{-1} and ζ_{23}^{-1} in Eq.(53) can drive CP-violating phases to vary almost abruptly. Corrections of this origin are distinguishable since they are enhanced by s_z^{-1} and thus are exactly the same for δ , ρ and σ . Only after θ_z has grown large enough, will the evolution of CP-violating phases slow down and acquire a strength comparable to that of mixing angles.

The evolution of CP-violating phases is also illustrated in Figure 6. In the figure, the difference between the RG evolution of mixing angles and that of CP-violating phases (as discussed above) is obvious.

2. RG Corrections in the MSSM when $\tan \beta \sim 10$

In this case, the contribution of y_{τ} is given in Eq.(51). From $M_{\rm Z}$ to M_3 , ζ_{12}^{-1} in Eq.(26) can enhance the correction to θ_x to be of $\mathcal{O}(50^\circ)$, while corrections to θ_y and θ_z (enhanced by ζ_{13}^{-1} and ζ_{23}^{-1}) are of $\mathcal{O}(2^\circ)$. Here, the $\mathcal{O}(50^\circ)$ correction to θ_x is usually negative and will be damped when θ_x is near zero. A positive correction to θ_x is also possible. But such a correction is always suppressed by factors of $\mathcal{O}(s_z)$. Considering the upper bound for $s_z (\leq 1/5)$, a positive correction to θ_x should be smaller than $\mathcal{O}(10^\circ)$. The smaller θ_z is, the smaller such a correction would be. If we take $\theta_z \sim 2^\circ$, the correction should be smaller

than $\mathcal{O}(2^{\circ})$. For θ_y , the situation is similar to that of θ_x . The dominant terms are usually negative, while a positive correction is possible but is always suppressed by s_z . For θ_z , the correction can be either positive or negative, depending on values of CP-violating phases and on the competition among the terms led by ζ_{13}^{-1} and ζ_{23}^{-1} .

From M_3 to Λ_{GUT} , y_{ν} dominates the corrections and the situation for mixing angles is quite similar to that in the SM. A notable difference is that, now one can start the evolution with a small θ_x at M_3 , since it has a good chance of being damped from M_{Z} to M_3 .

For CP-violating phases, ζ_{12}^{-1} alone can enhance the corrections to $\mathcal{O}(50^\circ)$, in the energy range from M_Z to M_3 . But this correction is strongly damped when $\rho \approx \sigma$. Alternative large corrections can come from terms led by ζ_{13}^{-1} and ζ_{23}^{-1} , in cases when θ_z is very small. The corresponding RGEs are the same as those in Eq.(53). Since dominant corrections to δ, ρ and σ are exactly the same, the relation $\rho \approx \sigma$ can be easily retained in this case. CPviolating phases are often driven to near their (pseudo-) fixed points when θ_z is extremely small at M_Z .

From M_3 to Λ_{GUT} , corrections to CP-violating phases are dominated by y_{ν} and the situation is quite similar to that in the SM.

For a numerical illustration and also as a check of the third remark at the beginning of this subsection, we stress that θ_x and the phases δ , ρ and σ cannot acquire their largest possible corrections simultaneously. For example, one needs $\rho \approx \sigma$ to make the ζ_{12}^{-1} term in Eq.(14) least suppressed, so that the correction to θ_x can be largest. However, in Table 1 and Eqs.(17)-(19), the condition $\rho \approx \sigma$ damps corrections (enhanced by ζ_{12}^{-1}) to CP-violating phases most strongly. Also, this $\rho \approx \sigma$ condition damps the RG correction to θ_z in the range from M_Z to M_3 . One can understand this point with the help of Eq.(16) and the fact that ζ_{13}^{-1} and ζ_{23}^{-1} are usually comparable to each other. These results are all illustrated in Figure 7. Note that, since we take $\rho \approx \sigma$ in the calculation, the correction to θ_z is largely damped and θ_z is kept unchanged until M_3 . As a result, CP-violating phases vary almost abruptly at the point where y_{ν} is turned on, just as in the case of the SM.

3. RG Corrections in the MSSM when $\tan \beta \sim 50$

In this case, the contribution of y_{τ} is given in Eq.(52). Along with the usual $\mathcal{O}(0.1)$ factor explained below Eq.(50), ζ_{12}^{-1} in Eq.(26) can enhance the correction to θ_x to $\mathcal{O}(1700^\circ)$, in the energy range from M_Z to M_3 . The other two factors ζ_{13}^{-1} and ζ_{23}^{-1} in Eq.(26) can enhance corrections to θ_y and θ_z to $\mathcal{O}(60^\circ)$. However, these values are overestimated. We have shown in the first subsection that, $|\zeta_{ij}^{-1}|$ usually are strongly reduced from M_Z to M_3 . In Figure 2, ζ_{12}^{-1} is more than an order smaller in magnitude than it is at M_Z , during a large part of the energy range from M_Z to M_3 . So in general, the correction to θ_x should also be an order of magnitude smaller than estimated above. The variation of $|\zeta_{13}^{-1}|$ and $|\zeta_{23}^{-1}|$ is more moderate and the strength of the reduction is more uniform in the whole energy range than those of $|\zeta_{12}^{-1}|$. For such reasons, we re-estimate that the correction to θ_x is roughly of $\mathcal{O}(170^\circ)$, while those to θ_y and θ_z are roughly of $\mathcal{O}(30^\circ)$.

Furthermore, (pseudo-) fixed points are always possible for mixing angles and CPviolating phases whenever corrections are very large. When certain angles or phases are near their fixed points, RG corrections to them are strongly damped. As a result, real corrections to mixing angles and CP-violating phases depend not only on values estimated above, but also on how (pseudo-) fixed points can be reached. This is also true for RG corrections above M_3 .

From M_3 to Λ_{GUT} , contributions from y_{ν} and y_{τ} are both important. As estimated in the first subsection, since $|\zeta_{12}^{-1}|$ is generally orders of magnitude smaller at M_3 than at M_Z , and since $|\zeta_{13}^{-1}|$ and $|\zeta_{23}^{-1}|$ are about 5 times smaller at M_3 than at M_Z , the correction to θ_x is roughly of $\mathcal{O}(100^\circ)$, while those to θ_y and θ_z are of $\mathcal{O}(10^\circ)$.

- For θ_x : From M_Z to M_3 , the dominant contribution in Eq.(14) (that enhanced by ζ_{12}^{-1} , but not suppressed by s_z) is always *negative*. But if θ_z is large, there can be a positive correction. The correction to θ_x becomes largest when $|c_{(\rho-\sigma)}| \sim 1$. Since $\rho = \sigma = 0$ is stable against RG corrections, this condition is easy to retain. Furthermore, $|c_{(\rho-\sigma)}| \sim$ 1 can also lead to a large correction to θ_x in the energy range from M_3 to Λ_{GUT} , if there is in addition $|c_{(\phi_1-\phi_2)}| \sim 1$. In contrast, if we want the correction to θ_x to be small, we need both $c_{(\rho-\sigma)} \sim 0$ and $c_\rho c_{\phi_1} \sim c_\sigma c_{\phi_2} \sim 0$. These requirements can only be partially satisfied by $\delta = \rho = \phi_1 = 0$ (up to π) and $\sigma = \phi_2 = \pi/2$ (up to π), which are also stable against RG corrections.
- For θ_y : From M_Z to M_3 , dominant contributions in Eq.(15) (those enhanced by ζ_{13}^{-1} , ζ_{23}^{-1} , but not suppressed by s_z) are also *negative*. Though a positive correction seems possible in Eq.(15) when θ_z is large, a numerical example is hard to find. The correction to θ_y becomes largest when $|c_{(\delta-\rho)}| \sim |c_{(\delta-\sigma)}| \sim 1$ below M_3 , and $|c_{(\delta-\rho)}c_{\phi_1}| \sim |c_{(\delta-\sigma)}c_{\phi_2}| \sim 1$ (these two terms should be in opposite signs) above M_3 . These conditions can be satisfied, e.g. by $\delta = \rho = \sigma = \phi_1 = 0$ and $\phi_2 = \pi$, which are stable against RG corrections. In contrast, the correction is smallest both at energies above and below M_3 when $c_{(\delta-\rho)} \sim c_{(\delta-\sigma)} \sim 0$. This condition can be satisfied, e.g. by $\delta = 0$ and $\rho = \sigma = \pi/2$, which are stable from M_Z to Λ_{GUT} if there is $\phi_1 = \phi_2 = \pi/2$ in addition.
- For θ_z: The sign of the dominant correction to θ_z depends on CP-violating phases and on the competition among terms led by ζ₁₃⁻¹ and ζ₂₃⁻¹ in Eq.(16). From M_Z to M₃, the O (30°) correction to θ_z is quite spectacular: it means that a too small value is no longer natural for θ_z. However, there is still a notable exception: if c_ρ ~ c_σ ~ 0 in Eq.(16), the correction to θ_z is strongly damped both at energies above and below M₃, and so θ_z can be kept at a small value. Only in this case, can a tiny θ_z be probable. However, for this to happen, ρ and σ must also be stable against RG corrections. In Eqs.(17)-(19) and Eqs.(C.4)-(C.6), we find that if s_(ρ-σ) ~ c_(δ-ρ) ~ c_(δ-σ) ~ c_{φ1} ~ c_{φ2} ~ 0, corrections to δ, ρ, σ, φ₁ and φ₂ can all be strongly damped, and so that δ, ρ, σ, φ₁ and φ₂ are all stable against RG corrections. Up to π, these conditions mean δ = 0 and ρ = σ = φ₁ = φ₂ = π/2. In contrast, the correction to θ_z can be large both at energies above and below M₃, if c_ρc_(δ-ρ) ≈ -c_σc_(δ-σ) and c_ρc_{φ1} ≈ c_σc_{φ2}, and all of them being of O(1) in magnitude. A simple but interesting phase configuration that can satisfy these conditions is δ = ρ + π/4 = σ π/4 = π/2 and φ₁ = φ₂ π = 0 (which, however, is not stable against RG corrections).

For CP-violating phases, possible corrections are roughly of $\mathcal{O}(100^\circ)$ both at energies above and below M_3 , if only the factors ζ_{ij}^{-1} are taken into account. But if the smallness of θ_z is retained during a small energy range, there can be extraordinarily large corrections to CP-violating phases. As in previous cases, such corrections often drive CP-violating phases to near their (pseudo-) fixed points dramatically. However, CP-violating phases usually need to be kept at special values if one wants to damp all large corrections to θ_z , just as we have mentioned above. So in such special cases, there should not be any large corrections to CP-violating phases, though θ_z may be tiny in a wide energy range such as from M_Z to Λ_{GUT} .

- For δ : Eq.(17) for the running of δ is rather complicated. But much simplified approximation can be obtained in the limit $\theta_z \to 0$, which is given in Eq.(53). A notable feature is that, only terms led by ζ_{13}^{-1} and ζ_{23}^{-1} are possibly enhanced by s_z^{-1} . The contribution from these terms can be dominant when θ_z is small enough ($\theta_z < |\zeta_{13}^{-1}/\zeta_{12}^{-1}| \sim |\zeta_{23}^{-1}/\zeta_{12}^{-1}| \sim \mathcal{O}(1^\circ)$).
- For ρ and σ : Eqs.(18) and (19) for the running of ρ and σ are also quite complicated. But the dominant contribution from y_{ν} is now simple. We can predict signs of corrections enhanced by different factors ζ_{ij}^{-1} with the help of Table 2, where the association of enhancing factors ζ_{ij}^{-1} with CP-violating phases is clearly shown. Furthermore, much simplified approximations of Eqs.(18) and (19) can be obtained in the limit $\theta_z \to 0$ and the results are also the same as given in Eq.(53).

In Figure 8, we illustrate cases in which the correction to a specific mixing angle is mostly enhanced or damped, just as discussed above. But the corrections shown are not largest in general. There can be larger corrections when specially chosen CP-violating phases are used. However, this is not our main concern here. What we want to demonstrate is that a good prediction of RG corrections can often be made with the help of Eqs.(14)-(19). For CPviolating phases, the situation is quite complicated and few general conclusions regarding their RG evolution can be reached. In Figure 8, we only give an example to illustrate how the RG evolution of CP-violating phases may be affected by θ_z .

D. The Inverted Hierarchy Case

What is special with the inverted hierarchy case is that only $|\zeta_{12}^{-1}|$ is moderately large among the factors $|\zeta_{ij}^{-1}|$ (for i < j; i, j = 1, 2, 3) defined in Eq.(12). So more interesting corrections can be expected than in the normal hierarchy case. However, since only $|\zeta_{12}^{-1}|$ is large, the situation will not be so complicated as the near degeneracy case. Similar to Eq.(48), we find in the limit $|\zeta_{12}^{-1}| >> 1$ and $s_z \to 0$:

$$\dot{\theta}_x = \frac{1}{\zeta_{12}} \left[C^l_\kappa y^2_\tau c^2_{(\rho-\sigma)} s^2_y - C^\nu_\kappa y^2_\nu c_{(\rho-\sigma)} c_{(\phi_1-\phi_2)} \frac{c_2 s_1 s_2}{c_x s_x} \right] c_x s_x + \cdots,$$
(54)

and

$$\dot{\delta} = \frac{1}{\zeta_{12}} \left[C_{\kappa}^{l} y_{\tau}^{2} c_{(\rho-\sigma)} s_{(\rho-\sigma)} s_{y}^{2} - C_{\kappa}^{\nu} y_{\nu}^{2} c_{(\phi_{1}-\phi_{2})} s_{(\rho-\sigma)} \frac{c_{2} s_{1} s_{2}}{c_{x} s_{x}} \right] + \mathcal{O}\left(s_{z}^{-1}\right) + \cdots;$$

$$\dot{\rho} = \frac{1}{\zeta_{12}} \left[C_{\kappa}^{l} y_{\tau}^{2} c_{(\rho-\sigma)} s_{(\rho-\sigma)} s_{y}^{2} - C_{\kappa}^{\nu} y_{\nu}^{2} c_{(\phi_{1}-\phi_{2})} s_{(\rho-\sigma)} \frac{c_{2} s_{1} s_{2}}{c_{x} s_{x}} \right] s_{x}^{2} + \mathcal{O}\left(s_{z}^{-1}\right) + \cdots;$$

$$\dot{\sigma} = \frac{1}{\zeta_{12}} \left[C_{\kappa}^{l} y_{\tau}^{2} c_{(\rho-\sigma)} s_{(\rho-\sigma)} s_{y}^{2} - C_{\kappa}^{\nu} y_{\nu}^{2} c_{(\phi_{1}-\phi_{2})} s_{(\rho-\sigma)} \frac{c_{2} s_{1} s_{2}}{c_{x} s_{x}} \right] c_{x}^{2} + \mathcal{O}\left(s_{z}^{-1}\right) + \cdots,$$
(55)

where the $\mathcal{O}(s_z^{-1})$ terms are exactly the same for $\dot{\delta}$, $\dot{\rho}$ and $\dot{\sigma}$, and have been given in Eq.(48). It is notable that terms led by ζ_{12}^{-1} are *not* enhanced by s_z^{-1} . As a result, there are no more extraordinary corrections in this case than in the normal hierarchy case with extremely small θ_z . However, the inclusion of a moderately large ζ_{12}^{-1} leads to two non-trivial consequences: (a) the correction to θ_x can now be much larger than in the normal hierarchy case, and (b) there can be much larger corrections to CP-violating phases than in the normal hierarchy case, when s_z^{-1} is not an efficient enhancing factor.

1. RG Correction in the SM

In the normal hierarchy case, enhancing factors $|\zeta_{ij}^{-1}| \sim \mathcal{O}(1)$ and RG corrections to mixing angles are negligible (except for θ_z when it is extremely tiny): $\mathcal{O}((10^{-4})^{\circ})$ in the energy range from M_Z to M_3 and $\mathcal{O}(0.1^{\circ})$ from M_3 to Λ_{GUT} . In the present case when $|\zeta_{12}^{-1}|$ is large, only the correction to θ_x is enhanced: $\mathcal{O}(0.01^{\circ})$ below M_3 and $\mathcal{O}(10^{\circ})$ above M_3 . Note that the correction to θ_x becomes largest when $|c_{(\rho-\sigma)}c_{(\phi_1-\phi_2)}| \sim 1$ in Eq.(54). Corrections to CP-violating phases are dominated by ζ_{12}^{-1} when $|\zeta_{12}^{-1}| >> s_z^{-1}$ (or equiv-

Corrections to CP-violating phases are dominated by ζ_{12}^{-1} when $|\zeta_{12}^{-1}| >> s_z^{-1}$ (or equivalently, $\theta_z >> |\zeta_{12}| \sim 0.01 \approx 0.5^{\circ}$). In this case, the situation for CP-violating phases is the same as that for θ_x . So the contribution from y_{τ} is negligible and large corrections of $\mathcal{O}(10^{\circ})$ are possible only in the energy range from M_3 to Λ_{GUT} . In Eq.(55), contributions to CP-violating phases become largest when $|c_{(\phi_1-\phi_2)}s_{(\rho-\sigma)}| \sim 1$.

When $\theta_z \ll 0.5^\circ$, corrections to CP-violating phases are dominated by s_z^{-1} and the situation is similar to the normal hierarchy case.

In Figure 9, we illustrate the typical evolution behavior of $\theta_x, \theta_z, \delta, \rho$ and σ in the SM. The competition between contributions from ζ_{12}^{-1} and s_z^{-1} is obvious.

2. RG Correction in the MSSM

In the MSSM when $\tan \beta \sim 10$, the contribution from y_{τ} is about $100 (\approx \tan^2 \beta)$ times larger than that in the SM but the contribution from y_{ν} is the same. So corrections to θ_y and θ_z are of $\mathcal{O}(0.01^\circ)$ in the energy range from M_Z to M_3 and of $\mathcal{O}(0.1^\circ)$ from M_3 to Λ_{GUT} , while the correction to θ_x (enhanced by ζ_{12}^{-1}) is of $\mathcal{O}(1^\circ)$ below M_3 and of $\mathcal{O}(10^\circ)$ above M_3 . Corrections to CP-violating phases are the same as that to θ_x when $|\zeta_{12}^{-1}| >> s_z^{-1}$ (or equivalently, $\theta_z >> 0.5^\circ$).

In the MSSM when $\tan \beta \sim 50$, the correction from y_{τ} is about 2500 times larger than that in the SM but the contribution from y_{ν} is the same. So corrections to θ_y and θ_z are of $\mathcal{O}(0.1^{\circ} \sim 1^{\circ})$ both at energies above and below M_3 , while the correction to θ_x is of $\mathcal{O}(30^{\circ})$ in the range from M_Z to M_3 but is still of $\mathcal{O}(10^{\circ})$ from M_3 to Λ_{GUT} . Note that in Eq.(54), the y_{τ} correction to θ_x is always *negative* in the MSSM, while the sign of the contribution from y_{ν} depends on simple phase factors. Corrections to CP-violating phases are the same as that to θ_x when ζ_{12}^{-1} is dominant. When $\theta_z \ll 0.5^{\circ}$, corrections to CP-violating phases are dominated by s_z^{-1} and the situation is similar to the normal hierarchy case. This is true no matter tan β is small or large.

In Figure 10, we illustrate the typical evolution behavior of θ_x , θ_z , δ , ρ and σ in the MSSM. In the calculation, we take $\rho \approx \sigma$ to make the correction to θ_x significant, and $|\rho - \sigma| \sim 90^\circ$ to make corrections to θ_z , δ , ρ and σ significant.

IV. SUMMARY

In this work, we have derived one-loop renormalization group equations for left-handed neutrino masses, leptonic mixing angles and CP-violating phases, both in the SM and the MSSM extended with three right-handed neutrinos. At energies above the seesaw threshold, we show explicitly the contribution from the Yukawa coupling matrix that connects leftand right-handed neutrinos. For simplicity, we have assumed hierarchical eigenvalues of this matrix in our derivation, so our analytical results may not be applicable when the eigenvalues are not hierarchical. And since we have also simplified the task by decoupling all right-handed neutrinos at a common scale, the discussion may have to be modified when the RG evolution between right-handed neutrino thresholds is important [15,28].

Based on these equations, we study possible RG corrections related to three typically interesting neutrino mass patterns: normal hierarchy, near degeneracy and inverted hierarchy.

We firstly study the RG evolution of the factors ζ_{ij} (for i < j; i, j = 1, 2, 3) defined in Eq.(12). We find that $|\zeta_{ij}^{-1}|$ can be significantly damped from M_Z to Λ_{GUT} , both in the near degeneracy case and in the inverted hierarchy case. It is also possible that $|\zeta_{ij}^{-1}|$ may develop extremely high and narrow peaks, so that the situation is nearly singular. However, signs of ζ_{ij} are not likely to be changed, neither is the order of sequence of left-handed neutrino masses.

In the normal hierarchy case, RG corrections from M_Z to Λ_{GUT} are always negligible for mixing angles, except for θ_z when it is extremely small. Appreciable or even significant RG corrections to CP-violating phases are possible only when $\theta_z < \mathcal{O}(1^\circ)$. In the SM, dominant RG corrections to CP-violating phases generally arise in the energy range from M_3 to Λ_{GUT} . In the MSSM when $\tan \beta$ is large, dominant RG corrections generally arise from M_Z to M_3 . Only in the MSSM when $\tan \beta$ is about 10, can large corrections to CP-violating phases arise both at energies above and below M_3 .

In the near degeneracy case, possible large corrections to mixing angles and CP-violating phases are plethora. Mixing angles and CP-violating phases are often driven to near their (pseudo-) fixed points, since corrections are usually very large. Interesting mixing angles at high energy scales are often possible. For example, it is natural to find a large θ_z (comparable to θ_x , θ_y) at Λ_{GUT} .

In the inverted hierarchy case, only ζ_{12}^{-1} and s_z^{-1} are significant enhancing factors. So the situation is much like that in the normal hierarchy case. However, because of the large ζ_{12}^{-1} , the correction to θ_x can be large, and significant RG corrections to CP-violating phases are possible even when θ_z^{-1} is not an efficient enhancing factor.

To conclude, since RG corrections play a significant role in relating the low- and highenergy physics, an analytical understanding of the RG evolution behavior of neutrino parameters is necessary and important. Following earlier works, we have extended this understanding beyond the seesaw threshold by deriving RGEs for left-handed neutrino masses, leptonic mixing angles and CP-violating phases running at energies above the heaviest right-handed neutrino mass, under a few reasonable simplifications. The significance of these equations are demonstrated by studying the RG correction related to three especially interesting neutrino mass patterns. We expect that our work will be very useful for building realistic neutrino mass models at high energy scales.

ACKNOWLEDGMENTS

The author is indebted to Professor Zhi-zhong Xing for reading the manuscript with great care and patience, and also for his valuable comments and numerous corrections. This work was supported in part by the National Nature Science Foundation of China.

Note added

When this work is being completed, we notice that another paper about the similar topic is released by S. Antusch et al [36]. However, our strategy and approach are apparently different from theirs. The two works are complementary to each other.

APPENDIX

APPENDIX A: ONE-LOOP RGES FOR LEPTONIC YUKAWA COUPLING MATRICES

Let Y_u , Y_d , Y_l and Y_{ν} denote Yukawa coupling matrices of up-type quarks, down-type quarks, charged leptons and the one that connects left- and right-handed neutrinos, respectively. At energies below the seesaw threshold, one-loop RGEs for Y_l and κ are $(H_f \equiv Y_f Y_f^{\dagger})$ for $f = u, d, l, \nu$:

$$\dot{Y}_l = \left(\hat{\alpha}_l + \hat{N}_l\right) Y_l; \qquad \qquad \hat{N}_l = C_l^l H_l; \qquad (A.1)$$

$$\dot{\kappa} = \hat{\alpha}_{\kappa}\kappa + \hat{N}_{\kappa}\kappa + \kappa\hat{N}_{\kappa}^{T}; \quad \hat{N}_{\kappa} = C_{\kappa}^{l}H_{l}.$$
(A.2)

For coefficients, we have in the MSSM

$$\hat{\alpha}_{l} = -\left(\frac{9}{5}g_{1}^{2} + 3g_{2}^{2}\right) + \operatorname{Tr}\left(3H_{d} + H_{l}\right); \quad C_{l}^{l} = 3;$$
$$\hat{\alpha}_{\kappa} = -\left(\frac{6}{5}g_{1}^{2} + 6g_{2}^{2}\right) + \operatorname{Tr}\left(6H_{u}\right); \qquad C_{\kappa}^{l} = 1;$$
(A.3)

and in the SM

$$\hat{\alpha}_{l} = -\left(\frac{9}{4}g_{1}^{2} + \frac{9}{4}g_{2}^{2}\right) + \operatorname{Tr}\left(3H_{u} + 3H_{d} + H_{l}\right); \qquad C_{l}^{l} = \frac{3}{2};$$
$$\hat{\alpha}_{\kappa} = -\left(3g_{2}^{2} - \lambda\right) + \operatorname{Tr}\left(6H_{u} + 6H_{d} + 2H_{l}\right); \qquad C_{\kappa}^{l} = -\frac{3}{2}.$$
(A.4)

At energies above the heaviest right-handed neutrino mass, one-loop RGEs for Y_l, Y_{ν} and M_R are

$$\dot{Y}_l = (\alpha_l + N_l) Y_l;$$
 $N_l = C_l^l H_l + C_l^{\nu} H_{\nu};$ (A.5)

$$\dot{Y}_{\nu} = (\alpha_{\nu} + N_{\nu}) Y_{\nu}; \qquad N_{\nu} = C_{\nu}^{l} H_{l} + C_{\nu}^{\nu} H_{\nu}; \qquad (A.6)$$

$$\dot{M}_R = N_R M_R + M_R N_R^T; \qquad N_R = C_R (Y_\nu^{\dagger} Y_\nu)^T.$$
 (A.7)

In the MSSM,

$$\alpha_l = -\left(\frac{9}{5}g_1^2 + 3g_2^2\right) + \text{Tr}\left(3H_d + H_l\right);$$
(A.8)

$$\alpha_{\nu} = -\left(\frac{3}{5}g_1^2 + 3g_2^2\right) + \text{Tr}\left(3H_u + H_{\nu}\right); \tag{A.9}$$

$$C_l^l = 3; \ C_l^\nu = 1; \ C_\nu^l = 1; \ C_\nu^\nu = 3; \ C_R = 2.$$
 (A.10)

In the SM,

$$\alpha_l = -\left(\frac{9}{4}g_1^2 + \frac{9}{4}g_2^2\right) + \text{Tr}\left(3H_u + 3H_d + H_l + H_\nu\right); \tag{A.11}$$

$$\alpha_{\nu} = -\left(\frac{9}{20}g_1^2 + \frac{9}{4}g_2^2\right) + \operatorname{Tr}\left(3H_u + 3H_d + H_l + H_\nu\right); \tag{A.12}$$

$$C_l^l = \frac{3}{2}; \ C_l^\nu = -\frac{3}{2}; \ C_\nu^l = -\frac{3}{2}; \ C_\nu^\nu = \frac{3}{2}; \ C_R = 1.$$
 (A.13)

If we make use of κ also at energies above the seesaw threshold, we can derive from Eqs.(4), (A.6) and (A.7)

$$\dot{\kappa} = \alpha_{\kappa}\kappa + N_{\kappa}\kappa + \kappa N_{\kappa}^{T} ; \qquad (A.14)$$

$$\alpha_{\kappa} = 2\alpha_{\nu} ; \quad N_{\kappa} = N_{\nu} - \tilde{N}_{R} = C_{\kappa}^{l} H_{l} + C_{\kappa}^{\nu} H_{\nu} ,$$
(A.15)

where $\tilde{N}_R = C_R H_{\nu}$, with C_R being the same as that in Eq.(A.7). Also, C_{κ}^l is the same as that in Eq.(A.2). In the MSSM, $C_{\kappa}^{\nu} = 1$; and in the SM, $C_{\kappa}^{\nu} = 1/2$.

APPENDIX B: DERIVATION OF RGES FOR INDIVIDUAL PARAMETERS

In the same way as in Refs. [18,27], we need to calculate $T_{\rm MNS} = U_{\rm MNS}^{\dagger} \dot{U}_{\rm MNS}$ to find out RGEs for leptonic mixing angles and CP-violating phases. From Eq.(8),

$$T_{\rm MNS} = U_{\rm MNS}^{\dagger} \left(U_l^{\dagger} \dot{U}_{\kappa} + \dot{U}_l^{\dagger} U_{\kappa} \right) = U_{\kappa}^{\dagger} \dot{U}_{\kappa} + U_{\rm MNS}^{\dagger} \dot{U}_l^{\dagger} U_l U_{\rm MNS}$$
$$= T_{\kappa} - U_{\rm MNS}^{\dagger} T_l U_{\rm MNS} , \qquad (B.1)$$

where

$$T_{\kappa} \equiv U_{\kappa}^{\dagger} \dot{U}_{\kappa} = -T_{\kappa}^{\dagger} ; \quad T_{l} \equiv U_{l}^{\dagger} \dot{U}_{l} = -T_{l}^{\dagger} .$$
 (B.2)

For T_{κ} , we find from Eqs.(A.14), (6), (7) and (10)

$$\dot{\kappa'} + T_{\kappa}\kappa' + \kappa'T_{\kappa}^{T} = 2\alpha_{\kappa}\kappa' + \tilde{N}_{\kappa}\kappa' + \kappa'\tilde{N}_{\kappa}^{T}; \qquad (B.3)$$

$$\tilde{N}_{\kappa} = U_{\kappa}^{\dagger} N_{\kappa} U_{\kappa} = C_{\kappa}^{l} U_{\text{MNS}}^{\dagger} H_{l}^{\prime} U_{\text{MNS}} + C_{\kappa}^{\nu} U_{\nu} H_{\nu}^{\prime} U_{\nu}^{\dagger} .$$
(B.4)

In Eq.(B.3), for diagonal elements (i = 1, 2, 3)

$$\dot{k}_i = 2\left(\alpha_{\kappa} + \operatorname{Re}\tilde{N}_{\kappa.ii}\right)k_i \; ; \quad \operatorname{Im}T_{\kappa.ii} = \operatorname{Im}\tilde{N}_{\kappa.ii} = 0 \; ; \tag{B.5}$$

and for off-diagonal elements (i, j = 1, 2, 3)

$$T_{\kappa.ij}k_j - k_i T^*_{\kappa.ij} = \tilde{N}_{\kappa.ij}k_j + k_i \left(\tilde{N}^T_{\kappa}\right)_{ij} = \tilde{N}_{\kappa.ij}k_j + k_i \tilde{N}_{\kappa.ji} ,$$

$$\Longrightarrow \begin{cases} \operatorname{Re}T_{\kappa.ij} = -\frac{k_i + k_j}{k_i - k_j} \operatorname{Re}\tilde{N}_{\kappa.ij} \equiv -\zeta_{ij}^{-1} \operatorname{Re}\tilde{N}_{\kappa.ij} ; \\ \operatorname{Im}T_{\kappa.ij} = -\frac{k_i - k_j}{k_i + k_j} \operatorname{Im}\tilde{N}_{\kappa.ij} \equiv -\zeta_{ij} \operatorname{Im}\tilde{N}_{\kappa.ij} . \end{cases}$$
(B.6)

For T_l , we find from Eq.(A.5)

$$\dot{H}_l = 2\alpha_l H_l + N_l H_l + H_l N_l . \tag{B.7}$$

Then from Eqs.(7) and (10)

$$\dot{H}'_{l} + T_{l}H'_{l} + H'_{l}T^{\dagger}_{l} = 2\alpha_{l}H'_{l} + \tilde{N}_{l}H'_{l} + H'_{l}\tilde{N}_{l} ; \qquad (B.8)$$

$$\tilde{N}_{l} = U_{l}^{\dagger} N_{l} U_{l} = C_{l}^{l} H_{l}^{\prime} + C_{l}^{\nu} \left(U_{\text{MNS}} U_{\nu} \right) H_{\nu}^{\prime} \left(U_{\text{MNS}} U_{\nu} \right)^{\dagger} .$$
(B.9)

In Eq.(B.8), for diagonal elements (i = 1, 2, 3)

$$\dot{H}_{l.ii}' = 2\left(\alpha_l + \tilde{N}_{l.ii}\right) H_{l.ii}' , \qquad (B.10)$$

and for off-diagonal elements (i, j = 1, 2, 3)

$$T_{l.ij}H'_{l.jj} + H'_{l.ii} \left(T_l^{\dagger}\right)_{ij} = \tilde{N}_{l.ij}H'_{l.jj} + H'_{l.ii}\tilde{N}_{l.ij}$$

$$\implies T_{l.ij} = -\frac{H'_{l.ii} + H'_{l.jj}}{H'_{l.ii} - H'_{l.jj}}\tilde{N}_{l.ij} .$$
(B.11)

Note that $T_{l,ii}$ (for i = 1, 2, 3) are arbitrary, since U_l is only determined up to a diagonal phase matrix on its right.

Furthermore, in order to derive RGEs for parameters in Y_{ν} , we find from Eq.(A.6)

$$\dot{H}_{\nu} = 2\alpha_{\nu}H_{\nu} + N_{\nu}H_{\nu} + H_{\nu}N_{\nu} .$$
(B.12)

Then from Eq.(10)

$$\dot{H}'_{\nu} + \left(T_{\nu} + U^{\dagger}_{\nu}T_{\kappa}U_{\nu}\right)H'_{\nu} + H'_{\nu}\left(T_{\nu} + U^{\dagger}_{\nu}T_{\kappa}U_{\nu}\right)^{\dagger} = 2\alpha_{\nu}H'_{\nu} + \tilde{N}_{\nu}H'_{\nu} + H'_{\nu}\tilde{N}_{\nu}; \qquad (B.13)$$

$$N_{\nu} = C_{\nu}^{l} \left(U_{\rm MNS} U_{\nu} \right)^{\dagger} H_{l}^{\prime} \left(U_{\rm MNS} U_{\nu} \right) + C_{\nu}^{\nu} H_{\nu}^{\prime} ; \qquad T_{\nu} \equiv U_{\nu}^{\dagger} U_{\nu} . \qquad (B.14)$$

In Eq.(B.13), for diagonal elements (i = 1, 2, 3)

$$H'_{\nu.ii} = 2\left(\alpha_{\nu} + \tilde{N}_{\nu.ii}\right) H'_{\nu.ii} ; \qquad (B.15)$$

and for off-diagonal elements (i, j = 1, 2, 3)

$$\left(T_{\nu} + U_{\nu}^{\dagger} T_{\kappa} U_{\nu} \right)_{ij} H_{\nu,ij}' - H_{\nu,ii}' \left(T_{\nu} + U_{\nu}^{\dagger} T_{\kappa} U_{\nu} \right)_{ij} = \tilde{N}_{\nu,ij} H_{\nu,ij}' + H_{\nu,ii}' \tilde{N}_{\nu,ij} ,$$

$$\Longrightarrow T_{\nu,ij} = -\frac{H_{\nu,ii}' + H_{\nu,jj}'}{H_{\nu,ii}' - H_{\nu,jj}'} \tilde{N}_{\nu,ij} - \left(U_{\nu}^{\dagger} T_{\kappa} U_{\nu} \right)_{ij} .$$
(B.16)

Just like the diagonal elements of T_l , $T_{\nu,ii}$ (for i = 1, 2, 3) are also arbitrary since U_{ν} is only determined up to a diagonal phase matrix on its right.

To calculate $\dot{\theta}_x$, $\dot{\theta}_y$, $\dot{\theta}_z$, $\dot{\delta}$, $\dot{\rho}$ and $\dot{\sigma}$ from $T_{\rm MNS}$, an auxiliary diagonal phase matrix is required on the left hand side of $U_{\rm MNS}$ as defined in Eq.(9), i.e. we have to use a more general parametrization of the MNS matrix in Eq.(B.1):

$$U_{\rm MNS} = P \cdot V \; ; \quad P \equiv \begin{pmatrix} e^{i\alpha} & & \\ & e^{i\beta} & \\ & & e^{i\gamma} \end{pmatrix} \; , \tag{B.17}$$

where V denoting the original $U_{\rm MNS}$ defined in Eq.(9). Then from Eq.(B.4)

$$\tilde{N}_{\kappa} = C^l_{\kappa} V^{\dagger} H^{\prime}_l V + C^{\nu}_{\kappa} U_{\nu} H^{\prime}_{\nu} U^{\dagger}_{\nu} . \qquad (B.18)$$

Together with Eqs.(B.5) and (B.6), this equation shows that T_{κ} is independent of the phase matrix P. Furthermore, from Eqs.(B.9) and (B.11), the product (P^*T_lP) is also independent of P. So in an equivalence of Eq.(B.1)

$$\dot{V}V^{\dagger} = VT_{\kappa}V^{\dagger} - P^{*}\left(T_{l} + P^{*}\dot{P}\right)P , \qquad (B.19)$$

the equations of off-diagonal elements are obviously independent of the matrix P. They are all together six linearly independent equations of $\dot{\theta}_x$, $\dot{\theta}_y$, $\dot{\theta}_z$, $\dot{\delta}$, $\dot{\rho}$ and $\dot{\sigma}$, and can thus determine these six quantities unambiguously. For the diagonal elements, $(P^*\dot{P})_{ii}$ (i = 1, 2, 3) are only determined up to arbitrary $T_{l.ii}$ (i = 1, 2, 3), but this is of no problem since α , β and γ are not physical by definition. We may choose whatever value for $T_{l.ii}$ (i = 1, 2, 3) as we like in the calculation, or may simply ignore the equations for the diagonal elements in Eq.(B.19).

In contrast, we can see from Eqs.(B.14) and (B.16) that to extract θ_1 , θ_2 , θ_3 , δ_{ν} , ϕ_1 and $\dot{\phi}_2$ from T_{ν} , an auxiliary phase matrix on the right hand side of U_{ν} (just as the phase matrix P on the left hand side of $U_{\rm MNS}$) is not necessary. We can use the U_{ν} defined in Eq.(11) directly during the calculation. There are totally six linearly independent equations of $\dot{\theta}_1$, $\dot{\theta}_2$, $\dot{\theta}_3$, $\dot{\delta}_{\nu}$, $\dot{\phi}_1$ and $\dot{\phi}_2$ in Eq.(B.16), so these quantities can also be determined unambiguously.

APPENDIX C: FULL EXPRESSIONS OF $\dot{\theta}_1, \dot{\theta}_2, \dot{\theta}_3, \dot{\delta}_{\nu}, \dot{\phi}_1$ AND $\dot{\phi}_2$

(Note that for what ever F: $\dot{F} \equiv 16\pi^2 \frac{dF}{dt}$; $t = \ln\mu$, with μ being the energy scale.)

$$\begin{split} \dot{\theta}_{1} &= C_{\kappa}^{l} y_{\tau}^{2} \left\{ \frac{C_{(\phi_{1}-\phi_{2})}}{\zeta_{12}} \left[c_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(c_{(\delta+\rho-\sigma)} c_{x}^{2} - c_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \frac{c_{1} s_{2}}{c_{2}} \right. \\ &\quad \left. - \zeta_{12} \cdot s_{(\phi_{1}-\phi_{2})} \left[s_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(s_{(\delta+\rho-\sigma)} c_{x}^{2} + s_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \frac{c_{1} s_{2}}{c_{2}} \right. \\ &\quad \left. + \left[\frac{c_{\phi_{1}}}{\zeta_{13}} \left(c_{(\delta-\rho)} s_{x} s_{y} - c_{\rho} c_{x} c_{y} s_{z} \right) + \zeta_{13} \cdot s_{\phi_{1}} \left(s_{(\delta-\rho)} s_{x} s_{y} + s_{\rho} c_{x} c_{y} s_{z} \right) \right] \frac{c_{y} c_{z} s_{1} s_{2}}{c_{2}} \\ &\quad \left. - \left[\frac{c_{\phi_{2}}}{\zeta_{23}} \left(c_{(\delta-\sigma)} c_{x} s_{y} + c_{\sigma} c_{y} s_{x} s_{z} \right) + \zeta_{23} \cdot s_{\phi_{2}} \left(s_{(\delta-\sigma)} c_{x} s_{y} - s_{\sigma} c_{y} s_{x} s_{z} \right) \right] c_{y} c_{z} \right\} \\ &\quad + C_{\kappa}^{\nu} y_{\nu}^{2} \left\{ - \left(\frac{c_{(\phi_{1}-\phi_{2})}^{2}}{\zeta_{12}} + \zeta_{12} \cdot s_{(\phi_{1}-\phi_{2})}^{2} \right) c_{1} s_{1} s_{2}^{2} \\ &\quad + \left(\frac{c_{\phi_{1}}^{2}}{\zeta_{13}} + \zeta_{13} \cdot s_{\phi_{1}}^{2} \right) c_{1} s_{1} s_{2}^{2} + \left(\frac{c_{\phi_{2}}^{2}}{\zeta_{23}} + \zeta_{23} \cdot s_{\phi_{2}}^{2} \right) c_{1} c_{2}^{2} s_{1} \right\} \\ &\quad + C_{\nu}^{l} y_{\tau}^{2} \left\{ c_{1} s_{1} \left[c_{x}^{2} s_{y}^{2} - c_{y}^{2} \left(c_{z}^{2} - s_{x}^{2} s_{z}^{2} \right] \right\} \\ &\quad + C_{\nu}^{l} y_{\tau}^{2} \left\{ c_{1} s_{1} \left[c_{x}^{2} s_{y}^{2} - c_{y}^{2} \left(c_{z}^{2} - s_{x}^{2} s_{z}^{2} \right] \right\} \\ &\quad + \left[c_{(\rho+\phi_{1})} c_{x} s_{1} s_{2} - c_{(\sigma+\phi_{2})} \left(c_{1}^{2} - s_{1}^{2} \right) c_{2} s_{x} \right] \frac{c_{y}^{2} c_{z} s_{z}}{c_{2}} \\ &\quad - \left[c_{(\delta-\rho-\phi_{1})} s_{1} s_{2} s_{x} + c_{(\delta-\sigma-\phi_{2})} c_{2} c_{x} \left(c_{1}^{2} - s_{1}^{2} \right) \right] \frac{c_{y} c_{z} s_{y} s_{z}}{c_{2}} \\ &\quad - \left[c_{(\delta+\rho-\sigma+\phi_{1}-\phi_{2})} \frac{c_{1} c_{x} s_{2} s_{x}}{c_{2}} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) \\ \\ &\quad + \left[c_{(\delta+\rho-\sigma+\phi_{1}-\phi_{2})} c_{x}^{2} - c_{(\delta-\rho+\sigma-\phi_{1}+\phi_{2})} s_{x}^{2} \right] \frac{c_{1} c_{y} s_{2} s_{y} s_{z}}{c_{2}} \\ \\ &\quad - \left[c_{(\delta+\rho-\sigma+\phi_{1}-\phi_{2})} c_{x}^{2} - c_{(\delta-\rho+\sigma-\phi_{1}+\phi_{2})} s_{x}^{2} \right] \frac{c_{1} c_{y} s_{2} s_{y} s_{z}}{c_{2}} \\ \\ &\quad - \left[c_{(\delta+\rho-\sigma+\phi_{1}-\phi_{2})} c_{x}^{2} - c_{(\delta-\rho+\sigma-\phi_{1}+\phi_{2})} s_{x$$

$$\begin{split} \dot{\theta}_{2} &= C_{\kappa}^{l} y_{\tau}^{2} \left\{ -\frac{c_{(\phi_{1}-\phi_{2})}}{\zeta_{12}} \left[c_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(c_{(\delta+\rho-\sigma)} c_{x}^{2} - c_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] s_{1} \right. \\ &+ \zeta_{12} \cdot s_{(\phi_{1}-\phi_{2})} \left[s_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(s_{(\delta+\rho-\sigma)} c_{x}^{2} + s_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] s_{1} \\ &+ \left[\frac{c_{\phi_{1}}}{\zeta_{13}} \left(c_{(\delta-\rho)} s_{x} s_{y} - c_{\rho} c_{x} c_{y} s_{z} \right) + \zeta_{13} \cdot s_{\phi_{1}} \left(s_{(\delta-\rho)} s_{x} s_{y} + s_{\rho} c_{x} c_{y} s_{z} \right) \right] c_{1} c_{y} c_{z} \right\} \\ &+ C_{\kappa}^{\nu} y_{\nu}^{2} \left\{ \left(\frac{c_{(\phi_{1}-\phi_{2})}}{\zeta_{12}} + \zeta_{12} \cdot s_{(\phi_{1}-\phi_{2})}^{2} \right) c_{2} s_{1}^{2} s_{2} + \left(\frac{c_{\phi_{1}}^{2}}{\zeta_{13}} + \zeta_{13} \cdot s_{\phi_{1}}^{2} \right) c_{1}^{2} c_{2} s_{2} \right\} \\ &+ C_{\nu}^{l} y_{\tau}^{2} \left\{ -c_{2} s_{2} \left[\left(c_{x}^{2} s_{1}^{2} - s_{x}^{2} \right) s_{y}^{2} + c_{y}^{2} \left(c_{1}^{2} c_{z}^{2} - \left(c_{x}^{2} - s_{1}^{2} s_{x}^{2} \right) s_{z}^{2} \right) \right] \\ &- 2 c_{\delta} c_{2} c_{x} c_{y} \left(1 + s_{1}^{2} \right) s_{2} s_{x} s_{y} s_{z} - \left[c_{(\rho+\phi_{1})} c_{x} \left(c_{2}^{2} - s_{2}^{2} \right) - 2 c_{(\sigma+\phi_{2})} c_{2} s_{1} s_{2} s_{x} \right] c_{1} c_{y}^{2} c_{z} s_{z} \\ &+ \left[c_{(\delta-\rho-\phi_{1})} \left(c_{2}^{2} - s_{2}^{2} \right) s_{x} + 2 c_{(\delta-\sigma-\phi_{2})} c_{2} c_{x} s_{1} s_{2} \right] c_{1} c_{y} c_{z} s_{y} \\ &- c_{(\rho-\sigma+\phi_{1}-\phi_{2})} c_{x}^{2} - c_{(\delta-\rho+\sigma-\phi_{1}+\phi_{2})} s_{x}^{2} \right] c_{y} s_{1} \left(c_{2}^{2} - s_{2}^{2} \right) s_{y} s_{z} \right\} . \tag{C.2}$$

$$\dot{\theta}_3 = C_{\kappa}^l y_{\tau}^2 \left\{ -\frac{c_{(\delta_{\nu}+\phi_1-\phi_2)}}{\zeta_{12}} \left[c_{(\rho-\sigma)} c_x s_x \left(s_y^2 - c_y^2 s_z^2 \right) - \left(c_{(\delta+\rho-\sigma)} c_x^2 - c_{(\delta-\rho+\sigma)} s_x^2 \right) c_y s_y s_z \right] \frac{c_1}{c_2} \right\}$$

$$\begin{aligned} +\zeta_{12} \cdot s_{(\delta\nu+\phi_1-\phi_2)} \left[s_{(\rho-\sigma)} c_x s_x \left(s_y^2 - c_y^2 s_z^2 \right) - \left(s_{(\delta+\rho-\sigma)} c_x^2 + s_{(\delta-\rho+\sigma)} s_x^2 \right) c_y s_y s_z \right] \frac{c_1}{c_2} \\ - \left[\frac{c_{(\delta\nu+\phi_1)}}{\zeta_{13}} \left(c_{(\delta-\rho)} s_x s_y - c_\rho c_x c_y s_z \right) + \zeta_{13} \cdot s_{(\delta\nu+\phi_1)} \left(s_{(\delta-\rho)} s_x s_y + s_\rho c_x c_y s_z \right) \right] \frac{c_y c_z s_1}{c_2} \right\} \\ + C_{\nu}^{\nu} y_{\nu}^2 \left\{ \left[\left(\zeta_{12}^{-1} + \zeta_{12} \right) c_{\delta\nu} + \left(\zeta_{13}^{-1} - \zeta_{12} \right) c_{(\delta\nu+2\phi_1-2\phi_2)} \right] \frac{c_1 s_1 s_2}{2} \right] \\ - \left[\left(\zeta_{13}^{-1} + \zeta_{13} \right) c_{\delta\nu} + \left(\zeta_{13}^{-1} - \zeta_{13} \right) c_{(\delta\nu+2\phi_1)} \right] \frac{c_1 s_1 s_2}{2} \right\} \\ + C_{\nu}^{l} y_{\tau}^2 \left\{ c_3 s_3 \left[c_2^2 \left(s_x^2 s_y^2 + c_x^2 c_y^2 s_z^2 \right) - \left(c_1^2 - s_1^2 s_2^2 \right) \left(c_x^2 s_y^2 + c_y^2 s_x^2 s_z^2 \right) - c_y^2 c_z^2 \left(s_1^2 - c_1^2 s_2^2 \right) \right] \\ - 2 c_{\delta c_3} c_x c_y \left(c_1^2 + c_2^2 - s_1^2 s_2^2 \right) s_3 s_x s_y s_z - 2 c_{\delta\nu} c_1 c_3^2 s_1 s_2 \left(c_x^2 s_y^2 - c_y^2 \left(c_z^2 - s_x^2 s_z^2 \right) \right) \\ + 2 \left[c_{(\rho+\phi_1)} c_2 c_x s_2 - c_{(\sigma+\phi_2)} s_1 \left(1 + s_2^2 \right) s_1 \left(c_1 c_3 c_y^2 c_z s_3 s_z \right) \\ + 2 c_{(\rho-\sigma+\phi_1-\phi_2)} c_2^2 c_3 c_x s_1 s_2 s_3 s_x \left(s_y^2 - c_y^2 s_z^2 \right) \\ - 2 \left[c_{(\delta+\rho-\sigma+\phi_1-\phi_2)} c_x^2 - c_{(\delta-\rho+\sigma-\phi_1+\phi_2)} s_x^2 \right] c_2 c_3 c_y s_1 s_2 s_3 s_y s_z \\ + c_{(\delta\nu+\rho+\phi_1)} \frac{c_x c_y^2 c_z s_1 s_2}{c_2} \left(c_2^2 \left(c_3^2 - s_3^2 \right) - s_2^2 \right) \\ + 2 \left[c_{(\delta\nu-\sigma-\phi_2)} c_1^2 - c_{(\delta\nu+\sigma+\phi_2)} s_1^2 \right] c_3^2 c_y^2 c_z s_2 s_x s_z \\ - c_{(\delta-\nu-\rho-\phi_1)} \frac{c_y c_z s_1 s_x s_y}{c_2} \left(c_2^2 \left(c_3^2 - s_3^2 \right) - s_2^2 \right) \\ + 2 \left[c_{(\delta+\delta\nu-\sigma-\phi_2)} c_1^2 - c_{(\delta-\delta\nu-\sigma-\phi_2)} s_1^2 \right] c_3^2 c_x c_y c_z s_2 s_y \\ + \left[c_{(\delta+\delta\nu-\sigma-\phi_2)} c_1^2 - c_{(\delta-\delta\nu-\sigma-\phi_2)} s_1^2 \right] c_3^2 c_x c_y c_z s_2 s_y \\ + \left[c_{(\delta+\delta\nu-\sigma-\phi_2)} c_1^2 - c_{(\delta-\delta\nu-\sigma-\phi_2)} s_1^2 \right] c_3^2 c_x c_y c_z s_2 s_y \\ + \left[c_{(\delta+\delta\nu+\sigma-\phi+\phi_1-\phi_2)} c_x^2 - c_{(\delta-\delta\nu-\sigma-\phi_2)} s_1^2 \right] c_3^2 c_x c_y c_z s_2 s_y \\ + \left[c_{(\delta+\delta\nu+\sigma-\phi+\phi_1-\phi_2)} c_x^2 - c_{(\delta-\delta\nu-\sigma-\phi+\sigma-\phi_1+\phi_2)} s_x^2 \right] \frac{c_1 c_y s_y s_z}{c_2} \left(c_2^2 \left(c_3^2 - s_3^2 \right) - s_2^2 \right) \\ - 4 c_{\delta} c_{\delta\nu} c_1 c_3^2 c_x c_y s_1 s_2 s_x s_y s_z \\ \right] .$$

$$\begin{split} \dot{\delta}_{\nu} &= C_{\kappa}^{l} y_{\tau}^{2} \left\{ -\frac{1}{\zeta_{12}} \left[s_{(\phi_{1}-\phi_{2})} c_{3} \left(s_{1}^{2} - c_{1}^{2} s_{2}^{2} \right) s_{3} - s_{(\delta_{\nu}+\phi_{1}-\phi_{2})} c_{1} s_{1} s_{2} \left(c_{3}^{2} - s_{3}^{2} \right) \right] \\ & \cdot \left[\left\{ c_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(c_{(\delta+\rho-\sigma)} c_{x}^{2} - c_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right\} \frac{1}{c_{2} c_{3} s_{1} s_{2} s_{3}} \right] \\ & - \zeta_{12} \left[c_{(\phi_{1}-\phi_{2})} c_{3} \left(s_{1}^{2} - c_{1}^{2} s_{2}^{2} \right) s_{3} - c_{(\delta_{\nu}+\phi_{1}-\phi_{2})} c_{1} s_{1} s_{2} \left(c_{3}^{2} - s_{3}^{2} \right) \right] \right. \\ & \cdot \left[\left\{ s_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(s_{(\delta+\rho-\sigma)} c_{x}^{2} + s_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right\} \frac{1}{c_{2} c_{3} s_{1} s_{2} s_{3}} \right] \\ & + \frac{1}{\zeta_{13}} \left[s_{\phi_{1}} c_{3} \left(c_{1}^{2} - s_{1}^{2} s_{2}^{2} \right) s_{3} + s_{(\delta_{\nu}+\phi_{1})} c_{1} s_{1} s_{2} \left(c_{3}^{2} - s_{3}^{2} \right) \right] \left(c_{(\delta-\rho)} s_{x} s_{y} - c_{\rho} c_{x} c_{y} s_{z} \right) \frac{c_{y} c_{z}}{c_{1} c_{2} c_{3} s_{2} s_{3}} \right] \\ & - \zeta_{13} \left[c_{\phi_{1}} c_{3} \left(c_{1}^{2} - s_{1}^{2} s_{2}^{2} \right) s_{3} + c_{(\delta_{\nu}+\phi_{1})} c_{1} s_{1} s_{2} \left(c_{3}^{2} - s_{3}^{2} \right) \right] \left(s_{(\delta-\rho)} s_{x} s_{y} + s_{\rho} c_{x} c_{y} s_{z} \right) \frac{c_{y} c_{z}}{c_{1} c_{2} c_{3} s_{2} s_{3}} \right] \\ & + \left[\frac{s_{\phi_{2}}}{\zeta_{23}} \left(c_{(\delta-\sigma)} c_{x} s_{y} + c_{\sigma} c_{y} s_{x} s_{z} \right) - \zeta_{23} \cdot c_{\phi_{2}} \left(s_{(\delta-\sigma)} c_{x} s_{y} - s_{\sigma} c_{y} s_{x} s_{z} \right) \right] \frac{c_{y} c_{z}}{c_{1} s_{1}} \right\} \end{aligned}$$

$$+ C_{\kappa}^{w} y_{\nu}^{2} \left\{ \frac{c_{(\phi_{1}-\phi_{2})}}{\zeta_{12}} \left[s_{(\phi_{1}-\phi_{2})} \left(s_{1}^{2} - c_{1}^{2} s_{2}^{2} \right) - s_{(\delta_{\nu}+\phi_{1}-\phi_{2})} \frac{c_{1} s_{1} s_{2}}{c_{3} s_{3}} \left(c_{3}^{2} - s_{3}^{2} \right) \right] \right. \\ \left. - \zeta_{12} \cdot s_{(\phi_{1}-\phi_{2})} \left[c_{(\phi_{1}-\phi_{2})} \left(s_{1}^{2} - c_{1}^{2} s_{2}^{2} \right) - c_{(\delta_{\nu}+\phi_{1}-\phi_{2})} \frac{c_{1} s_{1} s_{2}}{c_{3} s_{3}} \left(c_{3}^{2} - s_{3}^{2} \right) \right] \right. \\ \left. + \frac{c_{\phi_{1}}}{\zeta_{13}} \left[s_{\phi_{1}} \left(c_{1}^{2} - s_{1}^{2} s_{2}^{2} \right) + s_{(\delta_{\nu}+\phi_{1})} \frac{c_{1} s_{1} s_{2}}{c_{3} s_{3}} \left(c_{3}^{2} - s_{3}^{2} \right) \right] \right. \\ \left. - \zeta_{13} \cdot s_{\phi_{1}} \left[c_{\phi_{1}} \left(c_{1}^{2} - s_{1}^{2} s_{2}^{2} \right) + c_{(\delta_{\nu}+\phi_{1})} \frac{c_{1} s_{1} s_{2}}{c_{3} s_{3}} \left(c_{3}^{2} - s_{3}^{2} \right) \right] - \left(\zeta_{23}^{-1} - \zeta_{23} \right) c_{\phi_{2}} s_{\phi_{2}} c_{2}^{2} \right\} \\ \left. + C_{\nu}^{l} y_{\tau}^{2} \left\{ 2 s_{\delta_{\nu}} \left[c_{x}^{2} s_{y}^{2} - c_{y}^{2} \left(c_{z}^{2} - s_{x}^{2} s_{z}^{2} \right) + 2 c_{\delta} c_{x} c_{y} s_{x} s_{y} s_{z} \right] \frac{c_{1} c_{3} s_{1} s_{2}}{c_{1} s_{1} s_{1}} \\ \left. - s_{(\rho+\phi_{1})} \frac{c_{x} c_{y}^{2} c_{z} s_{x}}{c_{1} c_{2} c_{2} s_{z}} \left[s_{1}^{2} s_{2}^{2} + c_{1}^{2} \left(c_{2}^{2} - s_{2}^{2} \right) \right] + s_{(\sigma+\phi_{2})} \frac{c_{y}^{2} c_{x} s_{x} s_{z}}{c_{1} s_{1}} \left(c_{1}^{2} - s_{1}^{2} \right) \\ \left. - s_{(\rho-\sigma+\phi_{1}-\phi_{2})} \frac{c_{x} s_{x}}{c_{2} s_{1} s_{2}} \left[c_{2}^{2} s_{1}^{2} + c_{1}^{2} \left(c_{2}^{2} - s_{2}^{2} \right) \right] - s_{(\delta-\sigma-\phi_{2})} \frac{c_{x} c_{y} c_{x} s_{y}}{c_{1} s_{1}} \left(c_{1}^{2} - s_{1}^{2} \right) \\ \left. - s_{(\delta_{\nu+\rho-\phi+\phi_{1})} \frac{c_{x} c_{y}^{2} c_{x} s_{x} s_{z}}{c_{2} c_{3} s_{3}} \left[c_{2}^{2} - s_{2}^{2} \left(c_{3}^{2} - s_{3}^{2} \right) \right] - s_{(\delta_{\nu}-\sigma-\phi_{2})} c_{1}^{2} - s_{(\delta_{\nu}+\sigma+\phi_{2})} s_{1}^{2} \right] \frac{c_{3} c_{y}^{2} c_{z} s_{x} s_{x} s_{z}}{s_{3}} \\ \left. + s_{(\delta_{\nu+\rho-\sigma+\phi_{1}-\phi_{2})} c_{1}^{2} + s_{(\delta-\delta_{\nu}-\sigma-\phi-\phi_{2})} s_{1}^{2} \left] \frac{c_{3} c_{x} c_{y} c_{z} s_{x} s_{z}}{s_{3}} \right] \\ \left. - s_{(\delta_{\nu+\phi-\rho-\phi+\phi_{1}-\phi_{2})} c_{1}^{2} + s_{(\delta-\delta_{\nu}-\sigma-\phi-\phi_{2})} s_{1}^{2} \left] \frac{c_{2} c_{x} s_{z}^{2} s_{z} s_{z}}{s_{3}} \\ \left. - s_{(\delta_{\nu+\phi-\rho-\phi+\phi_{1}-\phi_{2})} c_{1}^{2} + s_{(\delta-\delta_{\nu}-\rho+\phi-\phi-\phi_{1}+\phi_{2})} s_{1}^{2}}$$

$$\begin{split} \dot{\phi}_{1} &= C_{\kappa}^{l} y_{\tau}^{2} \left\{ \frac{s_{(\phi_{1}-\phi_{2})}}{\zeta_{12}} \left[c_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(c_{(\delta+\rho-\sigma)} c_{x}^{2} - c_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \frac{c_{2} s_{1}}{s_{2}} \\ &+ \zeta_{12} \cdot c_{(\phi_{1}-\phi_{2})} \left[s_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(s_{(\delta+\rho-\sigma)} c_{x}^{2} + s_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \frac{c_{2} s_{1}}{s_{2}} \\ &- \left[\frac{s_{\phi_{1}}}{\zeta_{13}} \left(c_{(\delta-\rho)} s_{x} s_{y} - c_{\rho} c_{x} c_{y} s_{z} \right) - \zeta_{13} \cdot c_{\phi_{1}} \left(s_{(\delta-\rho)} s_{x} s_{y} + s_{\rho} c_{x} c_{y} s_{z} \right) \right] \frac{c_{y} c_{z} \left(c_{1}^{2} c_{2}^{2} - s_{2}^{2} \right)}{c_{1} c_{2} s_{2}} \\ &- \left[\frac{s_{\phi_{2}}}{\zeta_{23}} \left(c_{(\delta-\sigma)} c_{x} s_{y} + c_{\sigma} c_{y} s_{x} s_{z} \right) - \zeta_{23} \cdot c_{\phi_{2}} \left(s_{(\delta-\sigma)} c_{x} s_{y} - s_{\sigma} c_{y} s_{x} s_{z} \right) \right] \frac{c_{y} c_{z} s_{1}}{c_{1}} \right\} \\ &+ C_{\kappa}^{\nu} y_{\nu}^{2} \left\{ - \left(\zeta_{12}^{-1} - \zeta_{12} \right) c_{(\phi_{1}-\phi_{2})} s_{(\phi_{1}-\phi_{2})} c_{2}^{2} s_{1}^{2} - \left(\zeta_{13}^{-1} - \zeta_{13} \right) c_{\phi_{1}} s_{\phi_{1}} \left(c_{1}^{2} c_{2}^{2} - s_{2}^{2} \right) \right. \\ &+ \left(\zeta_{23}^{-1} - \zeta_{23} \right) c_{\phi_{2}} s_{\phi_{2}} c_{2}^{2} s_{1}^{2} \right\} \\ &+ C_{\nu}^{l} y_{\tau}^{2} \left\{ \left(s_{(\delta-\rho-\phi_{1})} s_{x} s_{y} + s_{(\rho+\phi_{1})} c_{x} c_{y} s_{z} \right) \frac{c_{y} c_{z} \left(c_{1}^{2} + s_{1}^{2} s_{2}^{2} \right)}{c_{1} c_{2} s_{2}} \\ &- \left(s_{(\delta-\sigma-\phi_{2})} c_{x} s_{y} - s_{(\sigma+\phi_{2})} c_{y} s_{x} s_{z} \right) \frac{c_{y} c_{z} \left(c_{1}^{2} + s_{1}^{2} s_{2}^{2} \right)}{c_{1} c_{2} s_{2}} \\ &- \left(s_{(\delta-\sigma-\phi_{2})} c_{x} s_{y} - s_{(\sigma+\phi_{2})} c_{y} s_{x} s_{z} \right) \frac{c_{y} c_{z} s_{1}}{c_{1}} + \left[s_{(\rho-\sigma+\phi_{1}-\phi_{2})} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) \right] \right]$$

$$-\left(s_{(\delta+\rho-\sigma+\phi_{1}-\phi_{2})}c_{x}^{2}+s_{(\delta-\rho+\sigma-\phi_{1}+\phi_{2})}s_{x}^{2}\right)c_{y}s_{y}s_{z}\left]\frac{c_{2}s_{1}}{s_{2}}\right\}$$
(C.5)

$$\begin{split} \dot{\phi}_{2} &= C_{\kappa}^{l} y_{\tau}^{2} \left\{ \frac{s_{(\phi_{1}-\phi_{2})}}{\zeta_{12}} \left[c_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(c_{(\delta+\rho-\sigma)} c_{x}^{2} - c_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \frac{s_{2}}{c_{2} s_{1}} \right\} \\ &+ \zeta_{12} \cdot c_{(\phi_{1}-\phi_{2})} \left[s_{(\rho-\sigma)} c_{x} s_{x} \left(s_{y}^{2} - c_{y}^{2} s_{z}^{2} \right) - \left(s_{(\delta+\rho-\sigma)} c_{x}^{2} + s_{(\delta-\rho+\sigma)} s_{x}^{2} \right) c_{y} s_{y} s_{z} \right] \frac{s_{2}}{c_{2} s_{1}} \\ &+ \left[\frac{s_{\phi_{1}}}{\zeta_{13}} \left(c_{(\delta-\rho)} s_{x} s_{y} - c_{\rho} c_{x} c_{y} s_{z} \right) - \zeta_{13} \cdot c_{\phi_{1}} \left(s_{(\delta-\rho)} s_{x} s_{y} + s_{\rho} c_{x} c_{y} s_{z} \right) \right] \frac{c_{y} c_{z} s_{2}}{c_{1} c_{2}} \\ &+ \left[\frac{s_{\phi_{2}}}{\zeta_{23}} \left(c_{(\delta-\sigma)} c_{x} s_{y} + c_{\sigma} c_{y} s_{x} s_{z} \right) - \zeta_{23} \cdot c_{\phi_{2}} \left(s_{(\delta-\sigma)} c_{x} s_{y} - s_{\sigma} c_{y} s_{x} s_{z} \right) \right] \frac{c_{y} c_{z} \left(c_{1}^{2} - s_{1}^{2} \right)}{c_{1} s_{1}} \right\} \\ &+ C_{\kappa}^{\nu} y_{\nu}^{2} \left\{ - \left(\zeta_{12}^{-1} - \zeta_{12} \right) c_{(\phi_{1}-\phi_{2})} s_{(\phi_{1}-\phi_{2})} s_{2}^{2} + \left(\zeta_{13}^{-1} - \zeta_{13} \right) c_{\phi_{1}} s_{\phi_{1}} s_{2}^{2} \right. \\ &- \left(\zeta_{23}^{-1} - \zeta_{23} \right) c_{\phi_{2}} s_{\phi_{2}} c_{2}^{2} \left(c_{1}^{2} - s_{1}^{2} \right) \right\} \\ &+ C_{\nu}^{l} y_{\tau}^{2} \left\{ \left(s_{(\rho-\phi_{1})} c_{x} s_{1} s_{2} + s_{(\sigma+\phi_{2})} c_{2} s_{x} \right) \frac{c_{y}^{2} c_{z} s_{z}}{c_{1} c_{2} s_{1}} - s_{(\rho-\sigma+\phi_{1}-\phi_{2})} \frac{c_{x} s_{2} s_{x} \left(s_{x}^{2} - c_{y}^{2} s_{z}^{2} \right)}{c_{2} s_{1}} \\ &+ \left(s_{(\delta-\rho-\phi_{1})} s_{1} s_{2} s_{x} - s_{(\delta-\sigma-\phi_{2})} c_{2} c_{x} \right) \frac{c_{y} c_{z} s_{y}}{c_{1} c_{2} s_{1}} \\ &+ \left(s_{(\delta+\rho-\sigma+\phi_{1}-\phi_{2})} c_{x}^{2} + s_{(\delta-\rho+\sigma-\phi_{1}+\phi_{2})} s_{x}^{2} \right) \frac{c_{y} s_{2} s_{y} s_{z}}{c_{2} s_{1}} \right\} .$$
(C.6)

REFERENCES

- Reviews, see, for example, H. Fritzsch, Z. Z. Xing, Prog. Part. Nucl. Phys. 45 (2000)
 hep-ph/9912358; G. Altarelli, F. Feruglio, Phys. Rept. 320 (1999) 295; and hep-ph/0206077; E. Ma, Mod. Phys. Lett. A 17 (2002) 2361, hep-ph/0211393; S. M. Barr,
 I. Dorsner, Nucl. Phys. B 585 (2000) 79, hep-ph/0003058; R. N. Mohapatra, hep-ph/0211252.
- [2] P. Minkowski, Phys. Lett. B 67 (1977) 421.
- [3] T. Yanagida, in proceedings of Workshop On The Baryon Number Of The Universe And Unified Theories, eds. O. Sawada and A. Sugamoto. KEK, Tsukuba, Japan, 1979. p95;
 M. Gell-Mann, P. Ramond, and R. Slansky, in Supergravity, eds. P. van Nieuwenhuizen and D. Z. Freedman, North Holland, Amsterdam, 1979, p. 315.
- [4] R. N. Mohapatra, G. Senjanovic, Phys. Rev. Lett. 44 (1980) 912.
- [5] C. Wetterich, Nucl. Phys. B 187 (1981) 343; G. M. Asatrian, A. N. Ioannisian, Sov. J. Nucl. Phys. 51 (1990) 115; Yad. Fiz. 51 (1990) 182.
- [6] P. H. Chankowski, Z. Pluciennik, Phys. Lett. B **316** (1993) 312, hep-ph/9306333; K. S. Babu, C. N. Leung, J. Pantaleone, Phys. Lett. B **319** (1993) 191, hep-ph/9309223.
- S. Antusch, M. Drees, J. Kersten, M. Lindner, M. Ratz, Phys. Lett. B 519 (2001) 238, hep-ph/0108225; Phys. Lett. B 525 (2002) 130, hep-ph/0110366; S. Antusch, M. Ratz, JHEP 0207 (2002) 059, hep-ph/0203027.
- [8] M. Tanimoto, Phys. Lett. B **360** (1995) 41, hep-ph/9508247.
- [9] J. R. Ellis, G. K. Leontaris, S. Lola, D. V. Nanopoulos, Eur. Phys. J. C 9 (1999) 389, hep-ph/9808251; J. R. Ellis, S. Lola, Phys. Lett. B 458 (1999) 310, hep-ph/9904279; S. Lola, hep-ph/9903203; Acta Phys. Polon. B 31 (2000) 1253, hep-ph/0005093.
- [10] J. A. Casas, J. R. Espinosa, A. Ibarra, I. Navarro, Nucl. Phys. B 556 (1999) 3, hep-ph/9904395; Nucl. Phys. B 569 (2000) 82, hep-ph/9905381; JHEP 9909 (1999) 015, hep-ph/9906281; E. Ma, J. Phys. G 25 (1999) L97, hep-ph/9907400; A. Ibarra, I. Navarro, JHEP 0002 (2000) 031, hep-ph/9912282; P. H. Chankowski, A. Ioannisian, S. Pokorski, J. W. F Valle, Phys. Rev. Lett. 86 (2001) 3488, hep-ph/0011150; M. K. Parida, C. R. Das, G. Rajasekaran, Pramana 62 (2004) 647, hep-ph/0203097.
- [11] N. Haba, N. Okamura, M. Sugiura, Prog. Theor. Phys. 103 (2000) 367, hep-ph/9810471;
 Y. L. Wu, J. Phys. G: Nucl. Part. Phys. 26 (2000) 1131, hep-ph/9905222; R. Barbieri, G. G. Ross, A. Strumia, JHEP 9910 (1999) 020, hep-ph/9906470; R. G. Felipe, F. R. Joaquim, JHEP 0109 (2001) 015, hep-ph/0106226; T. Miura, E. Takasugi, M. Yoshimura, Prog. Theor. Phys. 104 (2000) 1173, hep-ph/0007066; A. S. Dighe, A. S. Joshipura, hep-ph/0010079; A. S. Joshipura, Phys. Lett. B 543 (2002) 276, hep-ph/0205038; A. S. Joshipura, S. D. Rindani, N. N. Singh, Nucl. Phys. B 660 (2003) 362, hep-ph/0211378; S. Lavignac, I. Masina, C. A. Savoy, Nucl. Phys. B 633 (2002) 139, hep-ph/0202086; T. Fukuyama, N. Okada, JHEP 0211 (2002) 011, hep-ph/0205066; T. Miura, T. Shindou, E. Takasugi, Phys. Rev. D 66 (2002) 093002, hep-ph/0206207; Phys. Rev. D 68 (2003) 093009, hep-ph/0308109; G. Bhattacharyya, A. Raychaudhuri, A. Sil, Phys. Rev. D 67 (2003) 073004, hep-ph/0211074.
- [12] Z. Z. Xing, Phys. Rev. D 63 (2001) 057301, hep-ph/0011217.
- [13] K. R. S. Balaji, A. S. Dighe, R. N. Mohapatra, M. K. Parida, Phys. Rev. Lett. 84 (2000) 5034, hep-ph/0001310; Phys. Lett. B 481 (2000) 33, hep-ph/0002177; M. C. Chen, K.

T. Mahanthappa, Int. J. Mod. Phys. A 16 (2001) 3923, hep-ph/0102215; K. S. Babu,
R. N. Mohapatra, Phys. Lett. B 532 (2002) 77, hep-ph/0201176.

- [14] K. R. S. Balaji, R. N. Mohapatra, M. K. Parida, E. A. Paschos, Phys. Rev. D 63 (2001) 113002, hep-ph/0011263.
- [15] S. F. King, N. N. Singh, Nucl. Phys. B 591 (2000) 3, hep-ph/0006229; Nucl. Phys. B 596 (2001) 81, hep-ph/0007243.
- [16] G. Dutta, hep-ph/0203222.
- [17] P. H. Chankowski, W. Krolikowski, S. Pokorski, Phys. Lett. B 473 (2000) 109, hepph/9910231.
- [18] J. A. Casas, J. R. Espinosa, A. Ibarra, I. Navarro, Nucl. Phys. B 573 (2000) 652, hep-ph/9910420.
- [19] N. Haba, Y. Matsui, N. Okamura, Prog. Theor. Phys. 103 (2000) 807, hep-ph/9911481.
- [20] T. K. Kuo, J. Pantaleone, G. H. Wu, Phys. Lett. B **518** (2001) 101, hep-ph/0104131.
- [21] R. N. Mohapatra, M. K. Parida, G. Rajasekaran, Phys. Rev. D 69 (2004) 053007, hep-ph/0301234.
- [22] A. S. Joshipura, S. Mohanty, Phys. Rev. D 67 (2003) 091302, hep-ph/0302181; S. Antusch, P. Huber, J. Kersten, T.Schwetz, W. Winter, hep-ph/0404268; N. N. Singh, M. K. Das, hep-ph/0407206.
- [23] J. W. Mei, Z. Z. Xing, Phys. Rev. D 69 (2004) 073003, hep-ph/0312167.
- [24] C. Hagedorn, J. Kersten, M. Lindner, Phys. Lett. B 597 (2004) 63, hep-ph/0406103.
- [25] P. H. Chankowski, S. Pokorski, Int. J. Mod. Phys. A **17** (2002) 575, hep-ph/0110249.
- [26] J. A. Casas, J. R. Espinosa, I. Navarro, JHEP **0309** (2003) 048, hep-ph/0306243.
- [27] S. Antusch, J. Kersten, M. Lindner, M. Ratz, Nucl. Phys. B 674 (2003) 401, hepph/0305273.
- [28] S. Antusch, J. Kersten, M. Lindner, M. Ratz, Phys. Lett. B 538 (2002) 87, hepph/0203233.
- [29] S. Antusch, J. Kersten, M. Lindner, M. Ratz, Phys. Lett. B 544 (2002) 1, hep-ph/0206078; S. Antusch, M. Ratz, JHEP 0211 (2002) 010, hep-ph/0208136; S. Antusch, *Hamburg 2002, Supersymmetry and unification of fundamental interactions, vol. 2* 1018-1024, hep-ph/0208179.
- [30] J. W. Mei, Z. Z. Xing, Phys. Rev. D 70 (2004) 053002, hep-ph/0404081.
- [31] Z. Z. Xing, Int. J. Mod. Phys. A **19** (2004) 1, hep-ph/0307359.
- [32] Review of Particle Physics, Phys. Lett. B **592** (2004) 1.
- [33] M. Maltoni, T. Schwetz, M. A. Tortola, J. W. F. Valle, hep-ph/0405172.
- [34] R. N. Mohapatra et al., hep-ph/0412099.
- [35] C. Aalseth et al., hep-ph/0412300.
- [36] S. Antusch, J. Kersten, M. Lindner, M. Ratz, M. A. Schmidt, hep-ph/0501272.

TABLES

TABLE 1. Functions of δ , ρ and σ that can be factorized out together with ζ_{ij}^{-1} (for i < j; i, j = 1, 2, 3) defined in Eq.(12), both in contributions from y_{τ} and y_{ν} in Eqs.(14)-(19). Note that terms led by ζ_{13}^{-1} and ζ_{23}^{-1} in Eq.(17) are complicated, so that the last two rows in the table do not apply to $\dot{\delta}$.

	$\dot{ heta}_x$	$\dot{ heta}_y$	$\dot{ heta}_z$	$\dot{\delta}, \dot{ ho}, \dot{\sigma}$
ζ_{12}^{-1}	$c_{(ho-\sigma)}$	0	0	$s_{(ho-\sigma)}$
ζ_{13}^{-1}	$c_{ ho}$	$c_{(\delta- ho)}$	$c_{ ho}$	$s_{ ho}$
ζ_{23}^{-1}	c_{σ}	$\overline{c}_{(\delta-\sigma)}$	c_{σ}	s_{σ}

TABLE 2. Association of CP-violating phases with enhancing factors ζ_{ij}^{-1} (for i < j; i, j = 1, 2, 3) defined in Eq.(12), in the contribution from y_{ν} in Eqs.(14)-(19). Note that terms led by ζ_{13}^{-1} and ζ_{23}^{-1} in Eq.(17) are complicated, so that the last two rows in the table do not apply to $\dot{\delta}$.

	$\dot{ heta}_x$	$\dot{ heta}_y$	$\dot{ heta}_z$	$\dot{\delta},\dot{ ho},\dot{\sigma}$
ζ_{12}^{-1}	$-c_{(ho-\sigma)}c_{(\phi_1-\phi_2)}$	0	0	$-c_{(\phi_1-\phi_2)}s_{(\rho-\sigma)}$
ζ_{13}^{-1}	$-c_ ho c_{\phi_1}$	$c_{(\delta- ho)}c_{\phi_1}$	$-c_{ ho}c_{\phi_1}$	$c_{\phi_1}s_ ho$
ζ_{23}^{-1}	$c_\sigma c_{\phi_2}$	$-c_{(\delta-\sigma)}c_{\phi_2}$	$-c_{\sigma}c_{\phi_2}$	$c_{\phi_2}s_{\sigma}$

FIGURES



FIG. 1. The RG evolution of a left-handed neutrino mass $(m_1 \text{ of cases (i)} \text{ and (ii)}, \text{ and } m_3 \text{ of case (iii)})$ from M_Z to Λ_{GUT} , both in the SM and in the MSSM. At M_Z , $\theta_x = 33.2^\circ$, $\theta_y = 45^\circ$, $\theta_z = 5^\circ$, $\delta = 45^\circ$ and $\rho = \sigma = 5^\circ$. At M_3 , $y_\nu = 0.8$, $\theta_1 = \theta_2 = 45^\circ$ and $\theta_3 = \delta_\nu = \phi_1 = \phi_2 = 5^\circ$.



FIG. 2. The typical evolution behavior of enhancing factors $\left(-\zeta_{ij}^{-1}\right)$ defined in Eq.(12), both in the SM and in the MSSM. At M_Z , $\theta_x = 33.2^\circ$, $\theta_y = 45^\circ$, $\theta_z = 5^\circ$, $\delta = 45^\circ$ and $\rho = \sigma = 5^\circ$. At M_3 , $y_{\nu} = 0.8$, $\theta_1 = \theta_2 = 45^\circ$ and $\theta_3 = \delta_{\nu} = \phi_1 = \phi_2 = 5^\circ$.



FIG. 3. Typical peaks of enhancing factors $\left(-\zeta_{ij}^{-1}\right)$ defined in Eq.(12) in nearly singular situations. At $M_{\rm Z}$, $\theta_x = 33.2^{\circ}$, $\theta_y = 45^{\circ}$, $\theta_z = 5^{\circ}$, $\delta = 55^{\circ}$ and $\rho = \sigma = 5^{\circ}$. At M_3 , $y_{\nu} = 0.8$, $\theta_1 = \theta_2 = 89^{\circ}$ and $\theta_3 = \delta_{\nu} = \phi_1 = \phi_2 = 5^{\circ}$.



FIG. 4. The typical evolution behavior of the mixing angle θ_z and CP-violating phases (δ, ρ, σ) in the SM, in the normal hierarchy case. At M_Z , $\theta_x = 33.2^\circ$, $\theta_y = 45^\circ$, $\delta = 55^\circ$ and $\rho = \sigma = 45^\circ$. At M_3 , $y_\nu = 0.8$, $\theta_1 = \theta_2 = \phi_1 = \phi_2 = 45^\circ$ and $\theta_3 = \delta_\nu = 5^\circ$.



FIG. 5. The typical evolution behavior of CP-violating phases (δ, ρ, σ) in the MSSM, in the normal hierarchy case. At M_Z , $\theta_x = 33.2^\circ$, $\theta_y = 45^\circ$, $\delta = 55^\circ$ and $\rho = \sigma = 45^\circ$. At M_3 , $y_\nu = 0.8$, $\theta_1 = \theta_2 = \phi_1 = \phi_2 = 45^\circ$ and $\theta_3 = \delta_\nu = 5^\circ$.



FIG. 6. The typical evolution behavior of mixing angles $(\theta_x, \theta_y, \theta_z)$ and CP-violating phases (δ, ρ, σ) in the SM, in the near degeneracy case. At M_Z , $\theta_x = 33.2^\circ$, $\theta_y = 45^\circ$, $\delta = 55^\circ$, $\rho = 80^\circ$ and $\sigma = 5^\circ$. At M_3 , $y_\nu = 0.8$, $\theta_1 = \theta_2 = 45^\circ$ and $\phi_1 = \phi_2 = \theta_3 = \delta_\nu = 5^\circ$.



FIG. 7. The typical evolution behavior of mixing angles $(\theta_x, \theta_y, \theta_z)$ and CP-violating phases (δ, ρ, σ) in the MSSM when $\tan \beta$ is small, in the near degeneracy case. At M_Z , $\theta_x = 33.2^\circ$, $\theta_y = 45^\circ$, $\delta = 55^\circ$ and $\rho = \sigma = 45^\circ$. At M_3 , $y_\nu = 0.8$, $\theta_1 = \theta_2 = 45^\circ$ and $\phi_1 = \phi_2 = \theta_3 = \delta_\nu = 5^\circ$.



FIG. 8. The typical evolution behavior of mixing angles $(\theta_x, \theta_y, \theta_z)$ and CP-violating phases (δ, ρ, σ) in the MSSM when $\tan \beta$ is large, in the near degeneracy case. Figures A, B and C illustrate cases in which the correction to a specific mixing angle is mostly enhanced or damped. Figure A is devoted to θ_x , but we have also included the other two angles in the plot. Similarly, Figure B corresponds to θ_y , and Figure C corresponds to θ_z . Figure D serves to illustrate how θ_z may affect the RG evolution of CP-violating phases. At M_Z , $\theta_x = 33.2^\circ$, $\theta_y = 45^\circ$ and $\theta_z = 0.05^\circ$ (used in Figures A, B and C); at M_3 , $y_{\nu} = 0.8$, $\theta_1 = \theta_2 = 45^\circ$ and $\theta_3 = \delta_{\nu} = 5^\circ$. Initial values of CP-violating phases $(\delta, \rho \text{ and } \sigma \text{ at } M_Z, \text{ and } \phi_1 \text{ and } \phi_2 \text{ at } M_3)$ are marked on the graphs.



FIG. 9. The typical evolution behavior of mixing angles θ_x and θ_z and CP-violating phases (δ, ρ, σ) in the SM, in the inverted hierarchy case. At M_Z , $\theta_x = 33.2^\circ$, $\theta_y = 45^\circ$, $\delta = 55^\circ$, $\rho = 85^\circ$ and $\sigma = 5^\circ$; at M_3 , $y_\nu = 0.8$, $\theta_1 = \theta_2 = 45^\circ$ and $\phi_1 = \phi_2 = \theta_3 = \delta_\nu = 5^\circ$.



FIG. 10. The typical evolution behavior of mixing angles θ_x and θ_z and CP-violating phases (δ, ρ, σ) in the MSSM, in the inverted hierarchy case. At M_Z , $\theta_x = 33.2^\circ$, $\theta_y = 45^\circ$ and $\delta = 55^\circ$. Also at M_Z , $\rho = \sigma = 5^\circ$ in the plot of θ_x , but $\rho = 85^\circ$ and $\sigma = 5^\circ$ in the plot of θ_z, δ, ρ and σ . At $M_3, y_\nu = 0.8, \theta_1 = \theta_2 = 45^\circ$ and $\phi_1 = \phi_2 = \theta_3 = \delta_\nu = 5^\circ$.