

Charge Symmetry Breaking in Pion-Nucleon Coupling Constants Induced by Axial Anomaly

N.I.Kochelev

*School of Physics and Center for Theoretical Physics, Seoul National University,
Seoul 151-747, Korea*

*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research
Dubna 141980, Russia*

e-mail: kochelev@theor.jinr.ru

The contribution of axial anomaly to charge symmetry breaking in pion-nucleon coupling constants is calculated within instanton model for QCD vacuum. It has been demonstrated that the contribution is large and allows to explain Nolen-Schiffer anomaly.

The possibility of large violation of the charge symmetry breaking (CSB) in the strong NN interaction is widely discussed at last decade (see^{1,2} and for general review³). The so-called Nolen-Schiffer anomaly⁴ related to the mass difference between the mirror nuclei provides a well known example of CSB manifestation. In spite of the fact that large CSB effects are observed in various experiments, the full understanding of fundamental QCD mechanism behind them is absent so far.

One of the possible sources of CSB can be originated from the difference between the charged pion coupling constant to nucleon and the coupling constant of neutral pion. The evident mechanism for such difference is $\pi^0 - \eta - \eta'$ mixing. However, the estimation shows that this contribution to CSB is small due to the large differences between meson masses.

In this Letter we suggest a new mechanism for CSB based on the existence of the axial anomaly in QCD. We demonstrate that the large value of the topological charge matrix element between the vacuum and the neutral pion state coming from large difference of current masses of u- and d- quarks, $(m_d - m_u)/(m_d + m_u) \approx 0.4$,⁵

$$\langle 0 | \frac{\alpha_s}{4\pi} G_{\mu\nu}^a \tilde{G}_{\mu\nu}^a | \pi^0 \rangle = \frac{m_d - m_u}{m_d + m_u} f_\pi m_\pi^2, \quad (1)$$

where $f_\pi = 93$ MeV, produces significant CSB in pion-nucleon coupling constants.

The instanton liquid model for QCD vacuum^{6,7} is a very powerful tool to calculate non-perturbative effects in strong interactions. We will use this model to estimate the axial anomaly effect on CSB. Our starting point is the effective quark-gluon interaction induced by instantons⁸

$$\begin{aligned} \mathcal{L}_{eff} = & \int dU d\rho n(\rho) \prod_q \left[-\frac{2\pi^2 \rho^2}{m_q^*} \bar{q}_R (1 + \frac{i}{4} U_{ab} \tau^a \bar{\eta}_{b\mu\nu} \sigma_{\mu\nu}) q_L \right. \\ & \times \left. e^{-\frac{2\pi^2}{g} \rho^2 U_{cd} \bar{\eta}_{d\alpha\beta} G_{\alpha\beta}^c} + (R \leftrightarrow L) \right], \quad (2) \end{aligned}$$

where m_q^* is the effective quark mass in the instanton vacuum, $n(\rho)$ is the instanton density, and U is the orientation matrix of the instanton in $SU(3)_c$ color space. From




Figure 1: The π^0 meson coupling to nucleon, (a), through gluons and, (b), due to quark-quark interaction. The symbol I denotes the instanton.

the Lagrangian one can obtain the following quark-gluon and quark-quark interactions related to our consideration of CSB in pion-nucleon constants

$$\mathcal{L}_{ggq} = -i \frac{n_{eff} \pi^3 \rho_c^4}{4 \langle 0 | \bar{q}q | 0 \rangle} \alpha_s G_{\mu\nu}^a \tilde{G}_{\mu\nu}^a \{ \bar{u} \gamma_5 u + \bar{d} \gamma_5 d \}, \quad (3)$$

$$\mathcal{L}_{qq} = \frac{15 n_{eff}}{16 \langle 0 | \bar{q}q | 0 \rangle} \bar{q} \gamma_5 i \tau_3 q \bar{q} \gamma_5 i \tau_3 q, \quad (4)$$

and we have elaborated the version of Shuryak's instanton liquid model⁹:

$$n(\rho) = n_{eff} \delta(\rho - \rho_c), \quad m_q^* = -\frac{2}{3} \pi^2 \rho_c^2 \langle 0 | \bar{q}q | 0 \rangle, \quad (5)$$

where ρ_c is the average instanton size in QCD vacuum. To estimate the contribution of the effective interactions (3) and (4) to π^0 -nucleon coupling let us consider the diagrams presented in Fig.1. The straightforward calculation of the matrix elements $\langle N | \mathcal{L}_{ggq} | N \pi^0 \rangle$ and $\langle N | \mathcal{L}_{qq} | N \pi^0 \rangle$ using the vacuum dominance assumption, axial anomaly, (1), and PCAC relations

$$\langle 0 | \bar{q} \gamma_5 \tau_3 q | \pi^0 \rangle = \frac{2i f_\pi m_\pi^2}{m_u + m_d}, \quad (6)$$

gives the following result for the effective couplings of π^0 meson with proton and neutron

$$g_{\pi^0 pp, \pi^0 nn}^2 = g_{\pi^0 NN}^2 \left\{ 1 \pm \frac{4g_A^8}{15g_A^3} \frac{\pi^4 \rho_c^4}{\alpha_s^2} (m_d - m_u) \langle 0 | \bar{q}q | 0 \rangle \right\}, \quad (7)$$

where the valence quark approximation for the nucleon wave function has been used, $g_A^{3,8}$ are the axial couplings of nucleon, and we have absorbed all common factors, i.e n_c , $\langle 0 | \bar{q}q | 0 \rangle$ etc, into the effective pion-nucleon coupling $g_{\pi^0 NN}$ along the lines of the chiral quark model approach based on instantons^{7, 10}. For the value of CSB

$$\alpha = \frac{g_{\pi^0 nn}^2 - g_{\pi^0 pp}^2}{g_{\pi^0 NN}^2} \quad (8)$$

we derive the result

$$\alpha = -\frac{g_8}{g_3} \frac{8\pi^4 \rho_c^4}{15\alpha_s^2} (m_d - m_u) \langle 0|\bar{q}q|0 \rangle . \quad (9)$$

Using the value for the difference of current masses $m_d - m_u = 4.0$ MeV and value of the quark condensate $\langle 0|\bar{q}q|0 \rangle = -(260\text{MeV})^3$ from recent paper¹¹, and the values $g_A^3 = 1.267$ and $g_A^8 = 0.585$, and putting parameters of the instanton model from⁷ $\rho_c^{-1} = 600$ MeV, $2\pi/\alpha_s \approx 12$, we obtain finally

$$\alpha \approx 4\% . \quad (10)$$

So that, the parameter α of CSB is large and has positive sign. The sign is related to the sign of the quark condensate. It is well known that just the negative sign of the quark condensate leads to the sign flip the instanton induced interaction with odd or even number of incoming to instanton quark legs (see for example¹²). In our case this corresponds to the opposite signs of the contributions to $g_{\pi^0 pp}$ coupling constant produced from two diagrams presented in Fig.1. To explain the Nolen-Schiffer anomaly, for example for nuclei with $A = 41$, the parameter α of CSB should be larger than 2%¹³. Therefore, the axial anomaly contribution to CSB estimated above allows to resolve this longstanding problem in nuclear physics.

In summary, we have demonstrated that the charge symmetry breaking effect in pion-nucleon constants is significant due to the axial anomaly and large difference between current masses of d- and u- quarks,

Acknowledgments

The author thanks to A. Di Giacomo for numerous discussions on instanton effects in hadron physics and for his warm hospitality at beautiful Pisa. This work was supported by Brain Pool program of Korea Research Foundation through KOFST, grant 042T-1-1, and in part by grants of Russian Foundation for Basic Research, RFBR-03-02-17291 and RFBR-04-02-16445. The author also very grateful to Prof. Dong-Pil Min for warm hospitality at Seoul National University at the final stage of this work.

References

1. T. Nasu, M. Oka and S. Takeuchi, Phys. Rev. C **68**, 024006 (2003) [Erratum-ibid. C **69**, 029903 (2004)] [arXiv:nucl-th/0307002].
2. H. Machner and J. Niskanen, nucl-ex/0511027.
3. G. A. Miller, B. M. K. Nefkens and I. Slaus, Phys. Rept. **194** (1990) 1.
4. J. A. . Nolen and J. P. Schiffer, Ann. Rev. Nucl. Part. Sci. **19**(1969) 471.
5. D. J. Gross, S. B. Treiman and F. Wilczek, Phys. Rev. D **19** (1979) 2188.
6. T. Schäfer and E.V. Shuryak, Rev. Mod. Phys. **70** (1998) 1323.
7. D. Diakonov, Prog. Par. Nucl. Phys. **51** (2003) 173.
8. A.I. Vainshtein, V.I. Zakharov, V.A. Novikov, M.A. Shifman, Sov. Phys. Usp. **24** (1982) 195 [Usp. Fiz. Nauk **136** (1982) 553].
9. E.V. Shuryak, Nucl. Phys. **B203** (1982) 93; 116; 140.

10. A. E. Dorokhov, hep-ph/0212252.
11. B. L. Ioffe, Prog. Part. Nucl. Phys. **56**, 232 (2006) 232, hep-ph/0502148.
12. A. E. Dorokhov and N. I. Kochelev, hep-ph/0411362.
13. N. Auerbach, Phys. Rep. **98** (1983) 273.