On Goldstone bosons decoupling from high–lying hadrons

L. Ya. Glozman

Institute for Physics, University of Graz, Universitätsplatz 5, A-8010 Graz, Austria

A. V. Nefediev

Institute of Theoretical and Experimental Physics, 117218, B. Cheremushkinskaya 25, Moscow, Russia

In this paper, we discuss a decoupling of the Goldstone bosons from highly excited hadrons in relation to the restoration of chiral symmetry in such hadrons. We use a generalised Nambu–Jona-Lasinio model with the interaction between quarks in the form of an instantaneous Lorentz–vector confining potential. This model is known to provide spontaneous breaking of chiral symmetry in the vacuum via the standard selfenergy loops for valence quarks. For highly excited hadrons, where the typical momentum of valence quarks is large, the loop contributions represent only a small correction to the chiral–invariant classical contributions and asymptotically vanish. Consequently the chiral symmetry violating Lorentz–scalar dynamical mass of quarks vanishes. Then the conservation of the axial vector current in the chiral limit requires, via the Goldberger–Treiman relation, that the valence quarks decouple from the Goldstone boson. As a consequence, the whole hadron decouples from the Goldstone boson as well, which implies that its axial constant also vanishes.

PACS numbers: 12.38.Aw, 12.39.Ki, 12.39.Pn

I. INTRODUCTION

An approximate restoration of $SU(2)_L \times SU(2)_R$ and $U(1)_A$ symmetries in excited hadrons has recently become a subject of a significant theoretical effort [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14]. This *effective* restoration of chiral symmetry requires in particular that highly excited hadrons should gradually decouple from the Goldstone bosons [4, 7, 14]. There is an indirect phenomenological hint for such a decoupling. Indeed, the coupling constant for the process $h^* \to h + \pi$ decreases for high lying resonances, because the phase–space factor for such a decay increases with the mass of a resonance much faster than the decay width.

A coupling of the Goldstone bosons to the valence quarks is regulated by the conservation of the axial current (we consider for simplicity the chiral limit). This conservation results in a Goldberger–Treiman relation [15], taken at the "constituent quark" level, giving

$$g_{\pi} \propto m_q^{\text{eff}},$$
 (1)

where m_a^{eff} is the quark Lorentz–scalar dynamical mass which appears selfconsistently due to spontaneous breaking of chiral symmetry (SBCS) in the vacuum. Appearance of such a dynamical mass is a general feature of chiral symmetry breaking and has been studied in great detail in the context of different models like the Nambu and Jona-Lasinio model 16, 17, instanton liquid model 18, within the Schwinger–Dyson formalism with the quark kernel formed by the QCD string |19|, by the perturbative gluon exchange |20| or by the instantaneous Lorentz-vector confinement [21, 22, 23]. A general feature of this dynamical mass is that it results from quantum fluctuations of the quark field and vanishes at large momenta where the classical contributions dominate [5, 13]. Then, since the average momentum of valence quarks higher in the spectrum increases, the valence quarks decouple from the quark condensate and their dynamical Lorentz-scalar mass decreases (and asymptotically vanishes), so that chiral symmetry is *approximately* restored in the highly-excited hadrons [1, 12, 13]. This implies, via the Goldberger–Treiman relation (1), that valence quarks, as well as the whole hadron, decouple from the Goldstone bosons [4]. This, in turn, requires the axial coupling constant of the highly excited hadrons to decrease and to vanish asymptotically.

While this perspective was shortly outlined in the past, this has never been considered in detail microscopically. However, it is important to clarify this physics, especially because the origins of this phenomenon cannot be seen at the level of the effective Lagrangian approach, where the coupling constant of the Goldstone boson to the excited hadron is an input parameter and the decoupling looks unintuitive [14].

Even though the role of different gluonic interactions in QCD, which could be responsible for chiral and $U(1)_A$ symmetries breaking, is not yet clear, the most fundamental reason for the restoration of these symmetries in excited hadrons is universal [5]. Namely, both $SU(2)_L \times SU(2)_R$ and $U(1)_A$ symmetries breaking result from quantum fluctuations of the quark fields (that is, loops). However, for highly-excited hadrons, where the action of the intrinsic motion is large, a semiclassical regime necessarily takes place. Semiclassically, the contribution of quantum fluctuations is suppressed, relative to the classical contributions, by a factor \hbar/S , where S is the classical action of the intrinsic motion in the hadron in terms of the quark and gluon degrees of freedom. Since, for highly excited hadrons, $S \gg \hbar$, contributions of the quantum fluctuations of the quark fields are suppressed relative to the classical contributions. Consequently both chiral and $U(1)_A$ symmetries are approximately restored in this part of the spectrum.

Although this argument is quite general and solid, it does not provide one with any detailed microscopic picture of the symmetry restoration. Then in the absence of controllable analytic solutions of QCD such an insight can be obtained only through models.

It is instructive to outline the minimal set of requirements for such a model. It must be (i) relativistic, (ii) chirally symmetric, (iii) able to provide spontaneous breaking of chiral symmetry, (iv) it should contain confinement, (v) it must explain the restoration of chiral symmetry in excited states. There is a model which does incorporate all required elements. This is the generalised Nambu and Jona-Lasinio (GNJL) model with the instantaneous Lorentz-vector confining kernel [21, 22, 23]. In this model, confinement of quarks is guaranteed due to instantaneous infinitely rising (for example, linear) potential. Then chiral symmetry breaking can be described by the standard summation of the valence-quark selfinteraction loops giving rise to the Schwinger–Dyson equation for the quark selfenergy [21, 22]. Alternatively the model can be considered in the Hamiltonian approach using the BCS formalism [24]. In this case, chiral symmetry breaking in the vacuum happens via condensation of the ${}^{3}P_{0}$ quark-antiquark pairs and dressed quarks appear from the Bogoliubov–Valatin transformation applied to bare quarks. The mass–gap equation ensures absence of anomalous Bogoliubov terms in the Hamiltonian [23]. Finally, mesons are built using the Bethe–Salpeter equation for the quark–antiquark bound states [22, 23] or by a generalised bosonic Bogoliubov-like transformation applied to the operators creating/annihilating quark-antiquark pairs [25].

It was demonstrated in Ref. [12] that, for the low-lying states, where the typical momentum of valence quarks is not large and chiral symmetry breaking is important, this model leads to an *effective* Lorentz-scalar binding potential, while for high-lying states, such an effective potential becomes a pure Lorentz spatial vector. Then the discussed above quantum nature of chiral symmetry breaking in QCD as well as the transition to the semiclassical regime for excited states, with loop effects being suppressed relative to the classical contributions, has been illustrated within the same model in Ref. [13]. As a result, the model does provide chiral symmetry restoration for excited hadrons.

The purpose of this paper to give an insight into physics of decoupling of the Goldstone bosons from the excited hadrons which happens in line, and due to the same reason, with the approximate restoration of chiral symmetry for these hadrons. We resort to the GNJL model in view of its obvious advantage as a tractable model for QCD which can be used as a laboratory to get a microscopical insight into the restoration of chiral symmetry for excited hadrons.

II. GENERALISED NAMBU–JONA-LASINIO MODEL

A. Some generalities

In this chapter, we overview the GNJL chiral quark model [21, 22, 23] which is described by the Hamiltonian

$$\hat{H} = \int d^3x \bar{\psi}(\vec{x},t) \left(-i\vec{\gamma} \cdot \vec{\nabla} + m \right) \psi(\vec{x},t) + \frac{1}{2} \int d^3x d^3y \ J^a_\mu(\vec{x},t) K^{ab}_{\mu\nu}(\vec{x}-\vec{y}) J^b_\nu(\vec{y},t), \tag{2}$$

with the quark current–current $(J^a_{\mu}(\vec{x},t) = \bar{\psi}(\vec{x},t)\gamma_{\mu}\frac{\lambda^a}{2}\psi(\vec{x},t))$ interaction parametrised by an instantaneous confining kernel $K^{ab}_{\mu\nu}(\vec{x}-\vec{y})$ of a generic form. In this paper, we use the simplest form of the kernel compatible with the requirement of confinement,

$$K^{ab}_{\mu\nu}(\vec{x} - \vec{y}) = g_{\mu0}g_{\nu0}\delta^{ab}V_0(|\vec{x} - \vec{y}|).$$
(3)

We do not dwell at any particular form of the confining potential $V_0(|\vec{x} - \vec{y}|)$, though, if needed for an illustration purpose, we employ a power-like confining potential [21, 27],

$$V_0(|\vec{x}|) = K_0^{\alpha+1} |\vec{x}|^{\alpha}, \quad 0 \leqslant \alpha \leqslant 2, \tag{4}$$

for qualitative analysis, concentrating mostly at the case of the linear confinement ($\alpha = 1$) or, for numerical studies, resorting to the harmonic oscillator potential ($\alpha = 2$) [21, 22, 23].

$$\frac{1}{S} = \frac{1}{S_0} + \frac{1}{S_0} \underbrace{\Sigma}_{S_0} + \frac{1}{S_0} \underbrace{\Sigma}_{S_0} \underbrace{\Sigma}_{S_0} + \dots = \frac{1}{S_0} + \frac{1}{S_0} \underbrace{\Sigma}_{S_0} \underbrace{\Sigma}_{S_0} + \dots = \frac{1}{S_0} \underbrace{\Sigma}_{S_0} \underbrace{\Sigma}_{S_0} \underbrace{\Sigma}_{S_0} + \dots = \underbrace{\Sigma}_{S_0} \underbrace{\Xi}_{S_0} \underbrace{\Sigma}_{S_0} \underbrace{\Sigma}_{S_0} \underbrace{\Xi}_{S_0} \underbrace{\Xi$$

FIG. 1: Graphical representation of the equations for the dressed quark propagator, Eq. (5), and for the quark mass operator, Eq. (7).

B. Chiral symmetry breaking

Spontaneous breaking of chiral symmetry in the class of Hamiltonians (2) is described via the standard Dyson series for the quark propagator which takes the form, schematically (see Fig. 1):

$$S = S_0 + S_0 \Sigma S_0 + S_0 \Sigma S_0 \Sigma S_0 + \ldots = S = S_0 + S_0 \Sigma S,$$
 (5)

with S_0 and S being the bare– and the dressed–quark propagators, respectively, Σ is the quark mass operator. The Dyson–Schwinger Eq. (5) has the solution

$$S^{-1}(p_0, \vec{p}) = S_0^{-1}(p_0, \vec{p}) - \Sigma(\vec{p}), \tag{6}$$

where the mass operator independence of the energy p_0 follows from the instantaneous nature of the interaction. The expression for the mass operator through the dressed-quark propagator (see Fig. 1) reads:

$$i\Sigma(\vec{p}) = C_F \int \frac{d^4k}{(2\pi)^4} V_0(\vec{p} - \vec{k})\gamma_0 S(k_0, \vec{k})\gamma_0, \quad C_F = \frac{N_C^2 - 1}{2N_C},$$
(7)

with both quark-quark-potential vertices being bare momentum-independent vertices γ_0 . This corresponds to the so-called rainbow approximation which is well justified in the limit of the large number of colours N_C . We assume this limit in what follows. Then all nonplanar diagrams appear suppressed by N_C and can be consecutively removed from the theory. Eqs. (6) and (7) together produce a closed set of equations, equivalent to a single nonlinear equation for the mass operator,

$$i\Sigma(\vec{p}) = \int \frac{d^4k}{(2\pi)^4} V(\vec{p} - \vec{k})\gamma_0 \frac{1}{S_0^{-1}(k_0, \vec{k}) - \Sigma(\vec{k})}\gamma_0,$$
(8)

where the fundamental Casimir operator C_F is absorbed by the potential, $V(\vec{p}) = C_F V_0(\vec{p})$.

To proceed, we use the standard parametrisation of the mass operator $\Sigma(\vec{p})$ in the form:

$$\Sigma(\vec{p}) = [A_p - m] + (\vec{\gamma}\hat{\vec{p}})[B_p - p], \qquad (9)$$

so that the dressed–quark Green's function (6) becomes

$$S^{-1}(p_0, \vec{p}) = \gamma_0 p_0 - (\vec{\gamma} \vec{p}) B_p - A_p,$$
(10)

where, due to the instantaneous nature of the interquark interaction, the time component of the four-vector p_{μ} is not dressed.

It is easily seen from Eq. (10) that the functions A_p and B_p represent the scalar and the space-vectorial part of the effective Dirac operator, respectively. Notice, that it is the scalar part A_p that breaks chiral symmetry and hence it can be identified with the dynamical mass m_q^{eff} of the valence quark appearing in the Goldberger-Treiman relation (1). In the chiral limit, A_p vanishes, unless chiral symmetry is broken spontaneously. It is convenient, therefore, to introduce an angle, known as the chiral angle φ_p , according to the definition:

$$\tan\varphi_p = \frac{A_p}{B_p},\tag{11}$$

and varying in the range $-\frac{\pi}{2} < \varphi_p \leq \frac{\pi}{2}$, with the boundary conditions $\varphi(0) = \frac{\pi}{2}$, $\varphi(p \rightarrow \infty) \rightarrow 0$.

The selfconsistency condition for the parametrisation (9) of the nonlinear Eq. (8) requires that the chiral angle is subject to a nonlinear equation — the mass–gap equation,

$$A_p \cos \varphi_p - B_p \sin \varphi_p = 0, \tag{12}$$

with

$$A_p = m + \frac{1}{2} \int \frac{d^3k}{(2\pi)^3} V(\vec{p} - \vec{k}) \sin \varphi_k, \quad B_p = p + \frac{1}{2} \int \frac{d^3k}{(2\pi)^3} \left(\hat{\vec{p}k}\right) V(\vec{p} - \vec{k}) \cos \varphi_k.$$
(13)

For the given chiral angle φ_p the dispersive law of the dressed quark can be built as

$$E_p = A_p \sin \varphi_p + B_p \cos \varphi_p, \tag{14}$$

and it differs drastically from the free–quark energy $E_p^{(0)} = \sqrt{p^2 + m^2}$ in the low–momentum domain; E_p approaches this free–particle limit as $p \to \infty$.

It was demonstrated in the pioneering papers on the model (2) [21] that, for confining potentials, the mass–gap equation (12) always possesses nontrivial solutions which break



FIG. 2: Nontrivial solution to the mass-gap Eq. (12) with m = 0 and for the linear confinement (see, for example, Refs. [26, 27] for the details). The momentum p is given in the units of $\sqrt{\sigma}$; σ is the fundamental string tension.

chiral symmetry, by generating a nontrivial mass-like function A_p , even for a vanishing quark current mass. At Fig. 2, as an illustration, we show the numerical solution to the mass-gap equation (12) for the linearly rising potential $V(r) = \sigma r$ in the chiral limit. The chiral angle depicted at Fig. 2 shares all features of chiral symmetry breaking solutions to the mass–gap equation (12) for various confining quark kernels (for a comprehensive analysis of power-like potentials see Refs. [21, 27]), namely, it is given by a smooth function which starts at $\frac{\pi}{2}$ at the origin, with the slope inversely proportional to the scale of chiral symmetry breaking, generated by this solution. At large momenta it approaches zero fast. The latter property allows the reader to anticipate the principal conclusion of this paper. Indeed, since the Fourier transform of the potential is peaked at $\vec{p} \simeq \vec{k}$, whereas $\sin \varphi_k$ decreases with the increase of k, then, as $p \to \infty$, A_p is a decreasing function of the momentum p. In the chiral limit it vanishes asymptotically. Therefore, for highly excited hadrons, with the typical momentum of valence quarks being large, the dynamical Lorentz-scalar mass of such valence quarks decreases, and so does the coupling constant g_{π} , due to the Goldberger– Treiman relation (1). Hence highly excited hadrons decouple from the Goldstone bosons. Below we prove this general conclusion by a detailed analysis of the amplitude of the pion emission process $h \to h' + \pi$ with h (and perhaps h') being highly excited hadrons.



FIG. 3: Graphical representation of the Bethe–Salpeter Eq. (15) for the quark–antiquark bound state, in the ladder approximation.

III. PROPERTIES OF THE GOLDSTONE MODE

A. Mesonic Salpeter vertex; Bethe–Salpeter equation for the chiral pion

In this chapter we remind the reader the main steps to take in order to derive the Bethe– Salpeter equation for the generic quark–antiquark bound state, paying special attention to the case of the chiral pion. We follow the lines of Refs. [21, 23, 25, 28, 29].

We start from the homogeneous Bethe–Salpeter equation,

$$\chi(\vec{p},\vec{P}) = -i \int \frac{d^4k}{(2\pi)^4} V(\vec{p}-\vec{k}) \,\gamma_0 S(k_0,\vec{k}) \chi(\vec{k},\vec{P}) S(k_0-M,\vec{k}-\vec{P}) \gamma_0, \tag{15}$$

for the mesonic Salpeter amplitude $\chi(\vec{p}, \vec{P})$; M is the mass of the bound state and \vec{P} is its total momentum. Eq. (15) is written in the ladder approximation for the vertex which is consistent with the rainbow approximation for the quark mass operator and which is well justified in the large- N_C limit. For future convenience we introduce the matrix wave function of the meson as

$$\Psi(\vec{p}, \vec{P}) = \int \frac{dp_0}{2\pi} S(p_0, \vec{p}) \chi(\vec{p}, \vec{P}) S(p_0 - M, \vec{p} - \vec{P}),$$
(16)

and use the standard representation for the dressed quark propagator via Dirac projectors,

$$S(p_0, \vec{p}) = \frac{\Lambda^+(\vec{p})\gamma_0}{p_0 - E_p + i\epsilon} + \frac{\Lambda^-(\vec{p})\gamma_0}{p_0 + E_p - i\epsilon},$$
(17)

$$\Lambda^{\pm}(\vec{p}) = T_p P_{\pm} T_p^{\dagger}, \quad P_{\pm} = \frac{1 \pm \gamma_0}{2}, \quad T_p = \exp\left[-\frac{1}{2}(\vec{\gamma}\vec{p})\left(\frac{\pi}{2} - \varphi_p\right)\right]. \tag{18}$$

We also consider the bound-state in its rest frame setting $\vec{P} = 0$ and skipping this argument of the bound-state wave function for simplicity. It is easy to perform the integration in energy in Eq. (15) explicitly. Then, for the Foldy–rotated wave function $\tilde{\Psi}(\vec{p}) = T_p^{\dagger} \Psi(\vec{p}) T_p^{\dagger}$, the Bethe–Salpeter Eq. (15) reads:

$$\tilde{\Psi}(\vec{p}) = -\int \frac{d^3k}{(2\pi)^3} V(\vec{p} - \vec{k}) \left[P_+ \frac{T_p^{\dagger} T_k \tilde{\Psi}(\vec{k}) T_k T_p^{\dagger}}{2E_p - M} P_- + P_- \frac{T_p^{\dagger} T_k \tilde{\Psi}(\vec{k}) T_k T_p^{\dagger}}{2E_p + M} P_+ \right].$$
(19)

It is clear now that a solution to Eq. (19) is to have the form,

$$\tilde{\Psi}(\vec{p}) = P_+ \mathcal{A} P_- + P_- \mathcal{B} P_+, \qquad (20)$$

and, due to the obvious orthogonality property of the projectors P_{\pm} , $P_{+}P_{-} = P_{-}P_{+} = 0$, only matrices anticommuting with the matrix γ_{0} contribute to \mathcal{A} and \mathcal{B} . The set of such matrices is $\{\gamma_{5}, \gamma_{0}\gamma_{5}, \vec{\gamma}, \gamma_{0}\vec{\gamma}\}$ which can be reduced even more, down to $\{\gamma_{5}, \vec{\gamma}\}$, since the matrix γ_{0} can be always absorbed by the projectors P_{\pm} . The matrix wave function $\Psi_{n}(\vec{p})$ for the *n*-th generic mesonic state can be parametrised through the positive– and negative– energy components of the mesonic wave function $\varphi_{n}^{\pm}(p)$, and a bound–state equation in the form

$$\begin{cases} [2E_p - M_n]\varphi_n^+(p) = \int \frac{k^2 dk}{(2\pi)^3} [T_n^{++}(p,k)\varphi_n^+(k) + T_n^{+-}(p,k)\varphi_n^-(k)] \\ [2E_p + M_n]\varphi_n^-(p) = \int \frac{k^2 dk}{(2\pi)^3} [T_n^{-+}(p,k)\varphi_n^+(k) + T_n^{--}(p,k)\varphi_n^-(k)] \end{cases}$$
(21)

can be derived for such wave functions. The interested reader can find the details in Refs. [21, 23] or in Ref. [25], where also a Hamiltonian approach to the quark-antiquark bound-state problem is developed (as a generalisation of the method applied to the 't Hooft model for QCD in two dimensions in Ref. [29]) which, after a generalised Bogoliubov-like transformation leads to the same Eq. (21); $\varphi_n^{\pm}(p)$ play the role of the bosonic Bogoliubov amplitudes, so that their normalisation condition,

$$\int \frac{p^2 dp}{(2\pi)^3} \left[\varphi_n^+(p) \varphi_m^+(p) - \varphi_n^-(p) \varphi_m^-(p) \right] = \delta_{nm},$$

$$\int \frac{p^2 dp}{(2\pi)^3} \left[\varphi_n^+(p) \varphi_m^-(p) - \varphi_n^-(p) \varphi_m^+(p) \right] = 0,$$
(22)

should not come as a surprise.

Let us consider the case of the chiral pion in more detail. In this case only γ_5 contributes to \mathcal{A} and \mathcal{B} in Eq. (20) and one has:

$$\mathcal{A}_{\pi} = \gamma_5 \varphi_{\pi}^+(p), \quad \mathcal{B}_{\pi} = -\gamma_5 \varphi_{\pi}^-(p).$$
(23)

It is an easy task now to extract the amplitudes $T_{\pi}^{\pm\pm}$ (see Eq. (21)) from Eq. (19) using the explicit form of the matrix wave function $\tilde{\Psi}_{\pi}(\vec{p})$ and of the operator T_p . The result reads:

$$T_{\pi}^{++}(p,k) = T_{\pi}^{-+}(p,k) = -\int d\Omega_k V(\vec{p} - \vec{k}) \left[\cos^2 \frac{\varphi_p - \varphi_k}{2} - \frac{1 - (\hat{\vec{p}k})}{2} \cos \varphi_p \cos \varphi_k \right],$$

$$T_{\pi}^{+-}(p,k) = T_{\pi}^{--}(p,k) = -\int d\Omega_k V(\vec{p} - \vec{k}) \left[\sin^2 \frac{\varphi_p - \varphi_k}{2} + \frac{1 - (\hat{\vec{p}k})}{2} \cos \varphi_p \cos \varphi_k \right].$$
(24)

In the chiral limit, $\varphi_{\pi}^{+}(p) = -\varphi_{\pi}^{-}(p) \equiv \varphi_{\pi}(p)$, so that the bound-state Eq. (21) for the pion reduces to a single equation,

$$2E_p\varphi_{\pi}(p) = \int \frac{k^2 dk}{(2\pi)^3} [T_{\pi}^{++}(p,k) - T_{\pi}^{+-}(p,k)]\varphi_{\pi}(k) = -\int \frac{d^3k}{(2\pi)^3} V(\vec{p} - \vec{k})\varphi_{\pi}(k), \quad (25)$$

or, in the coordinate space, one arrives at the Schrödinger-like equation,

$$[2E_p + V(r)]\varphi_{\pi} = 0.$$
(26)

It is instructive to notice that Eq. (26) reproduces the mass-gap Eq. (12) for $\varphi_{\pi}(p) = \sin \varphi_{p}$. This is the wave function of the chiral pion whose dual nature is clearly seen from this consideration. Indeed, as a Goldstone boson, the pion appears, through the mass-gap equation, already at the level of the quark dressing, whereas the same entity reappears as the lowest pseudoscalar solution to the quark-antiquark bound-state Eq. (21). Below we study the properties of the pionic matrix wave function (16).

B. Bound-state equation in the matrix form: Gell-Mann-Oakes-Renner relation

Mesonic bound-state Eq. (21) admits a matrix form for the wave function $\Psi(\vec{p})$ and it can be derived directly from Eq. (19). Skipping the details of this derivation (the interested reader can find them in Ref. [29], for the two-dimensional 't Hooft model, whereas the generalisation to the model (2) is trivial), we give it here in the final form [25]:

$$M\Psi(\vec{p}) = [(\vec{\alpha}\vec{p}) + \beta m]\Psi(\vec{p}) + \Psi(\vec{p})[(\vec{\alpha}\vec{p}) - \beta m] + \int \frac{d^3q}{(2\pi)^3} V(\vec{p} - \vec{k}) \left\{ \Lambda^+(\vec{k})\Psi(\vec{p})\Lambda^-(-\vec{k}) - \Lambda^+(\vec{p})\Psi(\vec{k})\Lambda^-(-\vec{p}) - \Lambda^-(\vec{k})\Psi(\vec{p})\Lambda^+(-\vec{k}) + \Lambda^-(\vec{p})\Psi(\vec{k})\Lambda^+(-\vec{p}) \right\}.$$
(27)

$$\Psi_{\pi}(\vec{p}) = T_p \left[P_+ \gamma_5 \varphi_{\pi}^+ - P_- \gamma_5 \varphi_{\pi}^- \right] T_p = \gamma_5 G_{\pi} + \gamma_0 \gamma_5 T_p^2 F_{\pi}, \tag{28}$$

where $G_{\pi} = \frac{1}{2}(\varphi_{\pi}^{+} - \varphi_{\pi}^{-}), F_{\pi} = \frac{1}{2}(\varphi_{\pi}^{+} + \varphi_{\pi}^{-}).$

In order to normalise the pion wave function in its rest frame, one is to go slightly beyond the chiral limit and to consider pionic solutions to the bound-state Eq. (21) in the form [23, 25, 29]:

$$\varphi_{\pi}^{\pm}(p) = \mathcal{N}_{\pi} \left[\pm \frac{1}{\sqrt{m_{\pi}}} \sin \varphi_p + \sqrt{m_{\pi}} \Delta_p \right], \quad \mathcal{N}_{\pi}^{-2} = 4 \int_0^\infty \frac{p^2 dp}{(2\pi)^3} \Delta_p \sin \varphi_p, \tag{29}$$

where all corrections of higher order in the pion mass are neglected and the function Δ_p obeys a reduced m_{π} -independent equation (see, for example, Ref. [23] or the papers [29] where such an equation for Δ_p is discussed in two-dimensional QCD).

Furthermore, the pion norm \mathcal{N}_{π} can be easily related to the pion decay constant f_{π} . To this end we multiply the matrix bound-state equation (27) by $\gamma_0\gamma_5$, integrate its both parts over $\frac{d^3p}{(2\pi)^3}$, and, finally, take the trace. The resulting equation reads:

$$m_{\pi} \int \frac{d^3 p}{(2\pi)^3} F_{\pi} \sin \varphi_p = 2m \int \frac{d^3 p}{(2\pi)^3} G_{\pi},$$
(30)

and it is easy to recognise the celebrated Gell-Mann–Oakes–Renner relation [30] in Eq. (30). Indeed, using the explicit form of the pionic wave function beyond the chiral limit, Eq. (29), one can see that $G_{\pi} = (\mathcal{N}_{\pi}/\sqrt{m_{\pi}}) \sin \varphi_p$ and $F_{\pi} = (\mathcal{N}_{\pi}\sqrt{m_{\pi}})\Delta_p$ which, after substitution to Eq. (30), give the sought relation,

$$m_{\pi}^{2} \left[\frac{N_{C}}{\pi^{2}} \int_{0}^{\infty} dp \ p^{2} \Delta_{p} \sin \varphi_{p} \right] = -2m \langle \bar{q}q \rangle, \tag{31}$$

where the definition of the chiral condensate [21, 23],

$$\langle \bar{q}q \rangle = -\frac{N_C}{\pi^2} \int_0^\infty dp \ p^2 \sin \varphi_p, \tag{32}$$

was used. Therefore,

$$\mathcal{N}_{\pi} = \frac{\sqrt{2\pi N_C}}{f_{\pi}}.$$
(33)

Finally,

$$\Psi_{\pi}(\vec{p}) = T_p \left[P_+ \gamma_5 \varphi_{\pi}^+(p) - P_- \gamma_5 \varphi_{\pi}^-(p) \right] T_p = \frac{1}{f_{\pi}} \sqrt{\frac{2\pi N_C}{m_{\pi}}} [\gamma_5 \sin \varphi_p + O(m_{\pi})], \quad (34)$$

where the properties $T_p \gamma_5 = \gamma_5 T_p^{\dagger}$ and $T_p^{\dagger} T_p = T_p T_p^{\dagger} = 1$ of the Foldy operator T_p (see Eq. (18) for its definition) were taken into account.

C. Goldberger–Treiman relation and the pion emission vertex

The standard Goldberger-Treiman relation is the relation between the pion-nucleon coupling constant and the axial constant of the nucleon axial vector current, and its derivation is present in any textbook on hadronic physics (see, for example, Ref. [31]). This relation can be derived analytically for the GNJL model as well [32]. A similar relation holds in the GNJL models for the pion coupling to dressed quarks. Indeed, the model admits two representations: the dressed-quark and the mesonic representation which are interchangeable up to corrections suppressed by the large N_C number [25], so that, in the leading in the number of colours approximation, one has two equivalent representations for the axial-vector current complying with the PCAC theorem (for the sake of simplicity, we stick to the one-flavour theory and ignore the axial anomaly, generalisation to the flavour nonsinglet axial vector current in $SU(N_f)$, where there is no anomaly, is trivial)¹:

$$[J^5_\mu(x)]_\pi = f_\pi \partial_\mu \phi_\pi(x), \tag{35}$$

through the Goldstone boson, with $\phi_{\pi}(x)$ being the chiral pion wave function, and, generically,

$$[J^5_{\mu}(x)]_q = \bar{q}(x)[g_A\gamma_{\mu}\gamma_5 + h_A\partial_{\mu}\gamma_5]q(x), \qquad (36)$$

through the dressed–quark states, where $g_A = 1$ and $h_A = 0$. We evaluate now the matrix element of this current divergence between dressed quark states, $\langle q(p)|\partial_{\mu}J^{5}_{\mu}(x)|q(p')\rangle$, in the given two representations arriving at:

$$\langle q(p) | [\partial_{\mu} J^{5}_{\mu}(x)]_{\pi} | q(p') \rangle = f_{\pi} m_{\pi}^{2} \langle q(p) | \phi_{\pi}(x) | q(p') \rangle \propto f_{\pi} g_{\pi}(q^{2}) (\bar{u}_{p} \gamma_{5} u_{p'}), \quad q = p - p', \quad (37)$$

where the pion-quark-quark effective formfactor $g_{\pi}(q^2)$ is introduced, and

$$\langle q(p)|[\partial_{\mu}J^{5}_{\mu}(x)]_{q}|q(p')\rangle \propto m_{q}^{\text{eff}}(\bar{u}_{p}\gamma_{5}u_{p'}),$$
(38)

¹It was noticed long ago [21] that, in the given model, the pion decay constant in the temporal and in the spatial parts of the current in Eq. (35) may differ. This is a consequence of an explicit breaking of Lorentz covariance by the instantaneous interaction in the Hamiltonian (2). Although some improvements of the model can be made in order to get rid of this discrepancy (see, for example, Ref. [33]), its underlying origin cannot be removed by simple amends. Thus we consider the model (2) as it is, making emphasis on its qualitative predictions. Everywhere throughout this paper as f_{π} we denote the temporal constant extracted from the Gell-Mann–Oakes–Renner relation (31).



FIG. 4: The amplitude of the decay $h \to h' + \pi$.

where it is assumed that dressed quarks obey an effective Dirac equation with the dynamically generated effective Lorentz-scalar mass m_q^{eff} . Obviously this dynamical mass must be identified with the function A_p — see Eqs. (10) and (13), — and we discuss this issue in detail below.

Finally, Eqs. (37) and (38) together yield the needed relation,

$$f_{\pi}g_{\pi} = m_q^{\text{eff}}, \quad g_{\pi} \equiv g_{\pi}(m_{\pi}^2), \tag{39}$$

where, for the sake of simplicity, we absorb all coefficients into the definition of g_{π} .

This derivation of the Goldberger-Treiman relation (39) is based on quite general considerations of chiral symmetry breaking and the PCAC theorem. The only input is the requirement of the (partial) conservation of the total axial vector current in QCD and appearance of the Lorentz-scalar dynamical mass of quarks as a consequence of SBCS. It leaves a number of questions, such as the microscopic picture for the pion-quark-quark vertex and the dependence of the effective quark mass m_q^{eff} on the quark momentum. Below we give a microscopic derivation of the Goldberger-Treiman-like relation (39) for the pion-quark-quark vertex in the framework of the quark model (2).

Consider a hadronic process with an emission of a soft pion, for example, a decay $h \rightarrow h' + \pi$, with h and h' being hadronic states (below we consider them to be mesons) with the total momenta \vec{p} and $\vec{p'}$, respectively. The amplitude of this process is given by the sum of two triangle diagrams depicted at Fig. 4 and can be written as

$$M(h \to h' + \pi) = \int \frac{d^4k}{(2\pi)^4} Sp \left[\chi_h(\vec{k}, \vec{p}) S(k-p) \bar{\chi}_{h'}(\vec{k} - \vec{p}, \vec{p}') S(k-q) \bar{\chi}_{\pi}(\vec{k}, \vec{q}) S(k) \right] + \int \frac{d^4k}{(2\pi)^4} Sp \left[\chi_h(\vec{k}, \vec{p}) S(k-p) \bar{\chi}_{\pi}(\vec{k}, \vec{q}) S(k-q) \bar{\chi}_{h'}(\vec{k} - \vec{p}, \vec{p}') S(k) \right],$$
(40)

where the pion momentum is q = p - p'. Every vertex in these diagrams contains a Salpeter amplitude χ ($\bar{\chi}$ for outgoing vertices) which obeys a Bethe–Salpeter equation of the form of Eq. (15) which can be rewritten as

$$\chi(\vec{p}, \vec{P}) = \int \frac{d^3k}{(2\pi)^3} V(\vec{p} - \vec{k}) \gamma_0 \Psi(\vec{k}, \vec{P}) \gamma_0.$$
(41)

The incoming and outgoing matrix vertices for the given hadron are related to one another as

$$\bar{\chi}(\vec{p},\vec{P}) = \gamma_0 \chi^{\dagger}(\vec{p},\vec{P})\gamma_0.$$
(42)

Thence, considering Eqs. (34), (41), and (42) together, one arrives at the relation between the soft pion $(\vec{P}_{\pi} \equiv \vec{q} \rightarrow 0)$ emission vertex and the dressed quark effective dynamical mass:

$$f_{\pi}\bar{\chi}_{\pi}(\vec{p}) = \sqrt{\frac{2\pi N_C}{m_{\pi}}}\gamma_5 \int \frac{d^3k}{(2\pi)^3} V(\vec{p}-\vec{k})\sin\varphi_k = \text{const} \times \gamma_5 A_p, \tag{43}$$

where the definition of the function A_p , Eq. (13), was used. Thus, for the sake of transparency, introducing the formfactor $g_{\pi}(p)$ such that

$$\bar{u}_p \bar{\chi}_\pi(\vec{p}) u_p = \text{const} \times g_\pi(p) (\bar{u}_p \gamma_5 u_p), \tag{44}$$

with the same constant as in Eq. (43), one finally arrives at the Goldberger–Treiman relation,

$$f_{\pi}g_{\pi}(p) = A_p, \tag{45}$$

which explicitly relates the pion coupling to the dressed quarks with the effective chiral symmetry breaking Lorentz-scalar quark mass A_p . Notice an important difference between the naive Goldberger-Treiman relation (39) and the microscopic relation (45). In the former case, the pion coupling constant depends only of the momentum transfer in the pionic vertex (that is, the pion total momentum \vec{q}) squared, so that, for the on-shall pion, g_{π} is a constant. On the contrary, the pion emission vertex $\bar{\chi}_{\pi}(\vec{p},\vec{q})$ in the microscopic GNJL model depends on two arguments: one, as before, being the pion total momentum \vec{q} — as mentioned before, we treat it as for the standard derivation of the Goldberger-Treiman relation and continue the soft-pion vertex to the point $\vec{q} = 0$ — whereas the other argument being the momentum flow in the loop, which plays also the role of the momentum of the dressed quark the pion couples to. Without SBCS and beyond the chiral limit, $A_p = m$ and thus the coupling $g_{\pi}(p)$ turns to a constant. On the contrary, in the chiral limit and after SBCS, A_p is a rapidly decreasing function of the momentum p. Indeed, using the definition of A_p and that of the dressed quark dispersive law, one can find the relation:

$$A_p = E_p \sin \varphi_p. \tag{46}$$

At large momenta, $E_p \approx p$, whereas the chiral angle approaches zero fast — see Fig. 2. For a power-like confining potential (4) $\varphi_p \propto 1/p^{4+\alpha}$ as $p \to \infty$ (see, for example, Ref. [27]).

The decay amplitude of the process $h \to h' + \pi$ is given by an overlap of the three vertices — see Fig. 4 and Eq. (40). Each of them depends on the momentum circulating in the loop, and the maximal overlap is achieved for all three meson wave functions localised at comparable values of this momentum. Clearly the pion vertex (and the pion wave function $\sin \varphi_p$) is dominated by the low momenta and it decreases fast with the increase of the latter (see Fig. 2). In the meantime, for highly excited hadrons the momentum distribution in these hadrons as well as in the corresponding vertices in Fig. 4 is shifted to large momenta. Therefore, the amplitude (40) vanishes with the increase of the hadron h or/and h' excitation number and so does the effective coupling constant of the pion to such highly excited hadrons. We emphasise that it is the pion wave function $\sin \varphi_p$ that suppresses the Goldstone boson coupling to highly excited hadrons.

It was demonstrated recently that chiral symmetry restoration for excited hadrons happens due to the same reason — the effective interaction responsible for the splitting within a chiral doublet is also proportional to $\sin \varphi_p$ [12]. Therefore, if an effective constant is introduced in particular for the pion transition within the approximate parity doublet, this constant must be proportional to the splitting $\Delta M_{+-} = M_+ - M_-$ within the doublet and it has to vanish with ΔM_{+-} for highly excited hadrons.

D. Numerical estimates

In this chapter we present some quantitative estimates related to the chiral pion and its coupling to excited hadrons. For the sake of simplicity, we stick to the harmonic oscillator potential corresponding to the marginal choice of $\alpha = 2$ in Eq. (4). The mass–gap equation reduces in this case to a second–order differential equation studied in detail in Refs. [21, 22, 23],

$$p^{3}\sin\varphi_{p} = \frac{1}{2}K_{0}^{3}\left[p^{2}\varphi_{p}^{\prime\prime} + 2p\varphi_{p}^{\prime} + \sin 2\varphi_{p}\right] + mp^{2}\cos\varphi_{p},\tag{47}$$

where the dressed quark dispersive law is

$$E_p = m\sin\varphi_p + p\cos\varphi_p - K_0^3 \left[\frac{{\varphi'_p}^2}{2} + \frac{\cos^2\varphi_p}{p^2}\right].$$
(48)



FIG. 5: The ratio $g_{\pi}(p)/g_{\pi}(0)$ as a function of the dressed quark momentum.

The bound-state equation for the chiral pion, Eq. (21), takes the form

$$\left[\left[-K_0^3 \frac{d^2}{dp^2} + 2E_p \right] \left[\begin{array}{c} 1 & 0 \\ 0 & 1 \end{array} \right] + K_0^3 \left[\frac{\varphi_p'^2}{2} + \frac{\cos^2 \varphi_p}{p^2} \right] \left[\begin{array}{c} 1 & 1 \\ 1 & 1 \end{array} \right] - m_\pi \left[\begin{array}{c} 1 & 0 \\ 0 & -1 \end{array} \right] \right] \left[\begin{array}{c} \nu_\pi^+(p) \\ \nu_\pi^-(p) \end{array} \right] = 0,$$
(49)

where the radial wave functions $\nu_{\pi}^{\pm}(p) = p\varphi_{\pi}^{\pm}(p)$ were introduced for convenience, which are rescaled so as to obey the one-dimensional normalisation [23],

$$\int dp [\nu_{\pi}^{+2}(p) - \nu_{\pi}^{-2}(p)] = 1.$$
(50)

As noticed before, Eq. (49) reduces to the mass–gap Eq. (47) in the chiral limit, so that $\varphi_{\pi}^{\pm}(p) = \sin \varphi_p$ and the qualitative form of the solution for the chiral angle appears to be the same as for the linear confinement depicted in Fig. 2.

In Fig. 5, we plot the ratio $g_{\pi}(p)/g_{\pi}(0)$ defined with the help of Eqs. (45) and (46). This ratio also describes the actual decrease of the pion coupling to hadron with the increase of the average dressed quark momentum in it. Furthermore, for the sake of transparency, we measure this momentum in physical units, in MeV, using the chiral condensate to fix the value of the mass parameter K_0 . To this end, for the given solution to the mass–gap Eq. (47), we calculate the chiral condensate according to Eq. (32) which gives (we use the numerical results of Refs. [12, 25])

$$\langle \bar{q}q \rangle = -(0.51K_0)^3 \tag{51}$$

and, therefore, we fix $K_0 = 490 MeV$ to arrive at the standard value of the chiral condensate. From Fig. 5 one can see a fast decrease of the pion coupling to hadrons for large p's, as was discussed before using general qualitative arguments.

Now we estimate the average momentum of the valence quarks in excited mesons. We consider a radially excited S-wave light–light meson consisting of two light (massless) quarks. For highly excited bound states of such a system $(n \gg 1)$, the negative–energy component of the wave function $\varphi_n^-(p)$ vanishes and the bound–state equation reduces to a single equation for the positive–energy component $\varphi_n^+(p)$. This equation can be roughly approximated by the Schrödinger equation with the Hamiltonian

$$H = 2|p| + K_0^3 r^2, (52)$$

and we identify the corresponding selfenergies M_n as $M_n = \frac{1}{2}(M_{n+} + M_{n-})$, with $M_{n\pm}$ being the masses for the chiral doublet partners. Although the given naive Salpeter Hamiltonian is unable to correctly discriminate between M_{n+} and M_{n-} and a more sophisticated approach is required for this purpose (see Ref. [12] for a detailed discussion), it is sufficient to estimate the typical quark momentum in excited states. To make things simpler, we use the einbein field method (see the original paper [34] for the details and, for example, Ref. [35] for einbeins treated using the Dirac constraints formalism [36]) rewriting the Hamiltonian (52) as

$$H = \frac{p^2}{\mu} + \frac{\mu}{2} + K_0^3 r^2, \tag{53}$$

with the einbein μ treated as a variational parameter [37]. The spectrum of the Hamiltonian (53) minimised with respect to the parameter μ reads $M_n \approx 3K_0(2n+3/2)^{2/3}$ and the mean quark momentum can be easily estimated then using the virial theorem for the oscillator Hamiltonian (53) to be

$$\langle p \rangle \approx K_0 \left(2n + \frac{3}{2}\right)^{2/3},$$
(54)

which gives an approximate analytic dependence of $\langle p \rangle$ on the radial excitation number n. It is obvious therefore that already for $n \sim 1$ the quark momentum appears large enough and, as seen from Fig. 5, the corresponding coupling of the Goldstone boson to this hadron is suppressed as compared to its naive value of $g_{\pi}(0)$. An accurate systematic evaluation of the pion coupling to excited hadrons goes beyond the scope of the present qualitative paper — this work is in progress now and will be reported elsewhere.

IV. CONCLUSIONS

In this paper we considered in detail physics of the Goldstone bosons decoupling from highly excited hadrons, where chiral symmetry is approximately restored. This interesting physics can be summarised in a few words. The coupling of the Goldstone bosons to the valence quarks is regulated, via the Goldberger–Treiman relation, by the dynamical Lorentz– scalar mass of quarks. This dynamical mass which arises via the loop dressing of quarks represents effects of chiral symmetry breaking in the vacuum. A key feature of this dynamical mass is that it is strongly momentum-dependent and vanishes at large quark momenta. Hence at large momenta the valence quarks decouple from the quark condensates and from the Goldstone bosons. In this regime chiral symmetry breaking in the vacuum almost does not affect observables and physics is essentially such as if there had been no such chiral symmetry breaking in the vacuum. It is clear, therefore, that for a given hadron, the influence of SBCS on its properties is determined by the typical momentum of valence quarks in it. For low-lying hadrons this momentum lies below the chiral symmetry breaking scale, the valence quarks acquire a significant Lorentz–scalar dynamical mass, which results in a strong coupling of the low-lying hadrons to the Goldstone bosons. On the contrary, for high-lying hadrons, a typical momentum of valence quarks is large, their dynamical chiral symmetry breaking mass becomes small (and asymptotically vanishes), which implies that these high-lying hadrons decouple from the Goldstone bosons. Consequently the axial vector constant, g_A , of the highly excited hadrons appears suppressed and asymptotically vanishes.

We illustrate this physics in the framework of the GNJL model for QCD with the only interaction between quarks being the instantaneous Lorentz–vector confining potential. Such a model is tractable and contains all the required elements such as the spontaneous breaking of chiral symmetry via quantum fluctuations of the quark fields and confinement. While being only a model, it nevertheless provides a significant insight into physics of chiral symmetry restoration in excited hadrons.

Acknowledgments

L. Ya. G. thanks A. Gal for his hospitality at the Hebrew University of Jerusalem where this paper was prepared, and T. Cohen, M. Shifman and A. Vainstein for correspondence. He acknowledges the support of the P16823-N08 project of the Austrian Science Fund. A. V. N. would like to thank Yu. S. Kalashnikova and E. Ribeiro for reading the manuscript and critical comments and to acknowledge support of the grants DFG 436 RUS 113/820/0-1, RFFI 05-02-04012-NNIOa, NSh-843.2006.2 as well as of the Federal Programme of the Russian Ministry of Industry, Science, and Technology No 40.052.1.1112.

- [1] L. Ya. Glozman, Phys. Lett. B **475**, 329 (2000).
- [2] T. D. Cohen and L. Ya. Glozman, Phys. Rev. D 65, 016006 (2002); Int. J. Mod. Phys. A 17, 1327 (2002).
- [3] L. Ya. Glozman, Phys. Lett. B 539, 257 (2002); Phys. Lett. B 587, 69 (2004).
- [4] L. Ya. Glozman, Phys. Lett. B **541**, 115 (2002).
- [5] L. Ya. Glozman, hep-ph/0411281, Int. J. Mod. Phys. A., in press.
- [6] L. Ya. Glozman, Acta Phys. Polon. B 35, 2985 (2004); Nucl. Phys. A 755, 17 (2005).
- [7] T. D. Cohen and L. Ya. Glozman, hep-ph/0512185.
- [8] S. Beane, Phys. Rev. D 64, 116010 (2001); M. Golterman and S. Peris, Phys. Rev. D 67, 096001 (2003); S. S. Afonin et. al, JHEP 0404, 039 (2004).
- [9] E. Swanson, Phys. Lett. B **582**, 167 (2004).
- [10] T. DeGrand, Phys. Rev. D 69, 074024 (2004).
- [11] M. Shifman, hep-ph/0507246.
- [12] Yu. S. Kalashnikova, A. V. Nefediev, and J. E. F. T. Ribeiro, Phys. Rev. D 72, 034020 (2005).
- [13] L. Ya. Glozman, A. V. Nefediev, and J. E. F. T. Ribeiro, Phys. Rev. D 72, 094002 (2005).
- [14] R. L. Jaffe, D. Pirjol, and A. Scardicchio, hep-ph/0602010.
- [15] M. L. Goldberger and S. B. Treiman, Phys. Rev. 111, 354 (1958).
- [16] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961); **124**, 246 (1961).
- [17] U. Vogl and W. Weise, Progr. Part. Nucl. Phys. 27, 195 (1991); S. P. Klevansky, Rev. Mod.
 Phys. 64, 649 (1992); T. Hatsuda and T. Kunihiro, Phys. Rep. 247, 221 (1994).

- [18] T. Schäfer and E. Shuryak, Rev. Mod. Phys. 70, 323 (1998); D. Diakonov, Progr. Part. Nucl.
 Phys. 51, 173 (2003).
- [19] Yu. A. Simonov, Yad. Fiz. 60, 2252 (1997) (Phys. Atom. Nucl. 60, 2069 (1997)); Phys. Rev. D 65, 094018 (2002); Yad. Fiz. 67, 868 (2004) (Phys. Atom. Nucl. 67, 846 (2004)); J. A. Tjon and Yu. A. Simonov, Phys. Rev. D 62, 014501 (2000); A. V. Nefediev and Yu. A. Simonov, Pisma v ZhETF 82, 633 (2005) (JETP Lett 82 557 (2005)).
- [20] C. D. Roberts, A. G. Williams, Progr. Part. Nucl. Phys. 33, 477 (1994); R. Alkofer and L. von Smekal, Phys. Rep. 353, 281 (2001).
- [21] A. Amer, A. Le Yaouanc, L. Oliver, O. Pene, and J.-C. Raynal, Phys. Rev. Lett. 50, 87 (1983);
 A. Le Yaouanc, L. Oliver, O. Pene, and J.-C. Raynal, Phys. Lett. 134B, 249 (1984); Phys. Rev. D 29, 1233 (1984).
- [22] A. Le Yaouanc, L. Oliver, S. Ono, O. Pene and J.-C. Raynal, Phys. Rev. D 31, 137 (1985).
- [23] P. Bicudo and J. E. Ribeiro, Phys. Rev. D 42, 1611 (1990); *ibid.*, 1625 (1990); *ibid.*, 1635 (1990); P. Bicudo, Phys. Rev. Lett. 72, 1600 (1994).
- [24] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Phys. Rev. 106, 162 (1957); *ibid.* 108, 1175 (1957).
- [25] A. V. Nefediev and J. E. F. T. Ribeiro, Phys. Rev. D 70, 094020 (2004).
- [26] P. J. A. Bicudo, A. V. Nefediev, and J. E. F. T. Ribeiro, Phys. Rev. D 65, 085026 (2002).
- [27] P. J. A. Bicudo and A. V. Nefediev, Phys. Rev. D 68, 065021 (2003).
- [28] I. Bars, M. B. Green, Phys. Rev. D 17, 537 (1978).
- [29] Yu. S. Kalashnikova and A. V. Nefediev, Yad. Fiz. 62, 359 (1999) [Phys. Atom. Nucl. 62, 323 (1999)]; Usp. Fiz. Nauk 172, 378 (2002) [Phys. Usp. 45, 347 (2002)]; Yu. S. Kalashnikova, A. V. Nefediev, A. V. Volodin, Yad. Fiz. 63, 1710 (2000) [Phys. Atom. Nucl. 63, 1623 (2000)].
- [30] M. Gell-Mann, R. J. Oakes, and B. Renner, Phys. Rev. 175, 2195 (1968).
- [31] T.-P. Cheng and L.-F. Li, Gauge theories in elementary particles physics (Oxford University Press, London, England, 1984).
- [32] P. Bicudo, Phys. Rev. C 67, 035201 (2003).
- [33] P. Bicudo, Phys. Rev. C **60**, 035209 (1999).
- [34] L. Brink, P. Di Vecchia, and P. Howe, Nucl. Phys. B 118, 76 (1977).
- [35] Yu. S. Kalashnikova and A. V. Nefediev, Yad. Fiz. 60, 1529 (1997) (Phys. Atom. Nucl. 60, 1389 (1997)); Yad. Fiz. 61, 871 (1998) (Phys. Atom. Nucl. 61 785 (1998)).

- [36] P. A. M. Dirac, Lectures on Quantum Mechanics (Belter Graduate School of Science, Yeshiva University, New York, 1964).
- [37] A. Yu. Dubin, A. B. Kaidalov, and Yu. A. Simonov, Phys. Lett. B 323, 41 (1994); Phys. Lett. B 343, 310 (1995); E. L. Gubankova and A. Yu. Dubin, Phys. Lett. B 334, 180 (1994).