Hybrid meson properties in Lattice QCD and Flux Tube Models

T. J. Burns^{*} and F. E. $Close^{\dagger}$

Rudolf Peierls Centre for Theoretical Physics, University of Oxford, Keble Rd., Oxford, OX1 3NP, United Kingdom

(Dated: June 29, 2018)

Flux tube model predictions for hybrid meson decays are beginning to be confronted by Lattice QCD. We compare the two approaches for the S-wave decay of the exotic 1^{-+} , and find excellent agreement. Results suggest that in Strong QCD $q\bar{q}$ creation occurs with S = 1.

While Lattice QCD is now a mature guide for the masses of glueballs and hybrids, at least in the quenched approximation [1, 2], it is not yet able to determine hadronic decays extensively. Flux tube models of both spectra [3, 4] and decays[5, 6] have been developed, in part stimulated by attempts to model the lattice, and lattice work has confirmed their spectroscopy[2, 4]. The first study of the hybrid meson decays $1^{-+} \rightarrow \pi b_1$ and πf_1 has recently been made in lattice QCD[7] and shows features that had been anticipated in flux-tube models[5, 6]. In this paper we compare these results and assess the implications.

Lattice QCD enables the properties of strong QCD to be explored "experimentally" over a kinematic and parameter space that is richer than the "physical" values to which nature has restricted us. Thus, for example, ref [7] is restricted to S-wave decays, which it achieves with specially chosen masses so as to produce a decay at rest. To convert the results to widths (presented in Table I) a large extrapolation was made from the threshold, relying on an assumption that is beyond lattice QCD, thereby masking its primary results. It is tantalising that the ratio of widths to πb_1 and πf_1 in lattice QCD and the flux tube simulations agree whereas the absolute widths, as presented in ref[7], are rather larger than those of refs[5, 6]. The lattice QCD results for the known decay $b_1 \rightarrow \omega \pi$ also exceed data when the extrapolations of ref [7] are employed.

This prima facie suggests that the spin-dependent features of Strong QCD as revealed by the lattice are contained within the flux tube model, but that the momentum dependence of the assumed extrapolation differs. We show here that this is the case; that when the flux tube model is applied in the $k \to 0$ limit of the lattice the agreement is excellent; and that the results of lattice QCD reinforce the flux-tube hypothesis that $q\bar{q}$ creation is spin triplet.

Decay amplitudes in Lattice QCD and Flux Tube Models as $k \rightarrow 0$

The flux tube model has successfully described transitions among conventional mesons, $\mathcal{M} \to \mathcal{M} + \mathcal{M}[9]$, and also been applied to the decays of hybrid mesons, $\mathcal{H} \to \mathcal{M} + \mathcal{M}[5, 6, 10]$. A notable feature of the latter, which also emerges in some other models[11], is that the prominent decays are to excited mesons, notably S + P states[5]. In particular the exotic π_1 1⁻⁺ (called $\hat{\rho}$ in [7]) is expected to have prominent decays into πb_1 and πf_1 , with the former favoured by about a factor of four in both ${}^{3}P_0$ and ${}^{3}S_1$ flux-tube models[5, 6, 10]. For a decay of width Γ at momentum k, the coupling constant for the process $A \to B + C$ can be expressed

$$\Gamma/k = \frac{1}{\pi} (\mathbf{PS}) \langle BC | \sigma \cdot \nabla | A \rangle^2 \tag{1}$$

where **PS** denotes phase space and $\langle BC|\sigma \cdot \nabla |A\rangle$ is the overlap of the quark and string wavefunctions coupled by the ³P₀ string breaking operator. The overlap is explicitly momentum dependent, and assumes the same form in each of the cases of present interest. Defining

$$S(k) = \frac{\gamma_0 \pi^{3/4} 2^{5/2}}{\beta^{1/2} 3} \left(1 - \frac{2k^2}{9\beta^2} \right) e^{-k^2/12\beta^2}$$
(2)

^{*} e-mail: burns@thphys.ox.ac.uk

[†] e-mail: F.Close1@physics.ox.ac.uk

	FTM	FTM	Lattice
	Ref $[5]$	$\operatorname{Ref}[6]$	$\operatorname{Ref}\left[7\right]$
	$1.9 \mathrm{GeV}$	$2.0~{\rm GeV}$	$2.0~{\rm GeV}$
$\Gamma(\pi_1 \to b_1 \pi)_S$	100	70	400 ± 120
$\Gamma(\pi_1 \to b_1 \pi)_D$	30	30	
$\Gamma(\pi_1 \to f_1 \pi)_S$	30	20	90 ± 60
$\Gamma(\pi_1 \to f_1 \pi)_D$	20	25	

TABLE I: Comparison of flux tube and lattice predictions for π_1 decays.

the relevant S-wave amplitudes can be written [5, 6]

$$\langle \omega \pi | \sigma \cdot \nabla | b_1 \rangle = S(k) \tag{3}$$

$$\langle b_1 \pi | \sigma \cdot \nabla | \pi_1 \rangle = \sqrt{2} \frac{\kappa \sqrt{b}}{\beta} S(k)$$
 (4)

$$\langle f_1 \pi | \sigma \cdot \nabla | \pi_1 \rangle = \frac{1}{\sqrt{2}} \frac{\kappa \sqrt{b}}{\beta} S(k)$$
 (5)

for harmonic oscillator wavefunctions of equal width β and where γ_0 is an overall pair creation constant fit to data. We defer further discussion of the parameters temporarily, noting only that the relative scale of hybrid decays to that of conventional mesons is driven by the string tension b and a factor κ that emerges from the overlap of the excited flux tube with ground state flux tubes: thus in the flux tube model there is an immediate correlation of scale between the decay widths of conventional and hybrid states.

In Table I we present flux tube predictions for the π_1 width and compare with results extrapolated from lattice QCD in ref[7]. Ref [5] calculated the dominant decay amplitudes of exotic- J^{PC} hybrid states with a localised pair creation region, quoted results for a π_1 state at 1.9GeV. Ref [6] verified those results in an infinite flux tube approximation and calculated analytic forms for both exotic and non exotic hybrid decays as a function of mass. We quote the results for a π_1 at 2.0 GeV having corrected some numerical factors in ref [6], and present the amplitudes for arbitrary π_1 mass in Fig. 1. The model predicts widths for the 1⁻⁺ decays in both S and D waves and as a function of the hybrid mass, with the S-wave contributions dominating.

The lattice technique is to put a given decay channel at roughly the same energy as the decaying state so that the decay is just allowed while conserving energy in a two-point function [7]. For a lattice with spacing a and size L, the lattice transition amplitude xa gives the analogue of equations (3)–(5):

$$\langle BC|\text{Latt}|A\rangle = (L/a)^{3/2}(xa) \tag{6}$$

so that the coupling constant at threshold, with phase space \mathbf{PS} chosen appropriately with lattice masses and a decay at threshold, can be written

$$\Gamma/k = \frac{1}{\pi} (\mathbf{PS}) \langle BC | \text{Latt} | A \rangle^2.$$
(7)

In order to make a statement about physical widths, ref.[7] assumes that Γ/k doesn't vary with quark mass. This linear extrapolation leads to the large width of $\Gamma = 400 \pm 120$ MeV in Table I for a π_1 at 2.0GeV decaying to $b_1\pi$ with physical masses. Ref [7] also presented results for the conventional decay $b_1 \rightarrow \omega \pi$ noting that an equivalent extrapolation overestimates the data and that this could be generic. We note the result would be ~ 220MeV, significantly larger than the S-wave data ~ 130MeV [8].

On the other hand, the flux tube model extrapolation has been tested over a large range of k, predicting accurately the decays of both mesons and baryons [9, 12, 13]. For the physical b_1 at 1235MeV decaying to $\omega \pi$ the model reproduces very nicely the experimental data for both S-wave ~ 130MeV (using equation (3)) and D-wave ~ 10MeV. The successful phenomenology of this and a wide range of other conventional meson decays relies on momentumdependent form factors arising from the overlap of hadron wavefunctions. The need for such form factors is rather general, empirically supported as exclusive hadron decay widths do not show unrestricted growth with phase space[8]. Such phenomena are also expected for hybrid decays $\mathcal{H} \to \mathcal{M} + \mathcal{M}$ and appear explicitly in equations (4) and (5).

The extrapolation from the lattice limit $k \to 0$ assumed in ref [7] ignores any such k dependent suppression: thus the predicted widths are much larger and for $b_1 \to \omega \pi$ disagree with experiment. Such an assumption may apply for inclusive decays but is unphysical for exclusive channels as here. As the momentum k increases, individual channels

	b_1	$ ightarrow \omega \pi$	π_1	$\rightarrow b_1 \pi$	π_1	$\rightarrow f_1 \pi$
Lattice (C410)	2.3	± 0.1	2.9	± 0.4	1.5	± 0.4
Lattice $(U355)$	3.4	± 0.2	2.9	± 0.3	1.1	± 0.4
Flux tube (A)	2.7		2.9		1.4	
Flux tube (B)	3.3		3.9		1.9	

TABLE II: Transition amplitudes $\langle BC|\sigma \cdot \nabla |A\rangle$ and $\langle BC| \text{latt} |A\rangle$ in units of $\text{GeV}^{-1/2}$. The parameter sets (A) and (B) are (A) $\gamma_0 = 0.39, \beta = 0.40, \kappa = 0.7$

(B) $\gamma_0 = 0.45, \beta = 0.36, \kappa = 0.7$

fall at the expense of multi-body channels opening (this is the physics of exclusive form factors) even though the sum of channels may be k independent (scale invariant) [15]. More generally, the hadron size sets an explicit scale against which the momentum k of the exclusive process is weighed; a linear extrapolation ignores this.

Another difference is the functional parametrisation of phase space (\mathbf{PS}) in refs [5, 6] and [7], respectively

$$(\mathbf{PS})_{ft} = \frac{\widetilde{M}_B \widetilde{M}_C}{\widetilde{M}_H}, \qquad (\mathbf{PS})_{lat} = \frac{a E_B a E_C}{a E_B + a E_C}$$
(8)

where \widetilde{M} are meson masses calculated before spin interactions (for a discussion see Appendix A2 of ref [12]) and aE are the lattice masses. The latter phase space is for a decay at threshold: since the physical decay is far from threshold, $(\mathbf{PS})_{ft} < (\mathbf{PS})_{lat}$.

Thus there is no direct comparison between the widths of the lattice predictions in Table I and those of the flux tube model: the former works with unphysical masses and calculates the amplitude at threshold, extrapolating to the large momenta required for the physical masses; the latter calculates the amplitude with physical masses far from threshold, with dynamics at this momenta determined by the overlap of quark and string wavefunctions.

To compare the two approaches we evaluate the flux tube predictions for k = 0 and compare not the coupling constants Γ/k but the transition amplitudes $\langle BC | \sigma \cdot \nabla | A \rangle$ and $\langle BC | \text{latt} | A \rangle$. The lattice couplings follow from the slopes (xa) recorded in ref [7]; we list these couplings, for the two different codes, in the first two rows of Table II. The analagous flux tube couplings follow from equations (3)–(5) without further assumption, and these are shown in the remaining rows of Table II for two "standard" parameter sets (discussed below). The overall scale of decays in the flux tube model is driven by the pair creation constant γ_0 and the hadronic wavefunction width β , both of which are strongly constrained by data. The model then fixes the scale of decays involving hybrid mesons according to the overlap of the string degrees of freedom, exhibited in the ratio $\kappa \sqrt{b}/\beta$.

Before comparing the two approaches, we briefly discuss the nature of the parameter selection appropriate for the flux tube. The analytic expressions (3)–(5) are those appropriate to a radial hybrid wavefunction $\sim r^{\delta} e^{-\beta^2 r^2/2}$ with $\delta = 1$; as noted in [6] the results differ very little from the "true" radial wavefunction which, ignoring a term that raises/lowers the gluonic angular momentum, has $\delta \approx 0.6$. As they are presented, the expressions (3)–(5) correspond to setting $e^{-fby_{\perp}^2/2} = 1$ in the flux tube overlap term. The original flux tube formulation established that the inclusion of a localised pair creation region has very little effect on the predictions, on account of the asymptotic forms of the hadron wavefunctions automatically imposing a "flux tube"-like structure on the overlaps [9]. The approach of [6] uses an infinitely long flux tube, but for the case of equal wavefunction widths the flux tube information integrates out and merely rescales the parameter γ_0 . Thus in a sense the infinite flux tube approximation is equivalent to the absence of a $e^{-fby_{\perp}^2/2}$ flux tube term altogether: the naive ${}^{3}P_{0}$ model with equal probability pair creation everywhere in space has effectively been recovered, with a re-scaling of the overall pair creation parameter. Thus the appropriate choice of the γ_0 that appears in (3)–(5) is that which has been fit to the decays of mesons in a model with no flux tube supression away from the $q\bar{q}$ axis: several such fits are available and constrain both γ_0 and β rather strongly. The authors of ref [5] advocate $\gamma_0 = 0.39$ and $\beta = 0.4$; ref [13] fit to a wide range of higher quarkonia and settle on $\gamma_0 = 0.4 - 0.5$ and $\beta = 0.36$. As is evident Table II, these parameter sets correctly set the scale of the decay $b_1 \rightarrow \omega \pi$. The spread of values in the lattice calculation is reproduced by the spread of "phenomenologically-allowed" flux tube parameters.

The relative scale of hybrid decays is then set by the ratio $\kappa \sqrt{b}/\beta$. The string tension $b = 0.18 \text{GeV}^2$ is rather tightly constrained, thus the results hinge only on the factor κ that emerges directly from the overlap of the stringlike degrees of freedom of the hybrid and conventional mesons. Explicit calculation in the framework of the harmonic approximation [17] has κ as a function that depends on the longitudinal distance along the original $q\bar{q}$ axis at which the string breaks, varying from 0.8 - 0.9 at its peak in the centre of the meson to 0.4 - 0.6 at either end, depending on the degree of quantization. Refs [5] and [6] treated κ as a constant, choosing $\kappa = 1$ and 0.9 respectively. The




FIG. 1: Partial widths of π_1 as a function of mass in the flux tube model with parameter set (A).

aforementioned discussion would suggest $\kappa \sim 0.7$ may be a more realistic implementation of the model, and indeed such a choice reproduces the lattice calculations rather nicely.

The hadrons of the lattice are generally heavier than their experimental values, and as such it is not automatic that the β fit to the decays of experimental states should reproduce the decays on the lattice. However, it is only the pion that is drastically different from its true mass; moreover, this lattice-pion is no longer exceptional and in consequence may be described by the same scale as other hadrons.

Finally we consider the possibility of allowing the initial and final state wavefunction to have different widths. Ref [6] prefer a slightly smaller β for the initial hybrid state: such a choice manifests itself in a rescaling by β_A/β_B . The effect on the numbers in Table II is not drastic, though it improves the agreement somewhat, particularly for parameter set (B) with its larger γ_0 .

With the state of the lattice uncertainties at present, it is not appropriate to attempt a best fit for flux tube parameters. We are very much encouraged, however, that the "standard" choice of harmonic oscillator parameters reproduces the lattice results with remarkable accuracy. Even more encouraging is that the string overlap factor correctly sets the scale of hybrid decays relative to conventional meson decays: in this result we have at a very direct test of the flux tube dynamics. Agreement with the lattice is non-trivial: the relative strengths of hybrid and conventional decay amplitudes emerge naturally from the string-like description of the gluonic degrees of freedom and are determined by the same string tension that controls the conventional hadron spectrum. It is notable that the choice of κ suggested by the harmonic approximation calculations reproduces very nicely this relative scale. That this choice is smaller than that traditionally used suggests that the previous predictions for hybrid widths should be scaled by $\approx (0.7/0.9)^2 \approx 0.6$, which is encouraging from the point of view of the experimental hunt for these states. We also note that the same relative scale emerges from a continuum string picture, as can be seen from the plot of the continuum limit of κ in [17].

Emboldened by the good agreement, we suggest an extrapolation of the lattice results at threshold to the physical region. To the extent that the two approaches agree at k = 0, the relevant "extrapolation" is explicit in equations (4) and (5). The results are shown in Fig 1 with parameter set (A) (see the caption of Table II), spin-averaged masses in the phase space, and an f_1 octet-singlet mixing angle of 50° following [14]. The rather more modest widths than those of the linear lattice extrapolation offer encouragement for the experimental observation of such states. We note in passing, however, that if flux tube and lattice mass estimates are a guide, hybrids may couple strongly to many previously unconsidered modes such as 1P + vector and 1D + pseudoscalar. Although such modes appears not to spoil the rather narrow π_1 , they are important for other states such as 2^{+-} [18].

The predictions for isoscalar decays to πa_1 [7] also agree in relative scale with the flux tube. This is trivial as only Zweig connected diagrams are considered and the different modes are related primarily by flavour factors. Likewise the transition amplitudes for S-wave KK_1 directly follow from the aforementioned discussion, with some adjustments due to relative flavour factors: we find $\langle K^1 P_1 K | \sigma \cdot \nabla | \pi_1 \rangle = \langle b_1 \pi | \sigma \cdot \nabla | \pi_1 \rangle / \sqrt{2}$ and $\langle K^3 P_1 K | \sigma \cdot \nabla | \pi_1 \rangle = \langle f_1 \pi | \sigma \cdot \nabla | \pi_1 \rangle$. The ratio of πb_1 to πf_1 allows a direct test of the pair creation mechanism without entering into the nuances of parameters. The averaged lattice results of ref [7] suggest that the former is favoured by a factor of 2 in amplitude (4 in width), which is tantalising in that both the ${}^{3}P_{0}$ and ${}^{3}S_{1}$ decays models predict such a ratio, as can be read off from equations (4) and (5) in the ${}^{3}P_{0}$ case. Flavour immediately accounts for an enhancement of $b_{1}\pi$ by a factor of $\sqrt{2}$ in amplitude; the remaining enhancement observed in the lattice is therefore due to the spin and angular dynamics that differentiate the b_1 and f_1 (with quark spin 0 and 1 respectively).

Fundamental to the flux tube models has been the assumption that L and S factorise and that S is conserved in the decay. The latter assumption, coupled with spin 1 pair creation, forbids decays of the type $(S = 0) \rightarrow (S = 0) + (S = 0)$ by orthogonality of the spin wavefunctions and experiment seems to support it, for instance the absence of the decay $\pi_2(1670) \rightarrow b_1\pi$. On the other hand, S = 0 pair creation would forbid decays of the type $(S = 1) \rightarrow (S = 0) + (S = 0)$ and $(S = 0) \rightarrow (S = 1) + (S = 0)$; a possibility already excluded by data: $\rho \rightarrow \pi\pi$ and $b_1 \rightarrow \omega\pi$ are examples of such modes, both well known experimentally and on the lattice.

In flux tube and constituent gluon models, the 1^{-+} has $q\bar{q}$ in spin 1 and so would not decay to $b_1\pi$ if its decay were driven by spin 0 pair creation. Thus the lattice observation $\pi_1 \to b_1\pi$ supports the flux tube hypothesis that hybrid decays are driven by the same pair creation mechanism as that of conventionals, and that a hybrid can be described in terms of quark S = 0 or 1 coupled to L = 1 carried explicitly by the gluonic flux tube.

Spin 1 pair creation is natural in the flux tube model, with both ${}^{3}S_{1}$ and ${}^{3}P_{0}$ operators emerging from expansion of the strong coupling Hamiltonian. In the extremely strong coupled limit where the flux tube is straight, pair creation occurs along the interquark axis in ${}^{3}S_{1}$ via the operator $\sigma \cdot \hat{r}$. In the original formulation of the model it was argued that zero point oscillations of the flux tube will wipe out this term leaving instead ${}^{3}P_{0}$ pair creation via the operator $\sigma \cdot \nabla$, and this was found to give better agreement with experiment [3]. In either mechanism, there is a common feature that may well be verified by the lattice: the decay of $\pi_{1} \rightarrow \pi b_{1}$ is dominant over πf_{1} by a factor of four, independently of momentum [5, 6, 7, 10]. The origin of the dynamical effect is not readily explained by the lattice calculation; in the flux-tube model, as we now demonstrate, it emerges naturally from spin 1 pair creation by a scalar operator, and subsequent recoupling to final state mesons.

As shown in equations (4) and (5) of [6], the flux tube model decay amplitude can be written as a linear combination of spatial overlaps $I_{M_L^A M_L^B}$, that combination determined by angular momentum and partial wave recoupling of the initial and final states. The $I_{M_L^A M_L^B}$ are the matrix elements of the pair creation operator between the initial and final quark and string wavefunctions. Including flavour and spin wavefunction overlaps as in eqn. (3) of [6] gives

$$S(k) = \sqrt{3} \langle \phi_B \phi_C | \phi_A \phi_0 \rangle \sum_{M_L^A M_S^A \lambda} \langle 1 \ M_L^A, 1 \ M_S^A | 1 \ M_L^A + M_S^A \rangle \langle 1 \ \lambda, 1 - \lambda | 0 \ 0 \rangle \tag{9}$$

$$\langle 1 \ M_L^A + \lambda, S_B \ M_S^A - \lambda | 1 \ M_S^A + M_L^A \rangle \langle \ \chi_{S_B}^{M_S^A - \lambda} \chi_0^0 | \ \chi_1^{M_S^A} \chi_1^{-\lambda} \ \rangle I_{M_L^A, M_L^A + \lambda}(k) \tag{10}$$

where $\langle \phi_B \phi_C | \phi_A \phi_0 \rangle$ and $\langle \chi_{S_B}^{M_A^{-\lambda}} \chi_0^0 | \chi_1^{M_A^{-\lambda}} \chi_1^{-\lambda} \rangle$ are the overlaps of the flavour and spin wavefunctions of the initial meson $(q_1 \overline{q_2})$ and the emergent pair $(q_3 \overline{q_4})$ reordered to give final states $(q_1 \overline{q_4})$ and $(q_3 \overline{q_2})$, and the $q_3 \overline{q_4}$ emerge in either 3S_1 or 3P_0 with $|S, S_z\rangle = |1, -\lambda\rangle$. With $\phi_0 = (u\overline{u} + d\overline{d} + s\overline{s})/\sqrt{3}$ the flavour overlap is $1/\sqrt{3}$ for $b_1\pi$ and $1/\sqrt{6}$ for $f_1(n\overline{n})\pi$. The remaining algebra is tedious but straightforward giving, apart from common numerical factors,

$$\langle b_1 \pi | \sigma \cdot \nabla | \pi_1 \rangle = +2\sqrt{1/6}(I_{++} - I_{0+} + I_{+0}) \tag{11}$$

$$\langle f_1 \pi | \sigma \cdot \nabla | \pi_1 \rangle = -\sqrt{1/6} (I_{++} - I_{0+} + I_{+0})$$
 (12)

Crucial to the above result is the spin wavefunction of the emergent $q\bar{q}$ pair. For the $f_1\pi$ mode, the Clebsch-Gordan factor $\langle 1 \ M_L^A + \lambda, 1 \ M_S^A - \lambda | 1 \ M_S^A + M_L^A \rangle$ plays a role and brings $\pm \sqrt{1/2}$. This factor changes sign under $(M_L^A, M_S^A, \lambda) \rightarrow (-M_L^A, -M_S^A, -\lambda)$, matched by the changing sign of the spin overlap term $\langle \chi_1^{M_S^A - \lambda} \chi_0^0 | \chi_1^{M_S^A} \chi_1^{-\lambda} \rangle$ under the same operation. These compensating signs yield the same linear combination of I_{++}, I_{0+}, I_{+0} , scaled by $-\sqrt{1/2}$. The end result is that in spin triplet creation models

$$\frac{\Gamma(\pi_1 \to b_1 \pi)}{\Gamma(\pi_1 \to f_1 \pi)} = 4 \tag{13}$$

(apart from small phase space and k dependent corrections.) We stress that this result is independent of the spatial overlaps, and as such is independent of the detailed forms of the quark and string wavefunctions. Furthermore, the result is characteristic of both ${}^{3}S_{1}$ and ${}^{3}P_{0}$ models, being driven by the same angular momentum algebra, and depends

crucially on the spin 1 nature of the emergent $q\bar{q}$ pair. The same linear combination drives all of the S-wave decays of hybrids with negative parity to $1P + S_0$ modes, and the relevant recoupling coefficients can be read from Table 1 of [6], combined with the appropriate flavour overlaps.

Conclusions

These results show that near threshold lattice QCD and flux tube models are in excellent agreement. It is possible to compare quite directly the flux tube model with the underlying QCD, in part because one of the main uncertainties in any calculation of this type (phase space) has been removed. The standard quark model parameters give excellent agreement with the lattice results, and the encouraging agreement with the scale of hybrid decays supports the physical picture of the string wavefunction. For physical widths, momentum dependent form factors are crucial and we suggest that the widths of ref [7] are overestimates.

Any improvement of the uncertainties in the lattice ratio of $b_1\pi$: $f_1\pi$, or the calculation of any analogous ratios, would be a welcome advance. This will allow the decay mechanism to be probed rather directly: more general decay models, such as those trigerred by the emission of a single vector gluon with the possibility of spin flips, do not result automatically in the ratio of eq. (13). Finally we suggest that hybrid decays of the type $0^{+-}, 2^{+-} \rightarrow \rho\rho$ may be worth investigating on the lattice: these modes should not require drastic extrapolation to the physical regime, and are predicted to vanish exactly in several variants of quark models.

We acknowedge discussions with J Dudek. This work is supported, in part, by grants from the Particle Physics and Astronomy Research Council, the Oxford University Clarendon Fund and the EU-TMR program "Eurodice", HPRN-CT-2002-00311.

- C.J.Morningstar and M.Peardon, Phys. Rev. D60 (1999) 034509
- [2] P.Lacock, C. Michael, P.Boyle and P.Rowland, Phys. Lett. B401 (1997) 308
- [3] N. Isgur and J. Paton, Phys. Rev. D31 (1985) 2910
- [4] T. Barnes, F.E.Close, and E.S.Swanson, Phys. Rev. D52 (1995) 5242
- [5] N.Isgur, R.Kokoski and J.Paton, Phys.Rev.Lett. 54 (1985) 869
- [6] F.E.Close and P.R.Page, Nucl. Phys. B443 (1995) 233
- [7] C.Michael and C.McNeile, hep-lat/0603007
- [8] Particle Data Group, Phys. Lett. B592 (2004)
- [9] R. Kokoski and N. Isgur, Phys. Rev. D35 (1987) 907
- [10] P.R.Page, E.Swanson and A.P. Szczepaniak, Phys. Rev.D59 (1999) 034016
- [11] F.Iddir et al., Phys. Lett.B207 (1985) 325;
 P.R.Page, Phys. Lett. B402 (1997) 183
- [12] F.E.Close and E.S.Swanson, Phys. Rev. D72 (2005) 094004
- [13] T Barnes, F.E.Close, P.R.Page and E.S. Swanson, Phys. Rev. D55 (1997) 4157
- [14] F.E.Close and A.Kirk, Z.Phys. C76 (1997) 469
- [15] F.E.Close and N.Isgur, Phys. Lett. B509 (2001) 81
- [16] P. Geiger and E.S.Swanson, Phys. Rev. D50 (1994) 6855
- [17] N.Dowrick, J.Paton and S. Perantonis, J.Phys.G13 (1987) 423
- [18] T.J.Burns, in preparation

^[1] G.Bali et al., (UKQCD) Phys. Lett. B309 (1993) 378;