## Derivation of Covariant Dissipative Fluid Dynamics in the Renormalization-group Method

K. Tsumura

Department of Physics, Kyoto University, Kyoto 606-8502, Japan

T. Kunihiro and K. Ohnishi

Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

(Dated: September 16, 2018)

We derive generic relativistic hydrodynamical equations with dissipative effects from the underlying Boltzmann equation in a mechanical and systematic way on the basis of so called the renormalization-group method. A macroscopic frame vector is introduced to specify the frame on which the macroscopic dynamics is described. Our method is so mechanical with only few ansatz that our method give a microscopic foundation of the available hydrodynamical equations, and also can be applied to make a reduction of the kinetic equations other than the simple Boltzmann equation.

PACS numbers: 05.10.Cc,05.20.Dd,25.75.-q,47.75.+f

The relativistic hydrodynamical equation is widely used in various fields of physics, especially in high-energy nuclear physics and astrophysics [1, 2, 3], and it seems that the study of the *dissipative* hydrodynamical equations is now becoming a central interest in these fields [3, 4, 5, 6, 7, 8, 9, 10].

It is, however, noteworthy that we have not necessarily reached a full understanding of the theory of relativistic hydrodynamics for viscous fluids: (1) It is well known that the relativistic hydrodynamical equation with dissipation has a form depending on the choice of the Lorentz frame or the definition of the hydrodynamical flow. The typical frame is that on the particle or the energy flow, and the typical corresponding equations are the ones of Eckart [11] and Landau [12], which were derived phenomenologically on the basis of the number and energy-momentum conservation laws, the second law of thermodynamics and some specific assumptions on the choice of the flow. One might consider that the difference of the equations should be merely a kind of choice of coordinate to describe dynamics in a easy and practical way, and one might be able to go back and forth between the two reference frames, by a Lorentz transformation. However, it may not be the case. To be more specific, let  $\delta T^{\mu\nu}$  and  $\delta N^{\mu}$ , be the dissipative part of the energy-momentum tensor and the particle current, respectively. The point is that they are not determined uniquely only by the second law of thermodynamics without some physical constraints involving the four flow-vector  $u_{\mu}$  with  $u_{\mu}u^{\mu} = 1$ : The constraints of Eckart are expressed as [13], (i)  $u_{\mu}u_{\nu}\delta T^{\mu\nu} = 0$ , (ii)  $u_{\mu}\delta N^{\mu} = 0$ , (iii)  $\Delta_{\mu\nu}\delta N^{\nu} = 0$ , while those of Landau are (i)  $u_{\mu}u_{\nu}\delta T^{\mu\nu} = 0$ , (ii)  $u_{\mu}\delta N^{\mu} = 0$  and (iv)  $u_{\mu}\Delta_{\nu\rho}\delta T^{\mu\nu} = 0$ , where  $\Delta_{\mu\nu} = g_{\mu\nu} - u_{\mu}u_{\nu}$ . As one sees, the first two constraints are the same for the two theories, and the respective third condition specifies the frame of the coordinate. It is here noteworthy that there is a proposal by Stewart [14] for the constraints for the Eckart frame, i.e., the particle-flow frame, as given by (v)  $\delta T^{\mu}_{\ \mu} = 0$ , (ii)  $u_{\mu}\delta N^{\mu} = 0$ , (iii)  $\Delta_{\mu\nu}\delta N^{\nu} = 0$ , where the condition (i) of Eckart is replaced by a different one (v). One may ask if both the Eckart and Stewart constraints make sense or not. It has been scarcely examined [15] whether the constraints proposed phenomenologically hold internal consistency or match the underlying microscopic kinetic theories. (2) It is also known that these equations containing the time derivative only in the first order, unfortunately, suffer from the unphysical instability [13, 16]; the instability is attributed to the lack of the causality, and invented is a fluid dynamical equation which contains the second-order time-derivative as well as the first-order one [11, 12, 17]. The new equation called Israel-Stewart equation [20] contains new transport coefficients like the relaxation time yet to be determined microscopically or phenomenologically [14, 18, 19, 20].

There are some attempts to derive the phenomenological equations from the microscopic equation, i.e., the relativistic Boltzmann equation, with use of the Chapman-Enskog method [21] and the Grad's fourteen-moment method [22], and hence give a microscopic foundation to them [14, 18, 19, 20, 23]. We notice that such a microscopic approach has another merit that the hydrodynamical equations can be obtained, which are consistent with the underlying kinetic equation, because the theory gives the theory connect the kinetic and the hydrodynamical regimes and hence gives the natural initial condition for the hydrodynamical equation. This point should have a practical merit when one analyzes the freeze-out regime and also the hadron corona in the expanding system such as the intermediate stage of the relativistic heavy-ion collision [7, 9, 24].

One must say, however, that the past works in the microscopic approaches are not fully satisfactory in giving the foundation to the phenomenological equations: **a.** Although the past works succeeded in identifying the assumptions and/or approximations to reproduce the known hydrodynamical equations by Eckart, Landau and Israel, one must say that the physical meaning of these assumptions/approximations is somewhat obscure, and so is the uniqueness

of those hydrodynamical equations as the long-wavelength and low-frequency limit of the microscopic theory. **b.** It is also to be noted that the Chapman-Enskog method and the Grad's fourteen-moment method themselves have some ad-hoc part and may not be fully mechanical and systematic methods for the reduction of dynamics.

The purpose of this article is to derive the relativistic hydrodynamical equations with dissipative effects from the microscopic theory in a more natural and systematic way, and thereby establish the microscopic foundation of them. We shall show that when the particle-flow frame (Eckart frame) is chosen, the constraints by Stewart but not Eckart must be taken, while the resulting hydrodynamical equation which manifestly satisfies the second law of thermodynamics is different form that given by Stewart.

The problem is how to reduce the dynamics to a slower dynamics using a reliable reduction theory of the dynamics [25], which should be as mechanical as possible. In fact, van Kampen [23] applied his method [26] of the reduction to derive the relativistic hydrodynamic equations for the viscous fluid. However, his method is not formulated in a covariant way and hence failed in reproducing the phenomenological hydrodynamical equations with dissipation. We shall apply another powerful reduction theory known as the "renormalization-group(RG) method" [27, 28, 29, 30, 31, 32, 33] to reduce the relativistic Boltzmann equation to the hydrodynamical equations. Our theory is manifestly covariant and relies on almost no assumptions.

The RG method is a systematic reduction theory of the dynamics leading to the coarse-graining of temporal and spatial scales. This method is suitable for our present purpose to elucidate the physical meanings of each process of the reduction since the method is so mechanical and does not require any specific assumptions. Indeed, the RG method is already applied satisfactorily to the derivation of the Navier-Stokes equation from the (non-relativistic) Boltzmann equation [32, 33]. The present work is an attempt to extend these previous works to the relativistic case.

In this article, we shall treat, as a simplest example, the classical but relativistic system composed of identical particles; an extension to multi-component systems is possible and will be discussed in a separate paper [34]. Then the relativistic Boltzmann equation [18] describing such a system reads

$$p^{\mu} \partial_{\mu} f_p(x) = C[f]_p(x), \tag{1}$$

where  $f_p(x)$  denotes the one-particle distribution function defined in the phase space  $(x^{\mu}, p^{\mu})$  with  $p^{\mu}$  being the four momentum of the on-shell particle;  $p^{\mu} p_{\mu} \equiv p^2 = m^2$  and  $p^0 \ge 0$ . The right-hand side of Eq.(1) is called the collision integral,  $C[f]_p(x) \equiv \frac{1}{2!} \sum_{p_2} \frac{1}{p_2^0} \sum_{p_3} \frac{1}{p_3^0} \omega(p, p_1 | p_2, p_3) (f_{p_2}(x) f_{p_3}(x) - f_p(x) f_{p_1}(x))$ , where  $\omega(p, p_1 | p_2, p_3)$  denotes the transition probability owing to the microscopic two-particle interaction and contains the delta functions  $\delta^4(p + p_1 - p_2 - p_3)$  representing the energy-momentum conservation. The transition probability has the symmetric properties due to the indistinguishability of the particles and the time reversal invariance in the scattering process:  $\omega(p, p_1 | p_2, p_3) = \omega(p_2, p_3 | p, p_1) = \omega(p_1, p | p_3, p_2) = \omega(p_3, p_2 | p_1, p)$ . It should be stressed here that we have confined ourselves to the case in which the particle number is conserved in the collision process.

The particle-number and energy-momentum conservation in the collision process ensures the following important relations;  $\sum_{p} \frac{1}{p^0} C[f]_p(x) = \sum_{p} \frac{1}{p^0} p^{\mu} C[f]_p(x) = 0$ , which says that the five quantities  $(1, p^{\mu})$  are *collision invariants*. We have the continuity or balance equations for the particle current and the energy-momentum tensor,

$$\partial_{\mu} \sum_{p} \frac{1}{p^{0}} p^{\mu} f_{p}(x) \equiv \partial_{\mu} N^{\mu} = 0, \quad \partial_{\nu} \sum_{p} \frac{1}{p^{0}} p^{\mu} p^{\nu} f_{p}(x) \equiv \partial_{\nu} T^{\mu\nu} = 0, \tag{2}$$

respectively. Notice that while these equations have the same forms as the hydrodynamical equations, nothing about the dynamical properties is contained in these equations before the evolution of the distribution function  $f_p(x)$  is obtained by solving the microscopic Eq.(1).

The entropy current may be defined by  $S^{\mu} \equiv -\sum_{p} \frac{1}{p^{0}} p^{\mu} f_{p}(x) (\ln f_{p}(x) - 1)$ , which is conserved only if  $\ln f_{p}(x)$  is a collision invariant, or a linear combination of the basic collision invariants  $(1, p^{\mu})$  as  $\ln f_{p}(x) = \alpha(x) + p^{\mu} \beta_{\mu}(x)$  with  $\alpha(x)$  and  $\beta_{\mu}(x)$  being arbitrary functions.

Since we are interested in the hydrodynamical regime where the time- and space-dependence of the physical quantities are small, we solve Eq.(1) under the situation where the time variation of  $f_p(x)$  is slow with long wavelengths.

To retain the Lorentz covariance, and achieve a coarse-graining, it is found convenient to introduce a macroscopic Lorentz vector  $a_p^{\mu}$  characterizing the flow, which specifies the covariant coordinate system (frame) and we call the *macroscopic-frame vector*. Although  $a_p^{\mu}$  may depend on the momentum p and the space-time coordinate x, the time variation of it is supposed to be much smaller than that of the microscopic processes. Our point is that the separation of the scales can be nicely achieved by the RG method.

Keeping in mind the above scale difference, we define the covariant coordinate system  $(\tau, \sigma^{\mu})$  as  $d\tau \equiv a_p^{\mu}(x) dx_{\mu}$  and  $\varepsilon^{-1} d\sigma^{\mu} \equiv (g^{\mu\nu} - a_p^{\mu}(x) a_p^{\nu}(x)/a_p^2(x)) dx_{\nu} \equiv \Delta_p^{\mu\nu}(x) dx_{\nu}$ . We notice that the small quantity  $\varepsilon$  have been introduced to tag that the space derivatives are small for the system we are interested in; the  $\varepsilon$  may be identified with the ratio

of the average particle distance over the mean free path. In this coordinate system, Eq.(1) reads

$$\frac{\partial}{\partial \tau} f_p(\tau, \sigma) = \frac{1}{p \cdot \boldsymbol{a}_p(\tau, \sigma)} C[f]_p(\tau, \sigma) - \varepsilon \frac{1}{p \cdot \boldsymbol{a}_p(\tau, \sigma)} p \cdot \boldsymbol{\nabla} f_p(\tau, \sigma),$$
(3)

where  $a_p^{\mu}(\tau, \sigma) \equiv a_p^{\mu}(x)$ ,  $\Delta_p^{\mu\nu}(\tau, \sigma) \equiv \Delta_p^{\mu\nu}(x)$  and  $f_p(\tau, \sigma) \equiv f_p(x)$ . Since  $\varepsilon$  appears in front of  $\nabla^{\mu} \equiv \Delta_p^{\mu\nu}(\tau, \sigma) \frac{\partial}{\partial \sigma^{\nu}}$ , Eq.(3) has a form to which the perturbative expansion can be possible. In fact, a natural but significant assumption underlies this seemingly mere rewrite of the equation that only the spatial inhomogeneity over distances of the order of the mean free path is the origin of the dissipation. We notice that the RG method applied to non-relativistic Boltzmann equation with the corresponding assumption nicely leads to the Navier-Stokes equation [32, 33]; the present approach is simply a covariantization of the non-relativistic case.

Some remarks on  $a_p^{\mu}(\tau, \sigma)$  are in order: (i) In the perturbative expansion, we shall take the coordinate system where  $a_p^{\mu}(\tau, \sigma)$  has no  $\tau$  dependence, i.e.,  $a_p^{\mu}(\tau, \sigma) = a_p^{\mu}(\sigma)$ . (ii) In our coordinate system, the physical quantity transported per unit  $\tau$  is given by  $p \cdot a_p(\sigma)$ , we can control what is the flow represented in our theory by varying the specific expression of  $a_p^{\mu}(\sigma)$ . Owing to this freedom inherent in our coordinate system our theory may lead to various hydrodynamical equations, including the ones in the Eckart and Landau frames for non-ideal fluids.

In accordance with the general formulation of the RG method given in [28, 29, 31], we first try to obtain the perturbative solution  $\tilde{f}_p$  around the arbitrary initial time  $\tau = \tau_0$  with the initial value  $f_p(\tau_0, \sigma)$ ;  $\tilde{f}_p(\tau = \tau_0, \sigma; \tau_0) = f_p(\tau_0, \sigma)$ , where we have made explicit that the solution have the  $\tau_0$  dependence. The initial value is not yet specified, we suppose that the initial value is on an exact solution with  $\tau_0$  being varied. The initial value as well as the solution are expanded with respect to  $\varepsilon$  as follows;  $\tilde{f}_p(\tau, \sigma; \tau_0) = \tilde{f}_p^{(0)}(\tau, \sigma; \tau_0) + \varepsilon \tilde{f}_p^{(1)}(\tau, \sigma; \tau_0) + \varepsilon^2 \tilde{f}_p^{(2)}(\tau, \sigma; \tau_0) + \cdots$ , and  $f_p(\tau_0, \sigma) = f_p^{(0)}(\tau_0, \sigma) + \varepsilon f_p^{(1)}(\tau_0, \sigma) + \varepsilon^2 f_p^{(2)}(\tau_0, \sigma) + \cdots$ . The respective initial conditions at  $\tau = \tau_0$  are set up as  $\tilde{f}_p^{(l)}(\tau_0, \sigma; \tau_0) = f_p^{(l)}(\tau_0, \sigma)$  for  $l = 0, 1, 2, \cdots$ . In the expansion, the zeroth-order value  $\tilde{f}_p^{(0)}(\tau_0, \sigma; \tau_0) = f_p^{(0)}(\tau_0, \sigma)$  is supposed to be as close as possible to an exact solution.

Substituting the above expansions into Eq.(3) in the  $\tau$ -independent but  $\tau_0$ -dependent coordinate system with  $a_p^{\mu}(\tau, \sigma) \rightarrow a_p^{\mu}(\tau_0, \sigma) \equiv a_p^{\mu}(\sigma; \tau_0)$ , we obtain the series of the perturbative equations with respect to  $\varepsilon$ . The zeroth-order equation reads

$$\frac{\partial}{\partial \tau} \tilde{f}_p^{(0)} = \frac{1}{p \cdot \boldsymbol{a}_p} C[\tilde{f}^{(0)}]_p.$$
(4)

Since we are interested in the slow motion which may be realized asymptotically when  $\tau \to \infty$ , we should take the following stationary solution or the fixed point,  $\frac{\partial}{\partial \tau} \tilde{f}_p^{(0)} = 0$ , which is realized when  $C[\tilde{f}^{(0)}]_p = 0$  for arbitrary  $\sigma$ . This implies that  $\ln \tilde{f}_p^{(0)}$  is a linear combination of the five collision invariants  $(1, p^{\mu})$ , and hence  $\tilde{f}_p^{(0)}$  is found to be a local equilibrium distribution function or the Juettner function (the relativistic analog of the Maxwellian):  $\tilde{f}_p^{(0)}(\tau, \sigma; \tau_0) = (2\pi)^{-3} \exp\left[\left(\mu(\sigma; \tau_0) - p^{\mu} u_{\mu}(\sigma; \tau_0)\right) / T(\sigma; \tau_0)\right] \equiv f_p^{eq}(\sigma; \tau_0)$  with  $u^{\mu}(\sigma; \tau_0) u_{\mu}(\sigma; \tau_0) = 1$ , which implies that  $\tilde{f}_p^{(0)}(\tau, \sigma; \tau_0) = \tilde{f}_p^{(0)}(\tau_0, \sigma; \tau_0) = f_p^{(0)}(\tau_0, \sigma) = f_p^{eq}(\sigma; \tau_0)$ . It should be noticed that the five would-be integration constants  $T(\sigma; \tau_0), \mu(\sigma; \tau_0)$  and  $u_{\mu}(\sigma; \tau_0)$  is independent of  $\tau$  but may depend on  $\tau_0$  as well as  $\sigma$ .

Next the first-order equation reads

$$\frac{\partial}{\partial \tau} \tilde{f}_p^{(1)} = \sum_q A_{pq} \, \tilde{f}_q^{(1)} + F_p, \tag{5}$$

where the linear evolution operator A and the inhomogeneous term F are defined as  $A_{pq} \equiv (p \cdot a_p)^{-1} \frac{\partial}{\partial f_q} C[f]_p |_{f=f^{eq}}$ and  $F_p \equiv -(p \cdot a_p)^{-1} p \cdot \nabla f_p^{eq}$ , respectively. To obtain the solution which describes the slow motion, it is convenient to first analyze the properties of A, especially its spectra. For this purpose, we convert A to another linear evolution operator  $L \equiv f^{eq-1} A f^{eq}$  with the diagonal matrix  $f_{pq}^{eq} \equiv f_p^{eq} \delta_{pq}$ . Let us define the inner product between arbitrary vectors  $\varphi$  and  $\psi$  by

$$\langle \varphi, \psi \rangle \equiv \sum_{p} \frac{1}{p^{0}} \left( p \cdot \boldsymbol{a}_{p} \right) f_{p}^{\text{eq}} \varphi_{p} \psi_{p}.$$

$$\tag{6}$$

With this inner product, one can nicely show that L becomes self-adjoint  $\langle \varphi, L \psi \rangle = \langle L \varphi, \psi \rangle$  and non-positive definite  $\langle \varphi, L \varphi \rangle \leq 0$ , which means that the eigen values of L are zero or negative. The eigen vectors of the zero eigen value are  $\varphi_{0p}^{\alpha} = p^{\mu}$  for  $\alpha = \mu = 0 \sim 3$  and  $\varphi_{0p}^{4} = m$ , which span the kernel of L and satisfy  $L \varphi_{0}^{\alpha} = 0$ . Notice that these five vectors are the collision invariants. Following [31], we define the projection operator P onto the kernel

of L which is called the P space and the projection operator Q onto the Q space complement to the P space:

$$\left[P\,\psi\right]_{p} \equiv \sum_{\alpha\beta} \varphi^{\alpha}_{0p} \eta^{-1}_{\alpha\beta} \langle \varphi^{\beta}_{0}, \psi \rangle, \tag{7}$$

with  $Q \equiv 1 - P$  where  $\eta_{\alpha\beta}^{-1}$  is the inverse matrix of  $\eta^{\alpha\beta} \equiv \langle \varphi_0^{\alpha}, \varphi_0^{\beta} \rangle$ . In the following, we shall suppress the argument  $\sigma$  and the subscript p when no misunderstanding is expected. The solution to Eq.(5) with the initial condition  $\tilde{f}^{(1)}(\tau = \tau_0; \tau_0) = f^{(1)}(\tau_0)$  is found to be  $\tilde{f}^{(1)}(\tau; \tau_0) = e^{(\tau - \tau_0)A} \{ f^{(1)}(\tau_0) + e^{(\tau - \tau_0)A} \}$  $A^{-1}\bar{Q}F$  +  $(\tau - \tau_0)\bar{P}F - A^{-1}\bar{Q}F$ , where the modified projection operators  $\bar{P} \equiv f^{\text{eq}}Pf^{\text{eq-1}}$  and  $\bar{Q} \equiv f^{\text{eq}}Qf^{\text{eq-1}}$  have been introduced. Notice that the first term would be a fast motion coming from the Q space, which we can nicely eliminate by choosing the initial value  $f^{(1)}(\tau_0)$ , which has not yet been specified; thus we have the first-order solution

$$\tilde{f}^{(1)}(\tau;\tau_0) = (\tau - \tau_0) \,\bar{P} \,F - A^{-1} \,\bar{Q} \,F \tag{8}$$

with the initial value determined as  $\tilde{f}^{(1)}(\tau_0; \tau_0) = f^{(1)}(\tau_0) = -A^{-1}\bar{Q}F$ . We notice the appearance of the secular term proportional to  $\tau - \tau_0$ , which apparently invalidates the perturbative solution when  $|\tau - \tau_0|$  becomes large.

The second-order equation [36] is written as

$$\frac{\partial}{\partial \tau}\tilde{f}^{(2)} = A\,\tilde{f}^{(2)} + (\tau - \tau_0)\,H + I,\tag{9}$$

with  $H_p \equiv -(p \cdot \boldsymbol{a}_p)^{-1} p \cdot \nabla [\bar{P} F]_p$  and  $I_p \equiv (p \cdot \boldsymbol{a}_p)^{-1} p \cdot \nabla [A^{-1} \bar{Q} F]_p$ , the solution to which is found to be

$$\tilde{f}^{(2)}(\tau;\tau_0) = e^{(\tau-\tau_0)A} \left\{ f^{(2)}(\tau_0) + A^{-2} \bar{Q} H + A^{-1} \bar{Q} I \right\} + \frac{1}{2} (\tau-\tau_0)^2 \bar{P} F + (\tau-\tau_0) \left\{ \bar{P} I - A^{-1} \bar{Q} H \right\} - \left\{ A^{-2} \bar{Q} H + A^{-1} \bar{Q} I \right\}.$$
(10)

Again the would-be fast motion can be eliminated by the choice of the initial value  $f^{(2)}(\tau_0)$ , and thus we have the second-order solution

$$\tilde{f}^{(2)}(\tau;\tau_0) = \frac{1}{2} (\tau - \tau_0)^2 \,\bar{P} \,F + (\tau - \tau_0) \left\{ \bar{P} \,I - A^{-1} \,\bar{Q} \,H \right\} - \left\{ A^{-2} \,\bar{Q} \,H + A^{-1} \,\bar{Q} \,I \right\},\tag{11}$$

with the initial value determined as  $\tilde{f}^{(2)}(\tau_0;\tau_0) = f^{(2)}(\tau_0) = -A^{-2}\bar{Q}H - A^{-1}\bar{Q}I$ . We notice again the appearance of the secular terms.

Summing up the perturbative solutions up to the second order, we have an approximate solution around  $\tau \simeq \tau_0$  to this order;  $\tilde{f}(\tau; \tau_0) = \tilde{f}^{(0)}(\tau; \tau_0) + \varepsilon \tilde{f}^{(1)}(\tau; \tau_0) + \varepsilon^2 \tilde{f}^{(2)}(\tau; \tau_0) + O(\varepsilon^3)$ , containing the secular terms which apparently invalidates the perturbative expansion for  $\tau$  away from the initial time  $\tau_0$ .

The point of the RG method lies in the fact that we can utilize the secular terms to obtain a solution valid in a global domain. Now we may see that we have a family of curves  $f(\tau; \tau_0)$  parameterized with  $\tau_0$ . They are all on the exact solution  $f(\tau)$  at  $\tau = \tau_0$  up to  $O(\varepsilon^3)$ , but only valid locally for  $\tau$  near  $\tau_0$ . So it is conceivable that the envelope E of the family of curves which contacts with each local solution at  $\tau = \tau_0$  will give a global solution in our asymptotic situation. According to the classical theory of envelopes, the envelope which contact with any curve in the family at  $\tau = \tau_0$  is obtained by  $\frac{\mathrm{d}}{\mathrm{d}\tau_0} \tilde{f}_p(\tau, \sigma; \tau_0) \Big|_{\tau_0 = \tau} = 0$ , or explicitly

$$\frac{\partial}{\partial \tau} \left\{ f_p^{\rm eq} - \varepsilon \left[ A^{-1} \bar{Q} F \right]_p \right\} - \varepsilon \left[ \bar{P} F \right]_p - \varepsilon^2 \left\{ \left[ \bar{P} I \right]_p - \left[ A^{-1} \bar{Q} H \right]_p \right\} + O(\varepsilon^3) = 0.$$
(12)

This envelope equation is the basic equation in the RG method and gives the equation of motion governing the dynamics of the five slow variables  $T(\tau)$ ,  $\mu(\tau)$  and  $u^{\mu}(\tau)$  in  $f_{p}^{eq}$ . The global solution in the asymptotic region is given as the envelope function,

$$\tilde{f}_p(\tau,\sigma;\tau_0=\tau) \equiv f_p(\tau,\sigma) = f_p^{\rm eq} - \varepsilon \left[A^{-1}\bar{Q}F\right]_p - \varepsilon^2 \left\{ \left[A^{-2}\bar{Q}H\right]_p + \left[A^{-1}\bar{Q}I\right]_p \right\} + O(\varepsilon^3),\tag{13}$$

where the exact solution of Eq.(12) is inserted. Thus one sees that  $f_p(\tau)$  now describes the macroscopic-time evolution of the one-particle distribution function in Eq.(1), because the time-derivatives of the quantities in  $f_p(\tau)$  are all in the order of  $\varepsilon$ . We emphasize that we have derived the infrared effective theory of Eq.(1) in the form of the pair of Eq.'s (12) and (13). This is one of the main results in this article.

Now the RG/Envelop equation Eq.(12) is actually the hydrodynamic equation governing the the five slow variables  $T(\tau)$ ,  $\mu(\tau)$  and  $u^{\mu}(\tau)$ . To show this explicitly, we apply  $\bar{P}$  from the left and then take the inner product with the five zero modes  $\varphi_0^{\alpha}[37]$ . Putting back  $\varepsilon = 1$ , we arrive at  $\partial_{\mu} J_{\text{hydro}}^{\mu\alpha} = 0$  with

$$J_{\text{hydro}}^{\mu\alpha} \equiv \sum_{p} \frac{1}{p^{0}} p^{\mu} \varphi_{0p}^{\alpha} \left\{ f_{p}^{\text{eq}} - \left[ A^{-1} \bar{Q} F \right]_{p} \right\} = J_{0}^{\mu\alpha} + \delta J^{\mu\alpha}, \qquad (14)$$

where  $J_0^{\mu\alpha} \equiv \sum_p \frac{1}{p^0} p^{\mu} \varphi_{0p}^{\alpha} f_p^{\text{eq}}$  and  $\delta J^{\mu\alpha} \equiv -\sum_p \frac{1}{p^0} p^{\mu} \varphi_{0p}^{\alpha} \left[ A^{-1} \bar{Q} F \right]_p$ . Here  $J_0^{\mu\alpha}$  represents the currents of perfect-fluid part, while  $\delta J^{\mu\alpha}$  the dissipative part. In deriving Eq.(14), we have used the following relation :  $\sum_p \frac{1}{p^0} \left[ p \cdot a_p \right] \varphi_{0p}^{\alpha} \left[ \bar{P} \psi \right]_p = \sum_p \frac{1}{p^0} \left( p \cdot a_p \right) \varphi_{0p}^{\alpha} \psi_p$ , obtained from the definition  $\bar{P} = f^{\text{eq}} P f^{\text{eq}-1}$  and Eq.(7). Notice that the derivation of Eq.(14) is accomplished for arbitrary  $a_p^{\mu}$  which is now dependent on  $\tau$  as well as on  $\sigma$ ; in other words, we have a set of relativistic hydrodynamic equations for viscous fluids, which have still the freedom of choice of  $a_p^{\mu}$ . The is also one of the main results in the present work.

In the rest of the paper, we shall show how known relativistic dissipative hydrodynamic equations are derived with a choice of the macroscopic frame vector  $a_p^{\mu}$ .

As a simple but nontrivial choice, let us take the following set of the macroscopic frame vectors with  $\theta$  being a constant;  $a_p^{\mu} = (p \cdot u)^{-1} \{(p \cdot u) \cos \theta + m \sin \theta\} u^{\mu} \equiv \theta_p^{\mu}$ , which leads to the following relations  $\Delta_p^{\mu\nu} = g^{\mu\nu} - u^{\mu} u^{\nu} \equiv \Delta^{\mu\nu}$ ; then one finds that  $(p \cdot u)^{-1} \{(p \cdot u) \cos \theta + m \sin \theta\} \frac{\partial}{\partial \tau} = u^{\mu} \partial_{\mu} \equiv D$  and  $\nabla^{\mu} = \Delta^{\mu\nu} \partial_{\nu} \equiv \nabla^{\mu}$ . We notice that the factor m is introduced simply to make the expression dimensionless, so our method is also applicable to the case of massless particles.

In the reduction of the currents in terms of the fluid dynamical quantities, the central task is the evaluation of  $A^{-1}\bar{Q}F$ . A straightforward but tedious manipulation leads to the following formula,

$$\left[A^{-1}\bar{Q}F\right]_{p} = f_{p}^{\text{eq}}\sum_{q} \mathcal{L}_{pq}^{-1}\frac{1}{T}\left(\Pi_{q}X_{\theta} + Q_{q}^{\mu}X_{\theta\mu} + \Pi_{q}^{\mu\nu}X_{\mu\nu}\right),\tag{15}$$

where  $\mathcal{L}_{pq} \equiv (p \cdot \theta_p) L_{pq}$  being independent of  $\theta_p^{\mu}$ . Here we have introduced the following quantities: the dissipative currents, i.e., the bulk pressure,  $\Pi_p \equiv (4/3 - \gamma) (p \cdot u)^2 + [(\gamma - 1) T \hat{h} - \gamma T] (p \cdot u) - 1/3 m^2$ , the heat flow,  $Q_p^{\mu} \equiv -[(p \cdot u) - T \hat{h}] \Delta^{\mu\nu} p_{\nu}$  and the stress tensor,  $\Pi_p^{\mu\nu} \equiv 1/2 (\Delta^{\mu\rho} \Delta^{\nu\sigma} + \Delta^{\mu\sigma} \Delta^{\nu\rho} - 2/3 \Delta^{\mu\nu} \Delta^{\rho\sigma}) p_{\rho} p_{\sigma}$ . The coefficient of each dissipative current is the respective dissipative thermodynamic force, i.e.,  $X_{\theta} \equiv (-z^2 \cos 2\theta) \nabla \cdot u/\{z^2 \cos^2 \theta + z^2 (3\gamma - 4) \sin^2 \theta - 3z [1 - (\hat{h} - 1) (\gamma - 1)] \cos \theta \sin \theta\}$ ,  $X_{\theta\mu} \equiv \nabla_{\mu} \ln T - \cos \theta \nabla_{\mu} \ln(nT)/(\hat{h} \cos \theta + z \sin \theta)$ , and  $X_{\mu\nu} \equiv 1/2 (\Delta_{\mu\rho} \Delta_{\nu\sigma} + \Delta_{\mu\sigma} \Delta_{\nu\rho} - 2/3 \Delta_{\mu\nu} \Delta_{\rho\sigma}) \nabla^{\rho} u^{\sigma}$ , where the following definitions have been used; the reduced mass  $z \equiv \frac{m}{T}$ , the particle density  $n \equiv (2\pi)^{-3} 4\pi m^3 e^{\frac{\mu}{T}} \{z^{-1} K_2(z)\}$ , the energy density  $\epsilon \equiv m n \{K_3(z)/K_2(z) - z^{-1}\}$ , the reduced enthalpy per particle  $\hat{h} \equiv \frac{\epsilon + nT}{nT}$  and the ratio of the heat capacities  $\gamma \equiv 1 + \{z^2 + 3\hat{h} - (\hat{h} - 1)^2\}^{-1}$ , with the modified Bessel functions  $K_2(z)$  and  $K_3(z)$ .

Since the dissipative part of the currents has been obtained for arbitrary  $\theta$ , we can obtain the microscopic formulae for the transport coefficients, i.e., the bulk viscosity  $\zeta$ , the heat conductivity  $\lambda$  and the shear viscosity  $\eta$ , which are nicely given in forms of the Kubo formula as follows: Using the new inner product  $\langle \varphi, \psi \rangle_{eq} \equiv \sum_p \frac{1}{p^0} f_p^{eq} \varphi_p \psi_p$  and the evolved microscopic dissipative currents  $[\Pi(s)]_p \equiv \sum_q [e^{s\mathcal{L}}]_{pq} \Pi_q$  and so on, we have  $\zeta \equiv (1/T) \int_0^{\infty} ds \langle \Pi(0), \Pi(s) \rangle_{eq}$ ,  $\lambda \equiv -(1/3T^2) \int_0^{\infty} ds \langle Q^{\mu}(0), Q_{\mu}(s) \rangle_{eq}$  and  $\eta \equiv (1/10T) \int_0^{\infty} ds \langle \Pi^{\mu\nu}(0), \Pi_{\mu\nu}(s) \rangle_{eq}$ . It is noteworthy that these transport coefficients are independent of  $\theta$  while the dissipative thermodynamic forces are dependent on it.

Although we can now write down the generic form of the currents  $J^{\mu\alpha}_{\rm hydro}$  for arbitrary  $\theta$ , we shall give here them only for a few values of  $\theta$  for the sake of space, but show that our generic currents included the known currents for the relativistic fluid dynamics for a viscous fluid.

(A) With the simplest choice  $a_p^{\mu} = u^{\mu}$ , i.e.,  $\theta = 0$ , we have

$$J_{\text{hydro}}^{\mu\alpha} = \begin{cases} \epsilon \, u^{\mu} \, u^{\nu} - (p + \zeta \, X) \, \Delta^{\mu\nu} + 2 \, \eta \, X^{\mu\nu} & \alpha = \nu, \\ m \, n \, u^{\mu} - \frac{m}{\hat{h}} \, \lambda \, X^{\mu} & \alpha = 4, \end{cases}$$
(16)

where  $p \equiv nT$ ,  $X \equiv -\nabla \cdot u$  and  $X^{\mu} \equiv \nabla^{\mu} \ln T - \hat{h}^{-1} \nabla^{\mu} \ln(nT)$ . If we employ the constitutive equations;  $\Pi = \zeta X$ ,  $Q^{\mu} = T \lambda X^{\mu}$  and  $\Pi^{\mu\nu} = 2 \eta X^{\mu\nu} - \Pi \Delta^{\mu\nu}$ , it is possible to rewrite the equation in terms of the flows rather than the thermodynamic forces as

$$J_{\rm hydro}^{\mu\alpha} = \begin{cases} \epsilon \, u^{\mu} \, u^{\nu} - p \, \Delta^{\mu\nu} + \Pi^{\mu\nu} & \alpha = \nu, \\ m \, n \, u^{\mu} - \frac{m}{\hat{h}T} \, Q^{\mu} & \alpha = 4. \end{cases}$$
(17)

One should notice that  $J^{\mu\alpha}_{\rm hydro}$  in Eq.(16) or Eq.(17) agrees completely with the Landau theory for non-ideal fluids. This was actually anticipated: In fact, if we take a natural choice that  $\boldsymbol{a}_p^{\mu} = u^{\mu}$ , the physical quantity transported per unit $\tau$  becomes the microscopic thermal energy  $(p \cdot u)$ , which is reduced to the familiar form  $m + \frac{m}{2} |\frac{\boldsymbol{p}}{m} - \boldsymbol{u}|^2$  in the non-relativistic limit. Thus it is natural that the currents  $J^{\mu\alpha}_{\rm hydro}$  with the choice  $\boldsymbol{a}_p^{\mu} = u^{\mu}$  becomes those in the Landau frame.

(B) Another simple choice for the macroscopic frame vector is given with  $\theta = \pi/2$ ;  $a_p^{\mu} = \frac{m}{p \cdot u} u^{\mu}$ . A similar manipulation as the above gives the following currents

$$J_{\rm hydro}^{\mu\alpha} = \begin{cases} \left(\epsilon + 3\,\zeta\,\tilde{X}\right)u^{\mu}\,u^{\nu} - \left(p + \zeta\,\tilde{X}\right)\Delta^{\mu\nu} + \lambda\,T\,u^{\mu}\,\tilde{X}^{\nu} + \lambda\,T\,u^{\nu}\,\tilde{X}^{\mu} + 2\,\eta\,X^{\mu\nu} & \alpha = \nu, \\ m\,n\,u^{\mu} & \alpha = 4, \end{cases}$$
(18)

where  $\tilde{X} \equiv -\{1/3 (4/3 - \gamma)^{-1}\}^2 \nabla \cdot u$  and  $\tilde{X}^{\mu} \equiv \nabla^{\mu} \ln T$ . In terms of the flows, we have

$$J_{\rm hydro}^{\mu\alpha} = \begin{cases} \epsilon \, u^{\mu} \, u^{\nu} - p \, \Delta^{\mu\nu} + 3 \, \tilde{\Pi} \, u^{\mu} \, u^{\nu} + u^{\mu} \, \tilde{Q}^{\nu} + u^{\nu} \, \tilde{Q}^{\mu} + \tilde{\Pi}^{\mu\nu} & \alpha = \nu, \\ m \, n \, u^{\mu} & \alpha = 4, \end{cases}$$
(19)

where  $\tilde{\Pi} = \zeta \tilde{X}$ ,  $\tilde{Q}^{\mu} = \lambda T \tilde{X}^{\mu}$  and  $\tilde{\Pi}^{\mu\nu} = 2 \eta X^{\mu\nu} - \tilde{\Pi} \Delta^{\mu\nu}$ . Since the dissipative term with the heat conductivity is absent in  $J_{\text{hydro}}^{\mu4}$ , one might be tempted to identify the above equation with the Eckart one, but it is not the case: A detailed examination tells us that Eq.(18) is rather a modified or corrected version, we would say, of Stewart's[14]. This observation is based on the fact that Eq.(18) meets the Stewart's constraints, (v), (ii) and (iii) but not the Eckart's, (i), (ii) and (iii), mentioned in the Introduction. The reason why we call Eq.(18) the corrected version of Stewart equation will be shortly explained. A noteworthy point is the following: In the Eckart theory [38] used conventionally, the appearance of the terms with the bulk viscosity is the same as the Landau theory. In contrast, Eq.(18) shows the different appearance so that the bulk viscosity has an influence on the terms of not only the pressure p but also the energy density  $\epsilon$ . We can show that our equation (18) satisfies the second law of thermodynamics as will be explicitly shown in a separate paper [34, 35], although the original Stewart equation may not.

(C) The other interesting choice of  $\theta$  is  $\theta = -\pi/4$ , for which we have  $a_p^{\mu} = \frac{(p \cdot u) - m}{\sqrt{2} p \cdot u} u^{\mu}$ , and

$$J_{\rm hydro}^{\mu\alpha} = \begin{cases} \epsilon \, u^{\mu} \, u^{\nu} - p \, \Delta^{\mu\nu} - \lambda \, T \, u^{\mu} \, \bar{X}^{\nu} - \lambda \, T \, u^{\nu} \, \bar{X}^{\mu} + 2 \, \eta \, X^{\mu\nu} & \alpha = \nu, \\ m \, n \, u^{\mu} - \lambda \, T \, \bar{X}^{\mu} & \alpha = 4, \end{cases}$$
(20)

where  $\bar{X}^{\mu} \equiv z(\hat{h}-z)^{-1}\nabla^{\mu} \ln T - z(\hat{h}-z)^{-2}\nabla^{\mu} \ln(nT)$ . The alternative expression in terms of the flows is

$$J_{\rm hydro}^{\mu\alpha} = \begin{cases} \epsilon \, u^{\mu} \, u^{\nu} - p \, \Delta^{\mu\nu} - u^{\mu} \, \bar{Q}^{\nu} - u^{\nu} \, \bar{Q}^{\mu} + \bar{\Pi}^{\mu\nu} & \alpha = \nu, \\ m \, n \, u^{\mu} - \bar{Q}^{\mu} & \alpha = 4, \end{cases}$$
(21)

where  $\bar{Q}^{\mu} = \lambda T \bar{X}^{\mu}$  and  $\bar{\Pi}^{\mu\nu} = 2 \eta X^{\mu\nu}$ . In this frame,  $\bar{X} \equiv X_{\theta=-\pi/4} = 0$  so that the associated bulk pressure  $\bar{\Pi}(=\zeta \bar{X})$  disappears. The unique feature of this equation is the absence of terms with the bulk viscosity. As far as we know, this type of the dissipative fluid dynamical equation is written down for the first time, which was made possible by the introduction of the macroscopic frame vector in the powerful RG method.

Now we have seen that the fluid dynamical equation for dissipative fluids in the Eckart frame does not take the form of Eckart, if we start from the underlying Boltzmann equation. This fact is actually known for some time [15]. Nevertheless it is instructive to see that there can not exist the frame vector  $a_p^{\mu}$  that leads to the Eckart's constraints, (i), (ii) and (iii), because it would help to elucidate the physical meaning of the P- and Q-space in our formalism and their relation to the phenomenological constraints imposed to the dissipative part of the energy-momentum tensor  $\delta J^{\mu\nu}$  and the particle-number current  $\delta J^{\mu4}$ , which are defined just below Eq.(14). Eq.(15) tells us that the dissipative part of the one-particle distribution function  $f_p$ , i.e.,  $-[A^{-1}\bar{Q}F]_p$  can be written as  $f_p^{\text{eq}}\phi_p$  with  $\phi_p \equiv -[L^{-1}Q f^{\text{eq}-1}F]_p$ . Since the Q-space vector  $\phi_p$  is orthogonal to the five P-space vectors  $\varphi_{0p}^{\alpha}$  by definition, we have

$$\langle \varphi_0^{\alpha}, \phi \rangle = 0 \text{ for } \alpha = 0, 1, 2, 3, 4.$$
 (22)

Here the inner product is defined by Eq.(6) where  $a_p^{\mu}$  enters with the form  $(p \cdot a_p)$ . The notable point is that these five identical equations exactly correspond to the constraints on  $\delta J^{\mu\nu}$  and  $\delta J^{\mu4}$ . In fact, Eq.(22) with the choice of (A)  $a_p^{\mu} = u^{\mu}$  gives  $\sum_p \frac{1}{p^0} (p \cdot u) f_p^{eq} \varphi_{0p}^{\alpha} \phi_p = 0$ , which is equivalent to the set of equations  $u_{\nu} \delta J^{\mu\nu} = 0$  and  $u_{\mu} \delta J^{\mu4} = 0$ , because  $\delta J^{\mu\alpha} = \sum_p \frac{1}{p^0} p^{\mu} \varphi_{0p}^{\alpha} f_p^{eq} \phi_p$ . The former equation may be re-expressed by  $u_{\mu} u_{\nu} \delta J^{\mu\nu} = 0$  and

 $\Delta_{\mu\rho} u_{\nu} \delta J^{\mu\nu} = 0$ . Thus one can readily see that these equations coincide with the Landau's constraints, (i), (ii) and (iv). Similarly, with the choice of (B)  $a_p^{\mu} = \frac{m}{p \cdot u} u^{\mu}$ , we have  $\sum_p \frac{1}{p^0} m f_p^{eq} \varphi_{0p}^{\alpha} \phi_p = 0$ , which means that  $\delta J^{\mu 4} = 0$  and  $\delta J^{\mu}_{\mu} = 0$ ; the former equation is equivalent to the set of equations  $u_{\mu} \delta J^{\mu 4} = 0$  and  $\Delta_{\mu\nu} \delta J^{\mu 4} = 0$ . Thus one sees that the Stewart's constraints, (v), (ii) and (iii) are derived. Here we have used the on-shell condition  $m^2 = p^{\mu} p_{\mu}$ . Now it is easy to show that there exists no  $a_p^{\mu}$  leading to the Eckart's constraints, (i), (ii) and (iii), simultaneously. In order to lead to the constraints (ii) and (iii) on  $\delta J^{\mu 4}$ , Eq.(22) with  $\alpha = \mu$  requires that  $(p \cdot a_p) = \text{const.}$ , i.e., independent of  $p^{\mu}$ . On the other hand, in order to lead to the constraint (i) on  $\delta J^{\mu\nu}$ , Eq.(22) with  $\alpha = 4$  must lead to  $(p \cdot a_p) = \text{const.} \times (p \cdot u)^2$ , which is in contradiction with the condition for (ii) and (iii). Thus we can conclude that there exists no  $a_p^{\mu}$  leading to the Eckart's constraint (i) on  $\delta J^{\mu\nu}$ .

What we have seen is that whenever the Eckart frame is taken where the particle flow is constructed so as to have no dissipative part, the constraint on the dissipative part of the energy-momentum tensor must satisfy (v) but not (i). This fact suggests that the Eckart's theory may not be realized as the slow dynamics of the underlying Boltzmann equation, and hence nor have microscopic foundation [15].

One of the unique and important feature of the hydrodynamical equations obtained in the present work lies in the fact that the thermal force driving the heat flow contains  $\nabla^{\mu}T$  but no  $Du^{\mu}$  as in the Landau equation, which in contrast with the Stewart and Eckart equations. We shall briefly argue that the  $Du^{\mu}$  terms should not exist on the general ground on the basis of a natural assumption on the origin of the dissipation; the detailed argument will be presented in separate papers [35].

First of all, we can trace back the absence of  $Du^{\mu}$  terms in the thermal forces to our ansatz that the dissipation comes solely from the spatial inhomogeneity at the rest frame. This ansatz is clearly expressed in the choice of  $a_p^{\mu}$ : Although the generic form of  $a_p^{\mu}$  which keeps the Lorentz covariance can be given as  $a_p^{\mu} = \alpha(p \cdot u) u^{\mu} + \beta(p \cdot u) \Delta^{\mu\nu} p_{\nu}$ , where  $\alpha(x)$  and  $\beta(x)$  are the arbitrary Lorentz-scalar functions, we have intentionally put  $\beta(x) = 0$  so that the differential operator  $\nabla^{\mu}$  in the last term in Eq.(3), which gives rise to the deviation from the local equilibrium and hence the dissipation, agrees with the spatial derivative, i.e.,  $\nabla^{\mu} = \nabla^{\mu}$ . As mentioned before, the non-relativistic Navier-Stokes equation can be nicely derived by the same ansatz that the spatial inhomogeneity leads to the dissipation [32, 33], and the present approach is simply a covariantization of the non-relativistic case. On the other hand, the ansatz used by Landau or Stewart is that the dissipation comes from both of the spatial and the temporal inhomogeneity at the rest frame, although no  $Du^{\mu}$  terms appear in the Landau equation.

In a separate paper, we shall show that the covariant dissipative hydrodynamic equations constructed in the phenomenological way as was done by Eckart, Landau, Stewart so on but based on the same ansatz as that in the present work, completely agree with the equations derived in the present RG method for the Landau and Eckart frame, i.e., with the two choices for  $a_n^{\mu}$ .

In summary, we have derived generic covariant hydrodynamical equations for viscous fluids as a reduction of dynamics from relativistic Boltzmann equation in a mechanical way with no heuristic arguments on the basis of so called the renormalization-group method: This was made possible by introducing the macroscopic frame vector which defines the macroscopic Lorenz frame in which the slow dynamics is described. Our generic equation includes not only the Landau equation[12] but also a modified Eckart/Stewart equation[13, 14] and a novel one which does not contain the bulk viscosity term.

We notice that the derived equation is consistent with the underlying kinetic equation, so the equations and also the method developed here may be useful for the analysis of the system where the proper dynamics describing the system changes from the hydrodynamic to kinetic regime, as in the system near the freeze-out region in the RHIC experiment.

Our method is so mechanical and simple that it is successfully applicable to derive the dissipative hydrodynamics for the multi-component system in Landau frame [34]. In the present work, the fluid dynamical variables correspond to the conserved quantities in the collision process and span the five-dimensional P space, which is an invariant manifold in the terminology of the dynamical systems. One may suspect that the method could apply to derive the Israel-Stewart equation [20] with fourteen variables in accordance with Grad's moment method [22]. The answer is affirmative: We have found [34] that a simple extension of the P-space in our formalism gives the equation equivalent to Grad's fourteen-moment method [22]. These extensions will be reported elsewhere. In passing, we emphasize that our method itself has a universal nature and can be applied to derive a slow dynamics from other kinetic equations than the simple Boltzmann equation.

We thank Berndt Muller and Tetsufumi Hirano for their interest in this work and encouragement. T.K. is supported

by Grant-in-Aid for Scientific Research by Monbu-Kagakusyo(No. 17540250).

- [1] T. Hirano and K. Tsuda, Phys. Rev. C 66 (2002) 054905.
- See review articles, P. Huovinen, in "Quark gluon plasma", ed. R.C. Hwa and X.N. Wang, (World Scientific, Singapore), p.600 [arXiv:nucl-th/0305064];
- P. F. Kolb and U. W. Heinz, in "Quark gluon plasma", ed. R.C. Hwa and X.N. Wang, (World Scientific, Singapore), p.634.(arXiv:nucl-th/0305084).
- [2] M. Gyulassy and L. McLerran, Nucl. Phys. A 750 (2005) 30.
- [3] D. Teaney, Phys. Rev. C 68 (2003) 034913.
- [4] A. Muronga, Phys. Rev. Lett. 88 (2002) 062302; Phys. Rev. C 69 (2004) 034903.
- [5] A. Muronga and D. H. Rischke, nucl-th/0407114.
- [6] A. K. Chaudhuri and U. W. Heinz, nucl-th/0504022; U. W. Heinz, H. Song and A. K. Chaudhuri, nucl-th/0510014.
- [7] T. Hirano, U. W. Heinz, D. Kharzeev, P. Lacey and Y. Nara, Phys. Lett. B636 (2006) 299.
- [8] R. Baier, P. Romatschke and U. A. Wiedemann, hep-ph/0602249.
- [9] C. Nonaka and S. A. Bass, nucl-th/0607018.
- [10] D. Balsara, Astrophys. J. Suppl. 132 (2001) 83.
   J. C. Fabris, S. V. B. Goncalves and R. de Sa Ribeiro, Gen. Rel. Grav. 38 (2006) 495 [arXiv:astro-ph/0503362].
- [11] C. Eckart, Phys. Rev. 58 (1940) 919.
- [12] L. D. Landau and E. M. Lifshitz, Fluid Mechanics, (Pergamon Press, London, 1959).
- [13] N.G. van Kampen, J. of Stat. Phys. 46 (1987) 709.
- [14] J. M. Stewart, Non-Equilibrium Relativistic Kinetic Theory (Lecture Notes in Physics No. 10; Springer, Berlin, 1971).
- [15] Ch. G. van Weert, Some Problems in Relativistic Hydrodynamics, in Lecture Notes in Mathematics 1385, p.290, ed. by A. Anile and Y. Choquet-Bruhat (Springer, 1987).
- [16] W. A. Hiscock, L. Lindblom, Phys. Rev. D 31 (1985) 725.
- [17] W. Israel, Ann. Phys. 100 (1976) 310.
- [18] S. R. de Groot, W. A. van Leeuwen and Ch. G. van Weert, Relativistic Kinetic Theory (Elsevier North-Holland, 1980).
- [19] W. Israel, J. Math. Phys. 4 (1963) 1163.
- [20] W. Israel and J. M. Stewart, Ann. Phys. 118 (1979) 341.
- [21] S. Chapman and T. G. Cowling, The Mathematical Theory of Non-Uniform Gases, (Cambridge U.P., 1939).
- [22] H. Grad, Comm. Pure Appl. Math. 2 (1949) 331.
- [23] N.G.van Kampen, J. of Stat. Phys. 46 (1987) 709.
- [24] S.A.Bass and A.Dumitru, Phys.Rev. C61 (2000) 064909;
   D. Teaney, J. Lauret and E. V. Shuryak, Phys. Rev. Lett. 86 (2001)4783.
- [25] See for example, Y. Kuramoto, Prog. Theor. Phys. Suppl. **99** (1989) 244.
- [26] N. G. van Kampen, Phys. Rep. **124** (1985) 69.
- [27] L. Y. Chen, N. Goldenfeld and Y. Oono, Phys. Rev. Lett. 73 (1994) 1311; Phys. Rev. E 54 (1996) 376.
- [28] T. Kunihiro, Prog. Theor. Phys. 94 (1995) 503; 95(1996) 835 (E); Jpn. J. Ind. Appl. Math. 14 (1997) 51.
- [29] T. Kunihiro, Prog. Theor. Phys. 97 (1997) 179.
- [30] T. Kunihiro and J. Matsukidaira, Phys. Rev. E. 57 (1998) 4817.
- [31] S.-I. Ei, K. Fujii, and T. Kunihiro, Ann. Phys. 280 (2000) 236.
- [32] Y. Hatta and T. Kunihiro, Ann. Phys. 298 (2002) 24.
- [33] T. Kunihiro and K. Tsumura, J. Phys. A 39 (2006) 8089.
- [34] K. Tsumura, K. Ohnishi and T. Kunihiro, in preparation.
- [35] K. Tsumura and T. Kunihiro, in preparation.
- [36] Although the bilinear term of  $\tilde{f}^{(1)}(\tau)$  appears from the collision integral, here we neglect it. It is known that the neglected term produces the so-called Burnett terms which represents the higher-order non-equilibrium effects.
- [37] This is tantamount to inserting the solution  $f(\tau)$  into Eq.(2).
- [38] In the original paper [11], basically the bulk viscosity is not taken into account.