

# Topological charge screening and the infrared behavior of the instanton density

Nikolai Kochelev<sup>a,b1</sup>

(a) *School of Physics and Astronomy, Seoul National University,  
Seoul 151-747, Korea*

(b) *Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research,  
Dubna, Moscow region, 141980 Russia*

## Abstract

A new mechanism for suppression of the instanton density in the infrared is considered. This mechanism is based on the phenomenon of topological charge screening, which leads to an effective cutoff in the contribution of large instantons.

arXiv:hep-ph/0611198v2 20 Nov 2006

---

<sup>1</sup>kochelev@theor.jinr.ru

Instantons, strong topological fluctuations of gluon fields in QCD, are widely believed to play an important role in the physics of the strong interaction (for reviews see [1], [2]). In particular, instantons provide mechanisms for the violation of both  $U(1)_A$  and chiral symmetry in QCD, and may therefore be important in determining hadron masses and in the resolution of the famous  $U(1)_A$  problem. Furthermore, it was recently shown that instantons persist through the deconfinement transition, so that instanton-induced interactions between quarks and gluons may underlie the unusual properties of the so-called strongly coupled quark-gluon plasma recently discovered at RHIC [3].

One longstanding problem in the calculation of instanton contributions to physical processes is the uncertainty in their distribution over the full range of instanton length scales. For small instantons in the instanton dilute gas approximation this distribution is well known,

$$D_0(\rho) = \frac{d}{\rho^5} \left( \frac{2\pi}{\alpha_s(\rho)} \right)^{2N_c} e^{-S(\rho)} \quad (1)$$

where  $d$  is a renormalization scheme dependent constant, and

$$S(\rho) = S_0(\rho) = \frac{2\pi}{\alpha_s(\rho)} \quad (2)$$

is the Euclidean action of the instanton solution. This distribution increases rapidly with increasing instanton size, as  $D_0(\rho) \approx \rho^{\beta_0-5}$  (where  $\beta_0 = 11N_c/3 - 2N_f/3$ ). The resulting instanton density integral is divergent, so it is impossible to make definite quantitative predictions regarding instanton effects given this distribution. Shifman, Vainshtein and Zakharov (SVZ) [4] have shown that at an instanton length scale of  $\rho \approx 1 \text{ GeV}^{-1}$  the interaction between instantons and the gluon condensate becomes strong, which leads to a significant change in the instanton density at large instanton length scales,

$$D_{cond}(\rho) \approx D(\rho)_0 \exp \left\{ \frac{\pi^4 \rho^4}{8\alpha_s^2(\rho)} < 0 | \frac{\alpha_s}{\pi} G_{\mu\nu}^a G_{\mu\nu}^a | 0 > \right\} \quad (3)$$

where the gluon condensate is numerically [5]

$$< 0 | \frac{\alpha_s}{\pi} G_{\mu\nu}^a G_{\mu\nu}^a | 0 > = 0.009 \pm 0.007 \text{ GeV}^4. \quad (4)$$

Evidently this interaction leads to an even stronger divergence in the instanton density at large length scales than in Eq.(1). Recent lattice calculations however show a damping in the rate of increase of the instanton density for instanton lengths of  $\rho_c \sim 0.3 - 0.5 \text{ fm}$ , followed by a rapid decrease at larger instanton sizes [6]. Several scenarios have been suggested to explain the physics behind this phenomenon. One possible mechanism for this decrease is a repulsive interaction between large instantons. This idea was suggested by Ilgenfritz and Müller-Preussker in a papers [7] and was developed by Diakonov and Petrov later on [8]. The effect of confinement on the instanton distribution, considered by Shuryak, is another possible mechanism for the suppression of the instanton density at large  $\rho$  [9]. In a paper by Dorokhov *et al.* [10] it was shown that the deformation of instantons in external vacuum fields might also lead to a suppression of the density of large instantons.

In this Letter we consider another possible mechanism for the suppression of large instantons, which is the influence of topological charge screening on instanton density.

This screening effect is known to play an important role in the properties of the  $\eta'$  meson, and leads in particular to a vanishing topological susceptibility

$$\chi(k^2) = i \int d^4x e^{ikx} \langle 0|T\{Q_5(x)Q_5(0)\}|0 \rangle \quad (5)$$

as  $k^2 \rightarrow 0$ , in the chiral limit [1]. In Eq.(5),

$$Q_5(x) = \frac{\alpha_s}{8\pi} G_{\mu\nu}^a(x) \tilde{G}_{\mu\nu}^a(x) \quad (6)$$

is the topological charge density, and  $\tilde{G}_{\mu\nu}^a = 1/2\epsilon_{\mu\nu\alpha\beta}G_{\alpha\beta}^a$  is the dual gluon field strength tensor.

Our starting point is the SVZ interaction between an instanton and an external gluon field [4]:

$$\mathcal{L}_{eff}(x_0) = \int d\rho D_0(\rho) \exp\left(-\frac{2\pi^2\rho^2}{g_s} \bar{\eta}_{a\alpha\beta} G_{\alpha\beta}^a(x_0)\right). \quad (7)$$

Here  $\bar{\eta}_{a\alpha\beta}$  is 't Hooft's symbol, and  $x_0$  is the instanton's location. On expanding Eq.(7) in powers of the gluon field strength  $G_{\alpha\beta}$ , and keeping only terms with an even number of gluon fields, we find

$$\begin{aligned} \mathcal{L}_{eff} &= \int d\rho D_0(\rho) \left[1 + \frac{1}{2!} \left(\frac{2\pi^2\rho^2}{g_s}\right)^2 \bar{\eta}_{a\alpha\beta} \bar{\eta}_{b\mu\nu} G_{\alpha\beta}^a G_{\mu\nu}^b \right. \\ &\quad \left. + \frac{1}{4!} \left(\frac{2\pi^2\rho^2}{g_s}\right)^4 \bar{\eta}_{a\alpha\beta} \bar{\eta}_{b\mu\nu} \bar{\eta}_{c\rho\sigma} \bar{\eta}_{f\tau\gamma} G_{\alpha\beta}^a G_{\mu\nu}^b G_{\rho\sigma}^c G_{\tau\gamma}^f + \dots\right]. \end{aligned} \quad (8)$$

The effect of the gluon condensate on the instanton density at leading order is

$$\mathcal{L}_{eff} = \int d\rho D_0(\rho) \left[1 + \frac{\pi^4\rho^4}{8\alpha_s^2(\rho)} \langle 0|\frac{\alpha_s}{\pi} G_{\mu\nu}^a G_{\mu\nu}^a|0 \rangle + \dots\right], \quad (9)$$

which follows from the second term in Eq.(8), after using the color-singlet nature of the vacuum state

$$\langle 0|G_{\alpha\beta}^a G_{\mu\nu}^b|0 \rangle = \frac{1}{N_c^2 - 1} \delta^{ab} \langle 0|G_{\alpha\beta}^a G_{\mu\nu}^a|0 \rangle \quad (10)$$

and an identity for 't Hooft symbols in Minkowski space-time,

$$\bar{\eta}_{a\alpha\beta} \bar{\eta}_{a\mu\nu} = g_{\alpha\mu} g_{\beta\nu} - g_{\alpha\nu} g_{\beta\mu} - i\epsilon_{\alpha\beta\mu\nu}. \quad (11)$$

The expansion of Eq.(7) to high orders, combined with an assumption of factorization for the higher  $\langle 0|G^{2n}|0 \rangle$  condensates, leads to the exponential form given in Eq.(3).

We emphasize that in their original derivation of the effect of gluon condensates on the instanton density [4], SVZ ignored all terms involving the dual tensor  $\tilde{G}_{\mu\nu}^a$ . Actually only the leading term in the dual tensor expansion vanishes,

$$\langle 0|Q_5(x)|0 \rangle = 0, \quad (12)$$

due to the  $CP$  invariance of the QCD vacuum. Terms involving higher powers of  $\tilde{G}_{\mu\nu}^a$  generally do contribute to the expansion in Eq.(8). The leading dual tensor contribution arises from the third term in Eq.(8), and involves the matrix element

$$M = \langle 0|\bar{\eta}_{a\alpha\beta} \bar{\eta}_{b\mu\nu} \bar{\eta}_{c\rho\sigma} \bar{\eta}_{f\tau\gamma} G_{\alpha\beta}^a G_{\mu\nu}^b G_{\rho\sigma}^c G_{\tau\gamma}^f|0 \rangle. \quad (13)$$

By using the formula for the vacuum matrix element

$$\langle 0 | G_{\alpha\beta}^a G_{\mu\nu}^b G_{\rho\sigma}^c G_{\tau\gamma}^f | 0 \rangle \rightarrow \frac{\delta^{ab} \delta^{cf}}{N_c^4 - 1} \langle 0 | (G_{\alpha\beta}^d G_{\mu\nu}^d G_{\rho\sigma}^m G_{\tau\gamma}^m + perms.) | 0 \rangle, \quad (14)$$

one obtains

$$\Delta \mathcal{L}_{eff} \sim \int d\rho D_0(\rho) \frac{\pi^6 \rho^8}{5 \cdot 2^5 \alpha_s^4(\rho)} \langle 0 | \alpha_s G_{\mu\nu}^a(x_1) \tilde{G}_{\mu\nu}^a(x_2) \alpha_s G_{\alpha\beta}^b(x_3) \tilde{G}_{\alpha\beta}^b(x_4) | 0 \rangle, \quad (15)$$

where  $x_i \approx x_0$ . In Eq.(15) the nonlocality of the instanton is explicitly taken into account. This differs from the contribution of a pure gluon condensate, for which one can use the local approximation  $x_i = x_0$ . The difference follows from the singular behavior of the topological density at the origin. To illustrate this, note that in the instanton model of the QCD vacuum the correlator of topological density in the chiral limit is [1]

$$\langle 0 | Q_5(x) Q_5(0) | 0 \rangle \approx \frac{N}{V} \left[ \delta^4(x) - \frac{2N_f N}{f_\pi^2 V} \langle 0 | \eta'(x) \eta'(0) | 0 \rangle \right], \quad (16)$$

where  $N/V$  is the instanton density. The local term in Eq.(16) gives the contribution of a single instanton centered at the origin, and the second term is the so-called screening contribution. Evidently one should ignore the contribution of the first term in Eq.(16) in estimating the matrix element in Eq.(15), because it arises from the instanton under consideration. Therefore in the limit  $|x_i - x_j| \rightarrow 0$  we obtain

$$\langle 0 | \alpha_s G_{\mu\nu}^a \tilde{G}_{\mu\nu}^a \alpha_s G_{\alpha\beta}^b \tilde{G}_{\alpha\beta}^b | 0 \rangle^{scr} \approx -\frac{4}{\pi^2} \int d^4 k_E \bar{\chi}(k_E^2), \quad (17)$$

where  $\bar{\chi}(k_E^2)$  is the screening contribution to the topological susceptibility (for Euclidean momenta). It should be noted that the exact form of  $\chi(k^2)$  is unknown, and only model calculations of this function are available (see [1], [11], [12]). Here we are interested in the long range behavior of  $\chi$ , which can be represented by the expression

$$\chi(k_E^2) \approx -\chi'(0) m_{\eta'}^2 (1 - e^{-k_E^2/m_{\eta'}^2}), \quad (18)$$

where the value of the derivative  $\chi'(0)$  is known from a QCD sum rule calculation [11],  $\chi'(0) = (2.3 \pm 0.6) \times 10^{-3} \text{ GeV}^2$ . From Eq.(18), the screening part is

$$\bar{\chi}(k_E^2) \approx \chi'(0) m_{\eta'}^2 e^{-k_E^2/m_{\eta'}^2}. \quad (19)$$

Thus we find the estimate

$$\langle 0 | \alpha_s G_{\mu\nu}^a \tilde{G}_{\mu\nu}^a \alpha_s G_{\alpha\beta}^b \tilde{G}_{\alpha\beta}^b | 0 \rangle^{scr} \approx -4\chi'(0) m_{\eta'}^6. \quad (20)$$

It is easy to show that the higher order expansion in  $\tilde{G}_{\mu\nu}^a$  can be exponentiated, given the assumption of vacuum state dominance. Our final result for the screening correction to the instanton density is thus

$$\Delta \mathcal{L}_{eff} \approx \int d\rho D_0(\rho) \exp \left( -\frac{\pi^6 \rho^8}{40 \alpha_s^4(\rho)} \chi'(0) m_{\eta'}^6 \right). \quad (21)$$

In contrast to the  $\rho^4$  dependence in the exponential factor in Eq.(3), our novel correction evidently has a faster  $\rho^8$  behavior in the exponent. The sign of this contribution is very important; since it is negative, this effect leads to a *suppression* of the instanton density at large  $\rho$ . We emphasize that this sign is determined by the sign of the screening correction in Eq.(16), and follows from the general property of the positivity of state norms in Euclidean field theory [13].<sup>1</sup>




Figure 1: The ratio of our effective action to the free instanton action, after incorporating the gluon condensate contribution (dashed) and the screening correction (solid).

In Fig.1 we show the effective action ratio  $S_{eff}(\rho)^{cond,scr}/S_0(\rho)$ , where  $S_{eff}(\rho)^{cond}$  is the effective action including the gluonic condensate correction, Eq.(3), and  $S_{eff}(\rho)^{scr}$  is the action including the screening contribution, Eq.(21). To generate this figure the expression

$$\alpha_s(\rho) = \frac{4\pi}{9 \ln[(1/(\rho^2 \Lambda^2))]} \quad (22)$$

was used to estimate the strong coupling constant, where  $\Lambda \approx 300$  MeV. It is evident in Fig.1 that the effect of screening on the instanton density becomes dominant at rather small instanton length scales ( $\rho > 1$  GeV<sup>-1</sup>), and leads to a rapid increase in the effective instanton action at large instanton scales. One can therefore anticipate a corresponding cutoff in the instanton density relative to the dilute gas approximation of Eq.(1).

In summary, we have shown that topological charge screening strongly affects the length scale dependence of the distribution of instantons, and provides a natural infrared cutoff in the instanton density.

---

<sup>1</sup>We are grateful to A.Di Giacomo for clarifying this point.

We are happy to acknowledge useful discussions of various aspects of this work with A.Di Giacomo, A.E.Dorokhov, E.-M.Ilgenfritz, B.L.Ioffe, E.V.Shuryak and V.I.Zakharov. We would also like to thank T.Barnes for proofreading the manuscript. The author is very grateful to the School of Physics and Astronomy of Seoul National University, and especially Prof. Dong-Pil Min, for their kind hospitality. This work was supported in part by the Brain Pool program of the Korea Research Foundation through KOFST grant 042T-1-1, by the Russian Foundation for Basic Research through grant RFBR-04-02-16445.

## References

- [1] T. Schäfer and E.V. Shuryak, *Rev. Mod. Phys.* **70** (1998) 1323.
- [2] D. Diakonov, *Prog. Part. Nucl. Phys.* **51** (2003) 173.
- [3] E. V. Shuryak, arXiv:hep-ph/0608177.
- [4] M.A. Shifman, A.I.Vanshtein and V.I. Zakharov, *Nucl. Phys.* **B165** (1980) 45.
- [5] B. L. Ioffe, *Prog. Part. Nucl. Phys.* **56** (2006) 232.
- [6] B. Lucini, M. Teper and U. Wenger, *Nucl. Phys.* **B715** (2005) 461; A. Hasenfratz and C. Nieter, *Phys. Lett.* **B439** (1998) 366;
- [7] E.-M. Ilgenfritz and M. Muller-Preussker *Nucl. Phys.* **B184** (1981) 443; *Z.Phys.* **C16** (1983) 339, Erratum-ibid. **C20** (1983) 186.
- [8] D.I. Diakonov and V.Y. Petrov, *Nucl. Phys.* **B245** (1984) 259.
- [9] E.V. Shuryak, hep-ph/9909458; hep-th/0605219.
- [10] A. E. Dorokhov, S. V. Esaibegian, A. E. Maximov and S. V. Mikhailov, *Eur. Phys. J.* **C13** (2000) 331.
- [11] B. L. Ioffe, *Surveys High Energ. Phys.* **14** (1999) 89; B. L. Ioffe and A. V. Samsonov, *Phys. Atom. Nucl.* **63** (2000) 1448 [*Yad. Fiz.* **63** (2000) 1527]; B. V. Geshkenbein and B. L. Ioffe, *Nucl. Phys.* **B166**, 340 (1980).
- [12] A. E. Dorokhov and W. Broniowski, *Eur. Phys. J.* **C32** (2003) 79.
- [13] K. Osterwalder and R. Schrader, *Commun. Math. Phys.* **31** (1973) 83; **42** (1975) 281.