# Lepton Family Symmetry and Possible Application to the Koide Mass Formula

Ernest Ma

Physics and Astronomy Department University of California, Riverside Riverside, California 92521, USA

#### Abstract

A finite group generated by four  $Z_3$  transformations is applied to lepton families in a supersymmetric model, resulting in the charged-lepton masses  $m_i$  being proportional to  $v_i^2$ , where  $v_i$  are three vacuum expectation values. This may be relevant in obtaining the Koide formula  $m_e + m_\mu + m_\tau = (2/3)(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2$ . The Koide mass formula [1, 2, 3, 4]

$$m_e + m_\mu + m_\tau = \frac{2}{3} \left( \sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau} \right)^2$$
 (1)

is remarkably accurate, and cries out for a possible theoretical explanation. Koide himself has proposed [5] to understand it as

$$v_1^2 + v_2^2 + v_3^2 = \frac{2}{3}(v_1 + v_2 + v_3)^2.$$
 (2)

In that case, the condition  $m_i \propto v_i^2$  is required, and must be valid to great precision. To facilitate such a relationship, the following non-Abelian discrete family symmetry is proposed.

Consider the basis  $(a_1, a_2, a_3)$  and four separate  $Z_3$  transformations:

$$a_1 \to a_2 \to a_3 \to a_1, \quad a_i \to \omega a_i \to \omega^2 a_i \to a_i, \ (i = 1, 2, 3),$$
(3)

where  $\omega = \exp(2\pi i/3) = -1/2 + i\sqrt{3}/2$ . The resulting group, call it  $\Sigma(81)$ , has 81 elements, divided into 17 equivalence classes. It has 9 one-dimensional irreducible representations  $\mathbf{1}_{i}(i = 1, ..., 9)$  and 8 three-dimensional ones  $\mathbf{3}_{A}, \mathbf{\overline{3}}_{A}, \mathbf{3}_{B}, \mathbf{\overline{3}}_{B}, \mathbf{3}_{C}, \mathbf{\overline{3}}_{C}, \mathbf{3}_{D}, \mathbf{\overline{3}}_{D}$ . Its character table is given in the Appendix, together with the 81 matrices of the defining representation  $\mathbf{3}_{A}$ .

The key property of  $\Sigma(81)$  which allows this to work is that given just the representations  $\mathbf{3}_{\mathbf{A}}$  and  $\mathbf{\bar{3}}_{\mathbf{A}}$ , there are only three invariants:

 $a_1a_1a_1 + a_2a_2a_2 + a_3a_3a_3, \quad a_1\bar{a}_1 + a_2\bar{a}_2 + a_3\bar{a}_3, \quad \bar{a}_1\bar{a}_1\bar{a}_1 + \bar{a}_2\bar{a}_2\bar{a}_2 + \bar{a}_3\bar{a}_3\bar{a}_3. \tag{4}$ 

Consider now the supersymmetric standard model, with lepton superfields

$$L_i = (\nu_i, l_i), \quad l_i^c \sim \mathbf{3}_{\mathbf{A}}.$$
(5)

Add the following Higgs superfields:

$$\eta_i = (\eta_i^0, \eta_i^-), \quad \xi_i = (\xi_i^+, \xi_i^0) \sim \mathbf{3}_{\mathbf{A}},$$
(6)

$$\zeta_i = (\zeta_i^0, \zeta_i^-), \quad \psi_i = (\psi_i^+, \psi_i^0) \sim \bar{\mathbf{3}}_{\mathbf{A}}, \tag{7}$$

$$\sigma_i = \sigma_i^0 \sim \bar{\mathbf{3}}_{\mathbf{A}}, \quad \phi = (\phi^0, \phi^-) \sim \mathbf{1}.$$
(8)

The relevant superpotential is then given by

$$W = f L_i l_i^c \eta_i + M_1 \eta_i \psi_i + h_1 \zeta_i \psi_i \sigma_i + M_2 \zeta_i \xi_i + h_2 \xi_i \sigma_i \phi, \qquad (9)$$

resulting in the scalar potential

$$V = |M_1\psi_i + fL_i l_i^c|^2 + |M_2\xi_i + h_1\psi_i\sigma_i|^2 + |M_1\eta_i + h_1\zeta_i\sigma_i|^2 + |M_2\zeta_i + h_2\sigma_i\phi|^2.$$
(10)

Assuming  $\langle \phi^0 \rangle \neq 0$  and  $\langle \sigma_i \rangle \neq 0$  (to be discussed later), the supersymmetric minimum V = 0is obtained for

$$\langle \psi_i^0 \rangle = \langle \xi_i^0 \rangle = 0, \quad \langle \eta_i^0 \rangle = \frac{-h_1 \langle \zeta_i^0 \rangle \langle \sigma_i \rangle}{M_1}, \quad \langle \zeta_i^0 \rangle = \frac{-h_2 \langle \sigma_i \rangle \langle \phi^0 \rangle}{M_2}, \tag{11}$$

resulting thus in the charged-lepton mass relationship

$$m_i = f\langle \eta_i^0 \rangle = \frac{fh_1h_2\langle \sigma_i \rangle^2 \langle \phi^0 \rangle}{M_1M_2},\tag{12}$$

exactly as advertised.

To obtain Eq. (2), let [5]

$$\begin{pmatrix} \chi_0 \\ \chi_1 \\ \chi_2 \end{pmatrix} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{pmatrix} \begin{pmatrix} \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix},$$
(13)

so that

$$\sigma_1^2 + \sigma_2^2 + \sigma_3^2 = \chi_0^2 + 2\chi_1\chi_2, \tag{14}$$

$$\sigma_1^3 + \sigma_2^3 + \sigma_3^3 = \frac{1}{\sqrt{3}} (\chi_0^3 + \chi_1^3 + \chi_2^3 + 6\chi_0\chi_1\chi_2).$$
(15)

Consider the superpotential

$$W = \frac{1}{2}m_0\chi_0^2 + \frac{1}{2}m_1\chi_1^2 + \frac{1}{2}m_2\chi_2^2 + m_3\chi_1\chi_2 + \frac{1}{3}\lambda(\chi_0^3 + \chi_1^3 + \chi_2^3 + 6\chi_0\chi_1\chi_2), \quad (16)$$

which breaks  $\Sigma(81)$  only softly with its bilinear terms. The resulting scalar potential is given by

$$V = |m_0\chi_0 + \lambda(\chi_0^2 + 2\chi_1\chi_2)|^2 + |m_1\chi_1 + m_3\chi_2 + \lambda(\chi_1^2 + 2\chi_0\chi_2)|^2 + |m_2\chi_2 + m_3\chi_1 + \lambda(\chi_2^2 + 2\chi_0\chi_1)|^2.$$
(17)

The aim now is to find a supersymmetric minimum, i.e. V = 0, for which

$$\langle \chi_0 \rangle^2 = 2 \langle \chi_1 \rangle \langle \chi_2 \rangle = \frac{1}{2} (\langle \chi_0 \rangle^2 + 2 \langle \chi_1 \rangle \langle \chi_2 \rangle), \tag{18}$$

which is equivalent to [5]

$$\frac{1}{3}(\langle \sigma_1 \rangle + \langle \sigma_2 \rangle + \langle \sigma_3 \rangle)^2 = \frac{1}{2}(\langle \sigma_1 \rangle^2 + \langle \sigma_2 \rangle^2 + \langle \sigma_3 \rangle^2), \tag{19}$$

i.e. Eq. (2), and then use Eq. (12) to obtain the Koide formula of Eq. (1). This turns out to be extremely simple, i.e.

$$m_3 = m_0 = \sqrt{8m_1m_2},\tag{20}$$

for which

$$\langle \chi_0 \rangle = \frac{-m_0}{2\lambda}, \quad \langle \chi_1 \rangle = \frac{-m_1}{\lambda}, \quad \langle \chi_2 \rangle = \frac{-m_2}{\lambda}.$$
 (21)

The validity of Eq. (20) is of course not understood. It may be due to some underlying symmetry or dynamics, but whatever the origin, once it is established, Eq. (2) is protected by the unbroken supersymmetry of the theory. Assuming that  $\chi_i$  acquire vaccum expectation values not far above the supersymmetry breaking scale, the effective Yukawa couplings of the leptons at the electroweak scale will not deviate very much from their values which predict Eq. (1).

As for the neutrino sector [6], it is clear that having only superfields which transform as  $\mathbf{3}_{\mathbf{A}}$  and  $\mathbf{\overline{3}}_{\mathbf{A}}$  will not work, because that necessarily yields a diagonal mass matrix. To obtain a realistic neutrino mass matrix, consider for example scalar superfields  $(\Delta_i^{++}, \Delta_i^+, \Delta_i^0)$  which transform as  $\mathbf{3}_{\mathbf{A}}$  and  $\mathbf{3}_{\mathbf{B}}$  under  $\Sigma(81)$ . Using the multiplication rule

$$\mathbf{3}_{\mathbf{A}} \times \mathbf{3}_{\mathbf{A}} = \overline{\mathbf{3}}_{\mathbf{A}} + \overline{\mathbf{3}}_{\mathbf{B}} + \overline{\mathbf{3}}_{\mathbf{B}}, \tag{22}$$

it is possible to have  $\mathcal{M}_{\nu}$  of the most general form, i.e.

$$\mathcal{M}_{\nu} = \begin{pmatrix} a & f & e \\ f & b & d \\ e & d & c \end{pmatrix},\tag{23}$$

where (a, b, c) come from  $\mathbf{3}_{\mathbf{A}}$  and (d, e, f) from  $\mathbf{3}_{\mathbf{B}}$ . This makes no prediction, but if the vacuum structure can be restricted, say b = c and e = f, as for example in a model [7] based on  $S_4$ , then the approximate tribinaximal mixing of neutrinos may be obtained.

#### Acknowledgements

I thank Yoshio Koide for correspondence. This work was supported in part by the U. S. Department of Energy under Grant No. DE-FG03-94ER40837.

### Appendix

The proposed finite group  $\Sigma(81)$  contains  $\Delta(27)$  [8, 9]. The latter's character table and defining representation have been given elsewhere [10]. Here the  $17 \times 17$  character table of  $\Sigma(81)$  and the 81 matrices of its defining representation  $\mathbf{3}_{\mathbf{A}}$  are also given.

Table 1: Character table of  $\Sigma(81)$  for the 9 one-dimensional representations; n is the number of elements and h is the order of each element.

Class	n	h	$1_1$	$1_2$	$1_3$	$1_4$	$1_5$	$1_6$	$1_7$	$1_8$	$1_9$
$C_1$	1	1	1	1	1	1	1	1	1	1	1
$C_2$	1	3	1	1	1	1	1	1	1	1	1
$C_3$	1	3	1	1	1	1	1	1	1	1	1
$C_4$	3	3	1	1	1	1	1	1	1	1	1
$C_5$	3	3	1	1	1	1	1	1	1	1	1
$C_6$	9	3	1	ω	$\omega^2$	1	$\omega^2$	ω	1	ω	$\omega^2$
$C_7$	9	3	1	$\omega^2$	ω	1	ω	$\omega^2$	1	$\omega^2$	ω
$C_8$	3	3	1	1	1	$\omega^2$	$\omega^2$	$\omega^2$	ω	ω	ω
$C_9$	3	3	1	1	1	$\omega^2$	$\omega^2$	$\omega^2$	ω	ω	ω
$C_{10}$	3	3	1	1	1	$\omega^2$	$\omega^2$	$\omega^2$	ω	ω	ω
$C_{11}$	3	3	1	1	1	ω	ω	ω	$\omega^2$	$\omega^2$	$\omega^2$
$C_{12}$	3	3	1	1	1	ω	ω	ω	$\omega^2$	$\omega^2$	$\omega^2$
$C_{13}$	3	3	1	1	1	ω	ω	ω	$\omega^2$	$\omega^2$	$\omega^2$
$C_{14}$	9	9	1	ω	$\omega^2$	$\omega^2$	ω	1	ω	$\omega^2$	1
$C_{15}$	9	9	1	$\omega^2$	ω	$\omega^2$	1	ω	ω	1	$\omega^2$
$C_{16}$	9	9	1	$\omega^2$	ω	ω	$\omega^2$	1	$\omega^2$	ω	1
$C_{17}$	9	9	1	ω	$\omega^2$	ω	1	$\omega^2$	$\omega^2$	1	ω

Class	n	h	$3_{\mathbf{A}}$	$ar{3}_{\mathbf{A}}$	$3_{ m B}$	$ar{3}_{ ext{B}}$	$3_{ m C}$	$ar{3}_{\mathbf{C}}$	$3_{\mathrm{D}}$	$\bar{3}_{\mathrm{D}}$
$C_1$	1	1	3	3	3	3	3	3	3	3
$C_2$	1	3	$3\omega$	$3\omega^2$	$3\omega$	$3\omega^2$	$3\omega$	$3\omega^2$	3	3
$C_3$	1	3	$3\omega^2$	$3\omega$	$3\omega^2$	$3\omega$	$3\omega^2$	$3\omega$	3	3
$C_4$	3	3	0	0	0	0	0	0	$3\omega$	$3\omega^2$
$C_5$	3	3	0	0	0	0	0	0	$3\omega^2$	$3\omega$
$C_6$	9	3	0	0	0	0	0	0	0	0
$C_7$	9	3	0	0	0	0	0	0	0	0
$C_8$	3	3	$-i\sqrt{3}$	$i\sqrt{3}$	$-i\sqrt{3}\omega^2$	$i\sqrt{3}\omega$	$-i\sqrt{3}\omega$	$i\sqrt{3}\omega^2$	0	0
$C_9$	3	3	$-i\sqrt{3}\omega$	$i\sqrt{3}\omega^2$	$-i\sqrt{3}$	$i\sqrt{3}$	$-i\sqrt{3}\omega^2$	$i\sqrt{3}\omega$	0	0
$C_{10}$	3	3	$-i\sqrt{3}\omega^2$	$i\sqrt{3}\omega$	$-i\sqrt{3}\omega$	$i\sqrt{3}\omega^2$	$-i\sqrt{3}$	$i\sqrt{3}$	0	0
$C_{11}$	3	3	$i\sqrt{3}$	$-i\sqrt{3}$	$i\sqrt{3}\omega$	$-i\sqrt{3}\omega^2$	$i\sqrt{3}\omega^2$	$-i\sqrt{3}\omega$	0	0
$C_{12}$	3	3	$i\sqrt{3}\omega$	$-i\sqrt{3}\omega^2$	$i\sqrt{3}\omega^2$	$-i\sqrt{3}\omega$	$i\sqrt{3}$	$-i\sqrt{3}$	0	0
$C_{13}$	3	3	$i\sqrt{3}\omega^2$	$-i\sqrt{3}\omega$	$i\sqrt{3}$	$-i\sqrt{3}$	$i\sqrt{3}\omega$	$-i\sqrt{3}\omega^2$	0	0
$C_{14}$	9	9	0	0	0	0	0	0	0	0
$C_{15}$	9	9	0	0	0	0	0	0	0	0
$C_{16}$	9	9	0	0	0	0	0	0	0	0
$C_{17}$	9	9	0	0	0	0	0	0	0	0

Table 2: Character table of  $\Sigma(81)$  for the 8 three-dimensional representations.

The matrices of the  ${\bf 3_A}$  representation of  $\Sigma(81)$  are given by

$$\begin{array}{l} C_{1} : \left( \begin{array}{c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right), \quad C_{2} : \left( \begin{array}{c} \omega & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega \end{array} \right), \quad C_{3} : \left( \begin{array}{c} \omega^{2} & 0 & 0 \\ 0 & \omega^{2} & 0 \\ 0 & 0 & \omega^{2} \end{array} \right), \quad (24) \\ C_{4} : \left( \begin{array}{c} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^{2} \end{array} \right), \quad \left( \begin{array}{c} \omega^{2} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \omega \end{array} \right), \quad \left( \begin{array}{c} \omega^{2} & 0 & 0 \\ 0 & 0 & \omega \end{array} \right), \quad \left( \begin{array}{c} \omega & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^{2} \end{array} \right), \quad (25) \\ C_{5} : \left( \begin{array}{c} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & \omega & 0 \\ 0 & 0 & \omega \end{array} \right), \quad \left( \begin{array}{c} 0 & \omega^{2} & 0 \\ 0 & 0 & \omega^{2} \end{array} \right), \quad \left( \begin{array}{c} 0 & 1 & 0 \\ 0 & 0 & \omega^{2} \end{array} \right), \quad (26) \\ C_{6} : \left( \begin{array}{c} 0 & 1 & 0 \\ 0 & 0 & \omega^{2} \end{array} \right), \quad \left( \begin{array}{c} 0 & 1 & 0 \\ 0 & 0 & \omega^{2} \end{array} \right), \quad \left( \begin{array}{c} 0 & \omega^{2} & 0 \\ 0 & 0 & \omega^{2} \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0 & 0 & \omega^{2} \\ \omega^{2} & 0 & 0 \end{array} \right), \quad \left( \begin{array}{c} 0$$

$$\begin{array}{l} C_{12} : \begin{pmatrix} \omega^2 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, \begin{pmatrix} \omega^2 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, \begin{pmatrix} \omega & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega^2 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \omega^2 \end{pmatrix}, \begin{pmatrix} 0 & \omega^2 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & \omega^2 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & \omega^2 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & \omega^2 \\ 0 & 0 & \omega^2 \\ \omega^2 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & \omega^2 \\ 0 & 0 & \omega^2 \\ 0 & \omega & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & \omega^2 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \omega^2 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & \omega^2 \\ 0 & 0 & 0 \\ 0 & \omega^2 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \omega^2 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \omega^2 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \omega^2 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \omega & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \omega & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \omega & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \omega & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 &$$

## References

- [1] Y. Koide, Lett. Nuovo Cimento **34**, 201 (1982).
- [2] Y. Koide, Phys. Rev. **D28**, 252 (1983).
- [3] Y. Koide, Mod. Phys. Lett. A5, 2319 (1990).
- [4] For a recent review and other references, see Y. Koide, hep-ph/0506247.
- [5] Y. Koide, Phys. Rev. **D73**, 057901 (2006).
- [6] Y. Koide, hep-ph/0605074.
- [7] E. Ma, Phys. Lett. **B632**, 352 (2006).
- [8] G. C. Branco, J.-M. Gerard, and W. Grimus, Phys. Lett. 136B, 383 (1984).
- [9] I. de Medeiros Varzielas, S. F. King, and G. G. Ross, hep-ph/0607045.
- [10] E. Ma, Mod. Phys. Lett. **A21**, 1917 (2006).