WHY THE $SU(\infty)$ DECONFINING PHASE TRANSITION MIGHT BE OF SECOND ORDER ¹

ROBERT D. PISARSKI AND MICHEL TYTGAT

Dept. of Physics, Brookhaven National Lab Upton, NY 11973 USA

Abstract

Based upon what is known about the phase transition(s) of an SU(3) gauge theory, we argue that in a $SU(N_c)$ gauge theory without quarks, at nonzero temperature the deconfining phase transition is of second order when $N_c \geq 4$.

From 't Hooft, Witten, and others, it is known that at zero temperature, many properties of QCD can be understood by assuming that for a $SU(N_c)$ gauge theory, $N_c = 3$ is reasonably well described by the limit of $N_c = \infty$ [1]. This includes such features as the OZI rule, the approximate mass degeneracy between the ρ and ω vector mesons, *etc.* [2].

In this note we suggest that the large N_c expansion can also provide a way of understanding the phase diagram of a SU(3) gauge theory at nonzero temperature *if* the phase transition in a "pure" $SU(N_c)$ gauge theory (without dynamical quark fields) is of *second* order whenever $N_c \ge 4$, including $N_c = \infty$.

Simply counting the number of degrees of freedom allows one to make extremely strong statements about the thermodynamics of an $SU(N_c)$ gauge theory [3]-[6]. The basic point is simply that because gluons are in the adjoint representation, and quarks in the fundamental, at large N_c what happens to the $\sim N_c^2$ gluons totally dominates the $\sim N_c$ quarks. In the confined phase, confinement implies that all states are bound into colorless hadrons, so that the free energy is of necessity ~ 1 . At high temperature, the free energy is expected to be $\sim N_c^2$. Thus, as pointed out first by Thorn [3, 4], one can use the free energy itself to define the deconfining phase transition,

$$F(T) \sim 1$$
 , $T \le T_d$; $F(T) \sim N_c^2$, $T \ge T_d$. (1)

(See, however, [6].) In general, the deconfining phase transition is rigorously related to the global $Z(N_c)$ symmetry of a $SU(N_c)$ gauge theory, where the

¹To appear in the proceedings of the XXV Hirschegg Workshop on "QCD Phase Transitions", Jan. 1997.

order parameter is the Wilson line [7, 8],

$$L(\vec{x}) = tr\left(\mathcal{P}\exp\left(ig\int_{0}^{1/T}A_{0}(\vec{x},\tau)d\tau\right)\right) .$$
(2)

We assume that the $Z(N_c)$ symmetry is broken above T_d , $\langle L \rangle = 0$ for $T \leq T_d$, $\langle L \rangle \neq 0$ for $T \geq T_d$, which is most reasonable.

For n_f flavors of massless quarks, the analysis of the chiral phase transition proceeds as usual [9]. The only difference is that since the effects of the axial anomaly are $\sim g^2 n_f \sim (g^2 N_c) n_f / N_c$, if n_f and $g^2 N_c$ are held fixed as $N_c \to \infty$, then the effects of the anomaly vanish, and the global chiral symmetry is $S(U(n_f) \times U(n_f))$. We assume this is broken to $SU(n_f)$ at zero temperature [1], and restored at a temperature T_{χ} . Whatever the order of the chiral transition, however, since that part of the free energy is again $\sim N_c$, it cannot affect T_d . Thus at infinite N_c we can precisely characterize both the chiral and deconfining phase transitions. For the purposes of argument we take $T_{\chi} = T_d$.

Suppose now that the deconfining phase transition is of *first* order. Whatever the nature of the chiral transition, if we hold n_f finite as $N_c \to \infty$, the gluons dominate, and for any number of quark flavors, the first order deconfining transition always wins. Near T_d , the effective three dimensional theory for $L(\vec{x})$ is

$$\mathcal{L} = \frac{1}{2} |\partial_{\alpha}L|^2 + \frac{1}{2} m^2 |L|^2 + g_4 (|L|^2)^2 + g_6 (|L|^2)^3 .$$
(3)

A first order transition implies that the quartic coupling is negative, $g_4 < 0$. This is possible because the most general renormalizable theory in three dimensions includes a six-point coupling, which for stability must be positive, $g_6 > 0$.

At present, numerical simulations of an SU(3) gauge theory demonstrate the following [10]. In the pure gauge theory, if ϵ is the energy density, the latent heat is relatively weak [11],

$$\frac{\Delta\epsilon}{\epsilon}\Big|_{T_d} \le \frac{1}{3} \,. \tag{4}$$

If dynamical fermions are added, the deconfining transition can be washed out entirely; in the "Columbia" phase diagram, as a function of $m_u = m_d$ and m_s , there is a clean gap separating the regions of a first order deconfining phase transition, for large quark masses, and a first order chiral phase transition, for small quark masses [10].

These features of the $N_c = 3$ phase diagram are difficult to understand if the large N_c expansion is a reasonable guide, and if the deconfining phase transition for $N_c = \infty$ is strongly first order. One would expect that the deconfining phase transition would be strongly first order at $N_c = 3$, and that the deconfining transition would dominate for all quark masses. Thus there would be no need to draw the Columbia phase diagram — the transition would always be first order.

On the other hand, if the deconfining phase transition is of *second* order, then at least in a handwaving sense, everything seems to fit. Without dynamical fermions, at $N_c = 3$ the latent heat is small, down by $\sim 1/N_c \sim 1/3$. Further, since the deconfining phase transition is relatively weak to begin with, it is easily washed out by dynamical quarks. The chiral transition is of first order for three massless flavors, and about that point, but that is special to the chiral transition.

Moreover, the large N_c expansion does provide an understanding of one very familiar feature of the phase transition in an SU(3) gauge theory: the large increase in entropy. This is due, naturally, to the vast increase in the number of degrees of freedom between the hadronic and deconfined phases. But this terminology only makes sense if we can speak of a deconfined phase. Why there is such a large increase in entropy for small quark masses, when the transition is manifestly dominated by the chiral properties? No effective model of the chiral transition will produce such a large jump in entropy, simply because there is no great change in the number of (light) degrees of freedom. If we think of a second order transition for $N_c = \infty$, though, we automatically get a large increase in entropy. Not a jump, just an smooth but sharp increase.

We acknowledge that our suggestion contradicts known results from lattice gauge theory. Using the Eguchi-Kawai approximation to large N_c [12], under the assumption that the coupling between spacelike plaquettes can be neglected, Gocksch and Neri proved that the deconfining phase transition is of first order [13, 14]. (See, however, [15].) We note that a different approach to large N_c by Yaffe *et al.* [16] appears to indicate that the deconfining transition is of second order at $N_c = \infty$ [17].

Numerical simulations of a SU(4) lattice gauge theory have been done [18], and indicate a first order deconfining phase transition. Here we can only suggest that perhaps what was observed is a bulk transition, separate from the true deconfining phase transition at nonzero temperature.

In that regards, we would also like to make a technical aside about the deconfining phase transition for $N_c = 4$. For arbitrary N_c , the general effective lagrangian includes (3), which possesses a global O(2) symmetry, and the term

$$\mathcal{L}_Z = g_Z \left(L^{N_c} + (L^*)^{N_c} \right) \ . \tag{5}$$

which is only invariant under global $Z(N_c)$ transformations. For $N_c = 2$ this changes the symmetry from O(2) to Z(2). When $N_c = 3$ this is a cubic coupling, and drives the deconfining phase transition first order [7].

For $N_c = 4$, the coupling g_Z in \mathcal{L}_Z is as important as g_4 in \mathcal{L} , and one must be more careful. In particular, there is the possibility that having both g_Z and $g_4 \neq 0$ produces a fluctuation induced first order transition. For $N_c = 4$, the theory with the lagrangian $\mathcal{L} + \mathcal{L}_Z$ is equivalent to what is known as the n = 2 model of cubic anisotropy. In $4 - \epsilon$ dimensions, the O(2) fixed point is infrared stable [19]. This is supported by recent Monte Carlo simulations directly in three dimensions [20]. Hence, for $N_c = 4$, it does not appear as if the transition is fluctuation induced first order, and is of second order when g_4 and g_Z are positive. Of course the deconfining phase transition for SU(4)could still be first order because the couplings g_4 and/or g_Z are negative to begin with.

For $N_c = 5$, one expects that the pentic coupling can be neglected relative to the quartic. This must be qualified: precisely at the tricritical point, where $m^2 = g_4 = 0$, there is only a pentic and a hexic coupling. If $g_Z \neq 0$, the pentic coupling wins, and because it is odd in ϕ , drives the transition first order. This remains true in a region around the tricritical point, such as for $m^2 = 0$ and small $g_4 > 0$, when the dimensionless quantity $g_Z/\sqrt{g_4} \gg 1$. If $g_Z/\sqrt{g_4} \ll 1$, however, then the pentic coupling is negligible relative to the quartic, and the transition is of second order. This is in contrast to $N_c = 3$, where the cubic coupling $g_Z \neq 0$ drives the transition first order regardless of the magnitude of $g_Z/|g_4|^{3/2}$: it is weakly first order for $g_Z/|g_4|^{3/2} \ll 1$, but strongly first order when $g_Z/|g_4|^{3/2} \gg 1$. The difference is because the quartic coupling is less relevant than a cubic, but more relevant than a pentic.

For $N_c \geq 6$, we can certainly neglect the $Z(N_c)$ coupling. Thus we see that with some technical qualifications, that the conclusions of [7] remain: for $g_4 \geq 0$, when $N_c \geq 4$ the $SU(N_c)$ transition is second order, in the universality class of an O(2) model.

We conclude with two suggestions.

The first is to measure the coupling g_4 for SU(3) and see if it is positive. For a pure SU(2) gauge theory, it appears as if the deconfining phase transition is of second order [21] (see, however, [22]), which implies that the coupling g_4 is positive. Thus it would be interesting to know if $g_4 > 0$ for SU(3); if so, it would be reasonable to assume that $g_4 > 0$ for all N_c ; this implies that the deconfining phase transition is of second order whenever $N_c \ge 4$.

A second suggestion is simply to carefully measure again the deconfining phase transition for SU(4). Certainly $N_c = 4$ is closer to $N_c = \infty$ than $N_c = 3$. Of course our arguments are most indirect, with many obvious loopholes: large N_c may not describe thermodynamics for $N_c = 3$; the $N_c = \infty$ transition might be weakly first order (but then - why?); and so on. Still, a second order phase transition at $N_c = \infty$ helps one understand understand many qualitative features of the phase diagram at $N_c = 3$.

We thank Prof.'s Billó, Caselle, D'Adda, Panzeri, and Yaffe for discussions on large N_c , and Prof. Rajagopal for a comment on $N_c = 5$. This work is supported by a DOE grant at Brookhaven National Laboratory, DE-AC02-76CH00016.

References

- G. 't Hooft, Nucl. Phys. B72 (1974) 461; *ibid.* B75 (1974) 461; E. Witten, *ibid.* B156 (1979) 269; *ibid.* B160 (1979) 57; S. Coleman and E. Witten, Phys. Rev. Lett. 45 (1980) 100.
- [2] M. Teper, hep-lat/9701003, hep-lat/9701004.
- [3] C. B. Thorn, Phys. Lett. **B99** (1981) 458.
- [4] R. D. Pisarski, Phys. Rev. **D29** (1984) 1222.
- [5] J. J. Atick and E. Witten, Nucl. Phys. B310 (1988) 291; J. Polchinski, Phys. Rev. Lett. 68 (1992) 1267; T. H. Hansson and I. Zahed, Phys. Lett. B309 (1993) 385.
- [6] L. D. McLerran and A. Sen, Phys. Rev. **D32** (1985) 2794.
- [7] B. Svetitsky and L. G. Yaffe, Nucl. Phys. **B210** [FS6] (1982) 423.
- [8] H. Meyer-Ortmanns, Rev. Mod. Phys. 68 (1996) 473; A. Smilga, hepph/9612347.
- [9] R. D. Pisarski and F. Wilczek, Phys. Rev. D29 (1984) 338; K. Rajagopal and F. Wilczek, Nucl. Phys. B399 (1993) 395; *ibid.*, B404 (1993) 577; S. Gavin, A. Gocksch, and R. D. Pisarski, Phys. Rev. D49 (1994) R3079; K. Rajagopal, in *Quark-Gluon Plasma 2*, ed. R. Hwa (World Scientific, Singapore, 1995).
- [10] F. R. Brown, F. P. Butler, H. Chen, N. H. Christ, Z. Dong, W. Schaffer, L. I. Unger, A. Vaccarino, Phys. Rev. Lett. 65 (1990) 2491; F. Karsch, Nucl. Phys. A590 (1995) 367c; E. Laermann, proceedings of Quark Matter '96; A. Ukawa, hep-lat/9612011.
- [11] F. Karsch, B. Beinlich, J. Engels, R. Joswig, E. Laermann, A. Peikert, B. Petersson, hep-lat/9608047; F. Karsch, proceedings in this volume.

- [12] T. Eguchi and H. Kawai, Phys. Rev. Lett. 48 (1982) 1063; A. A. Migdal, Phys. Rep. 102 (1983) 199.
- [13] A. Gocksch and F. Neri, Phys. Rev. Lett. **50** (1983) 1099.
- [14] F. Neri and A. Gocksch, Phys. Rev. D28 (1983) 3147; S. R. Das and J. B. Kogut, Phys. Lett. 145B (1984) 375; Nucl. Phys. B257 [FS14] (1985) 141; A. Gocksch and M. Ogilvie, Phys. Rev. Lett. 54 (1985) 1772; S. R. Das, Rev. Mod. Phys. 59 (1987) 235; M. Caselle, A. D'Adda, and S. Panzeri, Phys. Lett. B293 (1992) 161; *ibid.* B302 (1993) 80; M. Billó, M. Caselle, A. D'Adda, L. Magnea, and S. Panzeri, Nucl. Phys. B435 (1995) 172; M. Billó, M. Caselle, A. D'Adda, S. Panzeri, hep-th/9610144; M. Billó, M. Caselle, A. D'Adda, hep-th/9701145.
- [15] F. R. Klinkhamer, Phys. Rev. **D29** (1984) 1795.
- [16] F. R. Brown, L. G. Yaffe, Nucl. Phys. B271 (1986) 267; T. A. Dickens,
 U. J. Lindqwister, W. R. Somsky, L. G. Yaffe, *ibid.* B309 (1988) 1.
- [17] W. R. Somsky, Princeton Univ. Thesis, 1988; L. G. Yaffe, private communication.
- [18] G. G. Batrouni and B. Svetitsky, Phys. Rev. Lett. **52** (1984) 2205; A. Gocksch and M. Okawa, *ibid.* **52** (1984) 1751; F. Green, Phys. Rev. **D29** (1984) 2986; J. F. Wheater and M. Gross, Phys. Lett. **B144** (1984) 409; Nucl. Phys. **B240** (1984) 253.
- [19] J. Rudnick, Phys. Rev. B11 (1975) 3397; D. J. Amit, Field Theory, the Renormalization Group, and Critical Phenomena, (World Scientific, Singapore, 1984), Part II, sec. 4.2; S. Midorikawa, H. So, and S. Yoshimoto, Z. Phys. C34 (1987) 307.
- [20] P. Arnold and L. G. Yaffe, hep-ph/9610447; P. Arnold and Y. Zhang, hep-ph/9610448; P. Arnold, S. R. Sharpe, L. G. Yaffe, Y. Zhang, hepph/9611201.
- [21] J. Engels, J. Fingberg, and D. E. Miller, Nucl. Phys. B387 (1992) 501;
 J. Fingberg, U. Heller, and F. Karsch, Nucl. Phys. *ibid.* (1993) 493; G. Cella, G. Curci, R. Tripiccione, A. Vicere, Phys. Rev. D49 (1994) 511; J. Engels and T. Scheideler, hep-lat/9610019.
- [22] R. V. Gavai, M. Grady, M. Mathur, Nucl. Phys. B423 (1994) 123;
 M. Mathur and R. V. Gavai, *ibid.* B448 (1995) 399; R. V. Gavai and M.Mathur, hep-lat/9512015.