Supersymmetry Breaking Without A Messenger Sector

Naoyuki Haba¹, Nobuhito Maru², and Takeo Matsuoka³

Department of Physics, Nagoya University Nagoya, JAPAN 464-01

Abstract

We propose a simple gauge-mediated supersymmetry breaking model in which suitable soft breaking masses are dynamically generated without relying on a messenger sector. This model is constructed as an extention of the 3 -2 model and needs no fine-tuning of parameters. The dynamical supersymmetry breaking sector contains non-renormalizable interactions and vector-like fields. Non-renormalizable terms are characterized by the couplings of $O(1)$ in units of the Planck scale. Hierarchy of various mass scales in the model arises from the non-renormalizable terms. The vacuum of this model conserves the color and electro-weak symmetry. We also discuss a possible solution to the μ problem in this framework.

¹E-mail: haba@eken.phys.nagoya-u.ac.jp, Address after April 1, 1997: Faculty of Engineering, Mie University, Mie, JAPAN 514

²E-mail: maru@eken.phys.nagoya-u.ac.jp

³E-mail: matsuoka@eken.phys.nagoya-u.ac.jp

1 Introduction

The dynamical supersymmetry breaking (DSB) provides an attractive explanation of the large hierarchy between the Planck scale and the electro-weak scale. Many modelswhich exhibit the DSB have been proposed $[1, 2, 3, 4]$ $[1, 2, 3, 4]$ $[1, 2, 3, 4]$ $[1, 2, 3, 4]$. Since these models by themselves have nothing to do with our visible world, we need some mechanisms of mediating supersymmetry breaking to the low-energy observable sector. For phenomenological implications it is of great importance to clarify how the supersymmetry breaking is communicated to the visible sector. Two ways of transmitting supersymmetry breaking to the visible sector have been proposed. The first is that gravitational interactions in $N = 1$ supergravity theory play its role. The second is that gauge interactions play a role of the messenger. As for the latter one, the scenario given in Ref. [\[1\]](#page-15-0) suffers from a problem that the QCD coupling becomes strong a few decades above the scale of the supersymmetry breaking. This difficulty originates from the situation of the model in which the standard model gauge group $G_{st} \equiv SU(3)_c \times SU(2)_L \times U(1)_Y$ is taken as the global symmetry of the DSB sector. In the model given in Refs.[[5\]](#page-15-0), this difficulty was avoided by introducing the "messenger sector" which connects the DSB sector with the standard model. In the messenger sector scenario the supersymmetry is broken at the energy as low as $O(100 \text{ TeV})$. In such a model several attractive features are derived. Namely, we obtain enough degeneracy among squarks and sleptons to ensure adequate suppression of flavor changing neutral currents (FCNC). Further, no new CP phases are induced in soft supersymmetry breaking parameters. This implies the absence of large particle electric dipole moment (EDM).

Recently, however, it has been pointed out that in the original model of the messenger sector scenario QCD color symmetry is not conserved[[6\]](#page-15-0). In Ref. [\[5\]](#page-15-0), the superpotential in the messenger sector is assumed to be

$$
W_{\text{mess}} = k_1 \phi^+ \phi^- S + \frac{1}{3} \lambda S^3 + k_3 S q \bar{q} + k_4 S l \bar{l}, \qquad (1.1)
$$

where the vector-like multiplets q, \bar{q} transform as 3, 3^{*} under color $SU(3)_C$ and l, \bar{l} transform as 2 under $SU(2)_L$. S is a singlet and ϕ^{\pm} have ± 1 charges of the messenger gauge group $U(1)_m$. Since the soft supersymmetry breaking masses squared for ϕ^+ and ϕ^- become negative, ϕ^+ and ϕ^- develop non-zero vacuum expectation values (VEVs). If the minimization of the scalar potential yields non-vanishing $\langle S \rangle$ and $\langle F_S \rangle$, the supersymmetry breaking is transmitted to the singlet S by the interaction $\phi^+ \phi^- S$ in Eq. (1.1). This is a crucial point for communicating supersymmetry breaking to the visible sector since gauginos, squarks and sleptons masses are radiatively induced by the interactions $Sq\bar{q}$ and $S_l\bar{l}$. However, in the true vacuum $\langle S \rangle$ and $\langle F_s \rangle$ vanish and at the same time QCD color is violated[[6](#page-15-0)].

Many attempts have been made in order to resolve the problem [\[7](#page-16-0), [8](#page-16-0)]. In the previous paper[[8\]](#page-16-0), the authors proposed a simple model in which the messenger gauge interaction is not necessarily needed and the vacuum does not break the color and electro-weak symmetry. So the model is free from the difficulty encountered in the original messenger sector scenario. In the model the effective theory from the DSB sector serves as the messenger sector and non-vanishing VEVs of S and F_S are ensured by the dynamics in the DSB sector. Thus, supersymmetry breaking is transmitted from DSB sector to the visible sector through the interactions $Sq\bar{q}$ and *Sll*. No elaborate messenger sector is needed to achieve the feed-down of the supersymmetry breaking. The strategy of this paper is to construct the model which satisfy the following requirements:

1. No messenger gauge interactions.

- 2. Communication of supersymmetry breaking to the visible sector without violating color symmetry.
- 3. The soft supersymmetry breaking masses coming predominantly from the standard model gauge interaction rather than the gravitational interaction.
- 4. No fine-tuning of parameters or very small coupling constants.
- 5. No μ problem.

Although the model proposed in the previous paper[[8\]](#page-16-0) satisfies the first three among the above requirements, the model relies on an extremely small coupling constant and a solution to the μ problem is not presented.

In this paper we propose a renewed model which is in line with the above five requirements. The superpotential in the present model consists of non-renormalizable interactions. Since the superpotential involves a scale explicitly, the present model, in a sense, seems to break the philosophy in Ref. [\[5](#page-15-0)] that all masses should arise through dimensional transmutation. However, in the case that the suppression factor of non-renormalizable terms is the Planck scale, the model does not contradict with the philosophy. This is because there exists the supergravity for the underlying theory. If we do not take the supergravity into account, models of the DSB suffer from a massless Goldstone fermion and a massless R-axion. A massless Goldstone fermion, which appears as a result of the spontaneous supersymmetry breaking, is absorbed by the gravitino. An R-axion can gain a mass by breaking the R symmetry explicitly to cancel the cosmological constant[[9](#page-16-0)]. Although soft masses are induced by both gravitational interactions and gauge interactions, the contribution of gauge interactions is dominant in the present model. The supersymmetry breaking scale is the intermediate scale of order $10^{10} \,\text{GeV}$.

This paper is organized as follows. After explaining the model proposed in the previous paper [\[8](#page-16-0)] in Section 2, we proceed in Section 3 to exhibit a renewed model. A possible solution to the μ problem is discussed in Section 4. Section 5 is devoted to summary.

2 Fine-tuning in the DSB models

To begin with, let us briefly review the parts of the 3 -2 model which are relevant to our study. The 3 -2 model is the simplest calculable model with the dynamical supersymmetrybreaking [[1\]](#page-15-0). The gauge group of this model is $SU(3) \times SU(2)$. This model contains chiral superfields

$$
Q(3,2)_{1/3}
$$
, $\overline{U}(3^*,1)_{-4/3}$, $\overline{D}(3^*,1)_{2/3}$, $L(1,2)_{-1}$, (2.1)

where the subscript denotes the quantum number of a global $U(1)$ symmetry of the model ⁴ . The superpotential of the 3 -2 model is given by

$$
W_{3-2} = \lambda_1 X_1 + \frac{\Lambda^7}{X_3},\tag{2.2}
$$

where X_i 's are gauge invariant combinations

$$
X_1 \equiv Q\overline{D}L \ , \ X_2 \equiv Q\overline{U}L \ , \ X_3 \equiv \det Q\overline{Q} \ . \tag{2.3}
$$

Here we use the notation $Q \equiv (U, D)$ and $\overline{Q} \equiv (\overline{U}, \overline{D})$. The first term of W_{3-2} is the most general renormalizable superpotential which is invariant under the symmetries. A coupling constant λ_1 is assumed to be very small so as $\lambda_1^{1/7} \ll 1$. The second term in Eq. (2.2) is dynamically generated by the one-instanton effect and yields the runaway behavior of the scalar potential. This runaway behavior is stabilized due to

 4 In the ordinary "messenger sector" scenario this $U(1)$ symmetry is localized to mediate the supersymmetry breaking [\[5](#page-15-0)]. The present model, however, does not need to localize it.

the lifting effect of the first term. The smallness of λ_1 ensures that the superfields develop large VEVs while maintaining only a small vacuum energy. Relying on a very small coupling is shared in common with a number of models which invoke the runaway-behaving potential. Now we consider the situation that the $SU(3)$ coupling is larger than the $SU(2)$ coupling. A denotes the scale at which the $SU(3)$ coupling becomeslarge ⁵. From the superpotential (2.2) (2.2) , VEVs of elementary fields $Q, L, ...$ are estimated to be of order $\Lambda/\lambda_1^{1/7} (\equiv v)$. Since we have the relation $v \gg \Lambda$, gauge couplings are effectively weak at the scale v . Therefore, effective fields X_i 's can be regarded as the product of elementary fields and Kähler potential has the canonical form for elementary fields. As F-terms of elementary fields turn out to be of $O(\lambda_1 v^2)$, the supersymmetry breaking occurs in the 3 -2 model. After integrating out the heavy degrees of freedom at the scale Λ , we can construct the effective theory below the scale Λ which is described in terms of the light fields X_i 's. The Kähler potential in theeffective theory is given by $[1, 9]$ $[1, 9]$ $[1, 9]$

$$
K = 3\left(t + \frac{B}{t}\right),\tag{2.4}
$$

where

$$
t \equiv (A + \sqrt{A^2 - B^3})^{1/3} + (A - \sqrt{A^2 - B^3})^{1/3},
$$

\n
$$
A \equiv \frac{1}{2} (X_1^{\dagger} X_1 + X_2^{\dagger} X_2),
$$

\n
$$
B \equiv \frac{1}{3} \sqrt{X_3^{\dagger} X_3}.
$$
\n(2.5)

In the previous paper [\[8\]](#page-16-0) it is assumed that the superpotential is of the form

$$
W = W_{3-2} + W_m + \frac{\lambda_2'}{M^5} X_2 \overline{E} X_3
$$
\n(2.6)

with

$$
W_m = \frac{h_1}{M^2} X_1 q \overline{q} + \frac{h_2}{M^2} X_1 l \overline{l} \ . \tag{2.7}
$$

⁵ If we consider the case where $SU(2)$ dynamics is dominant compared to $SU(3)$ dynamics, supersymmetry is broken due to the quantum deformation of the moduli space[[4\]](#page-15-0).

Here M is set equal to the reduced Planck scale M_{Planck} . Vector-like fields q, \overline{q} and l, \overline{l} are singlets under the 3-2 model gauge group $SU(3) \times SU(2)$ but charged under the standard model gauge group $G_{st} = SU(3)_c \times SU(2)_L \times U(1)_Y$ as

$$
q\ (3,1,-2/3),\ \ \overline{q}\ (3^*,1,2/3),\ \ l\ (1,2,1),\ \ \overline{l}\ (2,1,-1)\ .
$$
 (2.8)

Note that the global $U(1)$ symmetry offered in Eq. (2.1) is gauged and the singlet superfield \overline{E} is included to cancel the gauged $U(1)$ anomaly. Further, a non-anomalous $U(1)_R$ symmetry is introduced. However, the $U(1)$ gauge interaction and the last term in Eq.([2.6\)](#page-5-0) do not play an essential role in communicating the supersymmetry breaking.The fields X_1 and X_3 correspond to the singlet field S in Ref. [[5](#page-15-0)]. The above superpotential yields non-vanishing VEVs for X_1 , X_3 and F_{X_1} , F_{X_3} in the true vacuum, which are

$$
\langle X_1 \rangle = O(v^3), \qquad \langle F_{X_1} \rangle = O(\lambda_1 v^4), \tag{2.9}
$$

$$
\langle X_3 \rangle = O(v^4), \qquad \langle F_{X_3} \rangle = O(\lambda_1 v^5). \tag{2.10}
$$

Confronting the result with phenomenological constraints on gaugino, squark and slepton masses, the authors have obtained[[8\]](#page-16-0)

$$
10^{14} \,\text{GeV} < v < 10^{16} \,\text{GeV},\tag{2.11}
$$

$$
10^{-11} < \lambda_1 < 10^{-9}.\tag{2.12}
$$

This means that the model needs an unnatural fine-tuning of the parameter λ_1 . In order to find a phenomenologically viable model it is necessary to resolve this problem.

3 A new model

We now proceed to propose a new model, in which there is no fine-tuning of parameters. We introduce a global $U(1)_R$ symmetry but not the messenger gauge interaction. The $U(1)_R$ charges are assigned to each chiral superfield as shown in Table I. Under this $U(1)_R$ symmetry the first term in $W₃₋₂$ is forbidden. Instead, in the new model the superpotential is written as

$$
W = W_{SB} + W_m \tag{3.1}
$$

with

$$
W_{SB} = \frac{\lambda}{M^4} X_1 X_3 + \frac{\Lambda^7}{X_3},
$$
\n(3.2)

$$
W_m = \frac{h_1}{M^3} X_3 q \overline{q} + \frac{h_2}{M^3} X_3 l \overline{l} \,, \tag{3.3}
$$

where λ and h_i 's are coupling constants taken to be of $O(1)$ and $M = M_{\text{Planck}}$. The scale Λ at which the $SU(3)$ gauge coupling becomes strong is expected to be sufficiently small compared with M.

Table I

The D-flat direction of the $SU(3) \times SU(2)$ is given by

$$
\langle Q \rangle = \begin{pmatrix} b & 0 \\ 0 & a \\ 0 & 0 \end{pmatrix}, \qquad \langle \overline{U} \rangle = \begin{pmatrix} b & e^{i\theta} \\ 0 \\ 0 \end{pmatrix},
$$

$$
\langle \overline{D} \rangle = \begin{pmatrix} 0 \\ a & e^{i\delta} \\ 0 \end{pmatrix}, \qquad \langle L \rangle = (-\sqrt{a^2 - b^2} \ e^{i\eta}, 0).
$$
 (3.4)

Parameters a and b are real and positive with $a > b$. The symbols θ , δ , and η stand for relative phases. Along the D-flat direction the scalar potential is expressed as

$$
V = W_i \left(K^{-1} \right)_i \frac{1}{j} W_j^*, \tag{3.5}
$$

where $W_i \equiv \partial W / \partial X_i$ and $K_i \bar{j} \equiv \partial^2 K / \partial X_i \partial X_j^*$. The explicit form of V is

$$
V = \frac{2}{t} \left| \frac{2\lambda}{M^4} X_1 X_3 - \frac{\Lambda^7}{X_3} + \frac{h_1}{M^3} X_3 q \overline{q} + \frac{h_2}{M^3} X_3 l \overline{l} \right|^2
$$

$$
+ t^2 \left(\frac{\lambda}{M^4}\right)^2 |X_3|^2 + 6B t \left|\frac{\lambda}{M^4} X_1 - \frac{\Lambda^7}{X_3^2} + \frac{h_1}{M^3} q\overline{q} + \frac{h_2}{M^3} l\overline{l}\right|^2
$$

+
$$
\left(\frac{h_1}{M^3}\right)^2 |X_3|^2 \left(|q|^2 + |\overline{q}|^2\right) + \left(\frac{h_2}{M^3}\right)^2 |X_3|^2 \left(|l|^2 + |\overline{l}|^2\right). \tag{3.6}
$$

Here we denote the scalar components of the superfield by the same letters as the superfield itself. It can be easily shown that at the minimum point of V we have vanishing VEVs of q, \overline{q} and l, \overline{l} . This is due to the fact that

$$
\lambda \left(\frac{v}{M}\right)^4 \ll h_1 \left(\frac{v}{M}\right)^3, \quad h_2 \left(\frac{v}{M}\right)^3,\tag{3.7}
$$

where the scale v is redefined as

$$
v = \left(\frac{\Lambda^7 M^4}{\lambda}\right)^{1/11} \tag{3.8}
$$

in the new model and $\Lambda \ll v \ll M$. Further, since V takes the minimum value along the direction $\cos(2\theta + 3\delta + \eta) = 1$, we can put $\theta = \delta = \eta = 0$ without loss of generality. The minimization of V can be carried out numerically and results in

$$
a = 1.072 \times v, \t b = 0.897 \times v.
$$
 (3.9)

In addition, we have the VEVs

$$
\langle X_1 \rangle = a^2 \sqrt{a^2 - b^2} = 0.674 \times v^3, \tag{3.10}
$$

$$
\langle X_3 \rangle = a^2 b^2 = 0.924 \times v^4 \tag{3.11}
$$

and

$$
\langle F_{X_1} \rangle = 1.412 \times \lambda \frac{v^8}{M^4},\tag{3.12}
$$

$$
\langle F_{X_3} \rangle = -0.396 \times \lambda \frac{v^9}{M^4}.
$$
\n(3.13)

On the other hand, $\langle X_2 \rangle$ and $\langle F_{X_2} \rangle$ vanish. Then we obtain the ratios

$$
\frac{\langle F_{X_1} \rangle}{\langle X_1 \rangle} = 2.096 \times \lambda \frac{v^5}{M^4},\tag{3.14}
$$

$$
\frac{\langle F_{X_3} \rangle}{\langle X_3 \rangle} = -0.428 \times \lambda \frac{v^5}{M^4}.
$$
\n(3.15)

It is now understandable that the factor $\lambda(v/M)^4$ corresponds to the very small parameter λ_1 in the previous paper. The new model naturally explains the smallness of λ_1 .

Let us estimate gauge-mediated soft breaking masses of gauginos, squarks, and sleptons in the visible sector. Gaugino masses m_{λ_a} are induced at the one-loop level [[5\]](#page-15-0) and given by

$$
m_{\lambda_a} = \frac{g_a^2}{16\pi^2} \frac{\langle F_{X_3} \rangle}{\langle X_3 \rangle},\tag{3.16}
$$

where a denotes the index of G_{st} . Squark and slepton masses squared \tilde{m}^2 arise at the two-loop level [\[5](#page-15-0)] and are expressed as

$$
\tilde{m}^2 = \sum_a 2 C_F^a \left(\frac{g_a^2}{16\pi^2}\right)^2 \left(\frac{\langle F_{X_3} \rangle}{\langle X_3 \rangle}\right)^2, \tag{3.17}
$$

where C_F^a is 4/3 for $SU(3)_c$ triplets, 3/4 for $SU(2)_L$ doublets, and 3/5Y² for $U(1)_Y$. It is noteworthy that although the couplings $X_3q\overline{q}$ and $X_3l\overline{l}$ in W_m are suppressed by inverse powers of the scale M , soft breaking masses in the visible sector do not have any suppression factors of $1/M$. From Eqs. (3.14) and (3.16) , the relation

$$
\lambda \frac{v^5}{M^4} \simeq 10^{4 \sim 5} \,\text{GeV} \tag{3.18}
$$

should be satisfied so as to obtain appropriate soft masses of order $10^{2 \sim 3}$ GeV. By setting $\lambda = O(1)$ and $M = M_{Planck} = 10^{18.3}$ GeV, we are led to

$$
v \sim 10^{-2.8} \times M \simeq 10^{15.5} \,\text{GeV},\tag{3.19}
$$

which is around the GUT scale. This implies that

$$
\lambda \left(\frac{v}{M}\right)^4 \sim 10^{-11}, \qquad \frac{v}{\Lambda} \sim 40. \tag{3.20}
$$

Therefore, Λ is of order 10^{14} GeV and the perturbative picture is available.

In order to compare with the gauge-mediated contributions, we next estimate soft breaking masses of gauginos and sparticles transmitted by the gravitational interaction. Gaugino masses can arise from the operator

$$
\int d^2\theta \, \frac{X_1}{M^3} W_\alpha W^\alpha,\tag{3.21}
$$

where W^{α} is the field strength superfield of the standard model gauge group. Thus the gravitational contribution to gaugino masses becomes

$$
m_{\lambda} \simeq \frac{\langle F_{X_1} \rangle}{M^3} \simeq \lambda \frac{v^8}{M^7} \sim 10^{-4} \,\text{GeV}.\tag{3.22}
$$

Therefore, dominant contribution to gaugino masses comes from the standard model gauge interactions. On the other hand, soft masses squared of squarks and sleptons can arise from the operator

$$
\int d^4\theta \ c \frac{\phi^\dagger \phi}{M^2} \varphi_i^\dagger \varphi_i \tag{3.23}
$$

where ϕ is an elementary field of the DSB sector in Eq. ([2.1\)](#page-4-0), φ_i is the chiral superfield of the standard model and c is a constant with $c \leq 1$ which represents the elementary field content of the composite fields X_i . Thus, the gravitational contribution to squark and slepton masses are given by

$$
\tilde{m}^2 \sim \left(\frac{\langle F_{\phi} \rangle}{M}\right)^2 \simeq \left(\lambda \frac{v^6}{M^5}\right)^2 \sim \left(10^{1.5} \text{GeV}\right)^2,\tag{3.24}
$$

which are smaller than the gauge-mediated contribution by the factor $\sim 10^{-2}$.

In the present model mass spectra of other particles can be determined. Masses of vector-like fields q, \overline{q} and l, \overline{l} are of order $v^4/M^3 \sim 10^7 \,\text{GeV}$. Three scalars, two pseudoscalars, and one fermion of moduli fields X_i gain masses of order $v^5/M^4 \sim 10^4 \,\text{GeV}$. The fermion component of X_2 remains massless because the global $U(1)$ symmetry in the 3-2 sector is unbroken down to low energies. Since X_2 does not appear in the superpotential of Eq. (3.1) , the massless fermion of X_2 couples very weakly only through the Kähler potential. Thus the global $U(1)$ and the massless fermion are harmless to our visible sector. The mass of the R-axion is $O(10^3 \text{ GeV})$, which is free from the constraint of supernova astrophysical bounds. From the condition on vanishing cosmological constant, the gravitino mass is

$$
m_{3/2} \simeq \frac{\sqrt{|\langle F_{\phi} \rangle|^2}}{\sqrt{3}M} \simeq \frac{\lambda v^6}{\sqrt{3}M^5} \sim 10^{1.5} \,\text{GeV}.\tag{3.25}
$$

It is worth noting that the soft supersymmetry breaking parameters are blind to flavor at the intermediate scale $\Lambda \sim 10^{14}\,\text{GeV}$. As a consequence, enough degeneracy among squarks and sleptons is maintained at low energies and flavor changing neutral currents are naturally suppressed. As for the $SU(2)_L \times U(1)_Y$ breaking, it can be realized by the three-loop effects of the stop or by introducing another set of vectorlike fields in the visible sector [\[5](#page-15-0)].

Since the present model incorporates the Planck scale in non-renormalizable interactions, we should use the scalar potential of $N = 1$ supergravity for the correct discussion. However, even if we use the scalar potential of the supergravity, the above discussion remains unchanged.

4 The μ problem

In this section we explore a possible solution to the μ problem. As is well-known, phenomenological constraints on the Higgs potential including the vacuum stability condition require that the supersymmetric parameter μ and the soft supersymmetry breaking parameter B satisfy the condition

$$
B\mu \sim \mu^2 \sim (100 \,\text{GeV})^2. \tag{4.1}
$$

The minimal scheme of the messenger sector scenario has a difficulty in this regard. In the minimal messenger sector scenario, naively, the Higgs superpotential is written

$$
W_H = \lambda_H S H_u H_d, \qquad (4.2)
$$

where S is a gauge singlet superfield with the non-vanishing $\langle S \rangle$ and $\langle F_S \rangle$. It follows that $\mu = \lambda_H \langle S \rangle$ and $B\mu = \lambda_H \langle F_S \rangle$. Then we have $B = \langle F_S \rangle / \langle S \rangle$ irrespective of the magnitude of λ_H . However, the ratio $\langle F_S \rangle / \langle S \rangle$ has to be of order 10^{4∼5} GeV to generate appropriate soft masses in the minimal scenario. Thus we have too large B in disaccord with the above condition([4.1\)](#page-11-0).

Introducing an additional gauge singlet field N with the non-zero VEV but with the vanishing $\langle F_N \rangle$ as well as Higgs fields H_u and H_d in the present model, we assume an alternative superpotential for Higgs fields. Namely, we add the Higgs terms

$$
W_H = \left(\frac{k_1}{M}N^2 + \frac{k_2}{M^4}X_3N\right)H_uH_d\tag{4.3}
$$

to the superpotential, where constants k_1 and k_2 are $O(1)$. Under the condition $\langle N \rangle \gg \langle X_3 \rangle / M^3$, the superpotential W_H brings about

$$
\mu \simeq \frac{k_1}{M} \langle N \rangle^2, \tag{4.4}
$$

$$
B\mu \simeq \frac{k_2}{M^4} \langle F_{X_3} \rangle \langle N \rangle. \tag{4.5}
$$

If we take the value $\langle N \rangle \sim 10^{10.5}$ GeV as an ad hoc assumption here, then Eq. (4.4) yields $\mu = O(10^2 \text{ GeV})$. Substituting the relation $|\langle F_{X_3} \rangle| \simeq \lambda v^9 / M^4$ into Eq. (4.5), we obtain

$$
B\mu \simeq \frac{v^9}{M^8} \langle N \rangle. \tag{4.6}
$$

When we take Eq. [\(3.19\)](#page-9-0) and $\langle N \rangle \simeq 10^{10.5} \text{ GeV}$ into account, the numerical value becomes

$$
B\mu \sim \left(10^2 \,\text{GeV}\right)^2. \tag{4.7}
$$

This result is in accord with the above conditions (4.1) on μ and \ddot{B} . Further, the numerical value $\langle N \rangle \sim \sqrt{\mu M} \sim 10^{10.5} \text{ GeV}$ is also consistent with the condition

as

 $\langle N \rangle \gg \langle X_3 \rangle / M^3 (\sim 10^7 \text{ GeV})$. In this scenario it is essential that there exist not only the singlets X_1 and X_3 which communicate the supersymmetry breaking to the visible sector but also another singlet N with the non-zero VEV but with the vanishing $\langle F_N \rangle$.

Although in the above we put $\langle N \rangle \sim 10^{10.5} \, {\rm GeV}$ by hand, it is not unreasonable. This scale of $\langle N \rangle$ is nearly equal to $\langle X_1 \rangle/M^2 \sim 10^{10.5}$ GeV. It is plausible that the field N also corresponds to a composite operator. In the present model we have no such an operator fitting to the field N. However, when we take the gauge group larger than $SU(3) \times SU(2)$, the gauge singlet field fitting to the above N-field is possibly contained in the massless sector of the effective theory.

5 Summary

In this paper we proposed a new model in which suitable soft supersymmetry breaking masses are dynamically generated without the messenger sector. The model proposed here is free from the fine-tuning of coupling constants. The model is a simple extension of the 3 -2 model. The dynamical supersymmetry breaking sector contains extra vector-like quarks and leptons which have the gauge quantum number of the standard model. Non-renormalizable interactions play a crucial role of transmitting supersymmetry breaking to the visible sector through the gauge interaction. The non-renormalizable interactions are characterized by the couplings of $O(1)$ in units of the reduced Planck scale M_{Planck} . Hierarchy of various mass scales arises from the non-renormalizable terms. The scale of the DSB is of order $10^{14} \,\mathrm{GeV}.$ This model conserves the color symmetry. Although the supersymmetry breaking is transmitted by both gravity and gauge interactions, gaugino, squark, and slepton soft masses mainly come from the gauge-mediated contributions. Consequently, we obtain enough degeneracy between sfermions of the same gauge quantum numbers to ensure adequate suppression of FCNC. The gravitino mass is of order 10 GeV.

We discussed a possible solution to the μ problem in the present framework. It is necessary for us to introduce an additional gauge singlet N with the non-zero VEV but with the vanishing $\langle F_N \rangle$. When we take a large gauge group instead of $SU(3)\times SU(2),$ the massless sector of the effective theory may contain such a singlet. The present model will provide a useful guide to constructing the phenomenological viable models of gauge-mediated supersymmetry breaking.

Acknowledgements

The authors thank Prof. Kitakado for careful reading of the manuscript. They also thank C. Csáki and W. Skiba for a valuable comment. This work is supported in part by the Grant-in-Aid for Scientific Research, Ministry of Education, Science and Culture, Japan (No. 08640366).

References

(1988) 170.

- [1] I. Affleck, M. Dine and N. Seiberg, Nucl. Phys. B256 (1985) 557.
- [2] I. Affleck, M. Dine and N. Seiberg, Phys. Rev. Lett. 52 (1984) 1677. I. Affleck, M. Dine and N. Seiberg, Phys. Lett. B137 (1984) 187. I. Affleck, M. Dine and N. Seiberg, Phys. Lett. B140 (1984) 59. Y. Meurice and G. Venziano, Phys. Lett. B141 (1984) 69. D. Amati, K. Konishi, Y. Meurice, G. C. Rossi and G. Venziano, Phys. Rep. 162
	- K. Intriligator, N. Seiberg and S. H. Shenker, Phys. Lett. B342 (1995) 152.
	- E. Poppitz and S. P. Trivedi, Phys. Lett. B365 (1996) 125.
	- A. E. Nelson, Phys. Lett. B369 (1996) 277.
	- C. Csáki, L. Randall and W. Skiba, Nucl. Phys. **B479** (1996) 65.
	- C.-L. Chou, [hep-th/9605119.](http://arxiv.org/abs/hep-th/9605119)
	- E. Poppitz, Y. Shadmi and S. P. Trivedi, Phys. Lett. B388 (1996) 561.
	- C. Csáki, L. Randall, W. Skiba and R. G. Leigh, Phys. Lett. **B387** (1996) 791.
	- C. Csáki, W. Skiba and M. Schmaltz, [hep-th/9607210.](http://arxiv.org/abs/hep-th/9607210)
- [3] K.-I. Izawa and T. Yanagida, Prog. Theor. Phys. 95 (1996) 829.
- [4] K. Intriligator and S. Thomas, Nucl. Phys. **B473** (1996) 121.
- [5] M. Dine and A. E. Nelson, Phys. Rev. D48 (1993) 1277. M. Dine, A. E. Nelson and Y. Shirman, Phys. Rev. D51 (1995) 1362. M. Dine, A. E. Nelson, Y. Nir and Y. Shirman, Phys. Rev. D53 (1996) 2658.
- [6] I. Dasgupta, B. A. Dobrescu and L. Randall, Nucl. Phys. B483 (1997) 95. N. Arkani-Hamed, C. D. Carone, L. J. Hall and H. Murayama, Phys. Rev. D54 (1996) 7032.
- [7] see, for example
	- T. Hotta, K.-I. Izawa and T. Yanagida, Phys. Rev. D55 (1997) 415.
	- M. Dine, Y. Nir and Y. Shirman, Phys. Rev. D55 (1997) 1501.
	- S. P. Martin, Phys. Rev. D55 (1997) 3177.
	- S. Dimopoulos and G. F. Giudice, Phys. Lett. B393 (1997) 72.
	- S. Dimopoulos, S. Thomas and J. D. Wells, [hep-ph/9609434.](http://arxiv.org/abs/hep-ph/9609434)
	- J. Bagger, D. Pierce, K. Matchev and R.-J. Zhang, Phys. Rev. D55 (1997) 3188.
	- E. Poppitz and S. P. Trivedi, [hep-ph/9609529](http://arxiv.org/abs/hep-ph/9609529).
	- L. Randall, [hep-ph/9612426](http://arxiv.org/abs/hep-ph/9612426).
	- G. Dvali and M. Shifman, [hep-ph/9612490.](http://arxiv.org/abs/hep-ph/9612490)
	- N. Arkani-Hamed, J. March-Russell and H. Murayama, [hep-ph/9701286.](http://arxiv.org/abs/hep-ph/9701286)
	- R. N. Mohapatra and S. Nandi, [hep-ph/9702291](http://arxiv.org/abs/hep-ph/9702291).
	- S. Raby, [hep-ph/9702299.](http://arxiv.org/abs/hep-ph/9702299)
- [8] N. Haba, N. Maru and T. Matsuoka, [hep-ph/9612468.](http://arxiv.org/abs/hep-ph/9612468)
- [9] J. Bagger, E. Poppitz and L. Randall, Nucl. Phys. B426 (1994) 3.

Table Captions

Table I

The $U(1)_R$ -charges of matter superfields and their $SU(3) \times SU(2)$ gauge invariant operators. The parameter r is arbitrary other than $-5/3$.

Table I

