LA-UR-97-XXXX

Towards Resolution of the Scalar Meson Nonet Enigma

L. Burakovsky*

Theoretical Division, T-8 Los Alamos National Laboratory Los Alamos NM 87545, USA

Dedicated to Prof. L.P. Horwitz on the occasion of his 65th anniversary

Abstract

By the application of a linear mass spectrum to a composite system of both the pseudoscalar and scalar meson nonets, we find three relations for the masses of the scalar states which suggest the $q\bar{q}$ assignment for the scalar meson nonet: $a_0(1320)$, $K_0^*(1430)$, $f_0(1500)$ and $f'_0(980)$.

Key words: hadronic resonance spectrum, quark model, scalar mesons PACS: 12.39.Ki, 12.40.Ee, 12.40.Yx, 14.40.-n

^{*}E-mail: BURAKOV@PION.LANL.GOV

1 Introduction

The spectrum of the scalar meson nonet is a long-standing problem of light meson spectroscopy. The number of resonances found in the region of 1-2 GeV exceeds the number of states that conventional quark model can accommodate [1]. Extra states are interpreted alternatively as $K\bar{K}$ molecules, glueballs, multi-quark states or hybrids. In particular, except for a well established scalar isodoublet state, the $K_0^*(1430)$, the Particle Data Group (PDG) [1] lists two isovector states, the $a_0(980)$ and $a_0(1450)$. The latter, having mass and width 1450 ± 40 MeV, 270 ± 40 MeV, respectively, was discovered recently by the Crystal Barrel collaboration at LEAR [2]. The third isovector state (not included in [1]), $a_0(1320)$, having mass and width 1322 ± 30 MeV, 130 ± 30 MeV, respectively, was seen by GAMS [3]. There are four isoscalar states in [1], the $f_0(400 - 1200)$ (or σ), the interpretation of which as a particle is controversial due to a huge width of 600–1000 MeV, $f_0(980)$, $f_0(1370)$ (which stands for two separate states, $f_0(1300)$ and $f_0(1370)$, of a previous edition of PDG [4]), and $f_0(1500)$ (which also stands for two separate states, $f_0(1525)$ and $f_0(1590)$, of a previous edition of PDG), and two more possibly scalar states, the $f_J(1710), J = 0$ or 2, seen in radiative J/Ψ decays, and an η - η resonance X(1740)whose spin is also uncertain, produced in $p\bar{p}$ annihilation in flight and in chargeexchange. Recently several groups claimed different scalar isoscalar structures close to 1500 MeV, including a narrow state with mass 1450 ± 5 MeV and width 54 ± 7 MeV [5]. The lightest of the three states at 1505 MeV, 1750 MeV and 2104 MeV revealed upon reanalyzing of data on $J/\Psi \to \gamma 2\pi^+ 2\pi^-$ [6], and the $f_0(1450), f_0(1500), f_0(1520)$. The masses, widths and decay branching ratios of these states are incompatible within the errors quoted by the groups. We do not consider it as plausible that so many scalar isoscalar states exist in such a narrow mass interval. Instead, we take the various states as manifestation of one object which should be identified with the $f_0(1500)$ of recent PDG (which, however, may have a higher mass of about 1550 MeV).

It has been convincingly argued that the narrow $a_0(980)$, which has also been seen as a narrow structure in $\eta\pi$ scattering, can be generated by meson-meson dynamics alone [7]. This interpretation of the $a_0(980)$ leaves the $a_0(1320)$ or $a_0(1450)$ (which may be manifestations of one state having a mass in the interval 1350-1400 MeV) as the 1 ${}^{3}P_{0} q\bar{q}$ state. Similarly, it is mostly assumed that the $f_{0}(980)$ is a $K\bar{K}$ molecule. The mass degeneracy and their proximity to the $K\bar{K}$ threshold seem to require that the nature of both, $a_0(980)$ and $f_0(980)$, states should be the same. On the other hand, the $K\bar{K}$ interaction in the I = 1 and I = 0 channels is very different: the extremely attractive I = 0 interaction may not support a loosely bound state. Instead, it may just define the pole position of the $f_0(980) q\bar{q}$ resonance. Indeed, Morgan and Pennington [8] find the $f_0(980)$ pole structure characteristic for a genuine resonance of the constituents and not of a weakly bound system. The $I = 1 K\bar{K}$ interaction is weak and may generate a $K\bar{K}$ molecule. Alternatively, Törnqvist [9] interprets both the $f_0(980)$ and $a_0(980)$ as the members of the $q\bar{q}$ nonet with strong coupling to the decay channels, which, however, does not account for the recently discovered $a_0(1320)$ and $a_0(1450)$.

With respect to the $f_0(1370)$ (or two separate states, $f_0(1300)$ and $f_0(1370)$, according to a previous edition of PDG), we follow the arguments of Morgan and Pennington [8] and assume that the $\pi\pi$ interaction produces both very broad, $f_0(1000)$, and narrow, $f_0(980)$, states, giving rise to a dip at 980 MeV in the squared $\pi\pi$ scattering amplitude T_{11} . In this picture, the $f_0(1370)$ is interpreted as the high-mass part of the $f_0(1000)$ (the low-mass part may be associated with the σ of recent PDG). In experiments, the $f_0(1000)$ shows up at ~ 1300 MeV because of the pronounced dip in $|T_{11}|^2$ at ~ 1 GeV. The $f_0(1000)$ has an extremely large width; thus the resonance interpretation is questionable. It could be generated by *t*-channel exchanges instead of inter-quark forces [10].

The $f_0(1500)$ state has a peculiar decay pattern [11]

$$\pi\pi : \eta\eta : \eta\eta' : K\bar{K} = 1.45 : 0.39 \pm 0.15 : 0.28 \pm 0.12 :< 0.15.$$
 (1)

This pattern can be reproduced by assuming the existence of a further scalar state which is mainly $s\bar{s}$ and should have a mass of about 1700 MeV, possibly the $f_J(1710)$, and tuning the mixing of the $f_0(1500)$ with the $f_0(1370)$ $n\bar{n}$ and the (predicted) $f_0(1700)$ $s\bar{s}$ states [11]. In this picture, the $f_0(1500)$ is interpreted as a glueball state with strong mixing with the close-by conventional scalar mesons. The interpretation of the $f_0(1500)$ as a conventional $q\bar{q}$ state, as well as the qualitative explanation of its reduced $K\bar{K}$ partial width, were given by Klempt *et al.* [12] in a relativistic quark model with linear confinement and an instanton-induced interaction. The quantitative explanation of the reduced $K\bar{K}$ partial width of the $f_0(1500)$ was given in a very recent publication by the same authors [13].

The above arguments lead one to the following spectrum of the scalar meson nonet:

$$a_0(1320)$$
 or $a_0(1450)$, $K_0^*(1430)$,
 $f_0(980)$ or $f_0(1000)$, $f_0(1500)$. (2)

This spectrum agrees essentially with the $q\bar{q}$ assignments found by Klempt *et al.* [12], and Dmitrasinovic [14] who considered the Nambu–Jona-Lasinio model with a $U_A(1)$ breaking instanton-induced 't Hooft interaction. The spectrum of the meson nonet given in [12] is

$$a_0(1320), K_0^*(1430), f_0(1470), f_0'(980),$$
 (3)

while that suggested by Dmitrasinovic, on the basis of the sum rule

$$m_{f_0}^2 + m_{f'_0}^2 + m_{\eta}^2 + m_{\eta'}^2 = 2(m_K^2 + m_{K_0^*}^2)$$
(4)

derived in his paper, is [14]

$$a_0(1320), \ K_0^*(1430), \ f_0(1590), \ f_0'(1000).$$
 (5)

In this paper we show that similar results are reproduced by the application of the linear mass spectrum of a multiplet, discussed in a series of papers by Larry Horwitz and his collaborators [15, 16], to a composite system of the pseudoscalar and scalar meson nonets. The three mass relations that we obtain (one of these is the Dmitrasinovic sum rule (4), another its two-flavor version) enables one to calculate the unknown masses of the a_0 , f_0 and f'_0 states explicitly, in terms of the known masses of the pseudoscalar and scalar isodoublet states. The results turn out to agree essentially with the spectrum (2).

2 Hadronic resonance spectrum

The hadronic mass spectrum is an essential ingredient in theoretical investigations of the physics of strong interactions. It is well known that the correct thermodynamic description of hot hadronic matter requires consideration of higher mass excited states, the resonances, whose contribution becomes essential at temperatures ~ O(100 MeV)[17, 18]. The method for taking into account these resonances was suggested by Belenky and Landau [19] as considering unstable particles on an equal footing with the stable ones in the thermodynamic quantities; e.g., the formulas for the pressure and energy density in a resonance gas read¹

$$p = \sum_{i} p_{i} = \sum_{i} g_{i} \frac{m_{i}^{2} T^{2}}{2\pi^{2}} K_{2} \left(\frac{m_{i}}{T}\right), \qquad (6)$$

$$\rho = \sum_{i} \rho_i, \quad \rho_i = T \frac{dp_i}{dT} - p_i, \tag{7}$$

where g_i are the corresponding degeneracies (J and I are spin and isospin, respectively),

$$g_i = \frac{\pi^4}{90} \times \begin{bmatrix} (2J_i + 1)(2I_i + 1) & \text{for non} - \text{strange mesons} \\ 4(2J_i + 1) & \text{for strange } (K) \text{ mesons} \\ 2(2J_i + 1)(2I_i + 1) \times 7/8 & \text{for baryons} \end{bmatrix}$$

These expressions may be rewritten with the help of a *resonance spectrum*,

$$p = \int_{m_1}^{m_2} dm \ \tau(m) p(m), \quad p(m) \equiv \frac{m^2 T^2}{2\pi^2} K_2\left(\frac{m}{T}\right), \tag{8}$$

$$\rho = \int_{m_1}^{m_2} dm \ \tau(m)\rho(m), \quad \rho(m) \equiv T \frac{dp(m)}{dT} - p(m), \tag{9}$$

 1 For simplicity, we neglect the chemical potential and approximate the particle statistics by the Maxwell-Boltzmann one.

normalized as

$$\int_{m_1}^{m_2} dm \ \tau(m) = \sum_i g_i, \tag{10}$$

where m_1 and m_2 are the masses of the lightest and heaviest species, respectively, entering the formulas (6),(7).

In both the statistical bootstrap model [20, 21] and the dual resonance model [22], a resonance spectrum takes on the form

$$\tau(m) \sim m^a \ e^{m/T_0},\tag{11}$$

where a and T_0 are constants. The treatment of a hadronic resonance gas by means of the spectrum (11) leads to a singularity in the thermodynamic functions at $T = T_0$ [20, 21] and, in particular, to an infinite number of the effective degrees of freedom in the hadron phase, thus hindering a transition to the quark-gluon phase. Moreover, as shown by Fowler and Weiner [23], an exponential mass spectrum of the form (11) is incompatible with the existence of the quark-gluon phase: in order that a phase transition from the hadron phase to the quark-gluon phase be possible, the hadronic spectrum cannot grow with m faster than a power.

In ref. [16] a model for a transition from a phase of strongly interacting hadron constituents to the hadron phase described by a resonance spectrum, Eqs. (8),(9), was considered. The strongly interacting phase of hadron constituents was described by a manifestly covariant relativistic statistical mechanics which has been developed by Larry Horwitz and his collaborators and turned out to be a reliable framework in the description of realistic physical systems like strongly interacting matter [24]. An example of such a transition may be a relativistic high temperature Bose-Einstein condensation studied by the same authors in ref. [25], which corresponds, in the way suggested by Haber and Weldon [26], to spontaneous flavor symmetry breakdown, $SU(3)_F \rightarrow SU(2)_I \times U(1)_Y$, upon which hadronic multiplets are formed, with the masses obeying the Gell-Mann–Okubo formulas [27]

$$m^{\ell} = a + bY + c \left[\frac{Y^2}{4} - I(I+1) \right];$$
(12)

here I and Y are the isospin and hypercharge, respectively, ℓ is 2 for mesons and 1 for baryons, and a, b, c are independent of I and Y but, in general, depend on (p, q), where (p, q) is any irreducible representation of SU(3). Then only the assumption on the overall degeneracy being conserved during the transition is required to lead to the unique form of a resonance spectrum in the hadron phase:

$$\tau(m) = Cm, \quad C = \text{const.}$$
 (13)

Zhirov and Shuryak [28] have found the same result on phenomenological grounds. As shown in ref. [28], the spectrum (13), used in the formulas (8),(9) (with the upper

limit of integration infinity), leads to the equation of state $p, \rho \sim T^6$, $p = \rho/5$, called by Shuryak the "realistic" equation of state for hot hadronic matter [17], which has some experimental support. Zhirov and Shuryak [28] have calculated the velocity of sound, $c_s^2 \equiv dp/d\rho = c_s^2(T)$, with p and ρ defined in Eqs. (6),(7), and found that $c_s^2(T)$ at first increases with T very quickly and then saturates at the value of $c_s^2 \simeq 1/3$ if only the pions are taken into account, and at $c_s^2 \simeq 1/5$ if resonances up to $M \sim 1.7$ GeV are included.

The agreement of the results given by the linear spectrum (13) with those obtained directly from Eq. (6) for the actual hadronic species with the corresponding degeneracies, for all well-established multiplets, both mesonic and baryonic, was checked in ref. [15] and found to be excellent. Thus, the theoretical implication that a linear spectrum is the actual spectrum in the description of individual hadronic multiplets, is consistent with experiment as well.

The easiest way to see that a linear spectrum corresponds to the actual spectrum of a meson nonet is as follows. The average mass squared for a spin-s nonet is

$$\langle m^2 \rangle \equiv \frac{\sum_i g_i m_i^2}{\sum_i g_i} = \frac{3m_1^2 + 4m_{1/2}^2 + m_8^2 + m_9^2}{9},$$
 (14)

where m_1 , $m_{1/2}$, m_8 , m_9 are the masses of isovector, isodoublet, isoscalar octet and isoscalar singlet states, respectively, and the spin degeneracy, 2s + 1, cancels out. Since $m_9^2 = \langle m^2 \rangle$ [15], Eq. (14) may be rewritten as

$$\langle m^2 \rangle = \frac{3m_1^2 + 4m_{1/2}^2 + m_8^2}{8},$$
 (15)

which is the average mass squared of the octet. The same average mass squared can be also calculated with the help of the linear spectrum, as

$$\langle m^2 \rangle = \frac{\int_{m_1}^{m_8} dm \ m^3}{\int_{m_1}^{m_8} dm \ m} = \frac{(m_8^4 - m_1^4)/4}{(m_8^2 - m_1^2)/2} = \frac{m_1^2 + m_8^2}{2}.$$
 (16)

By equating (15) and (16), one obtains

$$4m_{1/2}^2 = 3m_8^2 + m_1^2, (17)$$

which is the Gell-Mann–Okubo mass formula for an octet (as follows from (12)). In general, the isoscalar octet and singlet states get mixed, because of SU(3) breaking, which results in the physical $\omega_{0'}$ and $\omega_{0''}$ states (the $\omega_{0'}$ is a mostly octet isoscalar):

$$\omega_{0'} = \omega_8 \cos \theta_M - \omega_9 \sin \theta_M,$$
$$\omega_{0''} = \omega_8 \sin \theta_M + \omega_9 \cos \theta_M,$$

where θ_M is a mixing angle. Assuming that the matrix element of the Hamiltonian between the states yields a mass squared, i.e., $m_{0'}^2 = \langle \omega_{0'} | H | \omega_{0'} \rangle$ etc., one obtains from the above relations [29],

$$m_{0'}^2 = m_8^2 \cos^2 \theta_M + m_9^2 \sin^2 \theta_M - 2m_{89}^2 \sin \theta_M \cos \theta_M,$$
(18)

$$m_{0''}^2 = m_8^2 \sin^2 \theta_M + m_9^2 \cos^2 \theta_M + 2m_{89}^2 \sin \theta_M \cos \theta_M.$$
(19)

Since $\omega_{0'}$ and $\omega_{0''}$ are orthogonal, one has further

$$m_{0'0''}^2 = 0 = (m_8^2 - m_9^2) \sin \theta_M \cos \theta_M + m_{89}^2 (\cos^2 \theta_M - \sin^2 \theta_M).$$
(20)

Eliminating m_9 and m_{89} from (18)-(20) yields

$$\tan^2 \theta_M = \frac{m_8^2 - m_{0'}^2}{m_{0''}^2 - m_8^2}.$$
(21)

It follows from (18),(19) that, independent of θ_M , $m_{0'}^2 + m_{0''}^2 = m_8^2 + m_9^2$, and therefore, as seen in Eq. (14), the average mass squared of the nonet does not change under the mixing of the ω_8 and ω_9 . This fact is easily understood in a manifestly covariant theory in which a total mass squared is rigorously conserved [24]: an average mass squared is equal to a total mass squared devided by the total number of degrees of freedom (e.g., 9 for a meson nonet); the conservation of the latter will mean that an average mass squared is conserved as well.

As also follows from (18),(19), $m_8^2 = m_{0'}^2 \cos^2 \theta_M + m_{0''}^2 \sin^2 \theta_M$, and therefore, Eq. (17) may be rewritten as

$$4m_{1/2}^2 - m_1^2 - 3m_{0'}^2 = 3\left(m_{0''}^2 - m_{0'}^2\right)\sin^2\theta_M,\tag{22}$$

which is the Sakurai mass formula [30]. For an octet-singlet mixing close to "ideal" one, for which $\omega_{0'} \simeq s\bar{s}$, $\omega_{0''} \simeq (u\bar{u} + d\bar{d})/\sqrt{2}$, $\sin\theta_M \simeq \sqrt{1/3}$; it then follows from (22) that

$$4m_{1/2}^2 \cong m_1^2 + m_{0''}^2 + 2m_{0'}^2, \tag{23}$$

which is a formula for the "ideal" structure of a nonet. As discussed in [15], this is the only mass relation for a mixed nonet consistent with the linear spectrum and the conservation of the average (and total) mass squared. This relation also appears in the algebraic approach to QCD [31] as the only solution to an infinite chain of constraints on the masses of the nonet states imposed by the charge algebras and the requirement of the asymptotic SU(3) flavor symmetry. One sees that the linear spectrum requires that the masses of the members of a given meson nonet satisfy the formula (23). We have found that Eq. (23) agrees with the experimentally established meson masses, with an accuracy of up to 3%, for all well-established meson nonets, except for the pseudoscalar and, possibly, scalar ones. As is generally believed, the reason for the invalidity of Eq. (23) for the pseudoscalar nonet is a large dynamical mass of the η_9 due to axial U(1) symmetry breakdown developed before it mixes with the η_8 to form the physical η and η' states. The well known solution to the $U_A(1)$ problem suggested by 't Hooft [32] is based on instanton effects. 't Hooft has shown that an expansion of the (euclidian) action around the one-instanton solutions of the gauge fields assuming dominance of the zero modes of the fermion fields leads to an effective $2N_f$ -fermion interaction (N_f being the number of fermion flavors) not covered by perturbative gluon exchange, which gives additional contribution to ordinary confinement quark-antiquark interaction. As shown in ref. [33], due to its point-like nature and specific spin structure, the instanton-induced interaction in the formulation of 't Hooft acts on the states with spin zero only. The masses of the other mesons with non-vanishing spin are therefore determined by the confinement interaction alone leading to the conventional splitting and an ideal mixing of the $q\bar{q}$ nonets which results in the mass spectra (23). The only two nonets whose mass spectra turn out to be affected by an instanton-induced interaction are spin zero pseudoscalar and scalar nonets. Quantitatively, an instanton-induced interaction for the scalar mesons is of the same magnitude but opposite in sign, in comparison with the pseudoscalars [12]. It, therefore, lowers the masses of the scalar states with a dominantly flavor singlet structure, and rises the masses of the scalar states with a dominantly flavor octet structure, just in contrast to the case of the pseudoscalar isoscalar flavor singlet and octet states whose masses are pushed up and down, respectively. Thus, the physical mass spectra of the pseudoscalar and scalar meson nonets are generated by the following mechanism provided by an instanton-induced interaction (in what follows, π stands for the mass of the pion, etc.):

for the pseudoscalar nonet:

$$(\pi, K, \eta_8, \eta_9) \longrightarrow (\pi, K, \eta'_8 \leq \eta_8, \eta'_9 \geq \eta_9)$$

 $\longrightarrow (\text{mixing of } \eta'_8 \text{ and } \eta'_9) \longrightarrow (\pi, K, \eta, \eta');$

for the scalar nonet:

$$(a_0, K_0^*, f_8, f_9) \longrightarrow (a_0, K_0^*, f_8' \ge f_8, f_9' \le f_9)$$

 $\longrightarrow (\text{mixing of } f_8' \text{ and } f_9') \longrightarrow (a_0, K_0^*, f_0, f_0').$

The parameters of the 't Hooft interaction may be fixed to reproduce the physical spectrum of the pseudoscalar states [34]; subsequent application to the scalar states yields the spectrum (3) [12]. In this paper we consider the spectra of the pseudoscalar and scalar states in a manifestly covariant framework, by the application of the linear mass spectrum to the composite system of both the pseudoscalar and scalar meson nonets.

3 Relations for the masses of the pseudoscalar and scalar states

We adopt the following pattern: both the pseudoscalar and scalar mesons emerge upon nucleation from the strongly interacting ensemble of hadron constituents to quasi-discrete (hadronic) mass levels, as the unique composite system which exhibits intermultiplet (instanton-induced) interaction. Let us denote the total shifts of the squared masses of the isoscalar states of the pseudoscalar and scalar nonets (which are now the subsystems of our composite system) which arise as a result of this interaction, as discussed above, as Δ_{PS} and Δ_S , respectively:

$$\Delta_{PS} \equiv \eta_8^{'2} + \eta_9^{'2} - \eta_8^2 - \eta_9^2, \qquad (24)$$

$$\Delta_S \equiv f_8^{\prime 2} + f_9^{\prime 2} - f_8^2 - f_9^2. \tag{25}$$

Since, independent of the mixing angle of the corresponding mixings,

$$\eta_8^{\prime 2} + \eta_9^{\prime 2} = \eta^2 + \eta^{\prime 2}, \tag{26}$$

$$f_8^{'2} + f_9^{'2} = f_0^2 + f_0^{'2}, (27)$$

Eqs. (24) and (25) may be rewritten as

$$\Delta_{PS} \equiv \eta^2 + \eta^{'2} - \eta_8^2 - \eta_9^2, \qquad (28)$$

$$\Delta_S \equiv f_0^2 + f_0^{\prime 2} - f_8^2 - f_9^2.$$
⁽²⁹⁾

It can be easily shown that \triangle_{PS} and \triangle_{S} are of the same magnitude but opposite in sign,

$$\Delta_{PS} = -\Delta_S,\tag{30}$$

just as the instanton-induced interaction for the scalar mesons as compared to that for the pseudoscalars, as discussed above. Indeed, independent of the details of the intermultiplet interaction (which is responsible only for the physical mass spectrum of a multiplet), the total mass squared of the composite system we are considering is rigorously conserved. Therefore, by equating the total mass squared of the bare (noninteracting) system and that of the physical pseudoscalar and scalar nonets, one obtains (taking into account the corresponding degeneracies)

$$3\pi^{2} + 4K^{2} + \eta_{8}^{2} + \eta_{9}^{2} + 3a_{0}^{2} + 4K_{0}^{*2} + f_{8}^{2} + f_{9}^{2} = 3\pi^{2} + 4K^{2} + \eta^{2} + \eta^{'2} + 3a_{0}^{2} + 4K_{0}^{*2} + f_{0}^{2} + f_{0}^{'2} + f_{0}$$

$$\eta^{2} + \eta^{'2} - \eta_{8}^{2} - \eta_{9}^{2} = f_{8}^{2} + f_{9}^{2} - f_{0}^{2} - f_{0}^{'2},$$

which reduces to Eq. (30), in view of (26)-(29). As discussed in [15] on the basis of the linear spectrum of a meson nonet, the squared mass of the isoscalar singlet state of a nonet is given by (16),

$$m_9^2 = \langle m^2 \rangle = \frac{m_1^2 + m_8^2}{2},$$

which reduces, through (17), to [15]

$$m_8^2 + m_9^2 = 2m_{1/2}^2.$$

In the case we are considering here, for the bare system of the two nonets one has, therefore, the two relations,

$$\eta_8^2 + \eta_9^2 = 2K^2,\tag{31}$$

$$f_8^2 + f_9^2 = 2K_0^{*2}, (32)$$

which for the interacting system transform into

$$\eta^2 + \eta'^2 = 2K^2 + \Delta_{PS},\tag{33}$$

$$f_0^2 + f_0^{\prime 2} = 2K_0^{*2} + \Delta_S, \tag{34}$$

in view of (24)-(27). Now, summing up (33) and (34) and using (30), one obtains

$$\eta^{2} + \eta^{'2} + f_{0}^{2} + f_{0}^{'2} = 2\left(K^{2} + K_{0}^{*2}\right), \qquad (35)$$

which is the Dmitrasinovich sum rule (4) and which represents the first of the three mass relations we need in order to determine the unknown masses of the scalar mesons. The remaining two relations are obtained by the application of the linear spectrum to the bare system and that of the two physical nonets, respectively.

For the bare system of the two nonets which occupies the mass interval (π, f_8) , by representing the average mass squared in two different ways, as was done in a previous section for the derivation of the Gell-Mann–Okubo mass formula (17),

$$\frac{3\pi^2 + 4K^2 + \eta_8^2 + \eta_9^2 + 3a_0^2 + 4K_0^{*2} + f_8^2 + f_9^2}{18} = \frac{\pi^2 + f_8^2}{2},$$
(36)

and using Eqs. (31),(32) for both, $\eta_8^2 + \eta_9^2$ and $f_8^2 + f_9^2$, in the l.h.s. of (36), and the Gell-Mann–Okubo formula

$$f_8^2 = \frac{4K_0^{*2} - a_0^2}{3} \tag{37}$$

in the r.h.s. of (36), one finally arrives at

$$K_0^{*2} - a_0^2 = K^2 - \pi^2, (38)$$

which is the second mass relations we need. A relation of this type may be anticipated on the basis of the relations

$$K^{*2} - \rho^2 = K^2 - \pi^2$$
, $K_2^{*2} - a_2^2 = K^2 - \pi^2$, etc.,

provided by either the algebraic approach to QCD developed in ref. [31] or phenomenological formulas

$$m_1^2 = 2Bm_n + C, \quad m_{1/2}^2 = B(m_n + m_s) + C$$

(where B is related to the quark condensate, C is a constant within a given meson nonet, and m_n and m_s are the masses of non-strange and strange quarks, respectively) motivated by the linear mass spectrum of a nonet and the collinearity of Regge trajectories of the corresponding I = 1 and I = 1/2 states, as discussed in refs. [15, 35], which reduce to

$$m_{1/2}^2 - m_1^2 = B(m_s - m_n),$$

independent of the quantum numbers of a nonet (for lower spin nonets, at least).

To obtain the third mass relation, we consider the system of the two physical nonets. As discussed above, in the case of a confinement interaction alone leading to an ideal mixture of a $q\bar{q}$ nonet, the physical mass spectrum of a nonet takes the form (23) which is, on the other hand, the manifestation of the conservation of the average (and total, in this case) mass squared. In the case we are considering here, the average squared masses of each of the two nonets are obviously not conserved under the mass shifts caused by an instanton-induced interaction, but the average mass squared of the composite system is conserved, due to the conservation of both the total mass squared and the total number of degrees of freedom. We, therefore, assume the validity of the two relations,

$$4K^2 = \pi^2 + \eta^{'2} + 2\eta^2 - \triangle^{'2}, \qquad (39)$$

$$4K_0^{*2} = a_0^2 + f_0^{'2} + 2f_0^2 + \Delta^{'2}, \tag{40}$$

which, upon summing up, lead to the formula

$$4\left(K^{2}+K_{0}^{*2}\right)=\pi^{2}+a_{0}^{2}+\eta^{'2}+f_{0}^{'2}+2\left(\eta^{2}+f_{0}^{2}\right),$$
(41)

consistent with the linear spectrum of the composite system of the two nonets and the conservation of the average mass squared of this composite system, just as Eq. (23) is consistent with the linear mass spectrum and the conserved average mass squared of an individual ideally mixed nonet. It then follows from (35) and (41) that

$$\eta^{'2} - \pi^2 = a_0^2 - f_0^{'2}, \tag{42}$$

which is the third relation we need in order to establish the unknown masses of the scalar mesons. The above relation was obtained by Dmitrasinovic in the twoflavor version of the Nambu–Jona-Lasinio model with the instanton-induced 't Hooft interaction [14]. It may be also derived by using the linear mass spectrum in the two-flavor case. Indeed, in this case a $q\bar{q}$ meson multiplet is a quadruplet composed with three isovector and one isoscalar states which are mass degenerate in the "ideal" case of the confinement interaction alone, since they are composed with the same nonstrange quarks (also, the standard procedure of representing the average mass squared in two different ways which we are using in this paper gives now $(3m_1^2 + m_0^2)/4 =$ $(m_1^2 + m_0^2)/2$, which reduces to $m_1 = m_0$). In the case of the instanton-induced interaction which perturbs the ideal structure of a multiplet, one has the physical mass spectrum of the composite system of the two pseudoscalar and scalar quadruplets, π, η', a_0, f'_0 ($\eta' \ge \pi, f'_0 \le a_0$). Now, the procedure of representing the average mass squared in two different ways gives

$$\frac{3\pi^2 + \eta'^2 + 3a_0^2 + f_0'^2}{8} = \frac{\pi^2 + a_0^2}{2},$$

which reduces to Eq. (42).

The three relations, (35),(38) and (42), obtained above enables one to determine the unknown masses of the scalar mesons explicitly, in terms of the known masses of the pseudoscalar and K_0^* mesons:

$$a_0^2 = K_0^{*2} - K^2 + \pi^2, (43)$$

$$f_0^2 = K_0^{*2} + 3K^2 - 2\pi^2 - \eta^2, (44)$$

$$f_0^{\prime 2} = K_0^{*2} - K^2 + 2\pi^2 - \eta^{\prime 2}.$$
(45)

With the physical masses of these states (in MeV) [1], $\pi = 138$, K = 495, $\eta = 547$, $\eta' = 958$, $K_0^* = 1429$, the formulas (43)-(45) give

$$a_0 = 1348,$$
 (46)

$$f_0 = 1562,$$
 (47)

$$f'_0 = 958.$$
 (48)

One sees that the obtained values for the masses of the scalar mesons agree essentially with the spectrum (2). The predicted mass of the isoscalar mostly singlet state (958) is within 2% of the physical mass of the $f_0(980)$ meson which is 980 ± 10 MeV [1], and is not too far from the mass of the $f_0(1000)$ meson. On the basis of the results obtained in this paper, we tentatively adopt the following $q\bar{q}$ assignment for the scalar meson nonet:

$$a_0(1320), \quad K_0^*(1430), \quad f_0(1550), \quad f_0'(980)$$

In order to obtain the mixing angle which corresponds to this assignment, we first calculate the mass of the f_8 using the Gell-Mann–Okubo mass formula and Eq. (43):

$$f_8^2 = \frac{4K_0^{*^2} - a_0^2}{3} = K_0^{*2} + \frac{K^2 - \pi^2}{3},$$
(49)

which gives

 $f_8 = 1455.$

Assuming now that the shift of the masses of the isoscalar octet states caused by an instanton-induced interaction is small compared to that of the masses of the isoscalar singlet states, i.e., $f'_8 \simeq f_8$, one has, with the help of (21),

$$\tan^2 \theta_M \simeq \frac{f_8^2 - f_0^2}{f_0'^2 - f_8^2},\tag{50}$$

which gives

 $\theta_M \simeq 26.5^o$.

This value is in good quantitative agreement with that predicted by Ritter *et al.* [13], $\theta_M \approx 25^{\circ}$, for which the partial widths of the $f_0(1500)$ calculated in their paper are

$$\pi\pi:\eta\eta:\eta\eta':KK=1.45:0.32:0.18:0.03$$

in excellent agreement with the experimentally observed partial widths, Eq. (1).

4 Concluding remarks

We have obtained three mass relations for the scalar states, by the application of the linear mass spectrum to a composite system of both the pseudoscalar and scalar meson nonets, which enable one to calculate the masses of the a_0 , f_0 and f'_0 mesons explicitly, in terms of the known masses of the pseudoscalar and K^*_0 mesons, and arrive at the $q\bar{q}$ assignment for the scalar meson nonet (2).

A question remains about which of the two, $f_0(980)$ or $f_0(1000)$, states should be associated with the predicted isoscalar mostly singlet state having a mass in the vicinity of 1 GeV. The corresponding decision is difficult to make on the basis of a naive quark model. However, two observations support the interpretation of the $f_0(980)$ as a $q\bar{q}$ state. First, the t-dependence of the $f_0(980)$ and the broad background produced in $\pi^- p \to \pi^0 \pi^0 n$ differ substantially [36]. The $f_0(1000)$ is produced in peripheral collisions only, while the $f_0(980)$ shows a t-dependence, as expected for a $q\bar{q}$ state. Second, since the scalar nonet is not ideally mixed, the both, $n\bar{n}$ and $s\bar{s}$, components of the isoscalar mostly singlet state should be appreciable. The $f_0(980)$ is seen strongly in $J/\Psi \to \phi f_0(980)$, but at most weakly in $J/\Psi \to \omega f_0(980)$. On the basis of quark diagrams, one must conclude that the $f_0(980)$ has a large $s\bar{s}$ component; its decay into $\pi\pi$ with the corresponding branching ratio 78% [1] underlines an appreciable $n\bar{n}$ component.

Although our results need experimental confirmation, the fact that the three different approaches developed in refs. [12, 14] and the present paper lead to essentially the same $q\bar{q}$ assignment for the scalar meson nonet should imply that we are not too far from the physics of a real world.

Acknowledgements

The author wishes to thank T. Goldman for very valuable discussions during the preparation of this paper.

References

- [1] Particle Data Group, Phys. Rev. D 54 (1996) 1
- [2] C. Amsler *et al.*, Phys. Lett. B **333** (1994) 277
- [3] M. Boutemeur, in Proceedings of the 3rd International Conference on Hadron Spectroscopy, "Hadron 89", Ajaccio, France, 1989; ed. F. Binon et al., Editions Frontieres (Gif-sur-Yvette) C29 (1990) 119
- [4] Particle Data Group, Phys. Rev. D 50 (1994) 1173
- [5] S. Abatzis *et al.*, Phys. Lett. B **324** (1994) 509
- [6] D.V. Bugg *et al.*, Phys. Lett. B **353** (1995) 378
- J. Weinstein and N. Isgur, Phys. Rev. D 41 (1990) 2236
 G. Janssen, B.C. Pierce, K. Holinde and J. Speth, KFA-Jülich preprint KFA-IKP-TH-1994-40; nucl-th/9411021
- [8] D. Morgan and M.R. Pennington, Phys. Rev. D 48 (1993) 1185, 5422
- [9] N.A. Törnqvist, Z. Phys. C 68 (1995) 647
- [10] B.S. Zou and D.V. Bugg, Phys. Rev. D 48 (1993) R3948
- [11] C. Amsler and F.E. Close, Phys. Lett. B **353** (1995) 385
- [12] E. Klempt, B.C. Metsch, C.R. Münz and H.R. Petry, Phys. Lett. B 361 (1995) 160

- [13] C. Ritter, B.C. Metsch, C.R. Münz and H.R. Petry, University of Bonn preprint Bonn TK-96-03; hep-ph/9601246
- [14] V. Dmitrasinovic, Phys. Rev. C 53 (1996) 1383
- [15] L. Burakovsky and L.P. Horwitz, Mass Spectrum of a Meson Nonet is Linear, Found. Phys. Lett., *in press*, Mass Spectrum of a Baryon Octet is Linear, Found. Phys. Lett., *in press*, Hadronic Resonance Spectrum May Help in Resolution of Meson Nonet Enigmas, Nucl. Phys. A, *in press*
- [16] L. Burakovsky, L.P. Horwitz and W.C. Schieve, Hadronic Resonance Spectrum: Power vs. Exponential Law. Experimental Evidence; to be published
- [17] E.V. Shuryak, The QCD Vacuum, Hadrons and the Superdense Matter, (World Scientific, Singapore, 1988)
- [18] C.R. Alcock, G.M. Fuller and G.J. Mathews, Astrophys. J. **320** (1987) 439
 H. Bebie, P. Gerber, J.L. Goity and H. Leutwyler, Nucl. Phys. B **378** (1992) 95
- [19] S.Z. Belenky and L.D. Landau, Sov. Phys. Uspekhi 56 (1955) 309; Nuovo Cim. Suppl. 3 (1956) 15
- [20] R. Hagedorn, Nuovo Cim. **35** (1965) 216; Nuovo Cim. Suppl. **6** (1968) 169, 311
- [21] S. Frautschi, Phys. Rev. D 3 (1971) 2821
- [22] S. Fubini and G. Veneziano, Nuovo Cim. A **64** (1969) 811
- [23] G.N. Fowler and R.M. Weiner, Phys. Lett. B 89 (1980) 394
- [24] L.P. Horwitz and C. Piron, Helv. Phys. Acta 46 (1973) 316
 L.P. Horwitz, W.C. Schieve and C. Piron, Ann. Phys. (N.Y.) 137 (1981) 306
 L.P. Horwitz, S. Shashoua and W.C. Schieve, Physica A 161 (1989) 300
 L. Burakovsky, Manifestly Covariant Relativistic Statistical Mechanics as a Framework for Description of Realistic Physical Systems, Ph.D. thesis (Tel-Aviv University, 1995), unpublished; L. Burakovsky, L.P. Horwitz and W.C. Schieve, Mass Proper Time Uncertainty Relation in a Manifestly Covariant Relativistic Statistical Mechanics, to be published
- [25] L. Burakovsky, L.P. Horwitz and W.C. Schieve, Phys. Rev. D 54 (1996) 4029
- [26] H.A. Haber and H.E. Weldon, Phys. Rev. D 25 (1982) 502
- [27] S. Okubo, Prog. Theor. Phys. 27 (1962) 949, 28 (1962) 24
 M. Gell-Mann and Y. Ne'eman, *The Eightfold Way*, (Benjamin, N.Y., 1964)
- [28] O.V. Zhirov and E.V. Shuryak, Sov. J. Nucl. Phys. **21** (1975) 443

- [29] D.H. Perkins, Introduction to High Energy Physics, 3rd ed., (Addison Wesley, Reading, MA, 1987), section 5.6
- [30] J.J. Sakurai, *Currents and Mesons*, (University of Chicago Press, Chicago, 1969)
- [31] S. Oneda and K. Terasaki, Prog. Theor. Phys. Suppl. 82 (1985) 1
- [32] G. 't Hooft, Phys. Rev. D 14 (1976) 3432
- [33] C.R. Münz, J. Resag, B.C. Metsch and H.R. Petry, Nucl. Phys. A 578 (1994) 418
- [34] W.H. Blask, U. Bohn, M.G. Huber, B.C. Metsch and H.R. Petry, Z. Phys. A 337 (1990) 327
- [35] L. Burakovsky and L.P. Horwitz, On the Thermodynamics of Hot Hadronic Matter, Nucl. Phys. A, *in press*
- [36] D. Alde *et al.*, Z. Phys. C **66** (1995) 375