

Pair production of charged Higgs bosons in gluon-gluon collisions

Jiang Yi^b, Han Liang^b, Ma Wen-Gan^{a,b}, Yu Zeng-Hui^b and Han Meng^b

^aCCAST (World Laboratory), P.O.Box 8730, Beijing 100080,
People's Republic of China

^bModern Physics Department, University of Science and
Technology of China, Anhui 230027,
People's Republic of China. *

Abstract

We studied the pair production rate of charged Higgs bosons via gluon-gluon collisions in the general two-Higgs-doublet model. The analysis of its production are made with possible parameter values of Higgs sectors in this model. We find that the production cross sections may be 4 to 41 fermibar in the future LHC pp collider with these parameters. The calculation also shows that this pair production process is not sensitive to the neutral Higgs masses, therefore its measurement could not give more imformations about the neutral Higgs bosons.

PACS number(s): 14.80.Cp, 12.15.Lk, 12.60.Cn

*Mailing address

1. Introduction

In the sense of Minimal Standard Model(MSM) [1][2], a neutral Higgs boson is predicted without giving its mass value. Therefore the search for Higgs boson is undoubtedly one of the important issues in testing the electroweak theory. Until now, the top quark mass is given as [3]: $m_t = 176 \pm 10^{+8}_{-10} GeV$. The absence of Higgs-boson signal events leads to a lower bound $M_{H^0} > 63.5 GeV$ at 95% C.L. obtained [4]. In order to investigate the sectors of the theory about which we know very little, the future multi-TeV hadron colliders such as the CERN Large Hadron Collider (LHC) and Next Linear Collider(NLC) are elaborately designed to devote to the precise characters of Higgs sector and top quark. Besides providing the first evidence of Higgs mechanism, the multi-TeV colliders offer the possibility of carrying on the research of other unknown particles in the hitherto unexplored mass range.

So far the extension of the MSM which includes a second Higgs doublet is far from being ruled out experimentally. One strong motivation for extending Higgs sector into two-Higgs-doublet model (THDM), comes from that it has a common feature of many extended models of the MSM, such as Minimal Supersymmetric Standard Model(MSSM). The MSSM model requires two SU(2) doublets to give masses to up and down quarks[5]. It is therefore of interest to determine whether THDM or MSSM is compatible with experiments, and what further extensions should be if it is not.

In future hadron colliders, several mechanisms can produce Higgs boson. Among them the Higgs pair productions via gluon-gluon collision, quark-antiquark annihilation or others, play a significant role in studying Higgs sectors. The on-mass shell charged Higgs boson pair

production with their subsequent cascade decays will provide a good opportunity of studying the two Higgs doublet model and a copious source of charged Higgs bosons. Of cause, the Higgs pair production has less rate comparing with the single production mechanism, but the Higgs pair production via gg and $q\bar{q}$ annihilation involves the trilinear and quartic couplings of the physical Higgs bosons. So the measurement of these couplings may be a good test of our theory. At the mean time, the pair production of Higgs boson in hadron collisions has a better signal-to-background ratio than one Higgs production process. Therefore it is worthwhile to study the precise features of the Higgs pair production processes at LHC, NLC and other future colliders which are designed operating in $\gamma\gamma, e^+e^-$, pp and $p\bar{p}$ collision modes. Especially when charged Higgs boson mass is not very heavy, its pair production will be more significant. Recently T. Plehn, M. Spira and P.M. Zerwas provided the calculations of pair production of neutral Higgs particles in gluon-gluon collisions in MSSM[6] and charged Higgs boson pair production at NLC in $\gamma\gamma$ modes are also investigated in references [7].

In this paper we represent the calculation of the charged Higgs boson pair production via gluon-gluon collisions at LHC in the frame of general THDM. The paper is organized as follows: The analytical evaluations are given in section 2. In section 3 there are numeriacal results, discussion and a short summary. Finally the explicit expressions of the relevant form factors are collected in Appendix.

2.Calculation

The general THDM contains six free parameters. Generally the four mass parameters,

i.e. the parameters for two scalar neutral Higgs boson masses m_{h^0} and m_{H^0} , one neutral pseudoscalar Higgs mass m_{A^0} , one pair charged Higgs boson masses m_{H^\pm} and the additional mixing angle α and β , are chosen as free parameters. The parameter α is defined as the mixing angle of the scalar Higgs bosons h^0 and H^0 . The angle β is a mixing parameter of charged and pseudoscalar Higgs bosons and is related directly to the vacuum expectation values of Higgs doublets (i.e. $\tan \beta = v_2/v_1$). In order to demonstrate the numerical features of the charged Higgs boson pair production, we specify $\alpha = \beta$ in our numerical calculation for simplicity.

The generic Feynman diagrams contributing to the subprocess $gg \rightarrow H^+ H^-$ which are involved in frame of THDM are depicted in figure 1. The relevant Feynman rules can be found in references[5]. Since we neglect the couplings between light-quarks and Higgs bosons, there are only the following three kinds of production mechanisms: One way is by exchanging a γ or a Z^0 boson to produce a charged Higgs boson pair(see Fig.1(a,b)), we call it as $\gamma - Z^0$ s-channel. The γ exchange s-channel diagrams have no contribution and only the axis-vector part of the coupling between Z^0 and the heavy quark in Z^0 exchange diagram, contributes to the cross section. The second mechanism is through producing the virtual neutral Higgs h^0 or H_0 bosons which is coupled to gluons by the usual heavy-quark triangle, and the charged Higgs boson pair is appeared by the subsequent decay of the virtual Higgs bosons (see Fig.1(c,d)), which are called the $h^0 - H^0$ s-channel. Another mechanism is named as heavy-quark box diagrams(see Fig.1(e ~ j)). The box-diagrams of exchanging the two initial gluons are not plotted in Fig.1. All the box diagrams can be divided into two groups. They are corresponding to t-channel and u-channel groups respectively. In this work, we used

n-dimensional regularization scheme in the one-loop diagram calculation.

We denote θ as the scattering angle between the one of the gluons and final H^+ boson.

Then in the system of center-of-mass (CMS) we can express all the four-momenta of initial and final particles by means of the total energy $\sqrt{\hat{s}}$, the scattering angle θ and the velocity of final charged Higgs bosons. In the CMS of the initial two gluons, the four-momenta of the final charged Higgs bosons read in term of following forms

$$\begin{aligned} p_1 &= \left(\frac{\sqrt{\hat{s}}}{2}, \frac{\sqrt{\hat{s}}}{2} \beta_{H^\pm} \sin \theta, 0, \frac{\sqrt{\hat{s}}}{2} \beta_{H^\pm} \cos \theta \right), \\ p_2 &= \left(\frac{\sqrt{\hat{s}}}{2}, \frac{-\sqrt{\hat{s}}}{2} \beta_{H^\pm} \sin \theta, 0, \frac{-\sqrt{\hat{s}}}{2} \beta_{H^\pm} \cos \theta \right). \end{aligned} \quad (1)$$

where the velocity of charged Higgs bosons in the CMS, β_{H^\pm} , can be represented as

$$\beta_{H^\pm} = \sqrt{1 - 4m_{H^\pm}^2/\hat{s}}. \quad (2)$$

p_3 and p_4 are the four-momenta of the initial gluons and expressed as

$$\begin{aligned} p_3 &= \left(\frac{\sqrt{\hat{s}}}{2}, 0, 0, \frac{\sqrt{\hat{s}}}{2} \right), \\ p_4 &= \left(\frac{\sqrt{\hat{s}}}{2}, 0, 0, \frac{-\sqrt{\hat{s}}}{2} \right). \end{aligned} \quad (3)$$

The corresponding matrix element for all the diagrams in figure 1 is written as

$$\begin{aligned} M &= M^{\hat{s}} + M^{box(\hat{t})} + M^{box(\hat{u})} \\ &= \frac{-1}{16\pi^2} T^a T^b g_s^2 \epsilon_\mu(p_3) \epsilon_\nu(p_4) \\ &\quad (f_1 g^{\mu\nu} + f_2 p_1^\mu p_1^\nu + f_3 p_1^\mu p_2^\nu + f_4 p_2^\mu p_1^\nu + f_5 p_2^\mu p_2^\nu + \\ &\quad f_6 \epsilon^{\mu\nu\alpha\beta} p_{1\alpha} p_{2\beta} + f_7 \epsilon^{\mu\nu\alpha\beta} p_{1\alpha} p_{3\beta} + f_8 \epsilon^{\mu\nu\alpha\beta} p_{2\alpha} p_{3\beta} + \\ &\quad f_9 \epsilon^{\nu\alpha\beta\gamma} p_{1\alpha} p_{2\beta} p_{3\gamma} p_1^\mu + f_{10} \epsilon^{\mu\alpha\beta\gamma} p_{1\alpha} p_{2\beta} p_{3\gamma} p_1^\nu + \end{aligned}$$

$$f_{11}\epsilon^{\nu\alpha\beta\gamma}p_{1,\alpha}p_{2,\beta}p_{3,\gamma}p_2^\mu + f_{12}\epsilon^{\mu\alpha\beta\gamma}p_{1,\alpha}p_{2,\beta}p_{3,\gamma}p_2^\nu. \quad (4)$$

The upper indexes \hat{s} , $\text{box}(\hat{t})$ and $\text{box}(\hat{u})$ represent the amplitudes corresponding to the s-channel diagrams, including all the $\gamma - Z^0$ and $h^0 - H^0$ diagrams, t-channal and u-channal box diagrams in Fig.1 respectively. The g_s is the strong coupling constant and $T^a(a = 1 \sim 8)$ are the $SU(3)_C$ generators introduced by Gell-Mann[8]. The variables \hat{s} , \hat{t} and \hat{u} are usual Mandelstam variables in the gluon-gluon center-of mass system with the definitions

$$\begin{aligned} \hat{s} &= (p_1 + p_2)^2 = (p_3 + p_4)^2, \\ \hat{t} &= (p_1 - p_3)^2 = (p_2 - p_4)^2, \\ \hat{u} &= (p_1 - p_4)^2 = (p_2 - p_3)^2. \end{aligned} \quad (5)$$

We collect the explicit expressions of the form factors $f_i(i = 1 \sim 12)$ which appear in equation (4) in Appendix. The total cross section for subprocess $gg \rightarrow H^+H^-$ can finally be written in the form

$$\hat{\sigma}(\hat{s}) = \frac{1}{16\pi\hat{s}^2} \int_{\hat{t}^-}^{\hat{t}^+} d\hat{t} |\bar{M}|^2. \quad (6)$$

where $|\bar{M}|^2$ is the initial spin and color averaged matrix element squared and $\hat{t}^\pm = (m_{H^\pm}^2 - \frac{1}{2}\hat{s}) \pm \frac{1}{2}\hat{s}\beta_{H^\pm}$. The color factor including in $|\bar{M}|^2$ is equal to 1/32. The total cross section for the charged Higgs pair production through gluon-gluon fusion in proton-proton collisions can be obtained by integrating the $\hat{\sigma}$ over the gluon luminosity.

$$\sigma(pp \rightarrow gg \rightarrow H^+H^- + X) = \int_{4m_{H^\pm}^2/s}^1 d\tau \frac{dL_{gg}}{d\tau} \hat{\sigma}(gg \rightarrow H^+H^- \text{ at } \hat{s} = \tau s). \quad (7)$$

Here we introduced a variable $\tau = x_1 x_2$ and x_1 and x_2 are defined as

$$x_1 = \frac{p_3}{P_1}, \quad x_2 = \frac{p_4}{P_2},$$

where P_1 and P_2 are the four momentum of both incident protons respectively. \sqrt{s} and $\sqrt{\hat{s}}$ are the proton-proton and gluon-gluon center-of-mass energies respectively, which can be expressed as

$$s = (P_1 + P_2)^2, \quad \hat{s} = (p_3 + p_4)^2 = x_1 x_2 s = \tau s. \quad (8)$$

The quantity of the gluon luminosity $\frac{dL_{gg}}{d\tau}$ can be calculated by using the following equation.

$$\frac{dL_{gg}}{d\tau} = 2 \int_\tau^1 \frac{dx_1}{x_1} \left[F_g(x_1, \mu) F_g\left(\frac{\tau}{x_1}, \mu\right) \right]. \quad (9)$$

The numerical results of the process $pp \rightarrow gg \rightarrow H^+ H^-$ are represented by adopting the MRS set G parton distribution function $F_g(x)$ [9].

3. Numerical results and discussion

In the numerical evolution we take the input parameters[10] as $m_b = 4.5 \text{ GeV}$, $m_Z = 91.1887 \text{ GeV}$, $m_W = 80.2226 \text{ GeV}$, $G_F = 1.166392 \times 10^{-5} (\text{GeV})^{-2}$ and $\alpha = \frac{1}{137.036}$. The top quark mass is just fixed to be the central value $m_t = 175 \text{ GeV}$. We also used a simple one-loop formula for the running strong coupling constant α_s as

$$\alpha_s(\mu) = \frac{\alpha_s(m_Z)}{1 + \frac{33-2n_f}{6\pi} \alpha_s(m_Z) \ln\left(\frac{\mu}{m_Z}\right)}. \quad (10)$$

where $\alpha_s(m_Z) = 0.117$ and n_f is the number of active flavors at scale μ . The ratio of the two vacuum expectation values $\tan \beta$ is chosen to be varied in our calculation. The coupling

strength of the charged Higgs boson with quarks is related to scalar and pseudo-scalar parts, each part has factor g_+ or g_- (The definitions of g_+ and g_- can be found in Appendix). These two parts of Yukawa coupling yield a crucial contribution alternatively depending on variate $\tan\beta$. Because the present experimental data disfavor the small values of $\tan\beta$ in general THDM. Therefore we limit its quantity in the representative range of 1.5 to 30 in the calculation.

The cross sections of the subprocess $gg \rightarrow H^+H^-$ with $m_{H^\pm} = 150 \text{ GeV}$ are depicted in Fig.2(1), on the conditions of $m_{h^0} = 100 \text{ GeV}$ and $m_{H^0} = 150 \text{ GeV}$. Fig.2(2) shows also the cross sections of this subprocess as the functions of CMS energy of incoming gluons with the same neutral Higgs boson masses, but $m_{H^\pm} = 300 \text{ GeV}$. In both figures, the full line curves and dashed line curves are corresponding to the $\tan\beta = 1.5$ and $\tan\beta = 30$ respectively. These graphs show that the cross section of the subprocess is approximately two times larger when the ratio of vacuum expectation values is 30 than that when $\tan\beta = 1.5$ in the energy region less than 2 TeV. In Fig.2(1) there is a peak around $\sqrt{\hat{s}} \sim 400 \text{ GeV}$. This peak comes from the $t\bar{t}$ threshold effects and enhanced by the top quark loop diagrams.

The calculation shows the result of the cross section $\hat{\sigma}$ is not sensitive to the masses of the related two neutral Higgs bosons m_{h^0} and m_{H^0} , especially when $\sqrt{\hat{s}}$ is far beyond the value regions of the two masses m_{h^0} and m_{H^0} . From the Feynman diagrams in Fig.1, we know that the diagrams, which are related to the neutral Higgs bosons and the trilinear Higgs couplings, are only appeared in the s-channel as shown in Fig.1(a,b), but their contribution is negligible quantitatively. This comes from two reasons. Firstly, the contribution is enhanced mainly by the Yukawa coupling which is proportional to m_t/m_W . Since the γ exchange s-channel

diagram has no contribution and Z^0 boson exchange s-channel diagram does not include Yukawa coupling, the contribution from these kinds of s-channel diagrams is rather small. Among the s-channel diagrams, Fig.1(b) has only one Yukawa coupling, whereas each box diagram in Fig.1($c \sim h$) has two such couplings, therefore the cross section of the subprocess is mainly contributed by the box diagrams. Secondly, when the energy of CMS system $\sqrt{\hat{s}}$ is not in the regions arround m_{h^0} and m_{H^0} , the propagators in s-channel diagrams will suppress their contributions to the cross section. So it is clear that the measurement of this process could not be expected to give constraint on the features of neutral Higgs bosons and the trilinear Higgs couplings at the future pp colliders.

In figure 3 we show the cross section $\hat{\sigma}$ as the function of $\tan\beta$ with $m_{H^\pm} = 150\text{ GeV}$, when $\sqrt{\hat{s}} = 400\text{ GeV}$ and $\sqrt{\hat{s}} = 800\text{ GeV}$ respectively. The cross section goes down when $\tan\beta$ varies from 1.5 to 6, and raises up when $\tan\beta$ varies from 6 to 30. The cross sections are extremely small in the region arround $\tan\beta = 6$. Fig.4 is the plot of cross section $\hat{\sigma}$ as the function of the charged Higgs boson mass with $\sqrt{\hat{s}} = 1\text{ TeV}$, $\tan\beta$ are 1.5 and 30 respectively. There the cross section increases with increasing mass of the charged Higgs boson m_{H^\pm} . In the region where $m_{H^\pm} < 300\text{ GeV}$, the curves are flat, whereas they raise up quickly when $m_{H^\pm} > 300\text{ GeV}$. That is because when the charged Higgs boson pair production threshold approaches the CMS energy $\sqrt{\hat{s}} = 1\text{ TeV}$, the cross section will be enhanced by the threshold effect of the charged Higgs boson pair production.

In Fig.5(1) and Fig.5(2) we show the cross section of $pp \rightarrow gg \rightarrow H^+H^- + X$ process with m_{H^\pm} being 150 GeV and 300 GeV respectively. The charged Higgs boson pair production

rates are read to be about 14 to 41 fermi-bar at the LHC energies when charged Higgs boson mass is 150 GeV and about 4.2 to 13.2 fermi-bar when $m_{H^\pm} = 300 \text{ GeV}$.

In conclusion, we have studied the pair production process of charged Higgs boson via gluon-gluon fusion in pp collider at LHC. The numerical analysis of its cross sections is carried out in the general two-Higgs- doublet model. With the possible parameters, the cross sections at future LHC collider can be 4 to 41 fermi-bar. These quantities are in the comparable range with that of the neutral Higgs boson pair production in gluon-gluon collisions as shown in Ref.[6]. The calculation shows also that the production rate of charged Higgs boson pair is not sensitive to the neutral Higgs boson masses and the trilinear Higgs couplings, but it is strongly related to the couplings of the charged Higgs boson to fermions.

This work was supported in part by the National Natural Science Foundation of China and the National Committee of Science and Technology of China. Part of this work was done when two of the authers, Ma Wen-Gan and Yu Zeng-Hui, visited the University Vienna under the exchange agreement(project number: IV.B.12).

Appendix

We adopt the same definitions of one-loop integral functions as in reference[11] and $d = 4 - \epsilon$. The integral functions are defined as

$$A_0(m) = -\frac{(2\pi\mu)^{4-D}}{i\pi^2} \int d^D q \frac{1}{[q^2 - m^2]},$$

$$\{B_0; B_\mu; B_{\mu\nu}\}(p, m_1, m_2) = \frac{(2\pi\mu)^{4-D}}{i\pi^2} \int d^D q \frac{\{1; q_\mu; q_{\mu\nu}\}}{[q^2 - m_1^2][(q + p)^2 - m_2^2]},$$

$$\{C_0; C_\mu; C_{\mu\nu}; C_{\mu\nu\rho}\}(p_1, p_2, m_1, m_2, m_3) =$$

$$\begin{aligned}
& -\frac{(2\pi\mu)^{4-D}}{i\pi^2} \int d^D q \frac{\{1; q_\mu; q_{\mu\nu}; q_{\mu\nu\rho}\}}{[q^2 - m_1^2][(q + p_1)^2 - m_2^2][(q + p_1 + p_2)^2 - m_3^2]}, \\
& \{D_0; D_\mu; D_{\mu\nu}; D_{\mu\nu\rho}; D_{\mu\nu\rho\alpha}\}(p_1, p_2, p_3, m_1, m_2, m_3, m_4) = \\
& \frac{(2\pi\mu)^{4-D}}{i\pi^2} \int d^D q \frac{\{1; q_\mu; q_{\mu\nu}; q_{\mu\nu\rho}; q_{\mu\nu\rho\alpha}\}}{[q^2 - m_1^2][(q + p_1)^2 - m_2^2][(q + p_1 + p_2)^2 - m_3^2][(q + p_1 + p_2 + p_3)^2 - m_4^2]},
\end{aligned}$$

The form factors involved in equation (4) are represented in terms of the above integral functions explicitly as:

$$\begin{aligned}
f_1 &= 2m_t(g_{h^0 H^\pm H^\mp} g_{h^0 tt} A_{h^0} + g_{H^0 H^\pm H^\mp} g_{H^0 tt} A_{H^0}) d B_0[p_1 + p_2, m_t, m_t] \\
&- 8m_t(g_{h^0 H^\pm H^\mp} g_{h^0 tt} A_{h^0} + g_{H^0 H^\pm H^\mp} g_{H^0 tt} A_{H^0}) \cdot \\
&(dC_{24} + (p_4 \cdot p_3)C_0)[-p_3, p_1 + p_2, m_t, m_t, m_t] + i f_1^- g_-^2 + i f_1^+ g_+^2,
\end{aligned} \tag{A-1}$$

where

$$f_1^- = f_{1a}^- + f_{1b}^- + f_{1c}^-, \tag{A-2}$$

$$\begin{aligned}
f_{1a}^- &= 2(2(m_b m_t C_0 - m_b^2 C_0 - m_{H^\pm}^2 C_{12}) + 2(p_1 \cdot p_3)(C_0 + C_{11} - C_{12}) - \\
&2(p_2 \cdot p_3)(C_0 + C_{11}) + 2(p_1 \cdot p_2)(C_{11} - C_{12}) - dC_{24})[-p_1 - p_3, -p_4, m_b, m_t, m_t] \\
&+ 2(2(p_1 \cdot p_3)(C_{11} - C_0 - C_{12}) - 2(p_1 \cdot p_2)C_{11} + 2(p_2 \cdot p_3)(C_0 + 3C_{11} - 2C_{12}) - \\
&dC_{24} + 2m_b(m_t - m_b)C_0 - 4m_{H^\pm}^2 C_{11})[-p_2 + p_3, -p_3, m_b, m_t, m_t] \\
&+ 4(2m_{H^\pm}^2(D_{27} + 2D_{311} - 2D_{312} + 2D_{313}) + 4m_t(m_t - m_b)D_{27} + \\
&2m_{H^\pm}^2 m_t^2(D_{11} - D_{12} + D_{13}) + 2(p_1 \cdot p_2)(D_{27} + 2D_{313} + m_t^2 D_{13}) + \\
&(p_1 \cdot p_3)(4D_{312} - 4D_{313} - 2D_{27} + m_{H^\pm}^2(D_{11} - D_{12} + 2D_{21} + 2D_{22} + \\
&2D_{23} - 4D_{24} + 4D_{25} - 4D_{26}) + m_b m_t(2D_{12} - D_0 - 2D_{11} - 2D_{13}) +
\end{aligned}$$

$$\begin{aligned}
& m_t^2(D_0 + 2D_{11})) + 2(p_1 \cdot p_2)(p_1 \cdot p_3)(D_{23} + D_{25} - D_{26}) + \\
& 2(p_1 \cdot p_3)^2(D_{24} - D_{22} - D_{23} - D_{25} + 2D_{26}) + \\
& (p_2 \cdot p_3)(m_{H^\pm}^2(D_{11} - D_{12} + 2D_{13} + 2D_{23} + 2D_{25} - 2D_{26}) - \\
& m_b m_t(D_0 + 2D_{13}) + m_t^2(D_0 + 2D_{13})) + 2(p_1 \cdot p_2)(p_2 \cdot p_3)(D_{13} + D_{23}) + \\
& 2(p_1 \cdot p_3)(p_2 \cdot p_3)(D_{26} - D_{23}))[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t] \\
+ & 4(4m_{H^\pm}^2 D_{311} + 2m_{H^\pm}^4(D_{11} + D_{21} + D_{24}) - 4m_b m_t D_{27} + \\
& 4m_t^2 D_{27} - 2m_b m_t m_{H^\pm}^2(D_0 + D_{11} + D_{12}) + 2m_{H^\pm}^2 m_t^2(D_0 + 2D_{11} + D_{12}) + \\
& 2(p_1 \cdot p_2)(2D_{312} + m_{H^\pm}^2(D_{11} + D_{12} + D_{21} + D_{22} + 2D_{24}) - m_b m_t(D_0 + D_{11} + D_{12}) + \\
& m_t^2(D_0 + D_{11} + 2D_{12})) + 2(p_1 \cdot p_2)^2(D_{12} + D_{22} + D_{24}) + (p_1 \cdot p_3)(2D_{27} - 4D_{312} + \\
& 4D_{313} + m_{H^\pm}^2(2D_{13} - D_{11} - D_{12} - 2D_{21} - 2D_{22} - 4D_{24} + 4D_{25} + 2D_{26}) + \\
& m_b m_t(D_0 + 2D_{11} + 2D_{12} - 2D_{13}) + m_t^2(4D_{13} - D_0 - 2D_{11} - 4D_{12})) + \\
& 2(p_1 \cdot p_2)(p_1 \cdot p_3)(D_{13} - D_{12} - 2D_{22} - 2D_{24} + D_{25} + 2D_{26}) + \\
& 2(p_1 \cdot p_3)^2(D_{22} + D_{23} + D_{24} - D_{25} - 2D_{26}) + \\
& (p_2 \cdot p_3)(m_{H^\pm}^2(2D_{25} - D_{11} - D_{12} - 4D_{24}) + m_b m_t(D_0 + 4D_{12} - 2D_{13}) + \\
& m_t^2(2D_{13} - D_0 - 4D_{12})) + 2(p_1 \cdot p_2)(p_2 \cdot p_3)(D_{26} - D_{12} - 2D_{22}) + \\
& 2(p_1 \cdot p_3)(p_2 \cdot p_3)(D_{23} + 2D_{22} - 3D_{26}))[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t], \quad (A-3)
\end{aligned}$$

$$f_{1b}^- = f_{1a}^- \quad (m_b \leftrightarrow m_t), \quad (A-4)$$

$$\begin{aligned}
f_{1c}^- = & 2d(B_0[-p_3, m_t, m_t] + B_0[-p_3, m_b, m_b] - B_0[-p_2, m_t, m_b]) \\
& + 4(m_{H^\pm}^2(C_{11} + C_{12}) + (p_1 \cdot p_2)(C_{11} + C_{12}) - (p_1 \cdot p_3)(2C_0 + 3C_{11}) - \\
& (p_2 \cdot p_3)(C_0 + C_{11} + 2C_{12}))[-p_1 + p_3, -p_2, m_b, m_t, m_b] \\
& + 4(m_{H^\pm}^2(C_{11} - C_0 - C_{12}) + (p_1 \cdot p_2)(C_{11} - C_0 - C_{12}) - (p_1 \cdot p_3)C_{11} + \\
& (p_2 \cdot p_3)(2C_{12} - C_0 - 3C_{11}))[p_2 - p_3, -p_2, m_b, m_t, m_b] \\
& + 4(2(m_b^2 - 2m_b m_t + m_t^2)D_{27} + 2m_{H^\pm}^2 m_t^2(D_{12} - D_{11}) + 2(p_1 \cdot p_2)(D_{27} + m_t^2 D_{13}) + \\
& (p_1 \cdot p_3)(2D_{27} + m_b^2(D_{11} - D_{12}) + m_{H^\pm}^2(D_{13} - 2D_{21} - 2D_{22} + 4D_{24}) + \\
& m_t^2(D_0 - D_{11} - D_{12}) - m_b m_t D_0) + 2(p_1 \cdot p_2)(p_1 \cdot p_3)(D_{25} - D_{26}) + \\
& 2(p_1 \cdot p_3)^2(D_{22} - D_{24}) + (p_2 \cdot p_3)(m_{H^\pm}^2(D_{11} - D_{12} + 2D_{25} - 2D_{26}) - \\
& m_b^2 D_{13} - m_b m_t D_0 + m_t^2(D_0 + D_{13})) + 2(p_1 \cdot p_3)(p_2 \cdot p_3)D_{26} - \\
& 2(p_1 \cdot p_2)(p_2 \cdot p_3)(D_{13} + D_{23})) [p_1, -p_1 + p_3, -p_2, m_t, m_b, m_t, m_b] \\
& + 4(2(m_b^2 + m_{H^\pm}^2 + m_t^2)D_{27} - 4m_b m_t D_{27} + m_b^2 m_{H^\pm}^2(D_{11} + D_{12} - D_{13}) + \\
& m_{H^\pm}^4(D_{13} - D_{11} - D_{12} - 2D_{21} - 2D_{24} + 2D_{25}) + \\
& m_{H^\pm}^2 m_t^2(D_{13} - 3D_{11} - D_{12}) + (p_1 \cdot p_2)(4D_{27} + m_b^2(D_{11} + D_{12} - D_{13}) + \\
& m_{H^\pm}^2(D_{13} - D_{11} - D_{12} - 2D_{21} - 2D_{22} - 2D_{23} - 4D_{24} + 4D_{25} + 4D_{26}) + \\
& m_t^2(3D_{13} - D_{11} - 3D_{12})) + 2(p_1 \cdot p_2)^2(D_{25} - D_{22} - D_{23} - D_{24} + 2D_{26}) + \\
& (p_1 \cdot p_3)(m_t^2(D_0 + D_{11} + 3D_{12}) - m_b m_t D_0 - m_b^2(D_{11} + D_{12}) + \\
& m_{H^\pm}^2(2D_{11} + 2D_{12} - D_{13} + 2D_{21} + 2D_{22} + 4D_{24} - 2D_{26} - 2D_{27})) + \\
& 2(p_1 \cdot p_2)(p_1 \cdot p_3)(D_{12} - D_{13} + 2D_{22} + 2D_{24} - D_{25} - 2D_{26}) -
\end{aligned}$$

$$\begin{aligned}
& 2(p_1 \cdot p_3)^2(D_{12} + D_{22} + D_{24}) + (p_2 \cdot p_3)(m_b^2(D_{13} - 2D_{12}) - m_b m_t D_0 + \\
& m_{H^\pm}^2(D_{11} + D_{12} + 4D_{24} - 2D_{25}) + m_t^2(D_0 + 2D_{12} - D_{13})) + \\
& 2(p_1 \cdot p_2)(p_2 \cdot p_3)(D_{23} + 2D_{22} - 3D_{26}) + \\
& 2(p_1 \cdot p_3)(p_2 \cdot p_3)(D_{26} - D_{12} - 2D_{22})) [p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b], \quad (A-5)
\end{aligned}$$

$$f_1^+ = f_1^- [m_b \rightarrow -m_b]. \quad (A-6)$$

$$\begin{aligned}
f_2 &= 8m_t(g_{h^0 H^\pm H^\mp} g_{h^0 tt} A_{h^0} + g_{H^0 H^\pm H^\mp} g_{H^0 tt} A_{H^0}) \cdot \\
&(C_{11} + 2C_{21} + 4C_{22} - 4C_{23})[-p_3, p_1 + p_2, m_t, m_t, m_t] + i f_2^- g_-^2 + i f_2^+ g_+^2, \quad (A-7)
\end{aligned}$$

where

$$f_2^- = f_{2a}^- + f_{2b}^- + f_{2c}^-, \quad (A-8)$$

$$\begin{aligned}
f_{2a}^- &= 4(C_0 + C_{11} - C_{12})[p_1 - p_3, -p_4, m_b, m_t, m_t] \\
&+ 4(C_0 + C_{11} - C_{12})[-p_2 + p_3, -p_3, m_b, m_t, m_t] \\
&+ (8(D_{27} + 4D_{311} - 2D_{312} + 2D_{313}) + 4m_{H^\pm}^2(D_{12} - D_{11} - 4D_{21} - 2D_{22} + 6D_{24} - \\
&4D_{25} + 2D_{26} - 4D_{31} + 8D_{310} + 8D_{34} - 8D_{35} - 4D_{36} - 4D_{37}) + \\
&4m_b m_t(D_0 + 4D_{11} - 4D_{12} + 4D_{13} + 4D_{21} - 4D_{24} + 4D_{25}) + 4m_t^2(4D_{12} - D_0 - 4D_{11} - \\
&4D_{13} - 4D_{21} + 4D_{24} - 4D_{25}) + 8(p_1 \cdot p_2)(D_{22} - D_{24} - D_{26} + 2D_{310} - 2D_{35} - 2D_{37}) + \\
&8(p_1 \cdot p_3)(D_{22} + D_{23} - D_{24} + D_{25} - 2D_{26} - 4D_{310} - 2D_{34} + 2D_{35} + 2D_{36} + 2D_{37}) + \\
&8(p_2 \cdot p_3)(D_{24} - D_{22} - D_{23} - D_{25} + 2D_{26}))[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t] \\
&+ 4(2(D_{27} + 4D_{311} - 2D_{312} + 2D_{313}) + m_{H^\pm}^2(D_{12} - D_{11} - 4D_{21} - 2D_{22} + \\
&6D_{24} - 4D_{25} + 2D_{26} - 4D_{31} + 4D_{34} - 4D_{35}) - m_t(m_t - m_b)(D_0 + 4D_{11} - 4D_{12} + \\
&4D_{13} + 4D_{21} - 4D_{24} + 4D_{25}) + 2(p_1 \cdot p_2)(D_{22} - D_{24} - D_{26} - 2D_{310} - 2D_{34} + 2D_{36}) + \\
&2(p_1 \cdot p_3)(D_{24} - D_{22} - D_{23} - D_{25} + 2D_{26} + 4D_{310} + 2D_{34} - 2D_{35} - 2D_{36} - 2D_{37}) + \\
&2(p_2 \cdot p_3)(D_{22} + D_{23} - D_{24} + D_{25} - 2D_{26}))[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t], \quad (A-9)
\end{aligned}$$

$$f_{2b}^- = f_{2a}^- \quad (m_b \leftrightarrow m_t), \quad (A-10)$$

$$\begin{aligned}
f_{2c}^- &= -4C_{11}[-p_1 + p_3, -p_2, m_b, m_t, m_b] - 4C_{11}[p_2 - p_3, -p_2, m_b, m_t, m_b] \\
&+ 4(m_b^2(D_{12} - D_{11} - 2D_{21} + 2D_{24}) + m_t^2(2D_{24} - D_0 - 3D_{11} + 3D_{12} - 2D_{21}) + \\
&\quad m_b m_t(D_0 + 4D_{11} - 4D_{12} + 4D_{21} - 4D_{24}) + m_{H^\pm}^2(2D_{26} - D_{13} - 2D_{25}) - 2D_{27} + \\
&\quad 2(p_1 \cdot p_2)(D_{22} - D_{24}) + 2(p_1 \cdot p_3)(D_{24} - D_{22}) + \\
&\quad 2(p_2 \cdot p_3)(D_{24} - D_{22} - D_{26}))[p_1, -p_1 + p_3, -p_2, m_t, m_b, m_t, m_b] \\
&+ 4(m_b^2(D_{12} - D_{11} - 2D_{21} + 2D_{24}) - 2D_{27} + m_{H^\pm}^2(2D_{12} - D_{13} + 4D_{24} - 2D_{25}) + \\
&\quad m_b m_t(D_0 + 4D_{11} - 4D_{12} + 4D_{21} - 4D_{24}) + m_t^2(3D_{12} - D_0 - 3D_{11} - 2D_{21} + 2D_{24}) + \\
&\quad 2(p_1 \cdot p_2)(D_{12} + D_{22} + D_{24} - D_{26}) - 2(p_1 \cdot p_3)(D_{12} + D_{22} + D_{24}) + \\
&\quad 2(p_2 \cdot p_3)(D_{26} - D_{12} - D_{22} - D_{24}))[p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b], \tag{A - 11}
\end{aligned}$$

$$f_2^+ = f_2^- [m_b \rightarrow -m_b]. \tag{A - 12}$$

$$\begin{aligned}
f_3 &= 8m_t(g_{h^0 H^\pm H^\mp} g_{h^0 tt} A_{h^0} + g_{H^0 H^\pm H^\mp} g_{H^0 tt} A_{H^0}) \cdot \\
&\quad (C_{11} + 2C_{21} + 4C_{22} - 4C_{23})[-p_3, p_1 + p_2, m_t, m_t, m_t] + i f_3^- g_-^2 + i f_3^+ g_+^2, \tag{A - 13}
\end{aligned}$$

where

$$f_3^- = f_{3a}^- + f_{3b}^- + f_{3c}^-, \tag{A - 14}$$

$$\begin{aligned}
f_{3a}^- &= 4(C_0 - C_{12})[-p_2 + p_3, -p_3, m_b, m_t, m_t] \\
&+ 4(2C_{12} - 3C_0 - 4C_{11} - 2C_{21} + 2C_{23})[p_1 - p_3, -p_4, m_b, m_t, m_t] \\
&+ 4(4(D_{27} + D_{312}) + m_{H^\pm}^2(D_{12} - D_{11} - 4D_{21} - 2D_{22} - 4D_{23} + 6D_{24} - \\
&\quad 8D_{25} + 6D_{26} - 8D_{310} - 4D_{32} - 4D_{34} + 8D_{36} + 8D_{38} - 4D_{39}) + \\
&\quad m_b m_t(D_0 + 4D_{11} - 4D_{12} + 4D_{13} - 4D_{22} + 4D_{24} + 4D_{26}) - m_t^2(D_0 + 4D_{11} - \\
&\quad 4D_{12} + 4D_{13} - 4D_{22} + 4D_{24} + 4D_{26}) + 2(p_1 \cdot p_2)(D_{22} - 2D_{23} - D_{24} - 2D_{25} + \\
&\quad D_{26} - 2D_{310} + 2D_{38} - 2D_{39}) + 4(p_1 \cdot p_3)(D_{22} + D_{23} - D_{24} + D_{25} - \\
&\quad 2D_{26} + D_{310} + D_{32} - D_{36} - 2D_{38} + D_{39})[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t] \\
&+ 4(2(2D_{313} - D_{27}) + m_{H^\pm}^2(2D_{26} - D_{11} - D_{12} - 4D_{25} - 4D_{35}) - \\
&\quad m_t(m_t - m_b)(D_0 + 4D_{13} + 4D_{25}) - 2(p_1 \cdot p_2)(D_{12} + D_{26} + 2D_{310}) + \\
&\quad 2(p_1 \cdot p_3)(D_{12} - D_{13} - D_{23} + D_{26} + 2D_{310} - 2D_{37}) + \\
&\quad 2(p_2 \cdot p_3)(D_{12} - D_{13} + D_{23} - D_{26}))[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t], \tag{A - 15}
\end{aligned}$$

$$f_{3b}^- = f_{3a}^- \quad (m_b \leftrightarrow m_t), \tag{A - 16}$$

$$\begin{aligned}
f_{3c}^- &= 4(C_0 - C_{12})[p_2 - p_3, -p_2, m_b, m_t, m_b] + 4(C_{12} + 2C_0)[-p_1 + p_3, -p_2, m_b, m_t, m_b] \\
&+ 4(m_b^2(D_{12} - D_{11} + 2D_{22} - 2D_{24} + 2D_{25} - 2D_{26}) + m_{H^\pm}^2(2D_{21} - D_{13} + \\
&\quad 2D_{22} - 4D_{24}) - 4D_{27} + m_b m_t(D_0 - 4D_{22} + 4D_{24} - 4D_{25} + 4D_{26}) + \\
&\quad m_t^2(D_{11} - D_0 - D_{12} + 2D_{22} - 2D_{24} + 2D_{25} - 2D_{26}) + \\
&\quad 2(p_1 \cdot p_2)(D_{22} - D_{24}) + 2(p_1 \cdot p_3)(D_{26} - D_{25} + 2D_{24} - 2D_{22}) + \\
&\quad 2(p_2 \cdot p_3)(D_{13} + D_{23} - D_{26}))[-p_1, -p_1 + p_3, -p_2, m_t, m_b, m_t, m_b] \\
&+ 4(m_{H^\pm}^2(2D_{13} + 2D_{25} - D_{11} - D_{12}) - 2D_{27} + m_b^2(D_{13} + 2D_{25}) + \\
&\quad m_b m_t(D_0 - 4D_{13} - 4D_{25}) + m_t^2(2D_{25} - D_0 + 3D_{13}) + 2(p_1 \cdot p_2)(D_{25} + D_{26} - \\
&\quad D_{12} + 2D_{13} - D_{23}) + 2(p_1 \cdot p_3)(D_{12} - D_{13} - D_{25} - D_{26}) + \\
&\quad 2(p_2 \cdot p_3)(D_{12} - D_{13} + D_{23} - D_{26}))[-p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b], \\
&\tag{A-17}
\end{aligned}$$

$$f_3^+ = f_3^- [m_b \rightarrow -m_b]. \tag{A-18}$$

$$\begin{aligned}
f_4 &= 8m_t(g_{h^0 H^\pm H^\mp} g_{h^0 tt} A_{h^0} + g_{H^0 H^\pm H^\mp} g_{H^0 tt} A_{H^0}) \cdot \\
&(C_{11} + 2C_{21} + 4C_{22} - 4C_{23})[-p_3, p_1 + p_2, m_t, m_t, m_t] + i f_4^- g_-^2 + i f_4^+ g_+^2, \tag{A-19}
\end{aligned}$$

where

$$f_4^- = f_{4a}^- + f_{4b}^- + f_{4c}^-, \tag{A-20}$$

$$\begin{aligned}
f_{4a}^- &= 4(C_0 - C_{12})[p_1 - p_3, -p_4, m_b, m_t, m_t] \\
&+ 4(2C_{12} - 3C_0 - 4C_{11} - 2C_{21} + 2C_{23})[-p_2 + p_3, -p_3, m_b, m_t, m_t] \\
&+ 4(2(2D_{313} - D_{27}) + m_{H^\pm}^2(D_{12} - D_{11} - 2D_{13} - 4D_{25} + 2D_{26} + \\
&\quad 4D_{310} - 4D_{35} - 4D_{37}) + m_b m_t(D_0 + 4D_{13} + 4D_{25}) - m_t^2(D_0 + 4D_{13} + 4D_{25}) - \\
&\quad 2(p_1 \cdot p_2)(D_{13} + D_{26} + 2D_{37}) + 2(p_1 \cdot p_3)(D_{23} - D_{26} - 2D_{310} + 2D_{37}) + \\
&\quad 2(p_2 \cdot p_3)(D_{26} - D_{23}))[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t] \\
&+ 4(4(D_{27} + D_{312}) + m_{H^\pm}^2(3D_{12} - D_{11} - 2D_{13} - 4D_{21} + 2D_{22} + 2D_{24} - \\
&\quad 4D_{25} - 2D_{26} - 4D_{310} - 4D_{34} + 4D_{36}) - m_t(m_t - m_b)(D_0 + 4D_{11} - 4D_{12} + \\
&\quad 4D_{13} - 4D_{22} + 4D_{24} + 4D_{26}) + 2(p_1 \cdot p_2)(D_{12} - D_{13} + 3D_{22} - 3D_{24} - 3D_{26} + \\
&\quad 2D_{32} - 2D_{36} - 2D_{38}) + 2(p_1 \cdot p_3)(D_{13} - D_{12} - 2D_{22} - 2D_{23} + 2D_{24} - 2D_{25} + \\
&\quad 4D_{26} - 2D_{310} - 2D_{32} + 2D_{36} + 4D_{38} - 2D_{39}) + \\
&\quad 2(p_2 \cdot p_3)(D_{13} - D_{12}))[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t], \\
&\tag{A-21}
\end{aligned}$$

$$f_{4b}^- = f_{4a}^- \quad (m_b \leftrightarrow m_t), \tag{A-22}$$

$$\begin{aligned}
f_{4c}^- &= 4(C_0 - C_{12})[-p_1 + p_3, -p_2, m_b, m_t, m_b] + 4(C_{12} + 2C_0)[p_2 - p_3, -p_2, m_b, m_t, m_b] \\
&+ 4(m_{H^\pm}^2(D_{12} - D_{11} + 2D_{23}) - 2D_{27} + m_b^2(D_{13} + 2D_{25}) + \\
&\quad m_b m_t(D_0 - 4D_{13} - 4D_{25}) + m_t^2(2D_{25} - D_0 + 3D_{13}) + 2(p_1 \cdot p_2)(D_{13} + D_{26}) + \\
&\quad 2(p_1 \cdot p_3)(D_{25} - D_{26}) - 2(p_2 \cdot p_3)(D_{23} + D_{26}))[-p_1, -p_1 + p_3, -p_2, m_t, m_b, m_t, m_b] \\
&+ 4(m_b^2(D_{12} - D_{11} + 2D_{22} - 2D_{24} + 2D_{25} - 2D_{26}) + \\
&\quad m_{H^\pm}^2(2D_{21} - D_{13} + 2D_{22} + 2D_{23} - 2D_{25} - 4D_{26}) + \\
&\quad m_b m_t(D_0 - 4D_{22} + 4D_{24} - 4D_{25} + 4D_{26}) - 4D_{27} + \\
&\quad m_t^2(D_{11} - D_0 - D_{12} + 2D_{22} - 2D_{24} + 2D_{25} - 2D_{26}) + \\
&\quad 2(p_1 \cdot p_2)(D_{22} + D_{23} + D_{24} - D_{25} - 2D_{26}) + 2(p_1 \cdot p_3)(D_{13} - 2D_{24} + D_{25} + D_{26}) + \\
&\quad 2(p_2 \cdot p_3)(3D_{26} - 2D_{22} - D_{23}))[-p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b], \\
&\tag{A-23}
\end{aligned}$$

$$f_4^+ = f_4^- [m_b \rightarrow -m_b]. \tag{A-24}$$

$$\begin{aligned}
f_5 &= 8m_t(g_{h^0 H^\pm H^\mp} g_{h^0 tt} A_{h^0} + g_{H^0 H^\pm H^\mp} g_{H^0 tt} A_{H^0}) \cdot \\
&(C_{11} + 2C_{21} + 4C_{22} - 4C_{23})[-p_3, p_1 + p_2, m_t, m_t, m_t] + i f_5^- g_-^2 + i f_5^+ g_+^2, \tag{A-25}
\end{aligned}$$

where

$$f_5^- = f_{5a}^- + f_{5b}^- + f_{5c}^-, \tag{A-26}$$

$$\begin{aligned}
f_{5a}^- &= 4(C_0 + C_{11} + 2C_{12} + 2C_{23})[p_1 - p_3, -p_4, m_b, m_t, m_t] \\
&+ 4(C_0 + C_{11} + 2C_{12} + 2C_{23})[-p_2 + p_3, -p_3, m_b, m_t, m_t] \\
&+ (4m_{H^\pm}^2(D_{12} - D_{11} - 2D_{13} - 4D_{23} - 4D_{25} + 2D_{26} - 4D_{310} + 4D_{38} - 4D_{39}) - \\
&\quad 4m_t(m_t - m_b)(D_0 + 4D_{13} + 4D_{26}) - 8(p_1 \cdot p_2)(D_{13} + 2D_{23} + D_{26} + 2D_{39}) + \\
&\quad 16(p_1 \cdot p_3)(D_{23} - D_{26} - D_{38} + D_{39}))[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t] \\
&+ 4(m_{H^\pm}^2(D_{12} - D_{11} - 2D_{13} - 4D_{25} - 2D_{26} - 4D_{310}) - \\
&\quad m_t(m_t - m_b)(D_0 + 4D_{13} + 4D_{26}) - 2(p_1 \cdot p_2)(D_{13} + 3D_{26} + 2D_{38}) + \\
&\quad 4(p_1 \cdot p_3)(D_{26} - D_{23} + D_{38} - D_{39}))[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t], \\
&\tag{A-27}
\end{aligned}$$

$$f_{5b}^- = f_{5a}^- \quad (m_b \leftrightarrow m_t), \tag{A-28}$$

$$\begin{aligned}
f_{5c}^- &= 4(C_0 + C_{11})[-p_1 + p_3, -p_2, m_b, m_t, m_b] + 4(C_0 + C_{11})[p_2 - p_3, -p_2, m_b, m_t, m_b] \\
&+ 4(m_b^2(D_{13} - 2D_{23} + 2D_{26}) + m_{H^\pm}^2(D_{12} - D_{11} - 2D_{25} + 2D_{26}) + \\
&\quad m_b m_t(D_0 + 4D_{23} - 4D_{26}) + m_t^2(2D_{26} - D_0 - D_{13} - 2D_{23}) + \\
&\quad 2(p_1 \cdot p_2)(D_{13} + D_{26}) - 2(p_1 \cdot p_3)(D_{13} + D_{26})) [p_1, -p_1 + p_3, -p_2, m_t, m_b, m_t, m_b] \\
&+ 4(m_{H^\pm}^2(D_{12} - D_{11} - 2D_{13} - 2D_{25}) + m_b^2(D_{13} - 2D_{23} + 2D_{26}) + \\
&\quad m_b m_t(D_0 + 4D_{23} - 4D_{26}) + m_t^2(2D_{26} - D_0 - D_{13} - 2D_{23}) + \\
&\quad 2(p_1 \cdot p_3)(D_{13} + D_{26})) [p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b], \\
\end{aligned} \tag{A-29}$$

$$f_5^+ = f_5^- [m_b \rightarrow -m_b]. \tag{A-30}$$

$$f_6 = f_{6a} + f_{6b} + f_{6c}, \tag{A-31}$$

$$\begin{aligned}
f_{6a} &= g_- g_+(8(C_{12} - C_{11})[p_1 - p_3, -p_4, m_t, m_b, m_b] - 8C_{11}[-p_2 + p_3, -p_3, m_b, m_t, m_t] \\
&+ 16((p_1 \cdot p_3)(D_{12} - D_{11} - D_{13} - D_{21} - D_{22} - D_{23} + 2D_{24} - 2D_{25} + 2D_{26}) + \\
&\quad (p_2 \cdot p_3)(D_{26} - D_{13} - D_{23} - D_{25}) - D_{27})[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t] \\
&+ 16(m_{H^\pm}^2(D_{12} + D_{22} + D_{24}) - 2D_{27} + (p_1 \cdot p_2)(D_{12} + D_{22} + D_{24}) + \\
&\quad (p_1 \cdot p_3)(D_{26} - D_{11} - D_{21} - D_{22}) + \\
&\quad (p_2 \cdot p_3)(D_{26} - D_{12} - D_{22} - D_{24}))[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t]), \\
\end{aligned} \tag{A-32}$$

$$f_{6b} = f_{6a} \quad (m_b \leftrightarrow m_t), \tag{A-33}$$

$$\begin{aligned}
f_{6c} &= 2ig_s^2 g_{ZH^\pm H^\mp} g_{Ztt}^a A_Z \cdot \{dB_0[-p_1 - p_2, m_t, m_t] - 4C_{24}[p_4, -p_1 - p_2, m_t, m_t, m_t]\} + \\
&\quad g_- g_+(8(C_{12} - C_{11})[-p_1 + p_3, -p_2, m_b, m_t, m_b] \\
&+ 8C_0 + C_{11} - C_{12})[p_2 - p_3, -p_2, m_b, m_t, m_b] \\
&- 16D_{27}[p_1, -p_1 + p_3, -p_2, m_t, m_b, m_b] + 8(m_{H^\pm}^2(D_{11} - D_{12} + D_{13} + 2D_{21} - \\
&\quad 2D_{24} + 2D_{25}) + m_b^2(D_{12} - D_{11} - D_{13}) + m_t^2(D_{11} - D_{12} + D_{13}) - 4D_{27} + \\
&\quad 2(p_1 \cdot p_2)(D_{24} - D_{22} - D_{23} - D_{25} + 2D_{26}) + \\
&\quad 2(p_1 \cdot p_3)(D_{22} - D_{24} - D_{26})) [p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b]). \\
\end{aligned} \tag{A-34}$$

$$f_7 = f_{7a} + f_{7b} + f_{7c}, \tag{A-35}$$

$$\begin{aligned}
f_{7a} = & g_- g_+ (8(C_0 + C_{11} - C_{12})[p_1 - p_3, -p_4, m_b, m_t, m_t] \\
& + 8(C_0 + C_{11} - C_{12})[-p_2 + p_3, -p_3, m_b, m_t, m_t] \\
& + 8(m_{H^\pm}^2(D_{11} - D_{12} + 2D_{13} + 2D_{21} + 2D_{22} + 4D_{23} - 4D_{24} + 6D_{25} - 6D_{26}) - \\
& \quad 2D_{27} - m_t^2 D_0 + 2(p_1 \cdot p_2)(D_{11} - D_{12} + D_{13} + D_{21} + D_{22} + 2D_{23} - 2D_{24} + 3D_{25} - \\
& \quad 3D_{26}) + 2(p_4 \cdot p_3)(D_{24} - D_{22} - D_{23} - D_{25} + 2D_{26}))[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t] \\
& + 8(m_{H^\pm}^2(D_{11} - D_{12} + 2D_{13} + 2D_{21} - 2D_{22} + 2D_{25} + 2D_{26}) - 2D_{27} - m_t^2 D_0 + \\
& \quad 2(p_1 \cdot p_2)(D_{11} - D_{12} + D_{13} + D_{21} - D_{22} + D_{25} + D_{26}) + \\
& \quad 2(p_4 \cdot p_3)(D_{22} + D_{23} - D_{24} + D_{25} - 2D_{26}))[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t]). \\
& \tag{A-36}
\end{aligned}$$

$$f_{7b} = f_{7a} \quad (m_b \leftrightarrow m_t), \tag{A-37}$$

$$\begin{aligned}
f_{7c} = & 2ig_s^2 g_{ZH^\pm H^\mp} g_{Ztt}^a A_Z \cdot \{-dB_0[-p_1 - p_2, m_t, m_t] + \\
& 4[(m_{H^\pm}^2 + p_1 \cdot p_2 - p_1 \cdot p_3 - p_2 \cdot p_3)C_{11} + dC_{24}][p_3, -p_1 - p_2, m_t, m_t, m_t]\} \\
& + g_- g_+ (-8(C_{11} + 2C_0)[-p_1 + p_3, -p_2, m_b, m_t, m_b] - \\
& \quad 8(C_{11} + 2C_0)[p_2 - p_3, -p_2, m_b, m_t, m_b] \\
& + 8(m_{H^\pm}^2(D_{13} - 2D_{21} - 2D_{22} + 4D_{24}) + m_b^2(D_{11} - D_{12}) + 6D_{27} + \\
& \quad m_t^2(D_0 - D_{11} + D_{12}) + 2(p_1 \cdot p_2)(D_{25} - D_{26}) + 2(p_1 \cdot p_3)(D_{22} - D_{24}) - \\
& \quad 2(p_2 \cdot p_3)D_{13})[p_1, -p_1 + p_3, -p_2, m_t, m_b, m_t, m_b] \\
& + 8(m_{H^\pm}^2(2D_{12} - D_{13} - 2D_{21} + 2D_{24}) + m_t^2(D_0 - D_{11} + D_{12}) + m_b^2(D_{11} - D_{12}) + \\
& \quad 6D_{27} + 2(p_1 \cdot p_2)(D_{12} - D_{13} + D_{22} - D_{24} + D_{25} - D_{26}) + 2(p_1 \cdot p_3)(D_{24} - \\
& \quad D_{12} - D_{22}) + 2(p_2 \cdot p_3)(D_{13} - D_{12}))[-p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b]). \\
& \tag{A-38}
\end{aligned}$$

$$f_8 = f_{8a} + f_{8b} + f_{8c}, \tag{A-39}$$

$$\begin{aligned}
f_{8a} = & g_- g_+ (8(C_0 - C_{11})[p_1 - p_3, -p_4, m_b, m_t, m_t] \\
& + 8(C_0 - C_{11})[-p_2 + p_3, -p_3, m_b, m_t, m_t] \\
& + 8(m_{H^\pm}^2(D_{12} - D_{11} + 4D_{23} + 2D_{25} - 2D_{26}) - 8D_{27} - m_t^2 D_0 + \\
& \quad 2(p_1 \cdot p_2)(D_{25} + 2D_{23} - D_{26}) + 2(p_1 \cdot p_3)(D_{24} - D_{22} - 2D_{23} - D_{25} + \\
& \quad 3D_{26}) + 4(p_2 \cdot p_3)(D_{26} - D_{23}))[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t] \\
& + 8(m_{H^\pm}^2(D_{12} - D_{11} + 2D_{22} + 2D_{24}) - 8D_{27} - m_t^2 D_0 + \\
& \quad 2(p_1 \cdot p_2)(D_{22} + D_{24}) + 2(p_1 \cdot p_3)(D_{25} + D_{26} - D_{22} - D_{24}) + \\
& \quad 4(p_2 \cdot p_3)(D_{26} - D_{22}))[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t]), \\
& \tag{A-40}
\end{aligned}$$

$$f_{8b} = f_{8a} \quad (m_b \leftrightarrow m_t), \quad (A - 41)$$

$$\begin{aligned}
f_{8c} = & 2ig_s^2 g_{ZH^\pm H^\mp} g_{Ztt}^a A_Z \cdot \{ dB_0[-p_1 - p_2, m_t, m_t] + \\
& 4[(p_1 \cdot p_3 + p_2 \cdot p_3 - p_1 \cdot p_2 - m_{H^\pm}^2)C_{11} - dC_{24}][p_3, -p_1 - p_2, m_t, m_t, m_t] \} \\
+ & g_- g_+ (-8(C_0 + C_{11})[-p_1 + p_3, -p_2, m_b, m_t, m_b] \\
- & 8(C_0 + C_{11})[p_2 - p_3, -p_2, m_b, m_t, m_b] \\
+ & 8(m_{H^\pm}^2(D_{11} - D_{12} + 2D_{25} - 2D_{26}) + m_t^2(D_0 + D_{13}) - m_b^2 D_{13} - \\
& 2(p_1 \cdot p_2)(D_{13} + D_{23}) + 2(p_1 \cdot p_3)(D_{13} + D_{26}))[-p_1, -p_1 + p_3, -p_2, m_t, m_b, m_t, m_b] \\
+ & 8(m_{H^\pm}^2(D_{11} - D_{12} + 2D_{13} + 2D_{25}) + m_t^2(D_0 + D_{13}) - m_b^2 D_{13} + \\
& 2(p_1 \cdot p_2)(D_{26} - D_{23}) - 2(p_1 \cdot p_3)(D_{13} + D_{26}))[-p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b]\}. \quad (A - 42)
\end{aligned}$$

$$f_9 = f_{9a} + f_{9b} + f_{9c}, \quad (A - 43)$$

$$\begin{aligned}
f_{9a} = & 16g_- g_+ ((D_{12} - D_{11} - D_{13} - D_{21} + D_{24} - D_{25})[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t] \\
+ & (D_{12} - D_{11} - D_{13} - D_{21} + D_{24} - D_{25})[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t]), \quad (A - 44)
\end{aligned}$$

$$f_{9b} = f_{9a} \quad (m_b \leftrightarrow m_t), \quad (A - 45)$$

$$\begin{aligned}
f_{9c} = & 32ig_s^2 g_{ZH^\pm H^\mp} g_{Ztt}^a A_Z \cdot (C_{23} - C_{22})[p_3, -p_1 - p_2, m_t, m_t, m_t] \\
+ & 16g_- g_+ ((D_{22} - D_{24} + D_{25} - D_{26})[p_1, -p_1 + p_3, -p_2, m_t, m_b, m_t, m_b] \quad (A - 46) \\
+ & (D_{13} + D_{25})[p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b]).
\end{aligned}$$

$$f_{10} = f_{10a} + f_{10b} + f_{10c}, \quad (A - 47)$$

$$\begin{aligned}
f_{10a} = & 16g_- g_+ ((D_{11} - D_{12} + D_{13} + D_{21} - D_{24} + D_{25})[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t] \\
+ & (D_{11} - D_{12} + D_{13} + D_{21} - D_{24} + D_{25})[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t]), \quad (A - 48)
\end{aligned}$$

$$f_{10b} = f_{10a} \quad (m_b \leftrightarrow m_t), \quad (A - 49)$$

$$\begin{aligned}
f_{10c} = & 16ig_s^2 g_{ZH^\pm H^\mp} g_{Ztt}^a A_Z \cdot (C_{11} + C_{21} + 2C_{22} - 2C_{23})[p_4, -p_1 - p_2, m_t, m_t, m_t] \\
& + 16g_- g_+(-(D_{13} + D_{25})[p_1, -p_1 + p_3, -p_2, m_t, m_b, m_t, m_b] \\
& + (D_{24} - D_{22} - D_{25} + D_{26})[p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b]).
\end{aligned} \tag{A-50}$$

$$f_{11} = f_{11a} + f_{11b} + f_{11c}, \tag{A-51}$$

$$\begin{aligned}
f_{11a} = & 16g_- g_+((D_{22} - D_{13} - D_{24} - D_{26})[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t] \\
& - (D_{13} + D_{25})[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t]),
\end{aligned} \tag{A-52}$$

$$f_{11b} = f_{11a} \quad (m_b \leftrightarrow m_t), \tag{A-53}$$

$$\begin{aligned}
f_{11c} = & 32ig_s^2 g_{ZH^\pm H^\mp} g_{Ztt}^a A_Z \cdot (C_{23} - C_{22})[p_3, -p_1 - p_2, m_t, m_t, m_t] \\
& + 16g_- g_+((D_{26} - D_{23})[p_1, -p_1 + p_3, -p_2, m_t, m_b, m_t, m_b] \\
& + (D_{26} - D_{23})[p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b]).
\end{aligned} \tag{A-54}$$

$$f_{12} = f_{12a} + f_{12b} + f_{12c}, \tag{A-55}$$

$$\begin{aligned}
f_{12a} = & 16g_- g_+((D_{13} - D_{22} + D_{24} + D_{26})[-p_1, p_1 - p_3, -p_4, m_t, m_b, m_t, m_t] \\
& + (D_{13} + D_{25})[-p_1, -p_2 + p_3, -p_3, m_t, m_b, m_t, m_t]),
\end{aligned} \tag{A-56}$$

$$f_{12b} = f_{12a} \quad (m_b \leftrightarrow m_t), \tag{A-57}$$

$$\begin{aligned}
f_{12c} = & 16ig_s^2 g_{ZH^\pm H^\mp} g_{Ztt}^a A_Z \cdot (C_{11} + C_{21} + 2C_{22} - 2C_{23})[p_4, -p_1 - p_2, m_t, m_t, m_t] \\
& + 16g_- g_+((D_{23} - D_{26})[p_1, -p_1 + p_3, -p_2, m_t, m_b, m_t, m_b] \\
& + (D_{23} - D_{26})[p_1, p_2 - p_3, -p_2, m_t, m_b, m_t, m_b]).
\end{aligned} \tag{A-58}$$

In the above form factor expressions we used the following definitions.

$$\begin{aligned}
g_{\gamma tt} &= \frac{-2i}{3}e, \\
g_{Ztt}^v &= \frac{-ig}{4\cos\theta}(1 - \frac{8}{3}\sin^2\theta),
\end{aligned}$$

$$\begin{aligned}
g_{Ztt}^a &= \frac{ig}{4 \cos \theta}, \\
g_{\gamma H^\pm H^\mp} &= -ie, \\
g_{ZH^\pm H^\mp} &= \frac{-ig \cos 2\theta}{2 \cos \theta}, \\
g_{H^0 tt} &= -ig \frac{m_t \sin \alpha}{2 m_W \sin \beta}, \\
g_{h^0 tt} &= -ig \frac{m_t \cos \alpha}{2 m_W \sin \beta}, \\
g_{H^0 H^\pm H^\mp} &= -ig(m_W \cos(\beta - \alpha) - m_Z \frac{\cos 2\beta}{2 \cos \theta_W} \cos(\beta + \alpha)), \\
g_{h^0 H^\pm H^\mp} &= -ig(m_W \sin(\beta - \alpha) + m_Z \frac{\cos 2\beta}{2 \cos \theta_W} \sin(\beta + \alpha)), \\
g_+ &= \frac{ig}{2\sqrt{2}m_W}(m_b \tan \beta + m_t \cot \beta), \\
g_- &= \frac{ig}{2\sqrt{2}m_W}(m_b \tan \beta - m_t \cot \beta), \\
A_\gamma &= \frac{-i}{s}, \\
A_Z &= \frac{-i}{s - m_Z^2}, \\
A_{H^0} &= \frac{i}{s - m_{H^0}^2 + i\Gamma_{H^0}m_{H^0}}, \\
A_{h^0} &= \frac{i}{s - m_{h^0}^2 + i\Gamma_{h^0}m_{h^0}}. \tag{A - 59}
\end{aligned}$$

The arguments of two-point, three-point and four-point integral functions are written at the end of formulae in parenthesis. The replacement $m_b \rightarrow -m_b$ is not performed on the arguments of loop intergral functions B, C and D.

References

- [1] P.W. Higgs, Phys.Lett 12(1964)132, Phys. Rev. Lett. 13 (1964)508; Phys.Rev. 145(1966)1156; F Englert and R.Brout, Phys. Rev. Lett. 13(1964)321.
- [2] G.S. Guralnik, C.R.Hagen and T.W.B. Kibble, Phys. Rev. Lett. 13 (1964)585; T.W.B. Kibble, Phys.Rev. 155(1967)1554.
- [3] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **74**, 2626(1995); D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **74**, 2632(1995).
- [4] J.Schwindling, in: Proc. Int. Europhysics Conference on High Energy Physics, Marseille, France, July 22-28, 1993, ed. by J. Carr and M. Perrottet (Editions Frontieres, Gif-sur-Yvette, 1993).
- [5] H. E. Haber, G. L. Kane, Phys. Rep. 117(1985) 75; J. F. Gunion, H. E. Haber, Nucl. Phys. **B272**(1986) 1.
- [6] T.Plehn, M.Spira and P.M. Zerwas, Nucl. Phys. **B479**(1996)46.
- [7] D.Bowser-Chao, K.Cheung, and S.Thomas, Phys. Lett **B315**, 399(1993); Ma Wen-Gan, Chong-Sheng Li and Hang Liang, Phys. Rev. **D53**,1304(1996).
- [8] M. Gell-Mann, Phys. Rev. **125**, 1067(1962).
- [9] A.D. Martin, W.J. Stirling and R.G. Roberts, Phys. Lett.**B354**, 155(1995).
- [10] Particle Data Group, Phys. Rev. **D 50**, No.3(1994); D. Schaile, CERN-PPE/94-162(11 October 1994).

[11] Bernd A. Kniehl, Phys. Rep. 240(1994)211 and the references therein.

Figure captions

Fig.1 The Feynman diagram of subprocess $gg \rightarrow H^+H^-$. (a,b,c,d)s-channel diagrams. (e ~ j)box diagrams including t- and u-channels.

Fig.2(1) Total cross sections of the subprocess $gg \rightarrow H^+H^-$ as the functions of the $\sqrt{\hat{s}}$ with $m_{h^0} = 100 \text{ GeV}$, $m_{H^0} = 150 \text{ GeV}$ and $m_{H^\pm} = 150 \text{ GeV}$. The full line is for $\tan \beta = 1.5$. The dashed line is for $\tan \beta = 30$.

Fig.2(2) Total cross sections of the subprocess $gg \rightarrow H^+H^-$ as the functions of the $\sqrt{\hat{s}}$ with $m_{h^0} = 100 \text{ GeV}$, $m_{H^0} = 150 \text{ GeV}$ and $m_{H^\pm} = 300 \text{ GeV}$. The full line is for $\tan \beta = 1.5$. The dashed line is for $\tan \beta = 30$.

Fig.3 Total cross sections of the subprocess $gg \rightarrow H^+H^-$ as the functions of the $\tan \beta$ with $m_{h^0} = 100 \text{ GeV}$, $m_{H^0} = 150 \text{ GeV}$ and $m_{H^\pm} = 150 \text{ GeV}$. The full line and dashed line correspond to $\sqrt{\hat{s}} = 400 \text{ GeV}$ and $\sqrt{\hat{s}} = 800 \text{ GeV}$ respectively.

Fig.4 Total cross sections of the subprocess $gg \rightarrow H^+H^-$ as the functions of the m_{H^\pm} with $m_{h^0} = 100 \text{ GeV}$, $m_{H^0} = 150 \text{ GeV}$ and $\sqrt{\hat{s}} = 1 \text{ TeV}$. The full line is for $\tan \beta = 1.5$. The dashed line is for $\tan \beta = 30$.

Fig.5(1) Total cross sections of the process $pp \rightarrow gg \rightarrow H^+H^- + X$ as the functions of

the \sqrt{s} with $m_{h^0} = 100 \text{ GeV}$, $m_{H^0} = 150 \text{ GeV}$ and $m_{H^\pm} = 150 \text{ GeV}$. The full line is for $\tan \beta = 1.5$. The dashed line is for $\tan \beta = 30$.

Fig.5(2) Total cross sections of the process $pp \rightarrow gg \rightarrow H^+H^- + X$ as the functions of the \sqrt{s} with $m_{h^0} = 100 \text{ GeV}$, $m_{H^0} = 150 \text{ GeV}$ and $m_{H^\pm} = 300 \text{ GeV}$. The full line is for $\tan \beta = 1.5$. The dashed line is for $\tan \beta = 30$.

Fig.3




Fig.4




Fig.2(1)




Fig.2(2)




Fig.5(1)




Fig.5(2)

