## CP-Violation in b Quark Radiative Inclusive Decays

H. M. Asatrian<sup>a</sup>

International Centre for Theoretical Physics, Strada Costiera 11, P.O. Box 586, 34100 Trieste, Italy e-mail: hrachia@moon.yerphi.am

G. K. Yeghiyan<sup>a</sup>

Deutches Electronen Synchrotron DESY, Hamburg, Germany

A. N. Ioannissian<sup>a</sup> Dept. of Physics, Technion - Israel Inst. of Tech., Haifa, Israel

## Abstract

The direct CP-violations in  $SU(2)_L \times SU(2)_R \times U(1)$  model and two-Higgs doublet extension of the standard model for  $b \to d\gamma$  and  $b \to s\gamma$  decays are investigated. The calculated value of CP-asymmetry for these two models and for  $b \to d\gamma$  and  $b \to s\gamma$  decays for the wide range of parameters may exceed the value, predicted by the standard model and has a sign opposite to that of the latter.

<sup>a</sup> Permanent address: Yerevan Physics Institute, 2 Alikhanyan Br., 375036, Yerevan, Armenia The investigation of rare B-meson decays can give an important information on new physics in the TeV region. The observation of direct CP asymmetry in B-meson decays will help to understand the CP breaking phenomenon.

The first experimental evidence for the exclusive  $\bar{B} \to K^* \gamma$  decay has been obtained at CLEO [1]. More recently, the branching ratio of the inclusive  $B \to X_s \gamma$  decay was measured [2].

The  $b \to s\gamma$  decay has been investigated theoretically for the standard model and its extensions in [3]-[12]. CP-violation in  $B - \bar{B}$  system in  $SU(2)_L \times$  $SU(2)_R \times U(1)$  model was considered in [13]. The problem of CP asymmetry for  $b \to s\gamma$  decay for standard model and its extensions was investigated in [14, 15, 16].

Although the expected decay rate for  $b \to d\gamma$  decay is about 10-20 times smaller than for the  $b \to s\gamma$  decay, the CP asymmetry for the first decay can be about 10 times larger [14, 17].

The aim of the present paper is to consider the direct CP decay asymmetry for the  $b \to d\gamma$  and  $b \to s\gamma$  decays for  $SU(2)_L \times SU(2)_R \times U(1)$  model and two Higgs doublet extension of the standard model.

In  $SU(2)_L \times SU(2)_R \times U(1)$  model the  $b \to d\gamma$  decay amplitude arises due to the interaction of quark charged weak current with the "left" and "right" W-bosons and charged Higgs field. This interaction has the following form [16]:

$$L^{ch} = \frac{1}{\sqrt{2}} \left( \bar{u}, \bar{c}, \bar{t} \right) \left[ \hat{W}_{1}^{+} \left[ -g_{L} \cos \beta K^{L} P_{-} - g_{R} \sin \beta e^{i\delta} K^{R} P_{+} \right] + \varphi^{+} \frac{g_{L}}{\sqrt{2} M_{W_{L}}} \left[ \left( -\tan 2\theta K^{L} M_{d} + e^{i\delta} \frac{1}{\cos 2\theta} M_{u} K_{R} \right) P_{+} + \left( \tan 2\theta M_{u} K_{L} - e^{i\delta} \frac{1}{\cos 2\theta} K^{R} M_{d} \right) P_{-} \right] \right] \begin{pmatrix} d \\ s \\ b \end{pmatrix}, \qquad (1)$$

where  $W_1$  is the "light" (predominantly left-handed) charged gauge boson and  $\beta$  is the mixing angle between left and right W - bosons,

$$\tan 2\beta = 2\sin 2\theta \frac{g_R}{g_L} \frac{M_{W_L}^2}{M_{W_R}^2} \quad \tan \theta = -\frac{k'}{k},\tag{2}$$

 $K^L$  and  $K^R$  are CKM mixing matrices for left and right charged currents respectively,  $P_{\pm} = (1 \pm \gamma_5)/2$ ,  $M_u$  and  $M_d$  are diagonal mass matrices for quarks with Q=2/3 and Q=-1/3 charges respectively. The matrices  $K^L$  and  $K^R$  can be expressed in a form, where  $K^L$  has only one complex phase and  $K^R$  has five complex phases [18]. Phase  $\delta$  in (2) takes his origin from the vacuum expectation value of Higgs field  $\Phi$ , connected with the  $SU(2)_L \times U(1)$ 

symmetry breaking:  $\Phi = \begin{pmatrix} k & 0 \\ 0 & -e^{i\delta}k' \end{pmatrix}$ . In (1) the term connected with the interaction with heavy (predominantly right) W- boson is omitted, since it is not relevant for  $b \to d\gamma$  decay.

The additional (compared with the standard model) phase factor  $exp(i\delta)$ in (1) leads to the existence of the new CP violation effects in  $SU(2)_L \times SU(2)_R \times U(1)$  model.

We define the direct CP asymmetry for  $b \to d\gamma$  decay as [14]:

$$a_{cp} = \frac{\Gamma(\bar{b} \to \bar{d}\gamma) - \Gamma(b \to d\gamma)}{\Gamma(\bar{b} \to \bar{d}\gamma) + \Gamma(b \to d\gamma)}$$
(3)

The direct CP asymmetry for  $b \to d\gamma$  decay arises only if the matrix element of decay has an absorptive part, which arises if the final state strong interaction effects are taken into account. In general case the amplitudes of the decays  $\bar{b} \to \bar{d}\gamma$  and  $b \to d\gamma$  can be expressed in the following form [14]:

$$A(\bar{b} \to \bar{d}\gamma) = \sum_{a} (A_a^r + iA_a^i) V_a^*$$
$$A(b \to d\gamma) = \sum_{a} (A_a^r + iA_a^i) V_a$$
(4)

where  $A_a^r$ ,  $A_a^i$  are real and absorptive parts of amplitudes,  $V_a$  are some phases and CKM-type factors, a=1,2,3.... Then CP asymmetry is given by:

$$a_{CP} = \frac{\sum_{a \neq b} (A_a^r A_b^i - A_a^i A_b^r) Im(V_a^* V_b)}{\sum_{a,b} (A_a^r A_b^r + A_a^i A_b^i) Re(V_a^* V_b)}$$
(5)

To take into account QCD- corrections to radiative decays matrix elements the effective Hamiltonian approach is used. We follow to [16] and use the results of [12] for the imaginary part of the amplitude, connected with the  $O_2$  operator:

$$Imr_{2} = \frac{16\pi}{81} \left[ -5 + (45 - 3\pi^{2} + 9L + 9L^{2})z + (-3\pi^{2} + 9L^{2})z^{2} + (28 - 12L)z^{3} \right]$$

where  $z = (m_c/m_b)^2$ , L=lnz. We take the ratio of c- and b- quark masses equal to 0.29 [12], then the ratio of imaginary parts of the amplitudes connected with  $\bar{c}c$  and  $\bar{u}u$  intermediate states is approximately equal to r=0.145. We obtain the following expression for effective Hamiltonian of  $b \to d\gamma$  decay in  $SU(2)_L \times SU(2)_R \times U(1)$  model:

$$H_{b\to d\gamma} = -\frac{e}{16\pi^2} \frac{2G_F}{\sqrt{2}} m_b \{ (K_{td}^{L*} K_{tb}^L A_{d\gamma}^{W_L} + e^{i\delta} K_{td}^{L*} K_{tb}^R A_{d\gamma}^R) O_7^L \\ + e^{-i\delta} K_{td}^{R*} K_{tb}^L A_{d\gamma}^R O_7^R + i [(\frac{2}{9} (K_{td}^{L*} K_{tb}^L A_{dg}^{W_L} + e^{i\delta} K_{td}^{L*} K_{tb}^L K_{tb}^R A_{dg}^R) (6) \\ + \frac{20}{81} (K_{ud}^{L*} K_{ub}^L y + r K_{cd}^{L*} K_{cb}^L) c_2) O_7^L + \frac{2}{9} e^{-i\delta} K_{td}^{L*} K_{tb}^L K_{td}^R A_{dg}^R O_7^R] \},$$

where

$$O_7^{L,R} = \bar{u}_d \sigma^{\mu\nu} (1 \pm \gamma_5) u_b F_{\mu\nu}$$
(7)

and the functions  $A_{d\gamma}^{W_L}$ ,  $A_{d\gamma}^R$ ,  $A_{dg}^{W_L}$ ,  $A_{dg}^R$ , which include leading logarithmic strong interaction corrections, were presented in [16].

Using (5) and (6), one obtains the following formula for CP asymmetry for  $b \to d\gamma$  decay in  $SU(2)_L \times SU(2)_R \times U(1)$  model:

$$a_{cp}(b \to d\gamma) = \frac{2\alpha_s}{(|C_7^L|^2 + |C_7^R|^2)v_t^*v_t} \left\{ (\operatorname{Im} v_t^* v_u + r \operatorname{Im} v_t^* v_c) \times (A_{d\gamma}^{W_L} + H \cos \alpha A_{d\gamma}^R) \frac{20c_2}{81} - (\operatorname{Re} v_t^* v_u + r \operatorname{Re} v_t^* v_c) \times (8) \times A_{d\gamma}^R H \sin \alpha \frac{20c_2}{81} + \frac{2}{9} H \sin \alpha v_t^* v_t (A_{d\gamma}^{W_L} A_{dg}^R - A_{d\gamma}^R A_{dg}^{W_L}) \right\},$$

where

$$He^{i\alpha} \equiv e^{i\delta} \frac{K_{Rtb}}{K_{Ltb}}, \quad C_7^L = A_{d\gamma}^{W_L} + \left(e^{i\delta} \frac{K_{tb}^R}{K_{tb}^L}\right) A_{d\gamma}^R, \quad C_7^R = e^{-i\delta} \frac{K_{td}^{R^*}}{K_{td}^L} A_{d\gamma}^R$$
$$v_t \equiv K_{Ltd}^* K_{Ltb}, \quad v_c \equiv K_{Lcd}^* K_{Lcb}, \quad v_u \equiv K_{Lud}^* K_{Lub} \tag{9}$$

The expression for CP asymmetry in  $b \to s\gamma$  (in the limit  $m_d=0, m_s=0$ ) decay can be obtained from (8) by making replacements  $v_u \to v'_u, v_c \to v'_c, v_t \to v'_t$ , where

$$v'_{t} \equiv K^{*}_{Lts}K_{Ltb}, \qquad v'_{c} \equiv K^{*}_{Lcs}K_{Lcb}, \qquad v'_{u} \equiv K^{*}_{Lus}K_{Lub} \tag{10}$$

We want to stress that in [14, 15, 16] the approximate value for r was used: r=0.12. Now we use the correct value r=0.145 [12], which is essential for the numerical results.

Let us proceed to the numerical results. We take  $\alpha_s = 0.212$ ,  $c_2 \simeq 1.1$ ,  $m_t = (175 \pm 9)$ GeV,  $m_b = 4.5 GeV$ ,  $m_b^* \equiv m_b(M_Z) = (3.5 \pm 0.5) GeV$  [19, 20]. For CKM matrix parameters we use Wolfenstein parametrization:

$$Im(v_t^*v_u) = -A^2\lambda^6\eta \qquad Re(v_t^*v_u) = A^2\lambda^6((1-\rho)\rho - \eta^2)$$
$$|v_t|^2 = A^2\lambda^6((1-\rho)^2 + \eta^2) \qquad |v_u|^2 = A^2\lambda^6(\rho^2 + \eta^2)$$
$$|v_t'|^2 = A^2\lambda^4((1+\lambda^2\rho)^2 + \lambda^8\eta^2) \qquad Im(v_t^{'*}v_u') = A^2\lambda^6\eta \qquad (11)$$
$$Re(v_t^{'*}v_u') = -A^2\lambda^6(\rho + \lambda^2(\rho^2 + \eta^2)) \qquad |v_u'|^2 = A^2\lambda^8(\rho^2 + \eta^2)$$

For parameters  $\lambda$ , A,  $\rho$ ,  $\eta$  in (11) we use values given in [20]. The CP asymmetry depends also on parameters of  $SU(2)_L \times SU(2)_R \times U(1)$ model:  $\alpha$ ,  $tan2\theta$ ,  $M_{W_R}$ ,  $M_{\varphi^+}$ ,  $|K_{tb}^R/K_{tb}^L|$ ,  $|K_{td}^R/K_{td}^L|$ . We will consider the following possibilities for the ratios  $|K_{tb}^R/K_{tb}^L|$ ,  $|K_{td(s)}^R/K_{td(s)}^L|$ :

(1) 
$$|K_{tb}^{R}/K_{tb}^{L}| = |K_{td(s)}^{R}/K_{td(s)}^{L}| = 1$$

(2) No restrictions on ratios  $|K_{tb}^R/K_{tb}^L|$ ,  $|K_{td(s)}^R/K_{td(s)}^L|$ , besides those which follow from the unitarity conditions for matrices  $K^L$  and  $K^R$ .

The case (1) corresponds to the pseudo-manifest left-right symmetry, when the absolute values of Kobayashi-Maskawa mixing matrices elements in left and right sectors ( $K_{ij}^L$  and  $K_{ij}^R$ , i,j=1,2,3 correspondingly) are equal to each other [18]. The case (2) corresponds to non-manifest left-right symmetry when  $|K_{ij}^L| \neq |K_{ij}^R|$  [18].

It is known that the experiment is in agreement with the standard model predictions for  $b \to s\gamma$  decay rate. Following to [16], we will consider that in  $SU(2)_L \times SU(2)_R \times U(1)$  model the branching of  $b \to s\gamma$  decay can differ from the standard model prediction no more than  $\Delta = 10\%$ . As for  $b \to d\gamma$ decay rate, there is no experimental restriction for it. However, if we assume that in  $SU(2)_L \times SU(2)_R \times U(1)$  model the  $b \to s\gamma$  decay rate is the same (with 10% accuracy) as in standard model, then for the case (1), the same condition will be satisfied for the  $b \to d\gamma$  decay rate also. For the case (2) we will consider two possibilities:

(2a)  $b \to d\gamma$  decay rate is equal with accuracy of  $\Delta = 10\%$  to that in the standard model

(2b)  $b \rightarrow d\gamma$  decay rate is arbitrary.

For case (1) and for a given  $M_{W_R}$ ,  $M_{\varphi^+}$  the decay asymmetry for  $b \to d\gamma$  and

 $b \to s\gamma$  decays depends on CKM parameters,  $\alpha$  and  $tan 2\theta$ . Taking into account (8) and the equivalence of decay rates (with 10% accuracy) in standard model and  $SU(2)_L \times SU(2)_R \times U(1)$  model, it is easy to understand, that for  $\alpha = 0$  the absolute value of decay asymmetry for both of decays can't exceed the standard model value more than 10% for all the values  $M_{W_R}$ ,  $M_{\varphi^+}$ ,  $tan2\theta$ . The sign of the decay asymmetry will be the same as in the standard model. When we have a new source of CP violation, i.e.  $\alpha \neq 0$ . the terms in (8) proportional to  $A_{d\gamma}^R$  and  $A_{d\gamma}^{W_L}A_{dg}^R - A_{d\gamma}^RA_{dg}^{W_L}$  contribute to decay asymmetry and one can expect less or more significant deviations from the standard model predictions. However, the restriction for decay rate here also plays the important role. In Fig 1 the  $tan 2\theta$  dependence of maximum and minimum values of  $a_{CP}$  for  $b \to d\gamma$  and  $b \to s\gamma$  decays for the case (1) are given for various values of  $M_{W_R}$ ,  $M_{\varphi^+}$ . Due to the presence of the terms proportional to  $sin\alpha$  in (8) there is a difference between standard model predictions  $(-a_{CP}(b \to s\gamma)/10^{-3} = 2.9 \div 6.4, a_{CP}(b \to d\gamma)/10^{-2} = 3.7 \div 16),$ which practically coincide with the results for  $M_{W_R} = M_{\varphi^+} = 50$  TeV and predictions of the  $SU(2)_L \times SU(2)_R \times U(1)$  model for  $M_{W_R}, M_{\varphi^+} \leq 20TeV$ ,  $tan2\theta \geq 1$ . The difference is most significant for  $b \to s\gamma$  decay. The sign of asymmetry can be different from those in the standard model for both decays. To illustrate the  $\alpha$  dependence of the decay asymmetry we give in the Table 1 minimum and maximum values of decay asymmetry for two decays for  $tan2\theta = 2$ ,  $M_{W_R} = M_{\varphi^+} = 10TeV$  and various values of  $\alpha$  (for  $2.30 \le |\alpha| \le 3.14$  and  $|\alpha| \le 1.9$  the condition for decay rate is not satisfied). As we have mentioned above, the difference between results for  $a_{CP}$  when taking r=0.12 or 0.145 in the expression (8) is non-negligible: for  $b \to s\gamma$ decay it can reach 30%. For this reason values of  $a_{CP}$  in Fig 1 are lower than those in [16].

Let us now proceed to the case (2). In Fig 2a we give the  $tan2\theta$  dependence of  $a_{CP}$  for  $b \to d\gamma$  decay for case (2a) and for  $b \to s\gamma$  decay for case (2). It is obvious that for the case (2), when there are no restrictions on right current mixing matrix (besides unitarity condition), the decay asymmetry in  $SU(2)_L \times SU(2)_R \times U(1)$  model is much more different from the standard model predictions, than for the case (1). Indeed, for the case (2) the minimum value of the asymmetry for  $b \to d\gamma$  decay can reach (for  $M_{W_R} = M_{\varphi^+} = 5TeV$ ) the value -0.18, while for the previous case the minimum value of  $a_{CP}$  for the same values  $M_{W_R}$  and  $M_{\varphi^+}$  is equal to -0.02. For  $b \to s\gamma$  decay the absolute value of decay asymmetry for the same values

 $M_{W_R}$  and  $M_{\varphi^+}$  is 1.5-2 times higher than for the previous case.

We give in Table 2 minimum and maximum values of decay asymmetry for two decays for  $tan2\theta = 3$ ,  $M_{W_R} = M_{\varphi^+} = 10TeV$  and various values of  $\alpha$  (for  $\alpha \leq 1.80$  the conditions for decay rate is not satisfied). It is clear that for the case (2a) the deviations from the standard model predictions are more significant and can take place for a larger parameter space, than for the case (1).

In Fig 3 for the case (2b) (when we have no restriction for  $b \to d\gamma$  decay rate) the decay asymmetry minimum and maximum values are given. We note, that for some values of  $tan2\theta$ ,  $M_{W_R}$ ,  $M_{\varphi^+}$  the decay asymmetry  $a_{CP}$  from (8) becomes abnomally large. This means, that for such values of  $tan2\theta$ ,  $M_{W_R}$ ,  $M_{\varphi^+}$  (8) becomes incorrect (imaginary part of the amplitude becomes non-small in comparison with the real part and it is necessary to take into account more terms of perturbation theory on  $\alpha_s$ ). Nevertheless, it is reasonably safe to suggest that in this case the difference from the standard model predictions for  $M_{W_R}, M_{\varphi^+} \leq 10TeV$  can be significant. Thus, for the case of non-manifest left-right symmetry for the large parameter space of the  $SU(2)_L \times SU(2)_R \times U(1)$  model  $(M_{W_R}, M_{\varphi^+} \leq (10-15)TeV,$  $\tan 2\theta \geq (1.5-2.5))$  one can expect a significant deviations from the standard model predictions for  $a_{CP}$  for both decays.

Let us proceed to the two Higgs doublet extension of the standard model. In general case Yukawa interaction of quarks with Higgs doublets  $\varphi_1$  and  $\varphi_2$  is:

$$L = \bar{q}_L(\gamma_1^d \varphi_1 + \gamma_2^d \varphi_2) d_R + \bar{q}(\gamma_1^u \varphi_1 + \gamma_2^u \varphi_2) u_R, \qquad (12)$$

where  $q_L$  is the quark doublet and  $d_R$  and  $u_R$  are quark singlets and  $\gamma_1^u$ ,  $\gamma_2^u$ ,  $\gamma_1^d$ ,  $\gamma_2^d$  are matrices in flavor space [15]. Usually two versions of this model are considered [15]:

model I, where only one doublet  $(\varphi_1)$  interacts with quarks:  $\gamma_2^u = \gamma_2^d = 0$ , model II, where one of doublets interacts with up-type quarks and the second one interacts with down-type quarks:  $\gamma_2^u = \gamma_1^d = 0$ .

In paper [15] the model was considered where both Higgs doublets interact with up and down quarks and all of the quantities  $\gamma_1^u$ ,  $\gamma_2^u$ ,  $\gamma_1^d$ ,  $\gamma_2^d$  are non-zero. Generally speaking, in this case the flavor changing neutral currents can arise [15]. The restrictions on Higgs particles masses and other parameters in such a model were considered in [21].

The last model (model III) is close in some respect to the  $SU(2)_L \times SU(2)_R \times U(1)$  model: for this model, as for the previous one, new CP-violating phase

arises. As for models I and II, there are no new sources of CP violation. Formula for CP asymmetry in  $b \rightarrow s\gamma$  decay for two Higgs doublet extension of the standard model is the following [15]:

$$a_{CP}(b \to s\gamma) = - 2\frac{2}{9} \frac{(C_{s\gamma}^{H} C_{sg}^{WH} - C_{s\gamma}^{WH} C_{sg}^{H})\alpha_{s}}{C_{s\gamma}^{2}} Im(\xi_{t}\xi_{b}) - 2\frac{20}{81} \frac{Re(v_{t}^{*}v_{u}) + Re(v_{t}^{*}v_{c})}{|v_{t}|^{2}} \frac{C_{s\gamma}^{H}c_{1}\alpha_{s}}{2C_{s\gamma}^{2}} Im(\xi_{t}\xi_{b})$$
(13)  
$$- 2\frac{20}{81} \frac{(1-r)Im(v_{t}^{*}v_{u})}{|v_{t}|^{2}} \frac{(C_{s\gamma}^{WH} + Re(\xi_{t}\xi_{b})C_{s\gamma}^{H})c_{1}\alpha_{s}}{2C_{s\gamma}^{2}}$$

where  $\xi_t$ ,  $\xi_b$ ,  $C_{s\gamma}^H$ ,  $C^{WH}$ ,  $C_{s\gamma}^{WH}$ ,  $C_{sg}^H$ ,  $C_{s\gamma}$  are given in [15]. We note, that there is a difference between formula (13) and the expression for CP asymmetry in [15]: in [15] the factor 2/9 is missed.

In Table 3 the numerical results for the model III for some values of charged Higgs boson masses are given. Generally speaking, values of  $a_{CP}$  for  $b \rightarrow s\gamma$  decay, given by Table 3, are lower than the results [15] for the reason, mentioned above, but the deviation from the standard model predictions for relatively low masses of charged Higgs boson  $\leq 200$ GeV (this is within the limits given in [21]) can be very large (more than 5 times). As for the  $b \rightarrow d\gamma$  decay (as it follows from the Table 3 a)) the restrictions on absolute value  $a_{CP}$  are close to the predictions of the standard model. The difference is that CP-asymmetry here can have opposite sign and the minimum value of  $a_{CP}$  can be very small. In Table 3 b) the minimum and maximum values of CP asymmetry for  $b \rightarrow d\gamma$  decay, for the case when we use the less severe condition  $\Delta < 50\%$  for the decay rate, are given. In this case the absolute value value of CP asymmetry can be 1.5 times larger than the standard model predictions.

As we have mentioned above, for the models I and II there is no new source of CP-violation and as it follows from the Tables 4 and 5, the values of  $a_{CP}$  for two decays are almost the same as for the standard model.

In conclusion, the direct CP-violation in  $SU(2)_L \times SU(2)_R \times U(1)$  model and two-Higgs doublet extension of the standard model for  $b \to d\gamma$  and  $b \to s\gamma$ decays was investigated. The calculated values of CP-asymmetry differ from the standard model predictions and can have a sign opposite to that of the latter. The difference is much stronger for non-manifest left-right symmetric model and two-Higgs doublet extension of the standard model (model III). Authors want to thank A. Ali for stimulating discussions. One of the authors (H. A.) wants to thank High Energy Group of ICTP for hospitality. The research described in this publication was made possible in part due to the contract INTAS-93-1630.

## References

- [1] R.Ammar et al. (CLEO Collaboration), Phys.Rev.Lett. 71 (1993) 674.
- [2] M. S. Alam et al. (CLEO Collaboration), Phys.Rev.Lett. 74 (1995) 2885.
- [3] B. Grinstein and M. B. Wise, Phys. Lett. B201, (1988), 274.
- [4] W. S. Hou, R. S. Willey, Phys. Lett. B202, (1988), 591.
- [5] R. Barbieri and G. F. Guidice, Phys. Lett. B309, (1993), 86.
- [6] S. Bertolini et al. Nucl. Phys. B353, (1991), 591.
- [7] L. Randal and R Sundrum, MIT preprint MIT-CTP-2211 (1993).
- [8] D.Cocolicchio et al, Phys.Rev. D40 (1989) 1477.
- [9] H.M.Asatrian and A.N.Ioannissian, Mod.Phys.Lett.A5 (1990) 1089; Sov. Journ. of Nucl. Phys. 51 (1990) 858.
- [10] K.S.Babu, K.Fujikawa and A.Yamada, Phys.Lett. B333 (1994) 196.
- [11] P.Cho and M.Misiak, Phys.Rev. D49 (1994) 5894.
- [12] C. Greub, T. Hurth, D. Wyler Phys. Rev. D54 (1996) 3350.
- [13] G. Ecker and W. Grimus, in: Moriond 86, v.1, p. 201.
- [14] J.Soares, Nucl.Phys.B 367 (1991) 575.
- [15] L.Wolfenstein and Y.L.Wu, Phys.Rev.Lett. 73 (1994) 2809.
- [16] H. M. Asatrian, A. N. Ioannissian, Phys. Rev. D54 (1996) 5242.

- [17] A. Ali DESY Report No. 96-106, hep-ph/9606324.
- [18] R.N.Mohapatra, in "CP Violation" ed. by C.Jarslkog, World Scientific, page 384.
- [19] M. Bando et al., Mod. Phys. Lett. A7, 3379 (1992)
- [20] A. Ali, D. London. DESY Report No.96-140, hep-ph/9607392.
- [21] D. Atwood, L. Reina, A. Soni. CEBAF preprint, JL-TH-96-15, hepph/9609279.

## **Figure Captions**

Fig. 1. Maximum and minimum values of  $a_{CP}$  in  $SU(2)_L \times SU(2)_R \times U(1)$ model for the case (1) and a) for  $b \to d\gamma$  decay, b) for  $b \to s\gamma$  decay for different values of  $M_{W_R}$  and  $M_{\varphi^+}$ :  $M_{W_R} = M_{\varphi^+} = 5$ TeV (curves 1 and 2);  $M_{W_R} = M_{\varphi^+} = 10$ TeV (curves 3 and 4);  $M_{W_R} = M_{\varphi^+} = 20$ TeV (curves 5 and 6);  $M_{W_R} = M_{\varphi^+} = 50$ TeV (curves 7 and 8).

Fig. 2. Maximum and minimum values of  $a_{CP}$  in  $SU(2)_L \times SU(2)_R \times U(1)$ model for the case (2a) and a) for  $b \to d\gamma$  decay, b) for  $b \to s\gamma$  decay for different values of  $M_{W_R}$  and  $M_{\varphi^+}$ :  $M_{W_R} = M_{\varphi^+} = 5$ TeV (curves 1 and 2);  $M_{W_R} = M_{\varphi^+} = 10$ TeV (curves 3 and 4);  $M_{W_R} = M_{\varphi^+} = 20$ TeV (curves 5 and 6);  $M_{W_R} = M_{\varphi^+} = 50$ TeV (curves 7 and 8).

Fig. 3. Maximum and minimum values of  $a_{CP}$  in  $SU(2)_L \times SU(2)_R \times U(1)$ model for the case (2b) and for  $b \to d\gamma$  decay for different values of  $M_{W_R}$  and  $M_{\varphi^+}$ :  $M_{W_R} = M_{\varphi^+} = 5$ TeV (curves 1 and 2);  $M_{W_R} = M_{\varphi^+} = 10$ TeV (curves 3 and 4);  $M_{W_R} = M_{\varphi^+} = 20$ TeV (curves 5 and 6);  $M_{W_R} = M_{\varphi^+} = 50$ TeV (curves 7 and 8).

Table 1: Minimum and maximum values of  $a_{CP}$  for case (1) for  $b \to d\gamma$  and  $b \to s\gamma$  decays for  $tan2\theta = 2.0$ ,  $M_{W_R} = M_{\varphi^+} = 10$  TeV and for various values of  $\alpha$ .

$\alpha$	$a_{CP}(b \to d\gamma)$	$a_{CP}(b \to s\gamma)$
-2.2	$0.043 \div 0.107$	$-0.0101 \div -0.0045$
-2.0	$0.041 \div 0.130$	$-0.096 \div -0.0040$
-1.8	$0.049 \div 0.123$	$-0.0087 \div -0.0039$
1.8	$0.0085 \div 0.162$	$-0.0037 \div 0.0019$
2.0	$-0.0024 \div 0.172$	$-0.0039 \div 0.0039$
2.2	$-0.0027 \div 0.160$	$-0.0018 \div 0.0044$

Table 2: Minimum and maximum values of  $a_{CP}$  for case (2a) for  $b \to d\gamma$ and  $b \to s\gamma$  decays for  $tan2\theta = 3.0$ ,  $M_{W_R} = M_{\varphi^+} = 10$ TeV and for various values of  $\alpha$ .

$\alpha$	$a_{CP}(b \to d\gamma)$	$a_{CP}(b \to s\gamma)$
-3.0	$-0.017 \div 0.044$	$-0.0033 \div -0.0045$
-2.5	$-0.035 \div 0.062$	$-0.0114 \div -0.0026$
-2.0	$0.008 \div 0.110$	$-0.0145 \div -0.0042$
2.0	$-0.031 \div 0.188$	$-0.0134 \div 0.0102$
2.5	$-0.036 \div 0.108$	$-0.0008 \div 0.0104$
3.0	$-0.012 \div 0.055$	$-0.0018 \div 0.0028$

a)		
$m_{H^+}, GeV$	$-a_{CP}(b \to s\gamma)/10^{-3}$	$a_{CP}(b \to d\gamma)/10^{-2}$
50	$-53 \div 53$	$-17 \div 17$
100	$-45 \div 45$	$-17 \div 17$
200	$-35 \div 35$	$-18 \div 18$
400	$-25 \div 25$	$-18 \div 18$
800	$-16 \div 16$	$-18 \div 18$
1600	$-9.7 \div 9.5$	$-17 \div 18$
3200	$0.6 \div 7$	$0.6 \div 17$
6400	$2 \div 7$	$3.1 \div 17$
12800	$2.6 \div 6.8$	$3.5 \div 17$
25600	$2.8 \div 6.5$	$3.6 \div 16$

Table 3: Values of CP-asymmetry for  $b \to s + \gamma$  and  $b \to d + \gamma$  for model III. a) for  $\Delta < 10\%$ , b) for  $\Delta < 50\%$ 

b)

$m_{H^+}, GeV$	$-a_{CP}(b \rightarrow s\gamma)/10^{-3}$	$a_{CP}(b \rightarrow d\gamma)/10^{-2}$
50	$-71 \div 71$	$-23 \div 23$
100	$-61 \div 61$	$-23 \div 23$
200	$-48 \div 48$	$-23 \div 23$
400	$-34 \div 34$	$-24 \div 24$
800	$-22 \div 21$	$-24 \div 24$
1600	$-12 \div 12$	$-22 \div 25$
3200	$-5 \div 11$	$-4.1 \div 25$
6400	$0.4 \div 8.3$	$2.4 \div 20$
12800	$2.3 \div 6.8$	$3.5 \div 17$
25600	$2.8 \div 6.5$	$3.6 \div 16$

a)		
$m_{H^+}, GeV$	$-a_{CP}(b \rightarrow s\gamma)/10^{-3}$	$a_{CP}(b \to d\gamma)/10^{-2}$
50	$2.9 \div 6.7$	$3.6 \div 17$
100	$2.9 \div 6.7$	$3.6 \div 17$
200	$2.9 \div 6.7$	$3.6 \div 17$
800	$2.9 \div 6.7$	$3.6 \div 17$
3200	$2.9 \div 6.5$	$3.6 \div 16$
1)		

Table 4:	Value	s of CF	<b>'</b> -asym	metry	for	the	Model	Ι
	a) for	$\Delta < 10$	0%, b)	for $\Delta$	< 5	0%		

1	1
h	1
$\mathbf{D}$	1

$m_{H^+}, GeV$	$-a_{CP}(b \rightarrow s\gamma)/10^{-3}$	$a_{CP}(b \to d\gamma)/10^{-2}$
50	$2.9 \div 8.6$	$3.6 \div 21$
100	$2.9 \div 8.9$	$3.6 \div 22$
200	$2.9 \div 8.7$	$3.6 \div 22$
800	$2.9 \div 7.6$	$3.6 \div 19$
3200	$2.9 \div 6.5$	$3.6 \div 17$

Table 5: Values of CP-asymmetry for the Model 2 a) for  $\Delta < 10\%,$  b) for  $\Delta < 50\%$ 

	<u>۱</u>
9	1
a	
	/

$m_{H^+}, GeV$	$-a_{CP}(b \to s\gamma)/10^{-3}$	$a_{CP}(b \to d\gamma)/10^{-2}$
400	-	-
650	-	-
1300	$2.8 \div 6.1$	$3.4 \div 15$
2600	$2.8 \div 6.2$	$3.5 \div 16$
5200	$2.9 \div 6.3$	$3.6 \div 16$

b)

$m_{H^+}, GeV$	$-a_{CP}(b \rightarrow s\gamma)/10^{-3}$	$a_{CP}(b \rightarrow d\gamma)/10^{-2}$
400	$2.4 \div 5.3$	$2.9 \div 13$
650	$2.5 \div 5.7$	$3.1 \div 14$
1300	$2.7 \div 6.1$	$3.4 \div 15$
2600	$2.8 \div 6.3$	$3.5 \div 16$
5200	$2.9 \div 6.3$	$3.6 \div 16$









