2/3 SPLITTING IN SUSY GUT — HIGGS AS GOLDSTONE BOSON

Z. BEREZHIANI

INFN Sezione di Ferrara, I-44100 Ferrara, Italy, and Institute of Physics, Georgian Academy of Sciences, 38077 Tbilisi, Georgia E-mail: berezhiani@axpfe1.fe.infn.it

ABSTRACT

We briefly review the GIFT (Goltstones instead of Fine Tuning) mechanism which provides a promising solution to the doublet-triplet splitting problem and some other puzzles of the supersymmetric grand unification as are the μ -problem, the fermion mass problem, etc. It can be naturally implemented by extending the minimal SUSY SU(5) model to the gauge SU(6) theory.

1. Introduction

The concepts of supersymmetry (SUSY) and grand unification theories (GUT) constitute the most promising ideas beyond the standard model. The present data on $\alpha_3(M_Z)$ and $\sin^2 \theta_W(M_Z)$ are in a remarkable agreement with the prediction of the SUSY SU(5) theory while exclude the non-supersymmetric SU(5).¹ On the other hand, the minimal supersymmetric standard model (MSSM) without GUT is also not in best shape: unification at the string scale gives too small $\sin^2 \theta_W(M_Z)$. In the MSSM the running gauge constants g_3 , g_2 and g_1 given at $\mu = M_Z$ within the present experimental error bars, in their evolution to higher energies join at the scale $M_5 \simeq 10^{16}$ GeV.¹ Hence, at this scale the $SU(3) \times SU(2) \times U(1)$ symmetry can be consistently embedded into SU(5). An additional hint is provided by the fact of the $b - \tau$ Yukawa unification at GUT energies². All these suggest a following paradigm:

At the scale $M_{\rm str} = g_{\rm str} M_P$ a basic 'string' theory reduces to a SUSY GUT given by a gauge group G (G = SU(5), SO(10), SU(6) etc.) with a gauge constant defined by condition $k_G g_G^2 = g_{\rm str}^2$, where k_G is a corresponding Kac-Moody level and $M_P \simeq 2 \cdot 10^{18}$ GeV is a reduced Planck mass. At some scale $M_G \ge M_5$ the gauge group breaks down to SU(5) subgroup, which then at $M_5 \simeq 10^{16}$ GeV breaks down to the $SU(3) \times SU(2) \times U(1)$. The value of the gauge constant at this scale is $g_5 \simeq 0.7$. In order to keep the gauge coupling in the perturbative regime up to the string scale, there should not be too many extra states with masses in the range from M_5 to M_P ,

Below the scale M_5 the theory is just MSSM containing the $SU(3) \times SU(2) \times U(1)$ gauge superfields, the chiral superfields of quarks and leptons q_i , l_i , u_i^c , d_i^c , e_i^c (i = 1, 2, 3), and two Higgs doublets $\phi_{1,2}$.^{*a*}Barring the gauge sector, its effective Lagrangian

^a In other words, no GUT debris is allowed below the scale M_G except the singlets. The extra complete degenerate SU(5) supermultiplets populating the intermediate scales would not affect the

is given by the superpotential terms:

$$W_{\text{Yuk}} = \lambda_{ij}^{u} q_i u_j^c \phi_2 + \lambda_{ij}^{d} q_i d_j^c \phi_1 + \lambda_{ij}^{e} l_i e_j^c \phi_1,$$

$$W_{\text{Higgs}} = \mu \phi_1 \phi_2$$
(1)

and the soft supersymmetry breaking (SSB) terms (tilde labels sfermions):

$$L_{A} = A_{ij}^{u} \tilde{q}_{i} \tilde{u}_{j}^{c} \phi_{2} + A_{ij}^{d} \tilde{q}_{i} \tilde{d}_{j}^{c} \phi_{1} + A_{ij}^{e} \tilde{l}_{i} \tilde{e}_{j}^{c} \phi_{1} + h.c.,$$

$$L_{\rm sf} = \sum m_{ij}^{2} \tilde{f}_{i}^{\dagger} \tilde{f}_{j}, \qquad f = q, u^{c}, d^{c}, l, e^{c}$$

$$L_{\rm Higgs} = m_{1}^{2} |\phi_{1}|^{2} + m_{2}^{2} |\phi_{2}|^{2} + (B_{\mu} \phi_{1} \phi_{2} + h.c.) \qquad (2)$$

with masses and dimensional constants of the order of supersymmetry breaking scale $m_S \sim m_Z$ (in the supergarvity folklore — the gravitino mass $m_{3/2}$). The VEVs $\langle \phi_1^0 \rangle = v_1 = v \cos \beta$ and $\langle \phi_2^0 \rangle = v_2 = v \sin \beta$, v = 174 GeV, break the electroweak symmetry and induce the fermion masses.

The softly broken SUSY is the only plausible idea that can support the GUT against the gauge hierarchy problem³. At the level of the standard model this is essentially a problem of the Higgs mass stability against radiative corrections (quadratic divergences). It is removed as soon as one appeals to SUSY, which links the scalar masses to those of their fermion superpartners while the latter are protected by the chiral symmetry. In the context of grand unification the gauge hierarchy problem concerns rather the origin of scales: why the weak scale $v \sim M_W$ is so small as compared to the GUT scale M_5 , which in itself is not far from the Planck scale M_P . This question is inevitably connected with the doublet-triplet (2/3) splitting puzzle³: the Higgs doublets ϕ_1, ϕ_2 embedded in the GUT multiplets are accompanied by the colour triplet partners \overline{T}, T . The latter would mediate unacceptably fast proton decay (via d = 5 operators⁴) unless they are superheavy, with mass $\sim M_5$.

Yet another problem is a so-called μ -problem⁵. The supersymmetric term $\mu \phi_1 \phi_2$ should be small, and at first sight $\mu = 0$ could be a most natural possibility: then the Higgs masses would emerge entirely from the soft SUSY breaking terms with $m_S \sim v$. However, $\mu = 0$ is excluded experimentally and it is required to be $\sim v$ as well. Therefore, one has to explain in a natural way, why the value of the supersymmetric mass μ is of the same order as that of the soft SUSY breaking mass m_S .

In the context of the realistic SUSY GUTs the 2/3 splitting and μ problems are intimately related to each other and to the other key problems as are the origin of the supersymmetry breaking, the fermion mass problem, proton stability, etc.

2. 2/3 splitting in SUSY SU(5)

In the minimal SUSY SU(5) model the Higgs sector consists of the chiral superfields in adjoint representation $\Sigma \sim 24$ and fundamental representations $H \sim 5$,

unification of the gauge constants. However, they would contribute the renormalization group (RG) running of the Yukawa constants from M_5 down to M_Z and thus affect the $b - \tau$ Yukawa unification.

 $\bar{H} \sim \bar{5}$: $H = (T + \phi_2)$ and $\bar{H} = (\bar{T} + \phi_1)$, where $\phi_{1,2}$ are the MSSM Higgs doublets and \bar{T}, T colour triplet partners. The most general superpotential terms involving these fields are the following:

$$\mathcal{W}_{\text{Higgs}} = M\Sigma^2 + h\Sigma^3 + M_H \bar{H} H + f \bar{H} \Sigma H \tag{3}$$

The SU(5) symmetry breaking down to $SU(3) \times SU(2) \times U(1)$ is provided by supersymmetric ground state $\langle \Sigma \rangle = (2M/3h) \text{diag}(2, 2, 2, -3, -3), \langle \bar{H}, H \rangle = 0$. In this vacuum the masses of the T and ϕ fragments are different: $M_3T\bar{T}$ and $\mu'\phi_1\phi_2$, where $M_3 = M_H - (3f/h)M$ and $\mu' = M_H + (2f/h)M$. So, the light doublet $(\mu \sim M_W)$ versus heavy triplet $(M_3 \sim M_X)$ requires that $hM_H \approx -2fM$, with the accuracy of about 10^{-14} . Supersymmetry renders this constraint stable against radiative corrections. However, such a *technical solution*³ is nothing but the *Fine Tuning* (FT) of the parameters in the superpotential.

The actual puzzle is to achieve the 2/3 splitting in a natural way, without FT. Several attempts have been done, which we briefly describe below:

• Missing doublet mechanism (MDM).⁶ In this scenario the 24-plet Σ is replaced by the 75-plet Φ of SU(5), and additional heavy superfields $\Psi \sim 50$ and $\bar{\Psi} \sim \overline{50}$ are introduced. The latter have a bare mass term $M_{\Psi}\bar{\Psi}\Psi$, $M_{\Psi} \sim M_X$, while $M_H H\bar{H}$ is absend in the superpotential. Instead the superpotential contains the terms $\bar{H}\Phi\Psi$ and $H\Phi\bar{\Psi}$. Then the triplet components T, \bar{T} in H, \bar{H} can get $\sim M_X$ mass via their mixing to triplet states in $\Psi, \bar{\Psi}$, while the doublets $\phi_{1,2}$, will remain massless since 50-plet does not contain doublet fragment. The MDM can be motivated by some additional symmetry reasons, among which the anomalous U(1) gauge symmetry seems to be a most compelling variant⁷.

New possibilities can emerge with introducing the extra singlet superfield I. Then the most general renormalizable Higgs superpotential reads:

$$\mathcal{W}_{\text{Higgs}} = M\Sigma^2 + h\Sigma^3 + \Lambda^2 I + M'I^2 + h'I^3 + h''I\Sigma^2 + M_H\bar{H}H + f\bar{H}\Sigma H + f'I\bar{H}H \quad (4)$$

The 2/3 splitting can be achieved under certain constraints on the coupling constants:

• Sliding singlet mechanism⁸ implies that I has no couplings in the superpotential except the last term in (4). Then at tree level, with taking into account also the SSB terms, the VEV of I can be adjusted to cancell the large contributions to the doublet mass from other terms. This mechanism, however, was shown to be unstable against radiative corrections⁹ – the tadpole diagrams induce the doublet mass $\sim \sqrt{m_S M_G}$.

• Missing VEV mechanism (MVM). One can assume that the Σ and I together form a reducible representation $\Sigma' \sim 25 = 24 + 1$ and the superpotential has a form:

$$\mathcal{W}_{\text{Higgs}} = M\Sigma'^2 + h\Sigma'^3 + f\bar{H}\Sigma'H, \quad (\Sigma')^{\alpha}_{\beta} = \Sigma^{\alpha}_{\beta} + \frac{1}{\sqrt{5}}I\delta^{\alpha}_{\beta} \tag{5}$$

Since Σ' is not traceless, its VEV can be chosen as $\langle \Sigma' \rangle = (2M/3h) \text{diag}(1, 1, 1, 0, 0)$. In this case the triplets get mass $M_3 = (2f/3h)M$ while the doublets remain massless. In the SU(5) framework it is difficult to justify such a situation, which in terms of the general expression (4) is equivalent to the following 5 conditions of the FT:

$$M' = M, \quad h' = \frac{1}{3}h'' = \frac{1}{\sqrt{5}}h, \quad \Lambda = M_H = 0$$
 (6)

Nevertheless, it can be realized in a natural way in models with a larger gauge symmetry where both 24 and 1 can be embbedded in one irreducible representation which is not constrained to be traceless. In particular, in SO(10) model the 45-plet Higgs of SO(10) containing both 24-plet and singlet fragments: 45 = 1 + 24 + 10 + 10, can have a VEV of the form¹² $\langle A \rangle \sim \text{diag}(1,1,1,0,0) \otimes i\sigma$. Since 45-plet is antisymmetric, the coupling HAH is not allowed. However, one can introduce the additional 10-plet H' with large mass term M'H'H' and mix with H through A.

• Goldstones instead of Fine Tuning (GIFT) mechanism.^{15,16} The superfields Σ , I, H and \overline{H} can be combined in the *irreducible* representation $\Sigma' \sim 35$ of the SU(6)group: $35 = 24 + 1 + 5 + \overline{5}$. Therefore, if the Higgs superpotential is a function of the combination Σ' :

$$\mathcal{W}_{\text{Higgs}} = M\Sigma'^2 + h\Sigma'^3, \qquad \Sigma' = \begin{pmatrix} -\frac{5}{\sqrt{30}}I & H\\ H & \Sigma + \frac{1}{\sqrt{30}}I \end{pmatrix}$$
(7)

then it possess a global symmetry SU(6), larger than the gauge SU(5) symmetry. The vacuum state can be chosen as $\langle \Sigma' \rangle = (M/h) \operatorname{diag}(1, 1, 1, 1, -2, -2)$, which breaks the global SU(6) symmetry down to $SU(4) \times SU(2) \times U(1)$ subgroup, while its gauged part SU(5) breaks down to standard $SU(3) \times SU(2) \times U(1)$. The corresponding Goldstone modes are in representations $(4, \bar{2}) + (\bar{4}, 2)$. Among those, the SU(3) triplet fragments $(3, \bar{2}) + (\bar{3}, 2)$ are eaten up by the SU(5) gauge supefields (X, Y bosons) due to Higgs mecghanism. As for the fragments $(1, \bar{2}) + (1, 2)$ they remain massless as Goldstone modes, and can be identified as the MSSM Higgs doublets $\phi_{1,2}$

In view of the general expression (4) this situation requires 9 FT constraints:

$$\Lambda = 0, \quad M = M' = \frac{1}{2}M_H, \quad h = -\sqrt{\frac{15}{8}}h' = \sqrt{\frac{10}{3}}h'' = \frac{1}{3}f = -\sqrt{\frac{5}{6}}f' \tag{8}$$

However, it can naturally emerge if one extends the gauge symmetry to SU(6).¹⁷ One can assume that at the scale $M_6 \gg M_5$ the local SU(6) symmetry is broken down to SU(5) by some Higgses which do not couple $\Sigma \sim 35$ Higgses in the Higgs superpotential. Then the superpotential of Σ would not 'feel' that the gauge symmetry is already reduced to SU(5) and maintain the global SU(6) invariance.

3. SUSY SU(6) Model

The $SU(6) \mod^{17}$ is a minimal extension of SU(5): the Higgs sector contains supermultiplets $\Sigma \sim 35$ and $H + \bar{H} \sim 6 + \bar{6}$ respectively in adjoint and fundamental representations, in analogy to 24 and $5 + \bar{5}$ of SU(5). However, this model drastically differs from the other GUTs where the Higgs sector usually consists of two different sets: one is for the GUT symmetry breaking (e.g. 24-plet in SU(5)), while another containing the Higgs doublets (like $5+\bar{5}$ in SU(5)) is just for the electroweak symmetry breaking. The SU(6) theory has no special superfields for the second purpose: 35 and $6 + \bar{6}$ constitute a minimal Higgs content needed for the local SU(6) symmetry breaking down to $SU(3) \times SU(2) \times U(1)$. As for the MSSM Higgs doublets $\phi_{1,2}$, they are contained in Σ and H, \bar{H} themselves, as the Goldstone modes of the accidental global symmetry $SU(6)_{\Sigma} \times U(6)_{H}$ related to independent transformations of Σ and H. This global symmetry arises if mixing terms of the form $\bar{H}\Sigma H$ are suppressed in the Higgs superpotential¹⁷. In particular, if the Higgs superpotential has a 'factorized' form

$$\mathcal{W}_{\text{Higgs}} = \mathcal{W}(\Sigma) + \mathcal{W}(H\bar{H}) \tag{9}$$

e.g. with (Y is some auxiliary singlet)

$$\mathcal{W}(\Sigma) = M\Sigma^2 + h\Sigma^3, \qquad \mathcal{W}(H\bar{H}) = \rho Y(\bar{H}H - \Lambda^2)$$
 (10)

Then it has an accidental global symmetry $SU(6)_{\Sigma} \times U(6)_{H}$.^b In the limit of exact SUSY (vanishing F and D terms), among the other degenerated vacua there is a following one:

$$\langle \Sigma \rangle = V_{\Sigma} \cdot \text{diag}(1, 1, 1, 1, -2, -2), \quad \langle H \rangle = \langle \bar{H} \rangle = V_{H} \cdot (1, 0, 0, 0, 0, 0)$$
(11)

where $V_{\Sigma} = M/h$ and $V_H = \Lambda$. After SUSY breaking, this configuration can indeed be a true vacuum state for a proper range of the soft parameters. Then H, \bar{H} break SU(6) down to SU(5) while Σ breaks SU(6) down to $SU(4) \times SU(2) \times U(1)$, and both channels together lead to the local symmetry breaking down to $SU(3) \times SU(2) \times$ U(1). At the same time, the global symmetry $SU(6)_{\Sigma} \times U(6)_{H}$ is broken down to $[SU(4) \times SU(2) \times U(1)]_{\Sigma} \times U(5)_{H}$. Most of the Goldstone modes are eaten up by the SU(6) gauge superfields through the Higgs mechanism. However, a couple of fragments survive and present in particle spectrum at lower energies as the Goldstone superfields. These constitute the MSSM Higgs doublets $\phi_{1,2}$ which in terms of the doublet (anti-doublet) fragments in Σ and H, \bar{H} are given as

$$\phi_2 = \cos \eta \phi_{\Sigma} - \sin \eta \phi_H , \qquad \phi_1 = \cos \eta \bar{\phi}_{\Sigma} - \sin \eta \bar{\phi}_{\bar{H}}$$
(12)

where $\tan \eta = 3V_{\Sigma}/V_H$. In the following we assume that $V_H \sim M_6$ is larger than $V_{\Sigma} \simeq M_5 \simeq 10^{16}$ GeV, as it is motivated by the SU(5) unification of the gauge

^b The global $SU(6)_{\Sigma} \times U(6)_{H}$ is an accidental symmetry of the Higgs superpotential and not a symmetry of the whole Lagrangian: the Yukawa terms and the gauge couplings (*D*-terms) do not respect it. However, in the exact supersymmetry limit it is effective for the field configurations on the vacuum valley where D = 0. Owing to non-renormalization theorem, it cannot be spoiled by the radiative corrections.

couplings. In this case the doublets $\phi_{1,2}$ dominantly come from Σ while in H, \bar{H} they are contained with small weight $\sim 3V_{\Sigma}/V_{H}$.

The scalar fields in $\phi_{1,2}$ then get mass from the SSB terms:

$$V_{SB} = (m_S [\theta^2 \mathcal{W}']_F + h.c.) + m_k^2 \sum_k |\varphi_k|^2$$
(13)

where φ_k imply all scalar fields involved, $m_k \sim m_S$ are their soft masses, and \mathcal{W}' repeates all structures present in the superpotential \mathcal{W} , but is not necessarily proportional to \mathcal{W} . Incidentally, the GIFT scenario naturally solves also the μ -problem. Taking into account the SSB terms (13) in minimization of the Higgs potential of Σ and H, \bar{H} , one observes that the VEV V_{Σ} is shifted by an amount of $\sim m_S$ as compared to the one of eq. (11) calculated in the exact SUSY limit. Then substituting these VEVs back in superpotential, this shift gives rise to the $\mu\phi_1\phi_2$ term, with $\mu \sim m_S$. Thus, in GIFT scenario the (supersymmetric) μ -term emerges as a consequence of the SUSY breaking. On the other hand, at the GUT scale one obtains for the soft parameters in L_{Higgs} :

$$m_1 = m_2 = m_{\Sigma}, \quad B_{\mu} = \mu^2 + m_{\Sigma}^2$$
 (14)

where m_{Σ} is a soft mass term of Σ in (13). In fact, this relation is a reflection of the fact that since \mathcal{W}' has the same form as \mathcal{W} , then it also has the global global symmetry $SU(6)_{\Sigma} \times U(6)_{H}$ and thus one combination $(\phi_1 + \phi_2^*)/\sqrt{2}$ of the scalar doublets remains massless as a true Goldstone boson.

However, in reality the SUSY breaking relaxes radiative corrections which lift the vacuum degeneracy (mainly due to the large top Yukawa coupling, origin of which we will clarify below) and fix the VEVs v_1 and v_2 . The effects of radiative corrections leading to the electroweak symmetry breaking were studied recently in refs.²⁴.

Thus, the SU(6) model naturally solves both the DT splitting and the μ problems. The Higgs doublets $\phi_{1,2}$ remain light, with $\mu \sim m_S$, and their triplet partners are superheavy: the triplets from Σ have masses $\sim M_5$, while the 'Goldstone' triplets from H, \bar{H} are eaten up by the SU(6) gauge superfields via the Higgs mechanism and thus acquire masses $\sim M_6$.

In order to built a consistent GIFT model, one has to find some valid symmetry reasons to forbid the mixed term $\bar{H}\Sigma H$: otherwise the global symmetry $SU(6)_{\Sigma} \times U(6)_{H}$ is not accidental and the mixed term should be put to zero by hands, which essentially would be a FT. It is natural to use for this purpose the discrete symmetries.^{18,20,21}

For example, one can introduce two 35-plets $\Sigma_{1,2}$ and impose the following discrete Z_3 symmetry²⁰: $\Sigma_1 \to e^{i\frac{\pi}{3}}\Sigma_1$ and $\Sigma_2 \to e^{-i\frac{\pi}{3}}\Sigma_2$, while H, \bar{H} and Y are invariant. Then the most general renormalizable superpotential of $\Sigma_{1,2}$ has a form:

$$W(\Sigma) = M\Sigma_1\Sigma_2 + h_1\Sigma_1^3 + h_2\Sigma_2^3$$
(15)

whereas the mixed terms $H\Sigma_{1,2}\overline{H}$ are forbidden by Z_3 symmetry, and thus the Higgs superpotential acquires an accidental global symmetry $SU(6)_{\Sigma} \times U(6)_{H}$.

However, problem arises if one includes also the higher order operators cutoff by M_P . Then the terms like $\frac{1}{M}H\Sigma_1\Sigma_2\bar{H}$ etc., allowed by the Z_3 symmetry would spoil the accidental global symmetry and Higgs bosons will get too large masses. Some possibilities avoiding the impact of the Planck scale terms were suggested in ref.²¹. A very intersting way to guarantee the accidental global symmetry at all orders in M_P^{-1} was suggested in ref.²² which makes use of the anomalous U(1) symmetry.

4. Fermion masses and mixing

In the minimal SUSY SU(5) model the quarks and leptons of each family fit into the multiplets $\overline{5}_i = (d_i^c + l_i)$ and $10_i = (u_i^c + q_i + e_i^c)$, i=1,2,3. The fermion masses are induced by the Yukawa terms:

$$\mathcal{W}_{\text{Yuk}} = \lambda_{ij}^u 10_i H 10_j + \lambda_{ij}^d 10_i \bar{H} \bar{5}_j \tag{16}$$

At the GUT scale M_5 these terms reduce to the MSSM couplings (1) with $\hat{\lambda}_{ij}^e = \hat{\lambda}_{ji}^d$, and hence $\lambda_{d,s,b} = \lambda_{e,\mu,\tau}$. The $\lambda_b = \lambda_{\tau}$ unification² is a definite success. After accounting for the RG running it translates into relation for physical masses $m_b/m_{\tau} \sim$ 3, while the more precise comparison of m_b and m_{τ} within the present experimental uncertainties implies that λ_t should be rather large (≥ 1), in which case the top mass is fixed by its infrared limit $M_t = (190 - 210) \sin \beta$ GeV = 140 - 210 GeV. Thus, the minimal SUSY SU(5) model explains the principal origin of the bottom quark mass and nicely links it to the large value of the top mass.

Unfortunately, the other predictions $\lambda_s = \lambda_{\mu}$ and $\lambda_d = \lambda_e$ are wrong: experimentally $m_s/m_d \simeq 20$ while $m_{\mu}/m_e = 200$. In addition, there is no explanation neither for the fermion mass hierarchy nor for the CKM mixing pattern: the Yukawa matrices λ^u and λ^d remain arbitrary and there is no reason neither for hierarchy of their eigenvalues nor for their allignment. Therefore, one is forced to go beyond the minimal SU(5) model and implement new ideas that could shed some more light on the origin of fermion masses and mixing¹¹.

In the GIFT SU(6) model the situation is drastically different. The fermion sector of the SU(6) theory consists of three families $(\bar{6}+\bar{6}'+15)_i$, i = 1, 2, 3, and one 20-plet (since 20 is a pseudo-real representation, its mass term is vanishing)^{19,20}. In terms of the SU(5) subgroup the fermions under consideration read

$$20 = 10 + \overline{10} = (q + u^c + e^c)_{10} + (Q^c + U + E)_{\overline{10}}$$

$$15_i = (10 + 5)_i = (q_i + u^c_i + e^c_i)_{10} + (D_i + L^c_i)_5$$

$$\bar{6}_i = (\bar{5} + 1)_i = (d^c_i + l_i)_{\bar{5}} + n_i, \quad \bar{6}'_i = (\bar{5} + 1)'_i = (D^c_i + L_i)_{\bar{5}'} + n'_i, \quad (17)$$

Only the following renormalizable Yukawa terms allowed by the SU(6) symmetry:

$$\mathcal{W}_{Yuk} = G \, 20\Sigma 20 \,+\, \Gamma \, 20H 15_3 \,+\, \Gamma_{ij} 15_i \bar{H} \bar{6}'_j \,, \qquad i, j = 1, 2, 3 \tag{18}$$

(all coupling constants are assumed to be ~ 1). Without loss of generality, the basis of 15-plets is chosen so that only the 15₃ state couples 20-plet in (18). Also, among six $\bar{6}$ -plets one can always choose three of them (denoted in eq. (18) as $\bar{6}'_{1,2,3}$) which couple 15_{1,2,3} while the other three states $\bar{6}_{1,2,3}$ have no Yukawa couplings.

Already at the scale V_H of the gauge symmetry breaking $SU(6) \rightarrow SU(5)$ the fermion content reduces to the one of the minimal SU(5). The extra vector-like fermions $\overline{10} + 10_3$ and $(5 + \overline{5}')_{1,2,3}$, get $\sim V_H$ masses from couplings (18) and thereby the light states remain as $\overline{5}_{1,2,3}$, $10_{1,2}$, 10 and singlets n_i, n'_i .

In the SU(6) model the Yukawa couplings of the Higgs doublets to fermions are very peculiar: should the Yukawa terms also respect the global $SU(6)_{\Sigma} \times U(6)_H$ symmetry, then $\phi_{1,2}$ being the Goldstone modes would have the vanishing Yukawa couplings to all fermions which remain massless after the GUT symmetry breaking down to the MSSM, that are ordinary quarks and leptons. Thus, the couplings relevant for fermion masses have to explicitly violate $SU(6)_{\Sigma} \times U(6)_H$. This leads to striking possibilities to understand the key features of the fermion mass and mixing without invoking the additional symmetry arguments. In particular, the only fermion which can get ~ 100 GeV mass through the renormalizable Yukawa coupling is the top quark, while other fermion masses can emerge only through the higher order operators and thus are suppressed by powers of the Planck scale M_P . Therefore, the observed hierarchy of fermion masses and mixings can be naturally explained in terms of small ratios $\varepsilon_{\Sigma} = V_{\Sigma}/V_H$ and $\varepsilon_H = V_H/M_P$.^{19,20,21}

Indeed, the couplings of 20-plet in (18) explicitly violate the global $SU(6)_{\Sigma} \times U(6)_H$ symmetry. Hence, the up-type quark from 20 (to be identified as top) has nonvanishing coupling with the Higgs doublet ϕ_2 . As far as $V_H \gg V_{\Sigma}$, it essentially emerges from $20\Sigma 20 \rightarrow qu^c \phi_2$. Thus, only the top quark can have ~ 100 GeV mass due to the large Yukawa constant $G: \lambda_t \sim 1$.

All other fermion masses can be induced only from the higher order operators scaled by the inverse powers of M_P . These operators can emerge effectively by integrating out some heavy fermion states.¹⁰ For example, the following d = 6 operator is responsible for the b and τ masses:

$$\mathcal{B} = \frac{B}{M_P^2} 20\bar{H}(\Sigma\bar{H})\bar{6}_i \tag{19}$$

At the MSSM level it reduces to the Yukawa couplings $\varepsilon_H^2 B(qd_3^c + e^c l_3)\phi_1$. Hence, though b and τ belong to the 20-plet as well as t, their Yukawa constants are by factor $\sim \varepsilon_H^2$ smaller than λ_t . In addition, the $b - \tau$ Yukawa constants are automatically unified at the GUT scale — up to $\sim \varepsilon_{\Sigma}^2$ corrections, $\lambda_b = \lambda_{\tau}$.

The second family fermions can get masses from the operators:

$$\mathcal{C} = \frac{C_{ij}}{M_P^2} 15_i H(\Sigma H) 15_j, \quad \mathcal{C}' = \frac{C_i}{M_P} 20\Sigma H 15_i$$
$$\mathcal{S} = \frac{S_{ik}}{M_P^2} 15_i (\Sigma^2 \bar{H}) \bar{6}_k, \quad \mathcal{S}' = \frac{S'_{ik}}{M_P^2} 15_i (\Sigma \bar{H}) (\Sigma \bar{6}_k) \tag{20}$$

 $(SU(6) \text{ indices are contracted so that combinations in the parentheses transform as effective <math>\overline{6} \text{ or } 6$). Since the $\overline{6}'$ and 15_3 states already have $\sim V_H$ masses, these operators are relevant only for the light states in 20, $15_{1,2}$ and $\overline{6}_{1,2,3}$. One can always redefine the basis of $\overline{6}$ -plets so that only the $\overline{6}_3$ couples 20 in eq. (20). In addition, we assume that constants B, C_{ij} , etc. all are order 1 as well as the constants in (18).

This allows to explain the observed hierarchy of fermion masses and mixing in terms of two small parameters, $\varepsilon_{\Sigma}, \varepsilon_{H} \sim 0.1$.^c More details on the fermion mass structures can be found in refs.^{19,20,21}. In the context of these one can obtain the following hierarchy for the Yukawa coupling eigenvalues at the GUT scale

$$\lambda_t \sim 1, \quad \lambda_b = \lambda_\tau \sim \lambda_c \sim \varepsilon^2, \quad \lambda_c \sim \varepsilon^2, \quad \lambda_s, \lambda_\mu \sim \varepsilon^3, \quad \lambda_u, \lambda_d, \lambda_e \sim \varepsilon^5$$
 (21)

and for the CKM angles

$$s_{12} \sim 1, \quad s_{23} \sim \varepsilon, \quad s_{13} \sim \varepsilon^2$$

$$\tag{22}$$

The lowest dimension operator relevant for the neutrino masses is $\mathcal{N} = \frac{N_{kl}}{M_P^2} \bar{6}_k H \Sigma H \bar{6}_l$. For $\varepsilon_{\Sigma}, \varepsilon_H \sim 0.1$ it induces the small Majorana masses $m_{\nu} \sim v^2/M_P \sim 10^{-5} \,\text{eV}$. This mass range (which as a matter of fact is also favoured by the minimal $SU(5) \,\text{model}^{25}$) is just what is needed for the long wavelength "just-so" oscillation solution to the solar neutrino problem.

5. discussion

Let us conclude with a brief summary of the simplest SUSY GUTs and appropriate mechanisms for the 2/3 splitting. Their main features are given in Table 1.

The only known solution of the 2/3 splitting problem in SUSY SU(5) is based on MDM⁶ or its variations (see ref.⁷ and references therein). It requires rather complicated Higgs sector involving large representations which in the string theory context can be allowed only at the Kac-Moody level $k_5 \ge 4$. The main problem of the MDM is that the extra states 50 and $\overline{50}$ with masses $\sim M_5$ contribute the RG running up of the SU(5) gauge constant and drive it out the perturbative regime well below the Planck scale M_P . Therefore, it is hard to imagine how such a model can be embedded in a string theory. Situation could be saved if the mass of 50-plets is $\sim M_P$. However, in this case the mass of the triplets T, \bar{T} from H, \bar{H} is $M_3 \sim M_5^2/M_P \sim 10^{14}$ GeV which would cause catastrophically fast proton decay via d = 5 operators.

The MVM solution¹² operative in SUSY SO(10) is also rather cumbersome and requires complicated Higgs sector. It also can be justified by symmetry reasons¹⁴.

^{*c*} As far as the scale $M_5 \simeq 10^{16} \text{ GeV}$ is fixed by the SU(5) unification of the gauge couplings, this corresponds to $M_6 \sim 10^{17} \text{ GeV}$ and $M_P \sim 10^{18} \text{ GeV}$, so that M_P is indeed close to the string or Planck scale. The relation $M_6 \sim \sqrt{M_5 M_P}$ could naturally emerge in the context of the models discussed in refs.^{20,21,22}.

Table 1. Summary of the known solutions to the 2/3 splitting problem in various SUSY GUTs: MDM in SU(5), MVM in SO(10) and GIFT in SU(6), with the minimal multiplet contents required for realization of these models. The lowest Kac-Moody levels at which these models can be in principle embeddable in the string theory are also shown.

GUT	Higgses	Fermions	μ and B_{μ}	Yukawas
SU(5) $(k=4)$	$5 + \overline{5} + 75 + 50 + \overline{50}$	$(\bar{5}+10)_i$?	$\lambda_b = \lambda_\tau (\tan\beta = ?)$
SO(10) $(k=2)$	$ \begin{array}{r} 10 + 10' + 45_{1,2,3} \\ +54 + 16 + \overline{16} \end{array} $	16_i	?	$\lambda_b = \lambda_\tau = \lambda_t \sim 1$ $(\tan\beta \sim 100)$
SU(6) $(k=2)$	$6 + \bar{6} + 35$	$27_i + 20$	$\begin{aligned} \mu &\sim m_S \\ B_\mu &= \mu^2 + m_{1,2}^2 \end{aligned}$	$\begin{array}{l} \lambda_t \sim 1, \lambda_b = \lambda_\tau \sim \varepsilon^2 \\ \lambda_c \sim \varepsilon^2, \lambda_{s,\mu} \sim \varepsilon^3 \\ (\tan \beta \sim 1) \end{array}$

Neither the MDM nor the MVM solutions for 2/3 splitting do not touch the μ -problem, and for its solution some additional ideas should be incorporated. In addition, the SU(5) or SO(10) models themselves give no understanding to the fermion mass hierarchy, though by involving the horizontal symmetry concept one can obtain the predictive mass textures.²⁶

The SU(6) model¹⁷ seems to be most favourable, and The GIFT mechanism can be justified by symmetry reasons^{21,22}. The Higgs sector of this theory is simple: it contains only the adjoint and fundamental representations. The fermion sector is also simple – three families of supermultiplets $(\bar{6} + \bar{6}' + 15)_i = 27_i$, i = 1, 2, 3, which in fact constitute the 27-plets of E_6 group, and one 20-plet. Therefore, there is no difficulty in maintaining the perturbative regime for the gauge coupling up to the scale M_P .

The SU(6) model provides a simultaneous solution to both 2/3 splitting and μ problems, leads to prediction for the soft parameters $m_{1,2}$ and B_{μ} pattern at the GUT scale, and, in addition, explains many features of the fermion mass spectrum and mixing by internal reasons, without involving additional ideas.

- U. Amaldi, W. de Boer and H. Fürstenau, Phys. Lett. B260 447;
 J. Ellis, S. Kelley and D.V. Nanopoulos, *ibid.* (1991) 131;
 - P. Langacker and M. Luo, Phys. Rev. D44 (1991) 817.
- 2. A. Buras et al., Nucl. Phys. B135 (1978) 66.
- 3. L. Maiani, in Comptes Rendus de l'Ecole d'Etè de Physique des Particules, Gif-sur-Yvette, 1979, IN2P3, Paris, 1980, p. 3;
 S. Dimopoulos and H. Georgi, Nucl. Phys. B150 (1981) 193;

M. Sakai, Z. Phys. C11 (1981) 153; E. Witten, Nucl. Phys. B188 (1981) 573

- 4. S. Weinberg, Phys. Rev. D26 (1982) 287;
 N. Sakai and T. Yanagida, Nucl. Phys. B197 (1982) 533.
- 5. J.E. Kim and H.P. Nilles, Phys. Lett. B138 (1984) 150.
- 6. H. Georgi, Phys. Lett. B108 (1982) 283;
 B. Grinstein, Nucl. Phys. B206 (1982) 387;
 A. Masiero *et al.*, Phys. Lett. B115 (1982) 380.
- 7. Z. Berezhiani and Z. Tavartkiladze, hep-ph/9611277.
- 8. E. Witten, Phys. Lett. **B105** (1981) 267;

D.V. Nanopoulos and K. Tamvakis, *ibid.* B113 (1982) 151.

- H.P. Nilles, M. Srednicki and D. Wyler, Phys. Lett. B124 (1982) 337;
 A. Lahanas, *ibid.* 341.
- C.D. Frogatt and H.B. Nielsen, Nucl. Phys. B147 (1979) 277;
 Z.G. Berezhiani, Phys. Lett. B129 (1983) 99; *ibid.* B150 (1985) 177;
 S. Dimopoulos, *ibid.* B129 (1983) 417.
- For a review, see Z. Berezhiani, hep-ph/9602325, to appear in Proc. ICTP Summer School (Trieste, July 1995), eds. E. Gava et al., World Scientific, Singapore, 1996 and references therein.
- 12. S. Dimopoulos and F. Wilczek, Preprint NSF-ITP-82-07 (1982).
- 13. K.S. Babu and S.M. Barr, Phys. Rev. **D48** (1993) 5354.
- 14. Z. Berezhiani and Z. Tavartkiladze, hep-ph/9612232.
- 15. K. Inoue, A. Kakuto and T. Takano, Progr. Theor. Phys. **75** (1986) 664.
- A. Anselm and A. Johansen, Phys. Lett. **B200** (1988) 331;
 A. Anselm, Sov. Phys. JETP 67 (1988) 663.
- 17. Z. Berezhiani and G. Dvali, Sov. Phys. Lebedev Inst. Reports 5 (1989) 55.
- 18. R. Barbieri, G. Dvali and M. Moretti, Phys. Lett. B322 (1993) 173.
- 19. R. Barbieri *et al.*, Nucl. Phys. **B432** (1994) 49.
- 20. Z.G. Berezhiani, Phys. Lett. **B355** (1995) 481.
- 21. Z. Berezhiani, C. Csaki and L. Randall, Nucl. Phys. **B444** (1995) 61.
- 22. G. Dvali and S. Pokorski, hep-ph/9610341
- Z. Berezhiani, Proc. Int. Workshop "Physics from Planck Scale to Electroweak Scale", eds. P. Nath, T. Taylor and S. Pokorski, World Scientific, Singapore, p. 240, hep-ph/9412372;

C. Csaki, Talk presented at DPF'96, hep-ph/9611336.

- 24. C. Csaki and L. Randall, Nucl. Phys. **B466** (1996) 41;
- B. Anantharayan and Q. Shafi, Phys. Rev. **D54** (1996) 3488.
- R. Barbieri, J. Ellis and M.K. Gaillard, Phys. Lett. **90B** (1980) 249;
 E. Akhmedov, Z. Berezhiani, G. Senjanović, Phys. Rev. Lett. **69** (1992) 3013.
- 26. G.W. Anderson et al., Phys. Rev. D49 (1994) 3660;
 K.S. Babu and R.N. Mohapatra, Phys. Rev. Lett. 74 (1995) 2418;
 Z. Berezhiani, Phys. Lett. B355 (1995) 178, and references therein.