

Quasiparticles and the quantum fluctuations of local observables

S. Ying

CCAST (World Laboratory), P. O. Box 8730, Beijing 100080, People's Republic of China
and

*Physics Department, Fudan University, Shanghai 200433, People's Republic of China**

(August 25, 2018)

The role that quasiparticles play in a strong interaction system with spontaneous symmetry breaking is examined. We find, using a non-perturbative cluster decomposition method, that the quasiparticles do not saturate the physical local observables at small distances. The fermion number density serves as a clearcut example. A component due to localized random quantum fluctuations of the order parameter(s) in the vacuum state and the contributions of “quasiparticles” corresponding to other local minima of the effective potential is needed. At large distances, the ordinary quasiparticle picture emerges in the response of the system to classical background fields but the above mentioned component acts as a source for them.

PACS number: 11.15.Tk, 11.30.Rd, 95.35.+d, 74.40.+k, 12.90.+b

In the self-consistent mean field and Hartree–Fock approximations, the behavior of a quasiparticle is similar to an ordinary particle. A particle in a free or weakly interacting theory has definite mass; observables of the theory are saturated by the contributions of these particles. As a consequence, the fermion number density of a system with massive fermions is non-zero only after the chemical potential μ exceeds the mass of the lightest fermions in the system. It is expected that the quasiparticles do the same at the mean field level.

Quasiparticles are however not particles. The essential difference between the mass of a particle and that of a quasiparticle lies in the fact that the mass of a particle is a macroscopic parameter of the theory, with no quantum fluctuations, and the “mass” of the quasiparticle, being proportional to the order parameter, is a local variable. Such a local variable contains quantum fluctuations that do not decouple in the thermodynamical limit. As a result the fermion number density is not saturated solely by the contributions of the quasiparticles but contains a component due to local quantum fluctuations. This component, being non-propagating and uncorrelated with each other in different spacetime region, is called the dark component.

Recent lattice studies of the spontaneous chiral symmetry breaking in QCD at finite density found the emergence of baryon number at chemical potential below the

expected threshold for particle production [1,2] even in the non-quenched approach [2]. The origin of this effect is not yet well understood. It is found here that the dark component is responsible for such a behavior.

For simplicity, the two flavor half bosonized NJL model is used for our investigation. The model Lagrangian density with a chemical potential μ is

$$\mathcal{L} = \bar{\psi} (i\not{\partial} + \mu\gamma^0 - \sigma - i\gamma^5 \vec{\tau} \cdot \vec{\pi}) \psi - \frac{1}{2G_0} (\sigma^2 + \vec{\pi}^2), \quad (1)$$

where G_0 is the coupling constant, σ and $\vec{\pi}$ are auxiliary fields and $\vec{\tau}$ is a set of three Pauli matrices. It has a $SU(2)_L \times SU(2)_R$ chiral symmetry.

After performing the integration over the fermion fields ψ and $\bar{\psi}$, the generating functional $W[J]$ can be written as

$$e^{W[J,\mu]} = \int D[\sigma, \vec{\pi}] e^{iS_{eff}[\sigma, \vec{\pi}, \mu] + i \int d^4x f \cdot J}, \quad (2)$$

where “ J ” and “ f ” represent, collectively, the external fields and auxiliary fields respectively. The effective action S_{eff} is given by

$$S_{eff}[\sigma, \vec{\pi}, \mu] = -i \frac{1}{2} Sp Ln S_F^{-1}[\sigma, \vec{\pi}, \mu] S_F[0, 0, 0] + \frac{1}{2G} \int d^4x (\sigma^2 + \vec{\pi}^2)$$

where Sp denotes the functional trace. The operator $iS_F^{-1}[\sigma, \vec{\pi}, \mu] = i\not{\partial} + \mu\gamma^0 - \sigma - i\gamma^5 \vec{\tau} \cdot \vec{\pi}$ is the inversed propagator of the fermions in the background auxiliary fields $\sigma(x)$ and $\vec{\pi}(x)$.

Let us define $\rho[\sigma, \vec{\pi}, \mu; x] \equiv \text{Tr} \gamma^0 \langle x | S_F[\sigma, \vec{\pi}, \mu] | x \rangle$, where the trace “Tr” is over the internal degrees of freedom of the fermions. The fermion number density of the model in a state $|\psi\rangle$ determined by the initial and final configurations of the auxiliary fields is

$$\langle \psi | \hat{\rho}(x) | \psi \rangle = \frac{1}{Z} \int D[\sigma, \vec{\pi}] \rho[\sigma, \vec{\pi}, \mu; x] e^{iS_{eff}[\sigma, \vec{\pi}, \mu]}, \quad (4)$$

where $Z = \int D[\sigma, \vec{\pi}] e^{iS_{eff}[\sigma, \vec{\pi}, \mu]}$. The Minkowski spacetime is not suitable to study the properties of the vacuum state using Eq. 4 since the initial and final auxiliary field configurations are not specified. For a given set of configurations, the contributing intermediate states are not merely the vacuum states but are contaminated by the excited states. The usual procedure to project out the contributions of the vacuum state is to go to the Euclidean spacetime in which the vacuum state has lowest energy.

*The present address.

In the mean field approximation, the vacuum phase is determined by minimizing the effective potential $V_{eff}(\sigma, \mu) = -S_{eff}[\sigma, 0, \mu]/\Omega$ with $\Omega \rightarrow \infty$ the spacetime volume of the system, σ spacetime independent and $\vec{\pi}$ assumed zero. Due to the chiral symmetry, the assumption $\vec{\pi} = 0$ does not loss generality. The phase of the system is determined by the condition $\delta V_{eff}(\sigma, \mu)/\delta\sigma = 0$. The solution for $\bar{\sigma}$ is non-zero after the coupling constant G_0 is greater than a critical value G_{0c} . A non-vanishing $\bar{\sigma}$ generates an effective mass for the fermions to form quasiparticles.

The mean field fermion number density for the vacuum is obtained from Eq. 4 by ignoring the functional integration over σ and $\vec{\pi}$ and let $\sigma = \bar{\sigma}$. The result is [3]

$$\bar{\rho}_{MF} = \rho(\bar{\sigma}, 0, \mu; x) = \theta(\mu - \bar{\sigma}) [\mu^2 - \bar{\sigma}^2]^{3/2}, \quad (5)$$

which is non-zero only when $\mu > \bar{\sigma}$ just like the fermion number density of free massive particles with mass $m = \bar{\sigma}$. Therefore, the quasiparticle contributions saturate the fermion number density in the mean field approximation.

The contributions of quantum fluctuations around the mean field $\bar{\sigma}$ are formally included in Eq. 4. The results are commonly expressed as loop corrections to the fermion number density vertex. It is not attempted here.

Instead of performing a loop expansion computation of the fermion number density, we evaluate non-perturbatively the effects of the quantum fluctuations by “doing” the path integral.

To proceed, the system under consideration is first putted in a Euclidean spacetime box of length L with periodic boundary conditions at its surfaces. The thermodynamical limit is defined as the limit of $L \rightarrow \infty$. In the thermodynamical limit, the extremal configuration dominates the path integral among those configurations of σ and $\vec{\pi}$ that give divergent action in the thermodynamical limit. The contributing finite action quantum fluctuation configurations, the number of which is proportional to the spacetime volume $\Omega = L^4$, are further classified into two categories: 1) correlated localized configurations, which are defined as the ones that approaches to the mean field configurations in the spacetime infinity and 2) correlated extended configurations, which are the ones that remain different from the mean field configurations at the spacetime infinity.

For a given system, whether or not a configuration is an correlated extended configurations or is of localized ones¹ is determined by dynamics. The on-shell amplitudes, are solutions of the “classical equation of motion” in the Euclidean spacetime

$$\frac{\delta S_{eff}^E[f]}{\delta f(x)} = 0, \quad (6)$$

¹In the sense that it can be decomposed into localized ones.

with S_{eff}^E the Euclidean effective action, f representing σ or $\vec{\pi}$ fields. The superscript E shall be suppressed in the following. The set of the extended solutions to Eq. 6 are correlated ones. Albeit there are plenty of extended on-shell amplitudes in the Minkowski spacetime, there is no known one in the Euclidean one. We shall assume the absence of them. The degree of correlation of an arbitrary extended configuration at different spacetime points is determined by the degree of their deviation from the extended on-shell amplitudes for propagating excitations of the system.

The correlation between the off-shell configurations at two different space time points decreases exponentially. For example, consider the field-field correlations or propagators

$$\langle 0 | T \hat{f}(x_1) \hat{f}(x_2) | 0 \rangle = \int \frac{d^4 p}{(2\pi)^4} e^{-ip \cdot (x_1 - x_2)} \frac{1}{p^2 + m^2} \quad (7)$$

with T denoting time ordering. The off-shellness of these configurations is measured by their mass m . The correlation of these configurations at two different location x_1 and x_2 (in the Euclidean spacetime) decreases as fast as $\exp(-|x_1 - x_2|m)$. So they can be decomposed into superposition of localized ones with a size of order $1/m$. For these set of configurations, one can divide the spacetime into cells with a dimension sufficiently larger than their correlation length. Then the contribution of this set of field configurations to the partition functional Eq. 2 can be cluster decomposed to

$$Z[J] \approx \prod_k z_k[J], \quad W[J] \approx \sum_k w_k[J] \quad (8)$$

with $w_k[J] \equiv \ln z_k[J]$, $z_k[J]$ and $w_k[J]$ the corresponding partition functional of the k th cell. The full partition of the k th cell is

$$z_k[0] \sim \int D_k[f] e^{-S_{eff}^{(k)}(\vec{f}+f)}, \quad (9)$$

where $S_{eff}^{(k)}[f]$ is the effective action of the k th cell and the path integration of f is over those ones that equals to the mean field value \vec{f} outside of the k th cell but with arbitrary amplitudes inside the finite volume. For a given theory, instead of arbitrary division of the spacetime, it is expected that there is an optimal one with minimum volume ϖ for each cell and yet has an error below a pre-determined one. We shall assume that such an optimal division of the spacetime into cells has already been found in the following discussion.

Let us evaluate the contributions of the uncorrelated localized quantum fluctuations of the σ and $\vec{\pi}$ fields to the vacuum fermion number density using Eqs. 4, 8 and 9 in the Euclidean spacetime.

In the phase where the chiral $SU(2)_L \times SU(2)_R$ symmetry is spontaneously broken down, the “chiral angle”

variable represented by $\vec{\pi}$ in the phase where $\langle 0|\vec{\pi}|0\rangle = 0$ becomes massless following the Goldstone theorem. The correlation length of the $\vec{\pi}$ field becomes divergent in the chiral symmetrical limit. Therefore the field configurations of the Goldstone boson degrees of freedom contain the dominating on-shell components that can be included by doing a loop expansion as usual. Such a loop expansion contains no infrared divergences. Because the fermion number density $\rho(\sigma, \vec{\pi}, \mu; x)$ under study is chiral symmetric, which means that it does not depend on a spacetime independent global “chiral angle”; it depends only on the derivatives of the “chiral angle” variables. The absence of a dependence of the fermion number density on a global “chiral angle” guarantees the absence of the infrared divergences in the quantum corrections due to the Goldstone bosons. The configurations of the “chiral angle”, being on the edge of their shell and extended configurations in nature, are uniform and infinitesimal in amplitudes since it contains no infrared divergences and has a number of distinct modes proportional to the volume Ω of the system. They can not modify the qualitative features of the quasiparticles. So, the $\vec{\pi}$ variable in the vacuum fermion number density can be eliminated. It is treated as zero in the following discussions.

The quantum fluctuation in the “mass” term, namely the chiral radius or order parameter represented by σ (when $\langle 0|\vec{\pi}|0\rangle = 0$) has different characteristics due to the fact that it contains no on-shell Euclidean configurations. These off-shell configurations have only short range correlations in spacetime.

If the spacetime is divided into cells with their dimension optimally determined, then the path integration within each cell can be done independently. This gives us a cluster decomposed partition functional of the form given by Eq. 8 with the partition functional for each cell computed independently of each other.

The cluster decomposition property of the partition functional of the system reduces the full fermion number density of the vacuum given by Euclidean form of Eq. 4 to

$$\rho_{vac} = \frac{1}{z_k[0]} \int D_k[\sigma'] \rho[\bar{\sigma} + \sigma', 0, \mu; x] e^{-S_{eff}^{(k)}[\bar{\sigma} + \sigma', \mu]} \quad (10)$$

where the cell labeled by k is the one that contains the spacetime point x , $\int D_k[\sigma']$ denotes integration over field configurations that approach the mean field value outside the cell and $S_{eff}^{(k)}$ is the Euclidean effective action of the cell.

Eq. 10 is still too complicated to evaluate analytically. We make a further simplification by assuming that the functional integration of σ' within a spacetime cell can be replaced by an ordinary integration over the spacetime averaged value of σ' within that cell. It can be achieved by keeping the average of σ' fixed while integrate over the rest degrees of freedom. The non-trivial part of it is in the assumption that after eliminating the rest of

the degrees of freedom, the resulting effective potential V_{eff} remains, at least in form, the same as the original one. This procedure is in the same spirit as the renormalization group analysis. In this way, Eq. 10 reduces to

$$\rho_{vac} = \frac{1}{z[0]} \int_{-\infty}^{\infty} d\delta\sigma \rho[\bar{\sigma} + \delta\sigma, 0, \mu; x] e^{-\varpi V_{eff}(\bar{\sigma} + \delta\sigma, \mu)}, \quad (11)$$

where $\delta\sigma = \langle \sigma' \rangle$ is the spacetime average of σ' within the cell and the same reduction is also made to $z[0]$. The finiteness of ϖ result in different qualitative behavior for ρ_{vac} as a function of μ . To explicitly see the difference, let us expand V_{eff} around $\bar{\sigma}$, keeping only the leading quadratic term

$$V_{eff}(\bar{\sigma} + \delta\sigma) = V_{eff}(\bar{\sigma}) + A(\bar{\sigma})\delta\sigma^2 + \dots \quad (12)$$

and using Eq. 5 for the trace term. If ϖA is sufficiently large, the result can be written as

$$\rho_{vac} = \frac{2}{\pi^2} \sqrt{\frac{\varpi A}{\pi}} \int_{-\mu-\bar{\sigma}}^{\mu-\bar{\sigma}} d\delta\sigma e^{-\varpi A \delta\sigma^2} [\mu^2 - (\bar{\sigma} + \delta\sigma)^2]^{3/2}. \quad (13)$$

It is clearly non-zero for any finite μ below $\bar{\sigma}$ since A is finite except in the limit of $G_0 \rightarrow 0$, which represents the free field case. The dependence of $\rho_{vac}^{1/3}$ on μ for a set of different values of $\alpha = \varpi A$ is plotted in Fig. 1. Instead of a sharp rise in $\rho_{vac}^{1/3}$ at $\mu = \bar{\sigma}$, ρ is non-vanishing all the way to $\mu = 0$. This component of the fermion number density can certainly not be attributed to the contributions of the quasiparticles. We find therefore that there is a dark component for the fermion number density that can not be accounted for by the quasiparticle contributions.

The NJL model has only one non-trivial vacuum phase since there is one independent absolute minimum in its effective potential. If the effective potential of the system contains a second local minimum with a higher energy density than the absolute minimum, then there are contributions from the local minimum to the fermion number density. The existence of such a contribution is another direct consequences of the cluster decomposability of the partition functional of the system. Models with a second minimum are studied in the literature. Those that have only one order parameter are represented by the Friedberg–Lee model [4] in which the two minima of the effective potential correspond to confinement and deconfinement phase of the model. Those that has two order parameters are introduced in Refs. [5,6] in which the β and ω phases of the massless fermion system are studied.

In the presence of a higher virtual phase that is separated from the actual phase of the system by a potential barrier, the fermion number density of the system is saturated by both the quasiparticles of the first phase and those ones of the second virtual phase together with their corresponding dark component discussed above. It has a form

$$\rho_{vac} = \frac{\rho_{vac_1} + e^{-\varpi\Delta}\rho_{vac_2}}{1 + e^{-\varpi\Delta}}, \quad (14)$$

where $\Delta = V_{eff}^{(2)} - V_{eff}^{(1)} > 0$ is the difference in energy density between the virtual phase and the actual phase, ρ_{vac_1} and ρ_{vac_2} are the vacuum fermion number density of the actual vacuum phase and that of the virtual vacuum phase respectively and ϖ is the optimal volume of the spacetime cell between which the order parameters of the system are decorrelated. The contributions of the virtual phase to the fermion number density or to any local physical observables are non-perturbative effects. Attempts had been made in Refs. [6,7] to search for possible other virtual phases [5,6] of the strong interaction vacuum.

The fermion number density discussed above are defined on a spacetime point. Such a precision is non-achievable in realistic observations. The physical observables can be represented by a coarse-grained averaging of the form $\bar{\rho}_{vac} = \Delta N_{vac}/\Delta\Omega$ with $\Delta\Omega$ the smallest volume in spacetime that the observation apparatus can resolve and ΔN_{vac} the average number of fermions *coherently* produced by an external classical field K within that volume. ΔN_{vac} is not in general identical to $\int_{\Delta\Omega} d^4x \rho_{vac}$ when $\Delta\Omega \gg \varpi$. It is obtained from the partition functional $W[J, \mu]$ by adding the external field K to μ . K has a constant non-zero value only within spacetime volume $\Delta\Omega$, then $\Delta N_{vac} = \partial W[0, \mu + K]/\partial iK|_{K=0}$. In case when $\Delta\Omega \ll \varpi$, namely the precision of the observation is much higher than the correlation length of the order parameter, the observed fermion number density $\bar{\rho}_{vac}$ behaves in the same way as ρ_{vac} . On the other hand, if $\Delta\Omega \gg \varpi$, then the smallest spacetime cell that contributes to ΔN_{vac} is a region of volume $\Delta\Omega$ rather than ϖ . In this case, $\bar{\rho}_{vac}$ is obtained from ρ_{vac} by substituting $\Delta\Omega$ for ϖ . The effects of the dark component in the observed fermion number density are reduced as a result. When the resolution is sufficiently low, which means $\Delta\Omega A \gg 1$, the quasiparticle picture with only one vacuum, namely the actual one, reemerges in the response of the system to K . Therefore the dark component of the fermion number density is of transient nature.

On the other hand, the K field radiated by the vacuum fermion number density is of the form $K(x) = \int d^4x' G(x, x') \rho_{vac}(x')$ with $G(x, x')$ the Green function for $K(x)$. It is ρ_{vac} that is the source for $K(x)$ rather than $\bar{\rho}_{vac}$ no matter how slow the resulting $K(x)$ varies in spacetime. Thus the effects of the dark component are indirectly observable even in low energy processes.

Albeit the fermion number density is discussed, the results of the letter are also applicable to other local observables like the energy density. The finding may have implications on the dark matter problem in cosmology since it implies that the apparent matterless space at the macroscopic level may reveal itself of matter effects in low energy gravitational processes even when μ is below

the baryonic particle production threshold and the $\mu = 0$ energy density subtracted. This is because gravitational fields couple locally to the source matter fields that contain the localized dark component.

In summary, it is found that the quasiparticle picture is insufficient in describing quantum processes of strong interaction at both small and large distance scales. The random quantum fluctuations of order parameters of the system can change the behavior of the system qualitatively. The finding may be relevant to our understanding of the dark matter problem in astronomy. Other implications in hadron physics are also worth exploring in the future.

-
- [1] R. A. Janik, M. A. Nowak, G. Papp, and I. Zahed, Phys. Rev. Lett. **77**, 4876 (1996); M. A. Stephanov, Phys. Rev. Lett. **76**, 4472 (1996); A. Gocksch, Phys. Rev. **D37**, 1014 (1988).
 - [2] E. Shuryak, Report No. hep-ph/9609249.
 - [3] S. Ying, Report No. hep-th/9611166; S. Ying, Submitted for a publication.
 - [4] R. Friedberg and T. D. Lee, Phys. Rev. **D15**, 1694 (1977); **D16**, 1096 (1977); **D18**, 2623 (1978).
 - [5] S. Ying, Ann. Phys. (N.Y.) **250**, 69 (1996); S. Ying, Phys. Lett. **B283**, 341 (1992).
 - [6] S. Ying, Report No. hep-ph/9604255 (1996); S. Ying, Submitted for a publication.
 - [7] S. Ying, J. Phys. **G22**, 293 (1996); S. Ying, Commun. Theor. Phys. to be published; Report No. hep-ph/9506290.

FIGURES

Fig. 1 The dependence of $\rho_{vac}^{1/3}$ on the chemical potential μ . Here $\alpha = \varpi A$. The unit for the dimensional quantities are GeV . Solid lines represent the case of free theory with mass 0 and 0.5 respectively. Other lines represent the results for the massless NJL model with different strength of local fluctuations characterized by the α of the order parameter.

