Supersymmetric Inflation, Baryogenesis and $\nu_{\mu} - \nu_{\tau}$ Oscillations

G. Lazarides ^{*a*}, Q. Shafi ^{*b*} and N. D. Vlachos ^{*c*}

a) Theory Division, CERN, 1211 Geneva 23,

Switzerland

and

Physics Division, School of Technology, Aristotle University of Thessaloniki, Thessaloniki GR 540 06, Greece.

> b) Bartol Research Institute, University of Delaware, Newark, DE 19716, USA.

c) Department of Physics, Aristotle University of Thessaloniki, Thessaloniki GR 540 06, Greece.

Abstract

In a supersymmetric left-right symmetric model, inflation, baryogenesis (via leptogenesis) and neutrino oscillations can become closely linked. A familiar ansatz for the neutrino Dirac masses and mixing of the two heaviest families, together with the MSW resolution of the solar neutrino puzzle, imply that $1 \text{ eV} \lesssim m_{\nu_{\tau}} \lesssim 9 \text{ eV}$. The predicted range for the mixing angle $\theta_{\mu\tau}$ will be partially tested by the Chorus/Nomad experiment. The CP violating phase $\delta_{\mu\tau}$ is also discussed.

The minimal supersymmetric standard model (MSSM) provides a particularly compelling extension of the $SU(3)_c \times SU(2)_L \times U(1)_Y$ gauge theory. Yet, it seems quite clear that MSSM, in turn, must be part of a larger picture. Let us list some reasons why: i) In MSSM, there is no understanding of how the supersymmetric μ term is ~ 10²-10³ GeV. In principle, it could be as large as the Planck mass. ii) An important (and undetermined) new parameter in MSSM is $\tan \beta$, the ratio of the vacuum expectation values of the two higgs doublets. Among other things, an understanding of this parameter can shed light on the mass of the Weinberg-Salam higgs. iii) It has become increasingly clear that a combination of both 'cold' and 'hot' dark matter (CHDM) provides [1] a good fit to the data on large scale structure formation, especially if the primordial density fluctuations are essentially scale invariant. In MSSM, there is no HDM candidate, even after including non-renormalizable terms. iv) It has been impossible, so far, to implement inflation within MSSM. v) Last, but not least, it is not easy to generate in MSSM the observed baryon asymmetry through the electroweak sphaleron processes.

Remarkably, all these challenges can be overcome in one fell swoop by considering a modest extension of the MSSM gauge symmetry to $H \equiv SU(3)_c \times SU(2)_L \times SU(2)_R \times$ $U(1)_{B-L}$. Of course, it is anticipated that H is embedded in a grand unified theory such as SO(10) or $SU(3)_c \times SU(3)_L \times SU(3)_R$. Apart from aesthetics, there are tantalizing hints from the extrapolation of low energy data for the existence of a supersymmetric unification scale $\sim 10^{16}$ GeV. The details on how the extension of MSSM to an H-based model can resolve the points above will not be discussed here, especially since an inflationary scenario based on H has been considered in some detail elsewhere [2,3]. This scenario gives rise to an essentially scale invariant spectrum (spectral index $n \simeq 0.98$), and contains both 'cold' (LSP) and 'hot' (massive neutrinos) dark matter candidates. The observed baryon asymmetry is generated through partial conversion of a primordial lepton asymmetry [4]. Finally, the parameter tan β is close to m_t/m_b , which also explains why the higgs boson of the electroweak theory has not been seen at LEP II. Its tree level mass is M_Z which, after radiative corrections, becomes $m_{h^o} \simeq 105 - 120$ GeV.

The inflationary phase is associated with the gauge symmetry breaking $SU(2)_R \times U(1)_{B-L} \to U(1)_Y$. Of course, since H is presumably embedded in some grand unified symmetry, there may well be more than one inflationary epoch. We concentrate on the last and most relevant one. The above breaking is achieved by a pair of $SU(2)_R$ doublet 'higgs' superfields which have the same gauge quantum numbers as the 'matter' right handed neu-

trino superfields. As a consequence, the inflaton decays primarily into 'matter' right handed neutrinos via quartic (or higher order) superpotential couplings. The 'reheat' temperature, T_R , turns out to be about one order of magnitude smaller than the mass of the heaviest right handed neutrino that the inflaton can decay into. The gravitino constraint on $T_R (\leq 10^9 \text{ GeV})$ allows us to restrict the second and third family right handed neutrino masses M_2, M_3 in a fairly narrow range. Our approach poses no obvious constraint on the first family right handed neutrino mass M_1 , except from $M_1 \leq M_2$. The constraints on M_2, M_3 , however, together with the leptogenesis scenario will enable us to restrict the oscillation parameters of the ν_{μ} - ν_{τ} system [5].

We consider the 2 × 2 'asymptotic' mass matrices $M^{\rm L}$, $M^{\rm D}$ and $M^{\rm R}$ in the weak basis, where the superscripts L,D and R denote the charged lepton, neutral Dirac, and right handed neutrino sectors respectively. $M^{\rm L}, M^{\rm D}$ are diagonalized by the biunitary rotations $L = U^{L}L', L^{c} = U^{L^{c}}L^{c'}, \nu = U^{\nu}\nu', \nu^{c} = U^{\nu^{c}}\nu^{c'}$:

$$M^L \to M^{L\,\prime} = \tilde{U}^{L^c} M^L U^L = \begin{pmatrix} m_\mu \\ m_\tau \end{pmatrix} \quad , \tag{1}$$

$$M^D \to M^{D'} = \tilde{U}^{\nu^c} M^D U^{\nu} = \begin{pmatrix} m_2^D \\ m_3^D \end{pmatrix} , \qquad (2)$$

where the diagonal entries are positive. This gives rise to the 'Dirac' mixing matrix $U^{\nu \dagger}U^{L}$ in the leptonic charged currents. Using the remaining freedom to perform arbitrary phase transformations on the components of L', ν' together with the compensating ones on the components of $L^{c'}, \nu^{c'}$ so that $M^{L'}, M^{D'}$ remain unaltered, we can bring this matrix to the form

$$U^{\nu \dagger} U^{L} \to \begin{pmatrix} \cos \theta^{D} & \sin \theta^{D} \\ -\sin \theta^{D} & \cos \theta^{D} \end{pmatrix}, \qquad (3)$$

where $\theta^D (0 \le \theta^D \le \pi/2)$ is the 'Dirac' (not the physical) mixing angle in the 2-3 leptonic sector.

In this basis, depicted with a double prime on the superfields, the Majorana mass matrix can be written as:

$$M^R = U^{-1} \ M_0 \ \tilde{U}^{-1} \ , \tag{4}$$

where $M_0 = diag(M_2, M_3)$, with M_2, M_3 (both positive) being the two Majorana masses, and U is a unitary matrix which can be parametrized as

$$U = \begin{pmatrix} \cos\theta & \sin\theta & e^{-i\delta} \\ -\sin\theta & e^{i\delta} & \cos\theta \end{pmatrix} \begin{pmatrix} e^{i\alpha_2} & \\ & e^{i\alpha_3} \end{pmatrix} , \qquad (5)$$

with $0 \le \theta \le \pi/2$ and $0 \le \delta < \pi$. The light neutrino mass matrix, to leading order in $M^{R-1}M^{D'}$, is

$$m = -\tilde{M}^{D'} \frac{1}{M^R} M^{D'} , \qquad (6)$$

where $M^{D'}$ is defined in eq.(2). We can express m as

$$m = \begin{pmatrix} e^{i\alpha_2} \\ e^{i\alpha_3} \end{pmatrix} \Psi(\theta, \delta) \begin{pmatrix} e^{i\alpha_2} \\ e^{i\alpha_3} \end{pmatrix} , \qquad (7)$$

where $\Psi(\theta, \delta)$ depends also on M_2 , M_3 , m_2^D , m_3^D . We diagonalize m by a unitary rotation $\nu'' = V\nu'''$ with

$$V = \begin{pmatrix} e^{i\beta_2} \\ e^{i\beta_3} \end{pmatrix} \begin{pmatrix} \cos\varphi & \sin\varphi e^{-i\epsilon} \\ -\sin\varphi e^{i\epsilon} & \cos\varphi \end{pmatrix}, \qquad (8)$$

where $0 \le \varphi \le \pi/2$, $0 \le \epsilon < \pi$. The 'Dirac' mixing matrix in eq.(3) is now multiplied by V^{\dagger} on the left and, after suitable phase absorptions, takes the form

$$\begin{pmatrix} \cos\theta_{23} & \sin\theta_{23} e^{-i\delta_{23}} \\ -\sin\theta_{23} e^{i\delta_{23}} & \cos\theta_{23} \end{pmatrix}, \qquad (9)$$

where $0 \le \theta_{23} \le \pi/2$, $0 \le \delta_{23} < \pi$. Here, θ_{23} (or $\theta_{\mu\tau}$) is the physical mixing angle in the 2-3 leptonic sector and its cosine equals the modulus of the complex number

$$\cos\varphi\cos\theta^D + \sin\varphi\sin\theta^D e^{i(\xi-\epsilon)} , \qquad (10)$$

where $-\pi \leq \xi - \epsilon = \beta_2 - \beta_3 - \epsilon \leq \pi$. Moreover, δ_{23} (or $\delta_{\mu\tau}$) is the associated CP violating phase which is given by $\delta_{23} = \xi + \rho - \lambda$ (modulo π), where λ ($-\pi \leq \lambda \leq \pi$) and $-\rho$ ($-\pi \leq \rho \leq \pi$) are the arguments of the complex numbers in eq.(10) and $\cos\varphi \sin\theta^D - \sin\varphi \cos\theta^D e^{i(\xi-\epsilon)}$ respectively. Since ξ remains undetermined (see below), the precise values of θ_{23} and δ_{23} cannot be found. However, we can determine the range in which θ_{23} lies:

$$|\varphi - \theta^D| \le \theta_{23} \le \varphi + \theta^D$$
, for $\varphi + \theta^D \le \pi/2$. (11)

The double valued function $\delta_{23}(\theta_{23})$ is also determined.

We will denote the two positive eigenvalues of the light neutrino mass matrix by m_2 (or $m_{\nu_{\mu}}$), m_3 (or $m_{\nu_{\tau}}$) with $m_2 \leq m_3$. Recall that all the quantities here (masses, mixings, etc.) are 'asymptotic' (defined at the GUT scale). The determinant and the trace invariance of $\Psi(\theta, \delta)^{\dagger}\Psi(\theta, \delta)$ provide us with two constraints on the (asymptotic) parameters which take the form:

$$m_2 m_3 = \frac{\left(m_2^D m_3^D\right)^2}{M_2 \ M_3} \quad , \tag{12}$$

$$m_2^2 + m_3^2 = \frac{\left(m_2^{D\ 2}c^2 + m_3^{D\ 2}s^2\right)^2}{M_2^2} +$$
(13)

$$\frac{\left(m_3^{D\ 2}c^2 + m_2^{D\ 2}s^2\right)^2}{M_3^2} + \frac{2(m_3^{D\ 2} - m_2^{D\ 2})^2c^2s^2\cos 2\delta}{M_2\,M_3},$$

where θ , δ are defined in eq.(5), $c = \cos \theta$, $s = \sin \theta$.

We now need information about the quantities m_2^D, m_3^D (the 'asymptotic' Dirac masses of the muon and tau neutrinos) as well as the 'Dirac' mixing angle θ^D . A plausible assumption, inspired by SO(10) for instance, is that these quantities are related to the quark sector parameters by the $SU(4)_c$ symmetry. In other words, we will assume the asymptotic relations:

$$m_2^D = m_c , \ m_3^D = \ m_t , \ \sin^D = |V_{cb}|$$
 (14)

Of course, the $SU(4)_c$ symmetry is not expected to hold in the down sector of the second family.

Contact with experiment can be made after renormalization effects have been taken into account. The pair of MSSM higgs doublets is assumed to belong to the (2, 2) representation of $SU(2)_L \times SU(2)_R$, implying that $\tan \beta \simeq m_t/m_b$ both 'asymptotically' and at low energies. The light neutrino masses, in this case, can be obtained by dividing the right hand side of eq.(6) by a factor of 2.44 [6]. Eqs.(12),(13) now hold with m_2,m_3 being the low energy neutrino masses and m^D 's replaced by their asymptotic values divided by 1.56. The latter turn out to be $m_2^D \simeq 0.23$ GeV and $m_3^D \simeq 116$ GeV (with $\tan \beta \simeq m_t/m_b$)[7]. Finally, using $SU(4)_c$ invariance, the asymptotic 'Dirac' mixing angle is calculated from $\sin \theta^D =$ $|V_{cb}|$ (asymptotic) $\simeq 0.03$. The renormalization of the mixing angle θ_{23} , with $\tan \beta \simeq m_t/m_b$, has been considered in ref.[6]. The net effect is that $\sin^2 2\theta_{23}$ increases by about 40% from M_{GUT} to M_Z .

In view of the lack of a compelling alternative theoretical framework, we will assume the hierarchy $m_1 \ll m_2 \ll m_3$. We will thus restrict m_2 in the range 1.7×10^{-3} eV $\lesssim m_2 \lesssim 3.5 \times 10^{-3}$ eV, as allowed by the small angle MSW solution [8]. We now recall a few salient features of the inflationary scenario associated with the breaking $SU(2)_R \times U(1)_{B-L} \rightarrow U(1)_Y$. As the inflaton $(SU(2)_R \text{ doublets } \phi, \bar{\phi})$ oscillates about its minimum, it decays into the appropriate 'matter' right handed neutrino (ν^c) via the effective superpotential coupling $\nu^c \nu^c \bar{\phi} \bar{\phi}$ permitted by the gauge symmetry. The 'reheat' temperature T_R is then related [3] to the mass M_H of the heaviest right handed neutrino the inflaton can decay into: $T_R \simeq M_H/9.2$. The inflaton decays predominantly into the heaviest of the two. Then, eq.(12) and the cosmological bound $m_3 \lesssim 23$ eV [9] require the smallest allowed mass of the heaviest right handed neutrino to be $\simeq 9.4 \times 10^{10}$ GeV giving $T_R \gtrsim 10^{10}$ GeV, in clear conflict with the gravitino constraint. Consequently, we find that

$$m_{infl}/2 \leq M_3 \lesssim 2.5 \times 10^{13} \text{ GeV},$$
 (15)

where the upper bound comes from the requirement that the coupling constant of the nonrenormalizable superpotential term $\nu^c \nu^c \bar{\phi} \bar{\phi}$, which provides mass for the right handed neutrinos, should not exceed unity. Thus, M_3 is restricted in a narrow range, and the inflaton decays into the second heaviest right handed neutrino.

The lepton asymmetry is generated by the subsequent decay of this neutrino and is given by [4]

$$\frac{n_L}{s} = -\frac{9 T_R}{8\pi m_{infl}} \frac{M_2}{M_3} \frac{Im(UM^{D'}M^{D'}^{\dagger}U^{\dagger})_{23}^2}{v^2(UM^{D'}M^{D'}^{\dagger}U^{\dagger})_{22}} , \qquad (16)$$

where v is the electroweak VEV at M_{GUT} . Substituting U from eq.(5), we get

$$\frac{n_L}{s} = \frac{9 T_R}{8\pi m_{infl}} \frac{M_2}{M_3} \frac{c^2 s^2 \sin 2\delta \ (m_3^{D\ 2} - m_2^{D\ 2})^2}{v^2 (m_3^{D\ 2} \ s^2 \ + \ m_2^{D\ 2} \ c^2)} \ . \tag{17}$$

Here we can again replace m_2^D , m_3^D by their 'asymptotic' values divided by 1.56 and v by 174 GeV. Assuming the MSSM spectrum between 1 TeV and M_{GUT} , the observed baryon asymmetry n_B/s is related [10] to n_L/s by $n_B/s = -28/79 (n_L/s)$.

We take a fixed value of M_3 in the allowed range (15) and, for any pair of values (m_2, m_3) , we calculate M_2 and T_R . The constraint $T_R \leq 10^9$ GeV yields a lower bound for the product m_2m_3 excluding the region below a hyperbola on the m_2,m_3 plot. Note that the gravitino constraint combined with the MSW restriction on m_2 yields a lower bound for m_3 . Inside the allowed m_2,m_3 region, we can use the trace condition (13) to solve for $\delta(\theta)$ in the interval $0 \leq \theta \leq \pi/2$. Given that we need a negative value for n_L/s so that $n_B/s > 0$, we see that there is at most one useful branch of the function $\delta(\theta)$ taken to lie in the region $-\pi/2 \leq \delta(\theta) \leq 0$. The expression for $\delta(\theta)$ is subsequently substituted in eq.(17) for the leptonic asymmetry, and the range of θ satisfying the constraint $0.02 \leq \Omega_B h^2 \leq 0.03$ is found. (This constraint is consistent with the low deuterium abundance as well as with structure formation in 'cold' plus 'hot' dark matter models). The m_2,m_3 pairs, for which this range of θ exists, are consistent with the observed baryon asymmetry. In Fig.1, we depict the areas on the m_2,m_3 plane which satisfy both the gravitino and baryogenesis constraints in the two extreme cases of $M_3 = m_{infl}/2$ (bounded by the thick solid line) and $M_3 = 2.5 \times 10^{13}$ GeV (bounded by the thick dashed line). The lower line, in both cases, corresponds to the gravitino constraint whereas the upper one comes from the baryogenesis constraint. Note that the baryogenesis constraint at $m_2 = 1.7 \times 10^{-3}$ eV provides us with an upper bound for m_3 . In summary, we get an interesting upper as well as a lower bound on m_3 (i.e. on Ω_{HDM} !), and, for each m_3 value, we know the allowed range of m_2 . Namely, for $M_3 = m_{infl}/2$, 1.3 eV $\lesssim m_3 \lesssim 8.8$ eV whereas, for $M_3 = 2.5 \times 10^{13}$ GeV, 0.9 eV $\lesssim m_3 \lesssim 5.1$ eV. Thus, m_{ν_7} is restricted in the range of 1 to 9 eV.

The discussion above can be extended to yield useful information for $\theta_{\mu\tau}$. For each allowed pair m_2, m_3 and, for every value of θ in the allowed range, we construct φ and ϵ in eq.(8). The phases α_2 and α_3 in eq.(5) remain undetermined by the conditions (12),(13) and, consequently, β_2, β_3 in eq.(8) and ξ in eq.(10) remain also undetermined. This fact does not allow us to predict the value of $\theta_{\mu\tau}$ for each value of θ , but only its allowed range given in eq.(11). The union of all these intervals, comprising all allowed values of θ and m_2 for a given m_3 , constitutes the allowed range of $\theta_{\mu\tau}$ for this m_3 . These ranges, after taking the renormalization of $\theta_{\mu\tau}$ into account, are depicted in Fig.2 for all possible values of m_3 , and constitute the allowed area of the neutrino oscillation parameters. The region bounded by the thick solid line is the allowed area for $M_3 = m_{infl}/2$, whereas the one bounded by the thick dashed line corresponds to $M_3 = 2.5 \times 10^{13}$ GeV. The areas tested (to be tested) by past (future) experiments are also indicated in Fig.2. The area excluded by E531 is depicted in Fig.1 and lies above the thin solid (dashed) E531 line, for $M_3 = m_{infl}/2$ (2.5 × 10¹³) GeV). CDHS does not appear to have any appreciable effect. Furthermore, if CHORUS gives negative results, we must further exclude the area above the corresponding thin solid (dashed) line in Fig.1, for $M_3 = m_{infl}/2$ (2.5 × 10¹³ GeV). A possibly negative CHORUS result implies that the upper bound for $m_{\nu_{\tau}}$ drops down to $\simeq 3.7$ eV. Notice that the upper limit on baryon asymmetry has no effect on Figs.1 and 2.

The CP violating phase $\delta_{\mu\tau}$ as a function of $\theta_{\mu\tau}$, for given values of M_3, m_2, m_3 and $\Omega_B h^2$, can now be constructed. We choose $M_3 = m_{infl}/2$, $m_2 = 2.6 \times 10^{-3}$ eV, $m_3 = 4$ eV, $\Omega_B h^2 = 0.025$ and solve for θ . We find two solutions and, for each one of them and any $\xi \ (-\pi \leq \xi - \epsilon \leq \pi)$, we calculate $\theta_{\mu\tau}$ and $\delta_{\mu\tau} = \xi + \rho - \lambda$. Eliminating ξ , we obtain the function $\delta_{\mu\tau}(\theta_{\mu\tau})$ in the region of eq.(11). This function turns out to be double valued-the sum of the two branches equals 2ϵ - and is depicted in Fig.3, for both values of θ , after renormalizing $\theta_{\mu\tau}$. The $\theta_{\mu\tau}$'s excluded by E531, for $m_{\nu\tau} = 4$ eV, are also indicated and lie to the right of the E531 line.

In conclusion, we find that a modest extension of MSSM to $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ can yield significant new results by tying together a number of apparently unrelated phenomena. In particular, inflation can be realized, the spectral index $n \simeq 0.98$, we get both 'cold' (essentially bino) and 'hot' (ν_{τ}) dark matter, while the ν_{τ} mass and ν_{μ} - ν_{τ} mixing is within reach of present and planned experiments. In the simplest scheme, the atmospheric neutrino anomaly remains a mystery.

We thank K.S. Babu for several discussions regarding the renormalization effects, and E. Tsesmelis for information about the experiments. TMR and NATO support under grant numbers ERBFMRXCT-960090 and CRG 970149 is gratefully acknowledged.

REFERENCES

- Q. Shafi and F.W. Stecker, Phys. Rev. Lett. 53, 1292 (1984). For a recent review and other refs. see Q. Shafi and R.K. Schaefer, hep-ph/9612428 (1996).
- [2] G. Dvali, Q. Shafi and R.K. Schaefer, Phys. Rev. Lett. 73, 1886 (1994). For related work, see E.J. Copeland, et al., Phys. Rev. D49, 6410 (1994); A. Linde and A. Riotto, hep-ph/9703209.
- [3] G. Lazarides, R.K. Schaefer and Q. Shafi, hep-ph/9608256, to appear in Phys. Rev. D.
- [4] M. Fukugita and T. Yanagida, Phys. Lett. B174, 45 (1986); G. Lazarides and Q. Shafi,
 Phys. Lett. B258, 305 (1991); L. Covi, E. Roulet and F. Vissani, Phys. Lett. B384,169 (1996).
- [5] For another approach relating baryogenesis and neutrino mixing, see W. Buchmuller and M. Plumacher, Phys. Lett. B389, 73 (1996).
- [6] K.S. Babu, C.N. Leung and J. Pantaleone, Phys. Lett. B319, 191 (1993).
- [7] K.S. Babu and Q. Shafi, Phys. Rev. D47, 5004 (1993); Phys. Lett. B357, 365 (1995).
- [8] A. Smirnov, hep-ph/9611465 and references therein.
- [9] For the consistency of the CHDM model, the Hubble constant is about 50 $kms^{-1}Mpc^{-1}$.
- [10] L. Ibanez and F. Quevedo, Phys. Lett. B283, 261 (1992).

Figure Captions

Fig.1.The allowed regions in the $m_{\nu_{\mu}}, m_{\nu_{\tau}}$ plane.

Fig.2.The allowed regions in the ν_{μ} - ν_{τ} oscillation plot.

Fig.3. The function $\delta_{\mu\tau}(\theta_{\mu\tau})$ for $M_3 = m_{infl}/2$ and 'central' values of $m_2, m_3, n_L/s$.

Fig. 1





Fig. 2

