WIMP AND AXION DARK MATTER^a

M. KAMIONKOWSKI

Department of Physics, Columbia University, 538 West 120th Street, New York, NY 10027 USA

There is almost universal agreement among cosmologists that most of the matter in the Universe is dark, and there are very good reasons to believe that most of this dark matter must be nonbaryonic. The two leading candidates for this dark matter are axions and weakly-interacting massive particles (WIMPs), such as the neutralino in supersymmetric extensions of the standard model. I discuss the arguments for these two dark-matter candidates and review techniques for discovery of these dark-matter particles.

1 Introduction

Almost all astronomers will agree that most of the mass in the Universe is nonluminous. The nature of this dark matter remains one of the great mysteries of science today. Dynamics of cluster of galaxies suggest a universal nonrelativistic-matter density of $\Omega_0 \simeq 0.1-0.3$. If the luminous matter were all there was, the duration of the epoch of structure formation would be very short, thereby requiring (in almost all theories of structure formation) fluctuations in the microwave background which would be larger than those observed. These considerations imply $\Omega_0 \gtrsim 0.3^{1}$. Second, if the current value of Ω_0 is of order unity today, then at the Planck time it must have been 1 ± 10^{-60} leading us to believe that Ω_0 is precisely unity for aesthetic reasons. A related argument comes from inflationary cosmology, which provides the most satisfying explanation for the smoothness of the microwave background². To account for this isotropy, inflation must set Ω (the total density, including a cosmological constant) to unity.

However, the most robust observational evidence for the existence of dark matter involves galactic dynamics. There is simply not enough luminous matter ($\Omega_{\rm lum} \lesssim 0.01$) observed in spiral galaxies to account for their observed rotation curves (for example, that for NGC6503 shown in Fig. 1 ³). Newton's laws imply galactic dark halos with masses that contribute $\Omega_{\rm halo} \gtrsim 0.1$.

^aTo appear in the proceedings of the 1997 ICTP Summer School on High Energy Physics and Cosmology, Trieste, Italy, June 2–July 4, 1997.




Figure 1: Rotation curve for the spiral galaxy NGC6503. The points are the measured circular rotation velocities as a function of distance from the center of the galaxy. The dashed and dotted curves are the contribution to the rotational velocity due to the observed disk and gas, respectively, and the dot-dash curve is the contribution from the dark halo.

On the other hand, big-bang nucleosynthesis suggests that the baryon density is $\Omega_b \lesssim 0.1^4$, too small to account for the dark matter in the Universe. Although a neutrino species of mass $\mathcal{O}(30\,\mathrm{eV})$ could provide the right dark-matter density, N-body simulations of structure formation in a neutrino-dominated Universe do a poor job of reproducing the observed structure ⁵. Furthermore, it is difficult to see (essentially from the Pauli principle) how such a neutrino could make up the dark matter in the halos of galaxies ⁶. It appears likely then, that some nonbaryonic, nonrelativistic matter is required.

The two leading candidates from particle theory are the axion ⁷, which arises in the Peccei-Quinn solution to the strong-CP problem, and a weaklyinteracting massive particle (WIMP), which may arise in supersymmetric (or other) extensions of the standard model⁸. As discussed below, there are good reasons to believe that if the Peccei-Quinn mechanism is responsible for preserving CP in the strong interactions, then the axion is the dark matter. Similarly, there are also excellent reasons to expect that if low-energy supersymmetry exists in Nature, then the dark matter should be composed of the lightest superpartner. The study of these ideas are no longer exclusively the domain of theorists: there are now a number of experiments aimed at discovery of axions and WIMPs. If axions populate the Galactic halo, they can be converted to photons in resonant cavities immersed in strong magnetic fields. An experiment to search for axions in this fashion is currently being carried out. If WIMPs populate the halo, they can be detected either directly in low-background laboratory detectors or indirectly via observation of energetic neutrinos from WIMPs which have accumulated and then annihilated in the Sun and/or Earth.

Here, I first show how the observed dynamics of the Milky Way indicate a local dark-matter density of $\rho_0 \simeq 0.4~{\rm GeV~cm^{-3}}$. I then review the arguments for axion and WIMP dark matter and the methods of detection. However, there is no way I can do this very active field of research justice in such short space. For background on early-Universe cosmology, I recommend the book by Kolb and Turner⁹. Readers with further interest in WIMPs should see the recent review article by Jungman, Griest and me ⁸. The first four sections of that article are meant to provide a general review of dark and supersymmetry, and the idea of WIMP dark matter. The remainder of that article provides technical details required by those interested in actively pursuing research on the topic. There are also several excellent axion reviews ⁷ and the recent book by Raffelt ¹⁰.

2 The Local Dark-Matter Density

The extent of the luminous disk of our Galaxy, the Milky Way, is roughly 10 kpc, and we live about 8.5 kpc from the center. Due to our location in it, the rotation curve of the Milky Way cannot be determined with the same precision as that of an external spiral galaxy, such as that shown in Fig. 1. However, it is qualitatively the same. The circular speed rises linearly from zero at the center and asymptotes to roughly 220 km sec⁻¹ somewhere near our own Galactocentric radius and remains roughly flat all the way out to ~ 25 kpc. Although direct measurements of the rotation curve are increasingly difficult at larger radii, the orbital motions of satellites of the Milky Way suggest that the rotation curve remains constant all the way out to radii of 50 kpc and perhaps even farther. According to Newton's laws, the rotation speed should fall as $v_c \propto r^{-1/2}$ at radii greater than the extent of the luminous disk. However, it is observed to remain flat to much larger distances. It therefore follows that the luminous disk and bulge must be immersed in an extended dark halo (or that Newton's laws are violated).

Our knowledge of the halo comes almost solely from this rotation curve. Therefore, we do not know empirically if the halo is round, elliptical, or perhaps flattened like the disk. However, there are good reasons to believe that the halo should be much more diffuse than the disk. The disk is believed to be flat because luminous matter can radiate photons and therefore gravitationally collapse to a pancake-like structure. On the other hand, dark matter (by definition) cannot radiate photons. There are also now empirical arguments which involve, e.g., the shape of the distribution of gas in the Milky Way, which suggest that the dark halo should be much more diffuse than the disk ¹¹.

Given that the halo is therefore nearly round, it must have a density distribution like

$$\rho(r) = \rho_0 \frac{r_0^2 + a^2}{r^2 + a^2},\tag{1}$$

where r is the radius, $r_0 \simeq 8.5$ kpc is our distance from the center, a is the core radius of the halo, and ρ_0 is the local halo density. Such a halo would give rise to a rotation curve,

$$v_h^2(r) = 4\pi G \rho_0(r_0^2 + a^2)[1 - (a/r)\tan^{-1}(r/a)], \tag{2}$$

where G is Newton's constant. If we know the rotation speed contributed by the halo at two points, we can determine ρ_0 and a. At large radii, the rotation curve of the Milky Way is supported entirely by this dark halo, so $v_h(r \gg 10\,\mathrm{kpc}) \simeq 220~\mathrm{km~sec^{-1}}$. However, the rotation curve locally is due in part to the disk, $v_c^2(r_0) = v_d^2(r_0) + v_h^2(r_0)$. The disk contribution to the

local rotation speed is somewhat uncertain but probably falls in the range $v_d(r_0) \simeq 118-155 \text{ km sec}^{-1}$, which means that the halo contribution to the local rotation speed is $v_h(r_0) \simeq 150-185 \; \mathrm{km \; sec^{-1}}$. Given the local and asymptotic rotation speeds, we infer that the local halo density is $\rho_0 \simeq$ $0.3-0.5 \,\mathrm{GeV} \,\mathrm{cm}^{-3}$. The particles which make up the dark halo move locally in the same gravitational potential well as the Sun. Therefore, the virial theorem tells us that they must move with velocities $v \sim v_c \sim 220 \text{ km sec}^{-1}$. Additional theoretical arguments suggest that the velocity distribution of these particles is locally nearly isotropic and nearly a Maxwell-Boltzmann distribution. To sum, application of Newton's laws to our Galaxy tells us that the luminous disk and bulge must be immersed in a dark halo with a local density $\rho_0 \simeq$ $0.4~{\rm GeV}~{\rm cm}^{-3}$ and that dark-matter particles (whatever they are) move with velocities comparable to the local circular speed. More careful investigations along these lines show that similar conclusions are reached even if we allow for the possibility of a slightly flattened halo or a radial distribution which differs from that in Eq. $(1)^{12}$.

3 Axions

Although supersymmetric particles seem to get more attention in the literature lately, we should not forget that the axion also provides a well-motivated and promising alternative dark-matter candidate ⁷. The QCD Lagrangian may be written

$$\mathcal{L}_{QCD} = \mathcal{L}_{pert} + \theta \frac{g^2}{32\pi^2} G\widetilde{G}, \tag{3}$$

where the first term is the perturbative Lagrangian responsible for the numerous phenomenological successes of QCD. However, the second term (where G is the gluon field-strength tensor and \widetilde{G} is its dual), which is a consequence of nonperturbative effects, violates CP. However, we know experimentally that CP is not violated in the strong interactions, or if it is, the level of strong-CP violation is tiny. From constraints to the neutron electric-dipole moment, $d_n \lesssim 10^{-25}$ e cm, it can be inferred that $\theta \lesssim 10^{-10}$. But why is θ so small? This is the strong-CP problem.

The axion arises in the Peccei-Quinn solution to the strong-CP problem ¹³, which close to twenty years after it was proposed still seems to be the most promising solution. The idea is to introduce a global $U(1)_{PQ}$ symmetry broken at a scale f_{PQ} , and θ becomes a dynamical field which is the Nambu-Goldstone mode of this symmetry. At temperatures below the QCD phase transition, non-perturbative quantum effects break explicitly the symmetry and drive $\theta \to 0$. The axion is the pseudo-Nambu-Goldstone boson of this near-global symme-

try. Its mass is $m_a \simeq \text{eV}(10^7 \,\text{GeV}/f_a)$, and its coupling to ordinary matter is $\propto f_a^{-1}$.

A priori, the Peccei-Quinn solution works equally well for any value of f_a (although one would generically expect it to be less than or of order the Planck scale). However, a variety of astrophysical observations and a few laboratory experiments constrain the axion mass to be $m_a \sim 10^{-4}$ eV, to within a few orders of magnitude. Smaller masses would lead to an unacceptably large cosmological abundance. Larger masses are ruled out by a combination of constraints from supernova 1987A, globular clusters, laboratory experiments, and a search for two-photon decays of relic axions 14 .

One conceivable theoretical difficulty with this axion mass comes from generic quantum-gravity arguments ¹⁵. For $m_a \sim 10^{-4}$ eV, the magnitude of the explicit symmetry breaking is incredibly tiny compared with the PQ scale, so the global symmetry, although broken, must be very close to exact. There are physical arguments involving, for example, the nonconservation of global charge in evaporation of a black hole produced by collapse of an initial state with nonzero global charge, which suggest that global symmetries should be violated to some extent in quantum gravity. When one writes down a reasonable ansatz for a term in a low-energy effective Lagrangian which might arise from global-symmetry violation at the Planck scale, the coupling of such a term is found to be extraordinarily small (e.g., $\leq 10^{-55}$). Of course, we have at this point no predictive theory of quantum gravity, and several mechanisms for forbidding these global-symmetry violating terms have been proposed ¹⁶. Therefore, these arguments by no means "rule out" the axion solution. In fact, discovery of an axion would provide much needed clues to the nature of Planck-scale physics.

Curiously enough, if the axion mass is in the relatively small viable range, the relic density is $\Omega_a \sim 1$ and may therefore account for the halo dark matter. Such axions would be produced with zero momentum by a misalignment mechanism in the early Universe and therefore act as cold dark matter. During the process of galaxy formation, these axions would fall into the Galactic potential well and would therefore be present in our halo with a velocity dispersion near 270 km sec⁻¹.

Although the interaction of axions with ordinary matter is extraordinarily weak, Sikivie proposed a very clever method of detection of Galactic axions ¹⁷. Just as the axion couples to gluons through the anomaly (i.e., the $G\widetilde{G}$ term), there is a very weak coupling of an axion to photons through the anomaly. The axion can therefore decay to two photons, but the lifetime is $\tau_{a\to\gamma\gamma}\sim 10^{50}\,\mathrm{s}\,(m_a/10^{-5}\,\mathrm{eV})^{-5}$ which is huge compared to the lifetime of the Universe and therefore unobservable. However, the $a\gamma\gamma$ term in the Lagrangian

is $\mathcal{L}_{a\gamma\gamma} \propto a\vec{E} \cdot \vec{B}$ where \vec{E} and \vec{B} are the electric and magnetic field strengths. Therefore, if one immerses a resonant cavity in a strong magnetic field, Galactic axions which pass through the detector may be converted to fundamental excitations of the cavity, and these may be observable ¹⁷. Such an experiment is currently underway ¹⁸. They have already begun to probe part of the cosmologically interesting parameter space (no, they haven't found anything yet) and expect to cover most of the interesting region parameter space in the next three years. A related experiment, which looks for excitations of Rydberg atoms, may also find dark-matter axions ¹⁹. Although the sensitivity of this technique should be excellent, it can only cover a limited axion-mass range.

It should be kept in mind that there are no accelerator tests for axions in the acceptable mass range. Therefore, these dark-matter axion experiment are actually our *only* way to test the Peccei-Quinn solution.

4 Weakly-Interacting Massive Particles

Suppose that in addition to the known particles of the standard model, there exists a new, yet undiscovered, stable (or long-lived) weakly-interacting massive particle (WIMP), χ . At temperatures greater than the mass of the particle, $T\gg m_\chi$, the equilibrium number density of such particles is $n_\chi\propto T^3$, but for lower temperatures, $T\ll m_\chi$, the equilibrium abundance is exponentially suppressed, $n_\chi\propto e^{-m_\chi/T}$. If the expansion of the Universe were so slow that thermal equilibrium was always maintained, the number of WIMPs today would be infinitesimal. However, the Universe is not static, so equilibrium thermodynamics is not the entire story.

At high temperatures $(T\gg m_\chi)$, χ 's are abundant and rapidly converting to lighter particles and $vice\ versa\ (\chi\bar\chi\leftrightarrow l\bar l$, where $l\bar l$ are quark-antiquark and lepton-antilepton pairs, and if m_χ is greater than the mass of the gauge and/or Higgs bosons, $l\bar l$ could be gauge- and/or Higgs-boson pairs as well). Shortly after T drops below m_χ the number density of χ 's drops exponentially, and the rate for annihilation of χ 's, $\Gamma=\langle\sigma v\rangle\,n_\chi$ —where $\langle\sigma v\rangle$ is the thermally averaged total cross section for annihilation of $\chi\bar \chi$ into lighter particles times relative velocity v—drops below the expansion rate, $\Gamma\lesssim H$. At this point, the χ 's cease to annihilate, they fall out of equilibrium, and a relic cosmological abundance remains.

Fig. 2 shows numerical solutions to the Boltzmann equation which determines the WIMP abundance. The equilibrium (solid line) and actual (dashed lines) abundances per comoving volume are plotted as a function of $x \equiv m_\chi/T$ (which increases with increasing time). As the annihilation cross section is increased the WIMPs stay in equilibrium longer, and we are left with a smaller




Figure 2: Comoving number density of a WIMP in the early Universe. The dashed curves are the actual abundance, and the solid curve is the equilibrium abundance.

relic abundance.

An approximate solution to the Boltzmann equation yields the following estimate for the current cosmological abundance of the WIMP:

$$\Omega_{\chi} h^2 = \frac{m_{\chi} n_{\chi}}{\rho_c} \simeq \left(\frac{3 \times 10^{-27} \,\mathrm{cm}^3 \,\mathrm{sec}^{-1}}{\sigma_A v}\right),\tag{4}$$

where h is the Hubble constant in units of 100 km sec⁻¹ Mpc⁻¹. The result is to a first approximation independent of the WIMP mass and is fixed primarily by its annihilation cross section.

The WIMP velocities at freeze out are typically some appreciable fraction of the speed of light. Therefore, from equation (4), the WIMP will have a cosmological abundance of order unity today if the annihilation cross section is roughly 10^{-9} GeV⁻². Curiously, this is the order of magnitude one would expect from a typical electroweak cross section,

$$\sigma_{\text{weak}} \simeq \frac{\alpha^2}{m_{\text{weak}}^2},$$
 (5)

where $\alpha \simeq \mathcal{O}(0.01)$ and $m_{\text{weak}} \simeq \mathcal{O}(100\,\text{GeV})$. The value of the cross section in equation (4) needed to provide $\Omega_\chi \sim 1$ comes essentially from the age of the Universe. However, there is no a priori reason why this cross section should be of the same order of magnitude as the cross section one would expect for new particles with masses and interactions characteristic of the electroweak scale. In other words, why should the age of the Universe have anything to do with electroweak physics? This "coincidence" suggests that if a new, yet undiscovered, massive particle with electroweak interactions exists, then it should have a relic density of order unity and therefore provides a natural dark-matter candidate. This argument has been the driving force behind a vast effort to detect WIMPs in the halo.

The first WIMPs considered were massive Dirac or Majorana neutrinos with masses in the range of a few GeV to a few TeV. (Due to the Yukawa coupling which gives a neutrino its mass, the neutrino interactions become strong above a few TeV, and it no longer remains a suitable WIMP candidate 20 .) LEP ruled out neutrino masses below half the Z^0 mass. Furthermore, heavier Dirac neutrinos have been ruled out as the primary component of the Galactic halo by direct-detection experiments (described below) 21 , and heavier Majorana neutrinos have been ruled out by indirect-detection experiments 22 (also described below) over much of their mass range. Therefore, Dirac neutrinos cannot comprise the halo dark matter 23 ; Majorana neutrinos can, but only over a small range of fairly large masses. This was a major triumph for experimental particle astrophysicists: the first falsification of a dark-matter candidate.

However, theorists were not too disappointed: The stability of a fourth generation neutrino had to be postulated *ad hoc*—it was not guaranteed by some new symmetry. So although heavy neutrinos were plausible, they certainly were not very well-motivated from the perspective of particle theory.

A much more promising WIMP candidate comes from supersymmetry (SUSY)^{8,24}. SUSY was hypothesized in particle physics to cure the naturalness problem with fundamental Higgs bosons at the electroweak scale. Coupling-constant unification at the GUT scale seems to be improved with SUSY, and it seems to be an essential ingredient in theories which unify gravity with the other three fundamental forces.

As another consequence, the existence of a new symmetry, R-parity, in SUSY theories guarantees that the lightest supersymmetric particle (LSP) is stable. In the minimal supersymmetric extension of the standard model (MSSM), the LSP is usually the neutralino, a linear combination of the supersymmetric partners of the photon, Z^0 , and Higgs bosons. (Another possibility is the sneutrino, but these particles interact like neutrinos and have been ruled out over most of the available mass range 25 .) Given a SUSY model, the cross section for neutralino annihilation to lighter particles is straightforward, so one can obtain the cosmological mass density. The mass scale of supersymmetry must be of order the weak scale to cure the naturalness problem, and the neutralino will have only electroweak interactions. Therefore, it is to be expected that the cosmological neutralino abundance is of order unity. In fact, with detailed calculations, one finds that the neutralino abundance in a very broad class of supersymmetric extensions of the standard model is near unity and can therefore account for the dark matter in our halo 26 .

If neutralinos reside in the halo, there are several avenues toward detection ⁸. One of the most promising techniques currently being pursued involves searches for the $\mathcal{O}(10\,\mathrm{keV})$ recoils produced by elastic scattering of neutralinos from nuclei in low-background detectors ^{27,28}. The idea here is simple. A particle with mass $m_\chi \sim 100~\mathrm{GeV}$ and electroweak-scale interactions will have a cross section for elastic scattering from a nucleus which is $\sigma \sim 10^{-38}\,\mathrm{cm}^2$. If the local halo density is $\rho_0 \simeq 0.4~\mathrm{GeV}$ cm⁻³, and the particles move with velocities $v \sim 300~\mathrm{km}~\mathrm{sec}^{-1}$, then the rate for elastic scattering of these particles from, e.g., germanium which has a mass $m_N \sim 70~\mathrm{GeV}$, will be $R \sim \rho_0 \sigma v/m_\chi/m_N \sim 1~\mathrm{event}~\mathrm{kg}^{-1}~\mathrm{yr}^{-1}$. If a 100-GeV WIMP moving at $v/c \sim 10^{-3}$ elastically scatters with a nucleus of similar mass, it will impart a recoil energy up to 100 keV to the nucleus. Therefore, if we have 1 kg of germanium, we expect to see roughly one nucleus per year spontaneously recoil with an energy nearly 100 keV.

Of course, this is only a very rough calculation. To do the calculation

more precisely, one needs to use a proper neutralino-quark interaction, treat the QCD and nuclear physics which takes you from a neutralino-quark interaction to a neutralino-nucleus interaction, and integrate over the WIMP velocity distribution. Even if all of these physical effects are included properly, there is still a significant degree of uncertainty in the predicted event rates. Although supersymmetry provides perhaps the most promising dark-matter candidate (and solves numerous problems in particle physics), it really provides little detailed predictive power. In SUSY models, the standard-model particle spectrum is more than doubled, and we really have no idea what the masses of all these superpartners should be. There are also couplings, mixing angles, etc. Therefore, what theorists generally do is survey a large set of models with masses and couplings within a plausible range, and present results for relic abundances and direct- and indirect-detection rates, usually as scatter plots versus neutralino mass.

After taking into account all the relevant physical effects and surveying the plausible region of SUSY parameter space, one generally finds that the predicted event rates seem to fall for the most part between 10^{-4} to 10 events kg⁻¹ day⁻¹, ⁸ although again, there may be models with higher or lower rates. Current experimental sensitivities in germanium detectors are around 10 events kg⁻¹ day⁻¹. ²¹ To illustrate future prospects, consider the CDMS experiment ³⁴ which expects to soon have a kg germanium detector with a background rate of 1 event day⁻¹. After a one-year exposure, their sensitivity would therefore be $\mathcal{O}(0.1 \, \text{event kg}^{-1} \, \text{day}^{-1})$; this could be improved with better background rejection. Future detectors will achieve better sensitivities, and it should be kept in mind that numerous other target nuclei are being considered by other groups. However, it also seems clear that it will be quite a while until a good fraction of the available SUSY parameter space is probed.

Another strategy is observation of energetic neutrinos produced by annihilation of neutralinos in the Sun and/or Earth in converted proton-decay and astrophysical-neutrino detectors (such as MACRO, Kamiokande, IMB, AMANDA, and NESTOR) 29 . If, upon passing through the Sun, a WIMP scatters elastically from a nucleus therein to a velocity less than the escape velocity, it will be gravitationally bound in the Sun. This leads to a significant enhancement in the density of WIMPs in the center of the Sun—or by a similar mechanism, the Earth. These WIMPs will annihilate to, e.g., c, b, and/or t quarks, and/or gauge and Higgs bosons. Among the decay products of these particles will be energetic muon neutrinos which can escape from the center of the Sun and/or Earth and be detected in neutrino telescopes such as IMB, Kamiokande, MACRO, AMANDA, or NESTOR. The energies of these muons will be typically 1/3 to 1/2 the neutralino mass (e.g., 10 to 100s of

GeV) so they will be much more energetic—and therefore cannot be confused with—ordinary solar neutrinos. The signature of such a neutrino would be the Cerenkov radiation emitted by an upward muon produced by a charged-current interaction between the neutrino and a nucleus in the rock below the detector.

The annihilation rate of these WIMPs is equal to the rate for capture of these particles in the Sun. This can be estimated in order of magnitude by determining the rate at which halo WIMPs elastically scatter from nuclei in the Sun. The flux of neutrinos at the Earth depends also on the Earth-Sun distance, WIMP annihilation branching ratios, and the decay branching ratios of the annihilation products. The flux of upward muons depends on the flux of neutrinos and the cross section for production of muons, which depends on the square of the neutrino energy. As in the case of direct detection, the precise prediction involves numerous factors from particle and nuclear physics and astrophysics, and on the SUSY parameters. When all these factors are taken into account, predictions for the fluxes of such muons in SUSY models seem to fall for the most part between 10⁻⁶ and 1 event m⁻² vr⁻¹, 8 although the numbers may be a bit higher or lower in some models. Presently, IMB and Kamiokande constrain the flux of energetic neutrinos from the Sun to be less than about $0.02 \text{ m}^{-2} \text{ yr}^{-1}$, 22,30 and the Baksan limit is perhaps a factor-oftwo better ³¹. MACRO expects to be able to improve on this sensitivity by perhaps an order of magnitude. Future detectors may be able to improve even further. For example, AMANDA expects to have an area of roughly 10⁴ m², and a 10⁶-m² detector is being discussed. However, it should be kept in mind that without muon energy resolution, the sensitivity of these detectors will not approach the inverse exposure; it will be limited by the atmospheric-neutrino background. If a detector has good angular resolution, the signal-to-noise ratio can be improved, and even moreso with energy resolution, so sensitivities approaching the inverse exposure could be achieved ³². Furthermore, ideas for neutrino detectors with energy resolution are being discussed ³³, although at this point these appear likely to be in the somewhat-distant future.

With two promising avenues toward detection, it is natural to inquire which is most promising. Due to the abundance of undetermined SUSY parameters and the complicated dependence of event rates on these parameters, the answer to this question is not entirely straightforward. Generally, most theorists have just plugged in SUSY parameters into the machinery which produces detection rates and plotted results for direct and indirect detection. However, another approach is to compare, in a somewhat model-independent although approximate fashion, the rates for direct and indirect detection ^{8,35,36}. The underlying observation is that the rates for the two types of detection are both

controlled primarily by the WIMP-nucleon coupling. One must then note that WIMPs generally undergo one of two types of interaction with the nucleon: an axial-vector interaction in which the WIMP couples to the nuclear spin (which, for nuclei with nonzero angular momentum is roughly 1/2 and not the total angular momentum), and a scalar interaction in which the WIMP couples to the total mass of the nucleus. The direct-detection rate depends on the WIMP-nucleon interaction strength and on the WIMP mass. On the other hand, indirect-detection rates will have an additional dependence on the energy spectrum of neutrinos from WIMP annihilation. By surveying the various possible neutrino energy spectra, one finds that for a given neutralino mass and annihilation rate in the Sun, the largest upward-muon flux is roughly three times as large as the smallest ³⁶. So even if we assume the neutralino-nucleus interaction is purely scalar or purely axial-vector, there will still be a residual model-dependence of a factor of three when comparing direct- and indirect-detection rates.

For example, for scalar-coupled WIMPs, the event rate in a kg germanium detector will be equivalent to the event rate in a $(2-6)\times 10^6$ m² neutrino detector for 10-GeV WIMPs and $(3-5)\times 10^4$ m² for TeV WIMPs ³⁶. Therefore, the relative sensitivity of indirect detection when compared with the direct-detection sensitivity increases with mass. The bottom line of such an analysis seems to be that direct-detection experiments will be more sensitive to neutralinos with scalar interactions with nuclei, although very-large neutrino telescopes may achieve comparable sensitivities at larger WIMP masses. This should come as no surprise given the fact that direct-detection experiments rule out Dirac neutrinos ²¹, which have scalar-like interactions, far more effectively than do indirect-detection experiments ³⁶.

Generically, the sensitivity of indirect searches (relative to direct searches) should be better for WIMPs with axial-vector interactions, since the Sun is composed primarily of nuclei with spin (i.e., protons). However, a comparison of direct- and indirect-detection rates is a bit more difficult for axially-coupled WIMPs, since the nuclear-physics uncertainties in the neutralino-nuclear cross section are much greater, and the spin distribution of each target nucleus must be modeled. Still, in a careful analysis, Rich and Tao found that in 1994, the existing sensitivity of energetic-neutrino searches to axially-coupled WIMPs greatly exceeded the sensitivities of direct-detection experiments ³⁵.

 and leptons is due primarily to Z^0 exchange with a coupling proportional to $\cos 2\beta$, where $\tan \beta$ is the usual ratio of Higgs vacuum expectation values in supersymmetric models. Therefore, the usually messy cross sections one deals with in a general MSSM simplify for this candidate, and the cross sections needed for the cosmology of this Higgsino depend only on the two parameters m_χ and $\cos 2\beta$. Furthermore, since the neutralino-quark interaction is due only to Z^0 exchange, this Higgsino will have only axial-vector interactions with nuclei.

The Earth is composed primarily of spinless nuclei, so WIMPs with axialvector interactions will not be captured in the Earth, and we expect no neutrinos from WIMP annihilation therein. However, most of the mass in the Sun is composed of nuclei with spin (i.e., protons). The flux of upward muons induced by neutrinos from annihilation of these light Higgsinos would be $\Gamma_{\rm det} \simeq 2.7 \times 10^{-2} \, \rm m^{-2} \, yr^{-1} \, cos^2 \, 2\beta$. ³⁹ On the other hand, the rate for scattering from ⁷³Ge is $R \simeq 300 \, \cos^2 2\beta \, \rm kg^{-1} \, yr^{-1}$. ^{37,39} For illustration, in addition to their kg of natural germanium, the CDMS experiment also plans to run with 0.5 kg of (almost) purified ⁷³Ge. With a background event rate of roughly one event kg⁻¹ day⁻¹, after one year, the 3σ sensitivity of the experiment will be roughly 80 kg⁻¹ yr⁻¹. Comparing the predictions for direct and indirect detection of this axially-coupled WIMP, we see that the enriched-⁷³Ge sensitivity should improve on the current limit to the upward-muon flux (0.02) m⁻² yr⁻¹) roughly by a factor of 4. When we compare this with the forecasted factor-of-ten improvement expected in MACRO, it appears that the sensitivity of indirect-detection experiments looks more promising. Before drawing any conclusions, however, it should be noted that the sensitivity in detectors with other nuclei with spin may be significantly better. On the other hand, the sensitivity of neutrino searches increases relative to direct-detection experiments for larger WIMP masses. It therefore seems at this point that the two schemes will be competitive for detection of light axially-coupled WIMPs, but the neutrino telescopes may have an advantage in probing larger masses.

A common question is whether theoretical considerations favor a WIMP which has predominantly scalar interactions or whether they favor axial-vector couplings. Unfortunately, there is no simple answer. When detection of supersymmetric dark matter was initially considered, it seemed that the neutralino in most models would have predominantly axial-vector interactions. It was then noted that in some fraction of models where the neutralino was a mixture of Higgsino and gaugino, there could be some significant scalar coupling as well ⁴⁰. As it became evident that the top quark had to be quite heavy, it was realized that nondegenerate squark masses would give rise to scalar couplings in most models ⁴¹. However, there are still large regions of supersymmetric

parameter space where the neutralino has primarily axial-vector interactions, and in fact, the Kane-Wells Higgsino candidate has primarily axial-vector interactions. The bottom line is that theory cannot currently reliable say which type of interaction the WIMP is likely to have, so experiments should continue to try to target both.

5 Discussion

There has been no shortage of proposed solutions since the advent of the dark-matter problem nearly seventy years ago. Moreover, there has been an explosion in the number of exotic solutions proposed in recent decades. However, among the numerous proposals, only WIMPs and axions have really survived extended theoretical scrutiny for close to twenty years. Unlike other proposed dark-matter candidates, neither WIMPs nor axions really require any exotic mechanisms in the early Universe to guarantee a relic density of order unity—the only speculation is that they exist, and if they do, they have a relic density near unity. Furthermore, it is important to keep in mind that these particles were invented by particle theorists to solve problems with the standard model and only later was it realized that they were dark-matter candidates; they were not introduced to solve the dark-matter problem.

Axions and WIMPs have not only intrigued theorists; a large community of experimentalists have devoted themselves to finding these particles. However, it should also be emphasized that although very attractive, these are still speculative ideas. There is still no direct evidence in accelerator experiments or otherwise for the existence of axions or of supersymmetry. The dark matter could be composed of something completely different. However, as argued here, the evidence for the existence of nonbaryonic dark matter is indeed extremely compelling, and the two particles discussed here provide our most promising candidates. Although it provides an enormous experimental challenge, it is clear that discovery of particle dark matter would be truly revolutionary for both particle physics and cosmology.

Acknowledgments

This work was supported by the D.O.E. under contract DEFG02-92-ER 40699, NASA under NAG5-3091, and the Alfred P. Sloan Foundation.

References

1. M. Kamionkowski and D.N. Spergel, Astrophys. J. 432, 7 (1994).

- A.H. Guth, *Phys. Rev.* D 28, 347 (1981); A.D. Linde, *Phys. Lett.* B 108, 389 (1982); A. Albrecht and P.J. Steinhardt, *Phys. Rev. Lett.* 48, 1220 (1982).
- 3. K.G. Begeman, A.H. Broeils, and R.H. Sanders, *Mon. Not. R. Astr. Soc.* **249**, 523 (1991).
- K.A. Olive, G. Steigman, D.N. Schramm, T.P. Walker, and H. Kang, Astrophys. J. 376, 51 (1991).
- 5. S.D.M. White, C.S. Frenk, and M. Davis, Astrophys. J. 274, L1-5 (1983).
- 6. S. Tremaine and J.E. Gunn, Phys. Rev. Lett., 42, 407 (1979).
- 7. For reviews, see, e.g., M.S. Turner, *Phys. Rep.* **197**, 67 (1990); G.G. Raffelt, *Phys. Rep.* **198**, 1 (1990). An illuminating pedagogical discussion has also been given recently by P. Sikivie, *Phys. Today* **49**, 22 (1996).
- 8. G. Jungman, M. Kamionkowski, and K. Griest, *Phys. Rep.* **267**, 195 (1996).
- 9. E.W.Kolb and M.S. Turner, *The Early Universe* (Addison-Wesley, Redwood City, 1989).
- 10. G.G. Raffelt, Stars as Laboratories for Fundamental Physics (University of Chicago Press, Chicago, 1996).
- 11. R. Olling and D. Merrifield, in preparation.
- 12. M. Kamionkowski and A. Kinkhabwala, hep-ph/9710337.
- R.D. Peccei and H.R. Quinn, *Phys. Rev. Lett.* 38, 1440 (1977); F. Wilczek, *Phys. Rev. Lett.* 40, 279 (1978); S. Weinberg, *Phys. Rev. Lett.* 40, 223 (1978).
- 14. M.A. Bershady, M.T. Ressell, and M.S. Turner, *Phys. Rev. Lett.* **66**, 1398 (1991).
- M. Kamionkowski and J. March-Russell, *Phys. Lett.* B **282**, 137 (1992);
 R. Holman *et al.*, *Phys. Lett.* B **282**, 132 (1992);
 S. M. Barr and D. Seckel, *Phys. Rev.* D **46**, 539 (1992);
 B.A. Dobrescu, *Phys. Rev.* D **55**, 5826 (1997).
- R. Holman et al., Phys. Lett. B 282, 132 (1992); N. Turok, Phys. Rev. Lett. 76, 1015 (1996); R. Kallosh et al., Phys. Rev. D 52, 912 (1995);
 E.A. Dudas, Phys. Lett. B 325, 124 (1994); K.S. Babu and S.M. Barr, Phys. Lett. B 300, 367 (1993).
- 17. P. Sikivie, *Phys. Rev. Lett.* **51**, 1415 (1983).
- 18. K. van Bibber *et al.*, *Int. J. Mod. Phys. Suppl.* D **3**, 33 (1994); C. Hagmann *et al.*, *Nucl. Phys. (Proc. Suppl.)* B **51**, 209 (1996).
- I. Ogawa, S. Matsuki, and K. Yamamoto, *Phys. Rev.* D **53**, 1740 (1996);
 S. Matsuki, I. Ogawa, and K. Yamamoto, *Phys. Lett.* B **336**, 573 (1994).
- 20. K. Griest and M. Kamionkowski, *Phys. Rev. Lett.* **64**, 615 (1990).
- 21. M. Beck, Nucl. Phys. (Proc. Suppl.) B 35, 150 (1994); M. Beck et al.,

- Phys. Lett. B **336**, 141 (1994); S.P. Ahlen et al., Phys. Lett. B **195**, 603 (1987); D.O. Caldwell et al., Phys. Rev. Lett. **61**, 510 (1988).
- 22. M. Mori et al. (Kamiokande Collaboration), Phys. Lett. B 289, 463 (1992); M. Mori et al. (Kamiokande Collaboration), Phys. Rev. D 48, 5505 (1993).
- 23. K. Griest and J. Silk, *Nature* **343**, 26 (1990); L.M. Krauss, *Phys. Rev. Lett.* **64**, 999 (1990).
- 24. H.E. Haber and G.L. Kane, *Phys. Rep.* **117**, 75 (1985).
- 25. T. Falk, K.A. Olive, and M. Srednicki, Phys. Lett. B 339, 248 (1994).
- 26. J. Ellis *et al.*, *Nucl. Phys. B* **238**, 453 (1984); K. Griest, M. Kamionkowski, and M.S. Turner, *Phys. Rev.* D **41**, 3565 (1990).
- M.W. Goodman and E. Witten, *Phys. Rev.* D **31**, 3059 (1986); I. Wasserman, *Phys. Rev.* D **33**, 2071 (1986); K. Freese, J. Frieman, and A. Gould, *Phys. Rev.* D **37**, 3388 (1988); A. Drukier, K. Freese, and D.N. Spergel, *Phys. Rev.* D **33**, 3495 (1986).
- J. Low Temp. Phys. 93 (1993); P.F. Smith and J.D. Lewin, Phys. Rep. 187, 203 (1990).
- J. Silk, K.A. Olive, and M. Srednicki, Phys. Rev. Lett. 55, 257 (1985); K. Freese, Phys. Lett. B 167, 295 (1986); L.M. Krauss, K. Freese, D.N. Spergel, and W.H. Press, Astrophys. J. 299, 1001 (1985); L.M. Krauss, M. Srednicki, and F. Wilczek, Phys. Rev. D 33, 2079 (1986); T. Gaisser, G. Steigman, and S. Tilav, Phys. Rev. D 34, 2206 (1986); M. Kamionkowski, Phys. Rev. D 44, 3021 (1991); S. Ritz and D. Seckel, Nucl. Phys. B 304, 877 (1988); F. Halzen, M. Kamionkowski, and T. Stelzer, Phys. Rev. D 45, 4439 (1992).
- 30. J.M. LoSecco et al. (IMB Collaboration), Phys. Lett. B 188, 388 (1987).
- M.M. Boliev et al., Bull. Acad. Sci. USSR, Phys. Ser. 55, 126 (1991)
 [Izv. Akad. Nauk. SSSR, Fiz. 55, 748 (1991)]; M.M. Boliev et al., in TAUP 95, proceedings of the Workshop, Toledo, Spain, September 17–21, 1995, ed. A. Morales, J. Morales and J.A. Villar, [Nucl. Phys. (Proc. Suppl.) B 48, 83 (1996)] (North-Holland, Amsterdam, 1996).
- 32. L. Bergstrom, J. Edsjö, and M. Kamionkowski, *Astropart. Phys.* **7**, 147 (1997).
- 33. Y. Ho (HANUL Collaboration), to appear in *Very High Energy Phenomena in the Universe*, proceedings of the XXXIInd Moriond Conference, ed. J. Tran Thanh Van (Editions Frontières, Gif-sur-Yvette, 1997).
- P.D. Barnes et al., J. Low Temp. Phys. 93, 79 (1993); T. Shutt et al., Phys. Rev. Lett. 61, 3425 (1992); ibid., 3531 (1992).
- 35. J. Rich and C. Tao, DAPNIA/SPP 95-01.
- 36. M. Kamionkowski, K. Griest, G. Jungman, and B. Sadoulet, Phys. Rev.

- Lett. 74, 5174 (1995).
- 37. G.L. Kane and J.D. Wells, *Phys. Rev. Lett.* **76**, 4458 (1996); S. Ambrosanio *et al.*, *Phys. Rev. Lett.* **76**, 3498 (1996).
- 38. S. Park, "Search for New Phenomena at CDF," 10th Topical Workshop on Proton-Antiproton Collider Physics, edited by R. Raja and J. Yoh (AIP Press, 1996).
- 39. K. Freese and M. Kamionkowski, Phys. Rev. D 55, 1771 (1997).
- K. Griest, Phys. Rev. D 38, 2357 (1988); FERMILAB-Pub-89/139-A
 (E).
- 41. M. Drees and M.M. Nojiri, *Phys. Rev.* D **47**, 4226 (1993); *ibid.* **48**, 3483 (1993).