

DYNAMICAL COSMOLOGICAL CONSTANT AND RELATIONS AMONG PSEUDO GOLDSTONE BOSONS

Juan Estrada Vigil¹ and Luis Masperi²

*Centro Atómico Bariloche and Instituto Balseiro, (Comisión Nacional de Energía
Atómica and Universidad Nacional de Cuyo) 8400 S.C. de Bariloche, Argentina*

Abstract

The present apparent cosmological constant is interpreted as the potential of the lightest pseudo Goldstone boson. Some numerical relations among cosmological parameters and particle masses are shown to arise from the mixture of this very light particle, whose interaction is of the gravitational order, with other pseudo Goldstone bosons like the pion and the axion which feel the strong interaction.

Evidence from clusters and superclusters of galaxies together with baryon abundance indicates that matter accounts for a fraction $\Omega_m \sim 0.3$ of the critical density of the universe[1]. For the formation of large scale structure, perhaps the age of the universe, the luminosity of galaxies and even the reionization of intergalactic medium[2] it is useful to add a contribution of the cosmological constant $\Omega_\Lambda \sim 0.7$ with the formal advantage of completing the critical density required by the standard version of inflation. However, for a correct normalization with the COBE anisotropies it is more convenient[3] that the apparent cosmological constant is caused by a field which is becoming dynamical near the present time due to the extremely light mass of the associated particle.

Cosmological parameters and typical particle masses are related by the Friedmann equation, that does not involve \hbar , which determines the expansion of the universe; and also by an intriguing relation[4] where \hbar appears, whose validity at any time would imply the variation of fundamental constants[5]. This latter relation has been derived recently[6] to calculate the Planck constant from classical considerations.

We show that with the dynamical cosmological constant interpretation, the Friedmann equation takes the form of the mass of a pseudo Goldstone boson produced by an explicit symmetry breaking term at the scale of a possible neutrino mass, and the second relation is consistent with the hypothesis that this very light pseudo Goldstone boson feels an interaction of the gravitational order. The fact that the latter relation includes \hbar is due, in our interpretation, to the present situation of balance between the microphysics term of the lightest pseudo Goldstone boson and the gravitational one corresponding to universe expansion.

¹E-mail: estradav@ib.cnea.edu.ar

²E-mail: masperi@cab.cnea.edu.ar

The Hubble constant H is related to the critical density ρ_c and the Newton constant G_N by the Friedmann equation

$$H^2 = \frac{8\pi}{3} G_N \rho_c \quad , \quad (1)$$

which comes from the Einstein equation and therefore has no quantum content. The variation of ρ_c with the universe scale determines the expansion rate through H .

On the other hand it has been noted that the following numerical relation holds

$$H_0 = G_N m_\pi^3 \quad (2)$$

between the present value of the Hubble constant H_0 and the pion mass. This equation is written in natural units $\hbar = c = 1$ but obviously involves quantum physics by making explicit a factor c/\hbar^2 on the right hand side. If one wishes that Eq.(2) holds for any time, since it contains the Hubble constant as the only cosmological parameter some fundamental constant should vary with time, e.g. the Newton one. A relation analogous to Eq.(2) has been obtained[6] calculating the Planck constant through the fluctuations of classical particles due to the background gravitational field.

If we adopt the dynamical cosmological constant approach, an extremely light particle should have a mass

$$m_\varphi \sim H_0 \quad , \quad (3)$$

so that for previous times the Hubble constant would prevail and the contribution to energy of the field φ would appear only through a potential term. φ should be therefore the lightest conceivable matter particle in our universe.

Applying Eq.(1) to present time we would have

$$m_\varphi^2 \sim \frac{1}{m_{Pl}^2} M^4 \quad (4)$$

where, due to the value of critical density, $M \sim 10^{-2} - 10^{-3} eV$. Eq.(4) suggests that φ is the Goldstone boson coming from a spontaneous symmetry breakdown at the Planck mass scale and which acquires a small mass through an explicit breaking of this symmetry at scale M . Since the numerical value of M may correspond to the mass of neutrinos which solves the solar problem, it has been thought[7] that the symmetry spontaneously broken at scale $f_\varphi \sim m_{Pl}$ is a chiral global one involving neutrinos.

The explicit breaking of this symmetry would produce an effective potential

$$V(\varphi) = M^4 \left(1 - \cos \frac{\varphi}{f_\varphi} \right) \quad , \quad (5)$$

due to the phase nature of φ , consistent with Eq.(4).

Regarding Eq.(2), it should be interpreted as

$$m_\varphi \sim \frac{1}{m_{Pl}^2} m_\pi^3 \quad , \quad (6)$$

which is satisfied for $m_\varphi \sim 10^{-32} eV$ that corresponds to Eq.(3). Now there is no time dependence and no need to think in changes of fundamental constants along the universe evolution, for which bounds exist[8].

The explanation of Eq.(6), that must be obviously consistent with Eq.(4), may be analogous to the one which suggests the mass of the axion. This well studied[9] light particle is the Goldstone boson of the spontaneous breaking of the Peccei-Quinn chiral symmetry of a quark which consequently becomes very heavy. Since the current corresponding to this symmetry has the QCD anomaly

$$\partial_\mu j_{PQ}^\mu = \frac{g_S^2}{16\pi^2} G_{\mu\nu} \tilde{G}^{\mu\nu} \quad (7)$$

in terms of the gluon field $G_{\mu\nu}$, one has to subtract, apart from a factor, the axial current of light quarks to define a current which is conserved in the chiral limit

$$j_a^\mu = j_{PQ}^\mu - (\bar{u}\gamma^\mu\gamma_5 u + \bar{d}\gamma^\mu\gamma_5 d + \bar{s}\gamma^\mu\gamma_5 s) \quad . \quad (8)$$

Therefore, this current will have a divergence structure analogous to that of the isospin axial current related to the symmetry whose spontaneous breaking at scale $f_\pi \sim 93 MeV$ gives the pion as Goldstone boson. Due to the explicit breaking of symmetry caused by the mass of light quarks, current algebra methods [10] allow to calculate the mass of pion and axion in terms of a typical quark q

$$m_\pi^2 \sim \frac{m_q}{f_\pi^2} \langle 0 | \bar{q}q | 0 \rangle \quad , \quad m_a^2 \sim \frac{m_q}{f_a^2} \langle 0 | \bar{q}q | 0 \rangle \quad (9)$$

apart from factors. Eq.(9) gives the relation

$$m_a \sim \frac{f_a}{f_\pi} m_\pi \quad . \quad (10)$$

This mass is equivalent to the one coming from the effective potential $V(a)$ originated by QCD which is of the type of Eq.(5) with a scale for the explicit breaking of Peccei-Quinn symmetry Λ_{QCD} replacing M . Eq.(10) gives the possible experimental window for axion mass $m_a \sim 10^{-5} eV$ if $f_a \sim 10^{12} GeV$.

For the Frieman et al. chiral symmetry, we may imagine that it applies to a neutrino which becomes very heavy through spontaneous breaking at the scale $f_\varphi \sim m_{Pl}$ and that its current j_F^μ has an anomaly due to fields corresponding to interactions of gravitational order. We again define the current associated to particle φ subtracting, apart from a factor, the axial current for a typical light neutrino to compensate this anomaly

$$j_\varphi^\mu = j_F^\mu - \bar{\nu}\gamma^\mu\gamma_5\nu \quad (11)$$




Figure 1: Mixture of pseudo Goldstone bosons

(a) axion with pion

(b) φ with pion

which will be conserved in the limit of massless neutrino. Again current algebra methods will give

$$m_\varphi^2 \sim \frac{m_\nu}{f_\varphi^2} \langle 0 | \bar{\nu} \nu | 0 \rangle \quad , \quad (12)$$

and the comparison with Eq.(9) will require

$$\frac{m_\nu \langle 0 | \bar{\nu} \nu | 0 \rangle}{m_q \langle 0 | \bar{q} q | 0 \rangle} \sim \left(\frac{m_\pi}{m_{Pl}} \right)^2 \quad (13)$$

to satisfy the relation (6). It is interesting to see that Eq.(13) holds if, together with the QCD values $m_q \sim 0.1 GeV$ and $\langle 0 | \bar{q} q | 0 \rangle \sim 10^{-3} GeV^3$, the estimation for the light neutrinos scale $m_\nu \sim 10^{-2} eV$ and $\langle 0 | \bar{\nu} \nu | 0 \rangle \sim 10^{-6} eV^3$ is introduced.

The previous result may be understood also in a pictorial way. The axion has an effective coupling with two gluons through a triangle of heavy quarks, of the order of the inverse of the scale f_a , which corresponds to the divergence of the Peccei-Quinn current. According to Fig.1(a), since the pion is coupled to the axial current through f_π , the axion lives a fraction of its time as a pion getting a mass as given by Eq.(10).

For the very light particle φ , due to the fact that its effective coupling is caused by the coupling with heavy neutrinos resulting from the spontaneous breakdown of chiral symmetry at scale f_φ , the interaction may be thought as of

gravitational order and the amplitude for the mixture with the pion according to Fig.1(b) needs an additional conversion factor. It seems reasonable that this factor must be the ratio of scales for strong and gravitational interactions i.e. of the order of the ratio of the Planck and Fermi lengths, or m_π/m_{Pl} . Therefore the mass of the φ particle will be

$$m_\varphi \sim \frac{1}{f_\varphi} \frac{m_\pi}{m_{Pl}} f_\pi m_\pi \simeq \frac{1}{m_{Pl}^2} m_\pi^3 \quad , \quad (14)$$

which is again Eq.(6).

The consistency of Eqs.(4) and (6) requires that the scale for the explicit breaking of the symmetry of which φ is the Goldstone boson, that should be of the order of the neutrino mass, is

$$M \sim \left(\frac{m_\pi}{m_{Pl}} \right)^{1/2} m_\pi \quad . \quad (15)$$

It is interesting to note that the derivation of Eq.(2) done in Ref.[6] starting from Friedmann Eq.(1) together with the expression of the energy contained in the present Hubble radius as given by N nonrelativistic particles of mass of the order of m_π , is expressed in our dynamical cosmological constant approach as the condition

$$m_\pi \sim \sqrt{N} m_\varphi \quad . \quad (16)$$

One might envisage the connection between classical and quantum mechanics of Ref.[6] by the fact that Eq.(3) indicates that the Compton length of the φ particle equals the radius of the observable universe. Only from this moment, i.e. around the present time, quantum mechanics may be applied to all particles.

Obviously the need of the variation with time of fundamental constants assuming the general validity of Eq.(2) is replaced in our scheme by the requirement that the field φ becomes dynamical at present which is a form of anthropic principle [11] even though the true cosmological constant may be zero because of some symmetry, perhaps supergravity[12]. Since our relations are valid only as an order of magnitude, φ might have become dynamical for a redshift z of some units and not necessarily for $z = 0$ and it is not relevant the precise value of H_0 .

Acknowledgments

We thank heartily Francesco Calogero for pointing us his derivation of Planck constant from gravitational effects and for stimulating discussions, and Carlos García Canal for extremely useful comments.

References

- [1] R. Carlberg et al., *Astrophys. J.* **462**, 32 (1996) and astro-ph 9706115;
N. Bahcall, astro-ph 9611080;
N. Bahcall, X. Fan and R. Cen, astro-ph 9706018;
L.M. Krauss, astro-ph 9706227.
- [2] J.P. Ostriker and P.J. Steinhardt, *Nature* **377**, 600 (1995);
L.M. Krauss and M.S. Turner, *Gen. Rel. Grav.* **27**, 1137 (1995);
T. Totani, Y. Yoshii and K. Sato, astro-ph 9705014;
L. Masperi and S. Savaglio, *Astronomy and Astrophysics* **321**, 1 (1997).
- [3] K. Coble, S. Dodelson and J.A. Frieman, *Phys. Rev.* **D 55**, 1851 (1997).
- [4] S. Weinberg, *Gravitation and Cosmology: Principles and applications of general theory of relativity* (Wiley, New York, 1972) p. 619.
- [5] P.A.M. Dirac, *Nature* **139**, 323 (1937), and *Proc. Roy. Soc.* **A 165**, 199 (1938).
- [6] F. Calogero, *Phys. Lett.* **A 238**, 335 (1997).
- [7] J.A. Frieman, C.T. Hill and R. Watkins, *Phys.Rev.* **D 46**, 1226 (1992);
J.A. Frieman, C.T. Hill, A. Stebbins and I. Waga, *Phys. Rev. Lett.* **75**, 2077 (1995).
- [8] P. Sisterna and H. Vucetich, Time variation of fundamental constants; bounds from local data, *Proc. of Conference on Mach's principle*, Ed. J. Babour and H. Pfister (Tübingen, 1993);
M.J. Drinkwater, J.K. Webb, J.D. Barrow and V.V. Flambaum, astro-ph 9709227.
- [9] R.D. Peccei, hep-ph 9606475;
M. Srednicki, *Nucl. Phys* **B 260**, 689 (1995);
W.A. Bardeen and S.-H.H. Tye, *Phys. Lett* **B 74** , 229 (1978).
- [10] R. Dashen, *Phys. Rev.* **183**, 1245 (1969).
- [11] S. Weinberg, *Phys. Rev. Lett.* **59**, 2067 (1987), and astro-ph 9610044;
H. Martel, P.R. Shapiro and S. Weinberg, astro-ph 9701099.
- [12] E. Witten, *Phys. Lett* **10**, 1247 (1995).