## $\theta$  VACUUM IN A RANDOM MATRIX MODEL

ROMUALD A. JANIK $^{1,2}$ , MACIEJ A. NOWAK $^{2,3}$ , GÁBOR PAPP $^4$ and ISMAIL ZAHED<sup>5</sup>

<sup>1</sup> Service de Physique Théorique, CEA Saclay, F-91191 Gif-Sur-Yvette, France. <sup>2</sup> Department of Physics, Jagellonian University, 30-059 Krakow, Poland.

 $4$  CNR Department of Physics, KSU, Kent, Ohio 44242, USA  $\mathcal{B}$ 

HAS Research Group for Theoretical Physics, Eötvös University, Budapest,

<sup>5</sup>Department of Physics and Astronomy, SUNY, Stony Brook, New York 11794.

Inspired by recent lattice calculations, we model certain aspects of the  $\theta$ -vacuum using a matrix model with gaussian weights. The vacuum energy exhibits a cusp at  $\theta < \pi$  that is sensitive to both the accuracy of the numerical analysis and the maximum density of winding modes present in a finite volume.

Recent lattice study of the  $\mathbb{CP}^3$  model on the lattice<sup>[1](#page-2-0)</sup> have revealed an intriguing result: the energy as a function of the CP breaking  $\theta$  angle levels off at large  $\theta$ . The result is very interesting, since in that model the string tension for small charges is related to the  $\theta$ -derivative of the vacuum energy. A plateau indicates the loss of confinement. The result was reexamined in  $2$  where the authors pointed out to some subtleties regarding the technical methods applied.

In this talk we will address these issues using a random matrix model with the partition function

$$
Z(\theta, N_f) = \left\langle \prod_{j=1}^{N_f} \det \left( \frac{im_j e^{i\theta/N_f}}{W^{\dagger}} \frac{W}{im_j e^{-i\theta/N_f}} \right) \right\rangle. \tag{1}
$$

where the averaging is carried over the matrices  $W, W^{\dagger}$  using the weight  $\exp(-n/2\text{Tr}W^{\dagger}W)$  and over the matrix size n and  $\sigma$ , with weights  $\exp(-\chi^2/2\chi_* V)$  and  $\exp(-\sigma^2/2\sigma_* V)$ , respectively. Here W is a complex asymmetric  $n_+ \times n_-$  matrix,  $n = n_+ + n_-$ , and  $\sigma \pm \chi = 2n_{\pm} - \langle n \rangle$ . The mean number of zero modes  $\langle n \rangle$  is either fixed from the outside or evaluated using the gaussian measure. Here and for simplicity we use the quenched measure without the fermion determinant to fix  $\langle n \rangle$ . Throughout, we set the value of the quark condensate to  $\Sigma = 1$  in the chiral limit.

Eq.  $(1)$  is borrowed from the effective instanton vacuum analysis  $3$  where  $n_{+}$  counts the number of right-handed zero modes, and  $n_{-}$  the number of lefthanded zero modes. The number of exact topological zero modes is commen-

*tr99-1: submitted to* World Scientific *on March 24, 2017* 1

<sup>3</sup> GSI, Planckstr. 1, D-64291 Darmstadt, Germany

surate with the net winding number carried by the instantons and antiinstan-tons. By analogy with <sup>[3](#page-2-0)</sup>,  $\chi_*$  and  $\sigma_*$  will refer to the unquenched topological susceptibility and particle compressibility, respectively, with  $\sigma_*^2 = 12\mathbf{n}_*/11N_c$ and  $\mathbf{n}_{*} = \langle n \rangle/V$  the mean density of zero modes. If the compressibility  $\sigma_{*}$  is assumed small in units of  $\Sigma = 1$ , then typically  $n \sim \langle n \rangle$ .

**quenched case:** In the case of the quenched partition function  $(N_f = 0)$ one probes the nature of the gaussian measure. We start discussing the case where  $\langle n \rangle = \infty$  with no restriction on the value of n, and hence no restriction on the value of  $\chi$ . Then, we discuss the case where  $\langle n \rangle$  is large but finite, so that  $|\chi| \leq n$  with typically  $n \sim \langle n \rangle$  for a peaked distribution in n.

When the sum is unrestricted and infinite, using Poisson resummation formula we have

$$
Z_Q(\theta) = \sum_{k=-\infty}^{+\infty} e^{-\frac{1}{2}V\chi_* (\theta - 2\pi k)^2} = \theta_3(\theta/2, e^{-\tau}).
$$
 (2)

with  $\theta$  being the third elliptic function and  $\tau = 1/(2V\chi_*)$ . The result is manifestly  $2\pi$  periodic. The vacuum energy,  $F_Q(\theta) = -\ln Z_Q(\theta)/V$  as  $V \to \infty$ is simply  $F_Q(\theta) = \min_{\frac{1}{2}} \chi_* (\theta + \text{mod } 2\pi)^2$  in agreement with the saddle-point approximation. This result is in agreement with the result using large  $N_c$ arguments  $4$ , and recent duality arguments  $5$ . We observe that the cusp at  $\theta = \pi \pmod{2\pi}$  sets in for  $V = \infty$ .

In the matrix model being considered the sum over  $\chi$  is restricted to  $|\chi| < N$ , with  $N = \max n$ . We denote by  $\mathbf{n} = N/V$  the maximum density of winding modes. While in general  $n \neq n<sub>*</sub>$ , for a peaked distribution in n (small compressibility  $\sigma_*$ ) we expect  $\mathbf{n} \sim \mathbf{n}_*$ . This will be assumed throughout unless indicated otherwise. Hence

$$
Z_Q(\theta) = \sum_{\chi = -(N-1)}^{N-1} e^{i\theta \chi} e^{-\chi^2/2V\chi_*}
$$
 (3)

Approximating the sum in (3) by an integral and evaluating it by saddle point we obtain  $Z_Q \sim e^{-V_{\chi_*} \theta^2/2}$  apparently in agreement with the infinite sum. However, the Euler-MacLaurin summation formula shows that deviation from this result in the case of a finite sum is expected for  $\theta > \theta_c \sim n/\chi_*$ <sup>[6](#page-2-0)</sup>. Indeed, this is confirmed by detailed numerical calculations of the vacuum energy as shown in Fig [1](#page-2-0) (left). For  $N = 250$  and  $\chi_* = 1$ , the double precision (16 digit) numerics (circles) breaks away from the saddle point approximation (solid line) at  $\theta/\pi \sim 0.2$  for both  $n = 1$  and  $n = 4$ , while the high-precision (64 digits) calculations (dashed line) agree with the saddle point result at

*tr99-1: submitted to* **World Scientific** *on March 24, 2017* 2

<span id="page-2-0"></span> $n = 4$  but break away at  $\theta/\pi \sim 0.3 \approx 1/\pi \chi_*$  at  $n = 1$ . On the right we show the numerical result for the  $\mathbb{CP}^3$  model <sup>1</sup> showing a similar behavior.



Figure 1. Free energy in a quenched model (left) and in the  $CP<sup>3</sup>$  model (right). See text.

unquenched case: Analytical and numerical studies of the unquenched matrix model for  $N_f = 1$  and  $N_f = 2^{-6}$  lead to results similar to the ones observed in the quenched case. The mean-field result breaks down for  $\theta$   $>$  $\theta_c \sim \mathbf{n}/\chi_{top}$ . For small quark masses  $\chi_{top} \sim m^{-7}$ , the breakdown is observed only for ensembles with very small maximal winding number.

In this talk we have shown that the vacuum energy of a chiral matrix model developes a cusp at  $\theta < \pi$  that is sensitive to the numerical accuracy. The cusp persits at high accuracy if the maximum winding number considered is low, in disagreement with mean-field results. Similar observations in current lattice simulations should therefore be interpreted with care.

## Acknowledgments

This work was supported by the US DOE grants DE-FG-88ER40388, by the Polish Government Project (KBN) grant 2P03B00814 and by the Hungarian grant OTKA-F026622.

## References

- 1. G. Schierholz, *Nucl. Phys. Proc. Suppl.* A37, 1994 (203).
- 2. J. C. Plefka and S. Samuel, *Phys. Rev.* D 56, 1977 (44).
- 3. M. Nowak, J. Verbaarschot and I. Zahed, *Phys. Lett.* B 228, 1989 (251).
- 4. E. Witten, *Annals of Phys.* 128, 1980 (363).
- 5. E. Witten, *Phys. Rev. Lett.* 81, 1998 (2862).
- 6. R.A. Janik, M.A. Nowak, G. Papp and I. Zahed, [hep-ph/9901390.](http://arxiv.org/abs/hep-ph/9901390)
- 7. R.A. Janik, M.A. Nowak, G. Papp and I. Zahed, *Nucl. Phys.* B 498, 1997 (313).

*tr99-1: submitted to* World Scientific *on March 24, 2017* 3