Single mass scale diagrams: construction of a basis for the ε -expansion.

J. Fleischer $^{a\ 1},\$ M. Yu. Kalmykov $^{a,b\ 2}$

- ^a Fakultät für Physik, Universität Bielefeld, D-33615 Bielefeld, Germany
- ^b BLTP, JINR, 141980, Dubna (Moscow Region), Russia

Abstract

Exploring the idea of Broadhurst on the "sixth root of unity" we present an ansatz for construction of a basis of transcendental numbers for the ε -expansion of single mass scale diagrams with two particle massive cut. As example, several new two- and three-loop master integrals are calculated.

Keywords: Feynman diagram.

 $PACS\ number(s)$: 12.38.Bx

¹E-mail: fleischer@physik.uni-bielefeld.de

² E-mail: misha@physik.uni-bielefeld.de

1 Introduction

One of the interesting cases of diagrams occurring in QCD and the Standard Model is of the propagator type with one mass and the external momentum on the mass shell (the bubble (vacuum) type diagrams also belong to this class). Only a limited number of analytical results is available in this case: the two- and three-loop diagrams occurring in QCD are given in Refs.[1, 2]; the analytical result for the two-loop bubble integral are presented up to the finite part in [1, 3, 4] and the $O(\varepsilon)$ term in [5]; the finite part of three-loop bubble master integrals is calculated in [6]; the results for all other two-loop on-shell propagator type integrals with one mass are collected in [7]. For some diagrams the results are available in terms of hypergeometric functions [8, 9]. However, the ε -expansion of hypergeometric functions is an independent and sometimes very complicated task.

At the same time a lot of different methods for numerical calculations of Feynman diagrams are available now [10, 11, 12, 13, 14]. From another point of view the development of numerical algorithms, in particular PSLQ [15], allows to determine with high confidence, whether or not particular numerical results for master-integrals can be expressed as a rational linear combination of several given transcendental constants. However, the problem arises of how to define the full set of the transcendental structure (the "basis") occurring in given diagrams. The structure of massless multiloop integrals is well understood at the present time. In contains mainly the ζ -function [16] and Euler sums [17]. Recently the connection between knot theory and the transcendental structure of Feynman diagrams was established [18]. In more complicated case, when dimensionless parameters exist, harmonic sums [19, 20] appeared. For studying the massive case, the differential equation method [21] and geometrical approach [22, 23] can be useful.

The aim of our investigation is to find the transcendental structure of single mass scale propagator type diagrams. In this paper we restrict ourselves to the consideration of diagrams with two particle massive cuts only. All our results are obtained empirically by carefully compiling and examining a huge data base on high precision (about 200 decimals) numerical calculation [15].

2 Basis

Our aim is to construct a basis (the full set of transcendental numbers) for single mass scale propagator type diagrams with two-particle massive cuts. The simplest diagrams of this type are shown in Fig.1.

An algebra for such a basis can be constructed as follows: the product of two basis elements having weights J_1 and J_2 , respectively, must belong to the basis of weight $J_1 + J_2$. The weight of basis elements can be connected, e.g., with powers of ε within the Laurent expansion of dimensionally regularized [24] Feynman diagrams.

In constructing the basis of weight J we will use Broadhurst's observation [6], that the sixth root of unity plays an important role in the calculation of the diagrams (see also [25]). Accordingly we conjecture, that the basis of weight J contains functions obtained by splitting of polylogarithms $\operatorname{Li}_J(z)$, $\operatorname{Li}_J(1-z)$ into real and imaginary parts, where z=




Figure 1: Examples of the simplest single mass scale integrals with two-particle massive cut. Bold and thin lines correspond to massive and massless propagators, respectively.

 $exp\left(i\frac{\pi}{3}\right), exp\left(i\frac{2\pi}{3}\right)$. The general case, $z = exp\left(\pm i\frac{\pi}{3} + i\pi m\right)$ can be reduced to these two values [26]. E.g., with $\operatorname{Li}_{2n}\left(e^{i\theta}\right) = \operatorname{Gl}_{2n}\left(\theta\right) + i\operatorname{Cl}_{2n}\left(\theta\right)$, the basis of weight 2n must contain $\operatorname{Gl}_{2n}\left(\frac{\pi}{3}\right), \operatorname{Gl}_{2n}\left(\frac{2\pi}{3}\right), \operatorname{Cl}_{2n}\left(\frac{\pi}{3}\right)$ and $\operatorname{Cl}_{2n}\left(\frac{2\pi}{3}\right)$. Not all of these functions are independent: $\operatorname{Gl}_{2n}\left(\omega\pi\right) \sim \zeta_{2n}$, where ω is an arbitrary rational number; $(1+2^{1-2n})\operatorname{Cl}_{2n}\left(\frac{2\pi}{3}\right) = \operatorname{Cl}_{2n}\left(\frac{\pi}{3}\right)$. The basis we are going to construct is supposed to hold for any number of loops and

The basis we are going to construct is supposed to hold for any number of loops and arbitrary order of the ε -expansion. Consequently we connect the lowest weight basis elements with Li₁(z) and Li₁(1-z), which results in π and ln 3. Comparing this with the one-loop diagram of Fig.1 (ONS11), the expression given in [23] allows us to find from its finite part the transcendental normalization factor $\frac{1}{\sqrt{3}}$, so that our lowest basis is given by $\frac{\pi}{\sqrt{3}}$ and ln 3. In general the normalization factor $\frac{1}{\sqrt{3}}$ is introduced according to our experience.

Using the above conjectures, we obtain the basis of higher weights: for weight **2**, e.g., we have $\frac{\pi}{\sqrt{3}} \ln 3$, ζ_2 , $\frac{1}{\sqrt{3}} \operatorname{Cl}_2\left(\frac{\pi}{3}\right)$ and $\ln^2 3$. The results up to weight **5** are presented in Table I. Some transcendental structures like, e.g., Ls₃ $\left(\frac{\pi}{3}\right)$ do not appear in the table since they could be eliminated by means of relations given in [26]. Including weight **3** the relations found in [26] were sufficient. For weight **4** we needed the new relation

$$\operatorname{Cl}_4\left(\frac{\pi}{3}\right) = \frac{2}{9}\operatorname{Ls}_4\left(\frac{\pi}{3}\right) - \frac{1}{9}\pi\zeta(3),\tag{1}$$

and for weight 5 we used the further relations

$$Ls_5^{(1)}\left(\pm\frac{\pi}{3}\right) = \frac{1}{3}\pi Ls_4\left(\frac{\pi}{3}\right) - \frac{19}{4}\zeta_5 - \frac{1}{2}\zeta_2\zeta_3,\tag{2}$$

$$Ls_5^{(2)} \left(\pm \frac{\pi}{3} \right) = \pm \left[\frac{2}{3} Ls_5 \left(\frac{\pi}{3} \right) + \frac{253}{54} \pi \zeta_4 \right], \tag{3}$$

in order to restrict the basis as much as possible.

Table I					
1	2	3	4	5^a	5^{b}
$\frac{\pi}{\sqrt{3}}$	$\frac{\pi}{\sqrt{3}} \ln 3$	$\frac{\pi}{\sqrt{3}}\ln^2 3$	$\frac{\pi}{\sqrt{3}} \ln^3 3$	$\frac{\pi}{\sqrt{3}}\ln^4 3$	$\frac{\pi}{\sqrt{3}}\zeta_4$
$\ln 3$	ζ_2	$\zeta_2 \ln 3$	$\zeta_2 \ln^2 3$	$\zeta_2 \ln^3 3$	$\pi \zeta_2 \operatorname{Cl}_2\left(\frac{\pi}{3}\right)$
	$\frac{\operatorname{Cl}_2\left(\frac{\pi}{3}\right)}{\sqrt{3}}$	$\frac{\operatorname{Cl}_2\left(\frac{\pi}{3}\right)}{\sqrt{3}}\ln 3$	$\frac{\operatorname{Cl}_2\left(\frac{\pi}{3}\right)}{\sqrt{3}}\ln^2 3$	$\frac{\operatorname{Cl}_2\left(\frac{\pi}{3}\right)}{\sqrt{3}}\ln^3 3$	$\frac{\pi}{\sqrt{3}} \left[\operatorname{Cl}_2\left(\frac{\pi}{3}\right) \right]^2$
	$\ln^2 3$	$\frac{\pi}{\sqrt{3}}\zeta_2$	$\frac{\pi}{\sqrt{3}}\zeta_2 \ln 3$	$\frac{\pi}{\sqrt{3}}\zeta_2\ln^23$	$\zeta_2 \frac{\operatorname{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}}$
		$\pi \operatorname{Cl}_2\left(\frac{\pi}{3}\right)$	$\pi \operatorname{Cl}_2\left(\frac{\pi}{3}\right) \ln 3$	$\pi \operatorname{Cl}_2\left(\frac{\pi}{3}\right) \ln^2 3$	$\operatorname{Cl}_2\left(\frac{\pi}{3}\right)\operatorname{Ls}_3\left(\frac{2\pi}{3}\right)$
		ζ_3	$\zeta_3 \ln 3$	$\zeta_3 \ln^2 3$	$\pi Ls_4\left(\frac{\pi}{3}\right)$
		$\frac{Ls_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}}$	$\frac{\operatorname{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\ln 3$	$\frac{Ls_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\ln^2 3$	$\pi Ls_4\left(\frac{2\pi}{3}\right)$
		$\ln^3 3$	$\zeta_2 \frac{\operatorname{Cl}_2\left(\frac{\pi}{3}\right)}{\sqrt{3}}$	$\zeta_2 \frac{\operatorname{Cl}_2\left(\frac{\pi}{3}\right)}{\sqrt{3}} \ln 3$	$\zeta_3 \frac{\operatorname{Cl}_2(\frac{\pi}{3})}{\sqrt{3}}$
			$\frac{\pi}{\sqrt{3}}\zeta_3$	$\frac{\pi}{\sqrt{3}}\zeta_3\ln 3$	$\zeta_2\zeta_3$
			$\pi Ls_3\left(\frac{2\pi}{3}\right)$	$\pi Ls_3\left(\frac{2\pi}{3}\right) \ln 3$	ζ_5
			$\left[\operatorname{Cl}_2\left(\frac{\pi}{3}\right)\right]^2$	$\left[\operatorname{Cl}_2\left(\frac{\pi}{3}\right)\right]^2\ln 3$	$\frac{\pi}{\sqrt{3}} \operatorname{Ls}_{4}^{(1)} \left(\frac{2\pi}{3} \right)$
			ζ_4	$\zeta_4 \ln 3$	$\frac{\operatorname{Cl}_5\left(\frac{\pi}{3}\right)}{\sqrt{3}}$
			$\frac{\operatorname{Ls}_4\left(\frac{\pi}{3}\right)}{\sqrt{3}}$	$\frac{\operatorname{Ls}_4\left(\frac{\pi}{3}\right)}{\sqrt{3}}\ln 3$	$\frac{\operatorname{Ls}_5\left(\frac{\pi}{3}\right)}{\sqrt{3}}$
			$\frac{\operatorname{Ls}_4\left(\frac{2\pi}{3}\right)}{\sqrt{3}}$	$\frac{\operatorname{Ls}_4\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\ln 3$	$\frac{\operatorname{Ls}_5\left(\frac{2\pi}{3}\right)}{\sqrt{3}}$
			$\operatorname{Ls}_4^{(1)}\left(\frac{2\pi}{3}\right)$	$Ls_4^{(1)}\left(\frac{2\pi}{3}\right)\ln 3$	$\operatorname{Ls}_{5}^{(1)}\left(\frac{2\pi}{3}\right)$
			$\ln^4 3$		$\frac{Ls_5^{(2)}\left(\frac{2\pi}{3}\right)}{\sqrt{3}}$
					$\ln^5 3$

The above table contains the following functions, which appear in the splitting of the polylogarithms into real and imaginary parts [26].

$$\operatorname{Ls}_{n}(\theta) = -\int_{0}^{\theta} \ln^{n-1} \left| 2 \sin \frac{\phi}{2} \right| d\phi,$$

$$\operatorname{Ls}_{n}^{(m)}(\theta) = -\int_{0}^{\theta} \phi^{m} \ln^{n-m-1} \left| 2 \sin \frac{\phi}{2} \right| d\phi.$$

The basis of weight $\mathbf{5}$ is sum of columns $\mathbf{5}^a$ and $\mathbf{5}^b$ in Table I. Some numerical values of these functions for special values of their argument are given in appendix A.

It should be noted, that the number N_J of basis elements with weight J satisfies the following relation:

$$N_J = 2N_{J-1} \equiv 2^J. \tag{4}$$

3 Examples

Let us present several examples of two- and three-loop diagrams expressible in terms of the basis structure elaborated in the previous section. Here we will use the notation $S_2 \equiv \frac{4}{9\sqrt{3}}\text{Cl}_2\left(\frac{\pi}{3}\right)$ and are working in in Euclidean space-time with dimension $N=4-2\varepsilon$. Moreover,

the normalization factor $\frac{\left(m^2\right)^{\varepsilon}(4\pi)^{\frac{N}{2}}}{\Gamma(1+\varepsilon)}$ for each loop is assumed.

As first example we consider the two-loop single mass scale bubble diagram (VL111 in Fig.1). Taking into account the 'magic connection' [5] between the one-loop off-shell massless triangle and two-loop bubble integral and using the one-fold integral representation [27] for the former, we have:

$$m^{-2}\mathbf{VL111}(1,1,1,m) = -\frac{3}{2}\frac{1}{(1-\varepsilon)(1-2\varepsilon)} \left\{ \frac{1}{\varepsilon^2} - \frac{1}{\varepsilon} \int_0^1 \frac{dx(1-x^{2\varepsilon})}{(1-x+x^2)^{1+\varepsilon}} \right\}.$$
 (5)

Another form of the integral representation of this diagram can be found in [9]. In this way, the calculation of the coefficients of the ε -expansion is reduced to the following integrals:

$$I_{a,b} = \int_{0}^{1} \frac{dx}{1 - x + x^{2}} \ln^{a} x \ln^{b} (1 - x + x^{2}).$$
 (6)

The values of these integrals for $a + b \le 2$ are know from [1, 4, 27]. The results for the next two indices (a + b = 3, 4) are given in appendix B. Combining all these results we have:

$$m^{-2}$$
VL111 $(1, 1, 1, m) = -\frac{3}{2} \frac{1}{(1 - \varepsilon)(1 - 2\varepsilon)} \left\{ \frac{1}{\varepsilon^2} - 9S_2 \right\}$

$$+2\varepsilon \left[\frac{9}{2} S_{2} \ln 3 - \frac{\pi}{\sqrt{3}} \zeta_{2} - 3 \frac{\text{Ls}_{3} \left(\frac{2\pi}{3} \right)}{\sqrt{3}} \right]$$

$$+2\varepsilon^{2} \left[-\frac{9}{4} S_{2} \ln^{2} 3 + \frac{\pi}{\sqrt{3}} \zeta_{2} \ln 3 + 3 \frac{\text{Ls}_{3} \left(\frac{2\pi}{3} \right)}{\sqrt{3}} \ln 3 + 2 \frac{\pi}{\sqrt{3}} \zeta_{3} - 2 \frac{\text{Ls}_{4} \left(\frac{2\pi}{3} \right)}{\sqrt{3}} \right]$$

$$+\varepsilon^{3} \left[\frac{3}{2} S_{2} \ln^{3} 3 - \frac{\pi}{\sqrt{3}} \zeta_{2} \ln^{2} 3 - 4 \frac{\pi}{\sqrt{3}} \zeta_{3} \ln 3 - \frac{19}{2} \frac{\pi}{\sqrt{3}} \zeta_{4} - 3 \frac{\text{Ls}_{3} \left(\frac{2\pi}{3} \right)}{\sqrt{3}} \ln^{2} 3 \right]$$

$$+4 \frac{\text{Ls}_{4} \left(\frac{2\pi}{3} \right)}{\sqrt{3}} \ln 3 - 2 \frac{\text{Ls}_{5} \left(\frac{2\pi}{3} \right)}{\sqrt{3}} \right] + \mathcal{O}(\varepsilon^{4}).$$
 (7)

This expression coincides with analytical results obtained before. The ε -part was obtained in [5] and all higher terms were obtained by Davydychev [28].

Let us consider another example: **J011** diagram of Fig.1 with the external momentum on the mass shell of the internal mass. The first two terms of the ε -expansion of this diagram has been calculated analytically in [7]. Here we present the next two terms of this expansion. It is well known that two diagrams with different sets of indices are needed as master integrals [29]. For the numerical investigation we choose the integral with indices 122 (the first index belonging to the massless line) and a linear combination investigated in [30]. To obtain the numerical results for these two integrals with high precision [31] we have constructed their Padé approximants from the small momentum Taylor expansion [13]. The Taylor coefficients were obtained by means of the package [32]. Within the above mentioned basis we have:

$$m^{2}\mathbf{J011}(1,2,2,m) = \frac{2}{3}\zeta_{2} - \varepsilon \frac{2}{3}\zeta_{3} + \varepsilon^{2}3\zeta_{4} - \varepsilon^{3} \left\{ 2\zeta_{5} + \frac{4}{3}\zeta_{2}\zeta_{3} \right\}$$

+ $\varepsilon^{4} \left\{ \frac{61}{6}\zeta_{6} + \frac{2}{3}\zeta_{3}^{2} \right\} - \varepsilon^{5} \left\{ 6\zeta_{7} + 4\zeta_{2}\zeta_{5} + 6\zeta_{3}\zeta_{4} \right\} + \mathcal{O}(\varepsilon^{6}),$ (8)

and

$$\begin{split} m^2 \bigg[\mathbf{J011}(1,2,2,m) + 2\mathbf{J011}(2,1,2,m) \bigg] &= -\frac{2}{3\varepsilon} \frac{\pi}{\sqrt{3}} + \left\{ -9S_2 + 2\frac{\pi}{\sqrt{3}} \ln 3 \right\} \\ &+ \varepsilon \bigg\{ -3\frac{\pi}{\sqrt{3}} \ln^2 3 + 27S_2 \ln 3 - \frac{28}{3} \frac{\pi}{\sqrt{3}} \zeta_2 - 18 \frac{\operatorname{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}} \right\} + \varepsilon^2 \bigg\{ -\frac{81}{8} S_2 \ln^2 3 \\ &+ 3\frac{\pi}{\sqrt{3}} \ln^3 3 + 28\frac{\pi}{\sqrt{3}} \zeta_2 \ln 3 + \frac{112}{3} \frac{\pi}{\sqrt{3}} \zeta_3 + 54 \frac{\operatorname{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}} \ln 3 - 36 \frac{\operatorname{Ls}_4\left(\frac{2\pi}{3}\right)}{\sqrt{3}} + \frac{32}{3} \frac{\operatorname{Ls}_4\left(\frac{\pi}{3}\right)}{\sqrt{3}} \bigg\} \\ &+ \varepsilon^3 \bigg\{ -\frac{9}{4} \frac{\pi}{\sqrt{3}} \ln^4 3 + \frac{81}{2} S_2 \ln^3 3 - 216 \zeta_2 \frac{\operatorname{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}} - 81 \frac{\operatorname{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}} \ln^2 3 - 32 \frac{\operatorname{Ls}_4\left(\frac{\pi}{3}\right)}{\sqrt{3}} \ln 3 \\ &+ 108 \frac{\operatorname{Ls}_4\left(\frac{2\pi}{3}\right)}{\sqrt{3}} \ln 3 + \frac{196}{3} \frac{\operatorname{Ls}_5\left(\frac{\pi}{3}\right)}{\sqrt{3}} - 54 \frac{\operatorname{Ls}_5\left(\frac{2\pi}{3}\right)}{\sqrt{3}} - 42 \frac{\pi}{\sqrt{3}} \zeta_2 \ln^2 3 - 112 \frac{\pi}{\sqrt{3}} \zeta_3 \ln 3 \end{split}$$




Figure 2: Three-loop master-integrals. Bold and thin lines correspond to massive and massless propagators, respectively.

$$-\frac{679}{6}\frac{\pi}{\sqrt{3}}\zeta_4 + 108\frac{\pi}{\sqrt{3}}Ls_4^{(1)}\left(\frac{2\pi}{3}\right) - 81\frac{Ls_5^{(2)}\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\right\} + \mathcal{O}(\varepsilon^4),\tag{9}$$

The values of the master integrals (with indices 111 and 112) can be obtained by means of recurrence relations [16] implemented in the package **ON-SHELL2** [33]. They are relatively lengthy and therefore will not be presented here.

Now we consider master integrals occurring in the package [34]: $D_3(0, 1, 0, 1, 1, 1, m)$, $E_3(1, 1, 1, 1, 1, m)$ and $D_5(1, 1, 1, 1, 1, 0, m)$ as shown in Fig.2, the notation being as in [34]. These integrals contribute in calculations like [35].

Results for $D_3(0, 1, 0, 1, 1, 1, m)$ and $E_3(1, 1, 1, 1, m)$ have not been published before. Two independent approaches have been applied to obtain high precision numerical values for these integrals. One method started from analytical results [1, 7, 19] for two-loop propagator diagrams. The finite three-loop diagrams were then obtained by multiplication with the massive propagator with power j > 1, resulting in a one-fold integral representation, which was integrated by MAPLE with accuracy more than 200 decimals. Alternatively the E_3 -integral was calculated by the method which was used earlier for the numerical calculation of $D_3(1, 1, 1, 1, 1, 1)$ [36]. Here the product of the two one-loop propagator diagrams were expanded in their small/large external momentum and the integration was performed piecewise. Details of such calculation were published in [37]. An accuracy of about 100 decimals was also obtained in this manner.

The analytical results are the following ones:

$$m^{-2}\mathbf{E}_{3}(1,1,1,1,1,m) = -\frac{2}{3\varepsilon^{3}} - \frac{11}{3\varepsilon^{2}} + \frac{1}{\varepsilon} \left\{ -14 + \frac{27}{2}S_{2} - \zeta_{2} \right\}$$

$$+ \left\{ -\frac{139}{3} + \frac{135}{2}S_{2} - \frac{27}{2}S_{2}\ln 3 - 5\zeta_{2} - \frac{1}{3}\zeta_{3} + \frac{5}{3}\frac{\pi}{\sqrt{3}}\zeta_{2} + 9\frac{\mathrm{Ls}_{3}\left(\frac{2\pi}{3}\right)}{\sqrt{3}} \right\}$$

$$+ \varepsilon \left\{ -\frac{430}{3} + \frac{459}{2}S_{2} - \frac{135}{2}S_{2}\ln 3 + \frac{27}{4}S_{2}\ln^{2} 3 + 45\frac{\mathrm{Ls}_{3}\left(\frac{2\pi}{3}\right)}{\sqrt{3}} - 9\frac{\mathrm{Ls}_{3}\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\ln 3 + \frac{80}{9}\frac{\mathrm{Ls}_{4}\left(\frac{\pi}{3}\right)}{\sqrt{3}} + 6\frac{\mathrm{Ls}_{4}\left(\frac{2\pi}{3}\right)}{\sqrt{3}} - 17\zeta_{2} + \frac{25}{3}\frac{\pi}{\sqrt{3}}\zeta_{2} - \frac{5}{3}\frac{\pi}{\sqrt{3}}\zeta_{2}\ln 3 - \frac{13}{3}\zeta_{3} - \frac{94}{9}\frac{\pi}{\sqrt{3}}\zeta_{3}$$

$$+\frac{3}{2}\zeta_4 + 9S_2\zeta_2 + \mathcal{O}(\varepsilon^2), \tag{10}$$

$$m^{-4}\mathbf{D_3}(0,1,0,1,1,1,m) = \frac{1}{\varepsilon^3} + \frac{15}{4\varepsilon^2} + \frac{65}{8\varepsilon} + \frac{135}{16} + \frac{81}{4}S_2 + \frac{1215}{8}S_2 - \frac{243}{8}S_2 \ln 3 + \frac{27}{2}\zeta_2 \frac{\pi}{\sqrt{3}} + \frac{81}{2}\frac{\mathrm{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\right\} + \mathcal{O}(\varepsilon^2).$$
(11)

Let us now consider the diagram $D_5(1, 1, 1, 1, 1, 0)$ shown in Fig.2. As shown by Broadhurst [6] it contains the new transcendental number $V_{3,1}$. We were able to verify this, but we also found the following decomposition in terms of our basis elements:

$$V_{3,1} = \frac{81}{16}S_2^2 - \frac{1}{4}\pi Ls_3\left(\frac{2\pi}{3}\right) + \frac{13}{24}\zeta_3 \ln 3 - \frac{259}{108}\zeta_4 + \frac{3}{8}Ls_4^{(1)}\left(\frac{2\pi}{3}\right). \tag{12}$$

4 Conclusion

In this work we have constructed the basis of transcendental numbers for single mass scale diagrams with two-particle massive cuts.

The basis, up to weight 5, is given in Table I. The conjectures which have been used for the construction of the basis are as follows: the set $\{b_J\}$ of transcendental numbers of weight J has the structure:

$$b_J = \{b_{J-K}b_K, \tilde{b}_J\}, K = 1, 2, \dots, J-1,$$

where new elements $\{\tilde{b}_J\}$ are connected with functions arising in the splitting of polylogarithms $\operatorname{Li}_J(z)$, $\operatorname{Li}_J(1-z)$ into real and imaginary parts, and $z = \exp\left(i\frac{\pi}{3}\right)$, $\exp\left(i\frac{2\pi}{3}\right)$. Up to the weight **5** we observed also, that the number N_J of basis elements with weight J satisfies the relation (4).

The fact that the 'sixth root of unity' [6] plays such a crucial role in the construction of our basis remains mysterious. The introduction of the 'normalization factor' $\frac{1}{\sqrt{3}}$ remains a matter of experience. Nevertheless our construction allows us to calculate several new diagrams (or obtain new terms in the ε -expansion) in an analytical form.

The interesting question remaining beyond our work is how to construct the basis for single-mass scale diagrams having several different massive cuts. From multiloop QED and QCD calculations [2, 6, 17, 38, 39] we know that other transcendental numbers, like π , $\ln 2$, G, $\text{Li}_n\left(\frac{1}{2}\right)$ or their products appear. In this case the structure of the basis elements is connected with polylogarithms of $z = \exp\left(i\frac{\pi}{2}\right)$, $\exp\left(i\pi\right)$ [6]. At the present moment we don't know how the mixing basis could look like.

Acknowledgments We are grateful to A. Davydychev and O. V. Tarasov for useful discussions and careful reading of the manuscript and to M. Tentyukov and O. Veretin for help in numerical calculation. We are very indebted to A. Davydychev for informing us about the results of [28] before publication. M.K's research has been supported by the DFG project FL241/4-1 and in part by RFBR #98-02-16923.

APPENDIX A. Numerical values of basic elements.

Here we present numerical values up to 40 decimals of functions, which for special values of their argument provide our basic elements. These numerical values were obtained by numerical integration of their integral representation by means of MAPLE.

$$\begin{array}{lll} \operatorname{Cl}_2\left(\frac{\pi}{3}\right) &=& 1.0149416064096536250212025542745202859417\cdots, \\ \operatorname{Ls}_3\left(\frac{2\pi}{3}\right) &=& -2.1447672125694943914826421122153512974711\cdots, \\ \operatorname{Ls}_4\left(\frac{\pi}{3}\right) &=& 6.0094975498188889162047887062032707405970\cdots, \\ \operatorname{Ls}_4\left(\frac{2\pi}{3}\right) &=& 5.9506682441397849307578101290139840215916\cdots, \\ \operatorname{Ls}_4^{(1)}\left(\frac{2\pi}{3}\right) &=& -0.4976755516066471960582989542950996307358\cdots, \\ \operatorname{Cl}_5\left(\frac{\pi}{3}\right) &=& 0.4800591458997082992274840215078861889153\cdots, \\ \operatorname{Ls}_5\left(\frac{\pi}{3}\right) &=& -24.0125331255169146150157139636316267950288\cdots, \\ \operatorname{Ls}_5\left(\frac{2\pi}{3}\right) &=& -24.0394144085686386066499683097012073111734\cdots, \\ \operatorname{Ls}_5^{(1)}\left(\frac{2\pi}{3}\right) &=& 0.2708016246010793737741869955751992363356\cdots, \\ \operatorname{Ls}_5^{(2)}\left(\frac{2\pi}{3}\right) &=& -0.5181087868296801173472656387316967550219\cdots. \end{array}$$

APPENDIX B. Analytical results of integrals occurring in the ε -expansion of the two-loop bubble.

Here we present the values of the integrals (6). Their numerical values were obtained by MAPLE with accuracy of 230 decimal and again the analytical results were obtained by PSLQ.

$$I_{3,0} = -\frac{8}{3} \frac{\text{Ls}_4\left(\frac{\pi}{3}\right)}{\sqrt{3}} + \frac{4}{3} \frac{\pi}{\sqrt{3}} \zeta_3,$$

$$I_{2,1} = \zeta_2 S_2 - \frac{248}{81} \frac{\text{Ls}_4\left(\frac{\pi}{3}\right)}{\sqrt{3}} + \frac{20}{27} \frac{\pi}{\sqrt{3}} \zeta_2 \ln 3 + \frac{280}{81} \frac{\pi}{\sqrt{3}} \zeta_3,$$

$$I_{1,2} = 2\zeta_{2}S_{2} - \frac{9}{2}S_{2}\ln^{2}3 + 6\frac{\text{Ls}_{3}\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\ln 3 - \frac{208}{81}\frac{\text{Ls}_{4}\left(\frac{\pi}{3}\right)}{\sqrt{3}} - 4\frac{\text{Ls}_{4}\left(\frac{2\pi}{3}\right)}{\sqrt{3}} + \frac{94}{27}\frac{\pi}{\sqrt{3}}\zeta_{2}\ln 3 + \frac{740}{81}\frac{\pi}{\sqrt{3}}\zeta_{3},$$

$$I_{0,3} = \frac{2}{3}\frac{\pi}{\sqrt{3}}\ln^{3}3 - 18S_{2}\ln^{2}3 + 24\frac{\text{Ls}_{3}\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\ln 3 - 16\frac{\text{Ls}_{4}\left(\frac{2\pi}{3}\right)}{\sqrt{3}} + 12\frac{\pi}{\sqrt{3}}\zeta_{2}\ln 3 + 24\frac{\pi}{\sqrt{3}}\zeta_{3},$$
(B.1)

and

$$I_{4,0} = \frac{340}{27} \frac{\pi}{\sqrt{3}} \zeta_4,$$

$$I_{3,1} = -6S_2 \zeta_3 - 18\zeta_2 \frac{\text{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}} - \frac{8}{3} \frac{\text{Ls}_4\left(\frac{\pi}{3}\right)}{\sqrt{3}} \ln 3 + \frac{49}{9} \frac{\text{Ls}_5\left(\frac{\pi}{3}\right)}{\sqrt{3}}$$

$$+ \frac{4}{3} \frac{\pi}{\sqrt{3}} \zeta_3 \ln 3 + \frac{5957}{216} \frac{\pi}{\sqrt{3}} \zeta_4 + 9 \frac{\pi}{\sqrt{3}} \text{Ls}_4^{(1)} \left(\frac{2\pi}{3}\right) - \frac{27}{4} \frac{\text{Ls}_5^{(2)}\left(\frac{2\pi}{3}\right)}{\sqrt{3}},$$

$$I_{2,2} = 2S_2 \zeta_2 \ln 3 - \frac{46}{3} S_2 \zeta_3 + 9 \frac{\pi}{\sqrt{3}} S_2^2 - \frac{128}{3} \zeta_2 \frac{\text{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}} - \frac{496}{81} \frac{\text{Ls}_4\left(\frac{\pi}{3}\right)}{\sqrt{3}} \ln 3 + \frac{986}{81} \frac{\text{Ls}_5\left(\frac{\pi}{3}\right)}{\sqrt{3}} + \frac{20}{27} \frac{\pi}{\sqrt{3}} \zeta_2 \ln^2 3 + \frac{560}{81} \frac{\pi}{\sqrt{3}} \zeta_3 \ln 3 + \frac{16963}{324} \frac{\pi}{\sqrt{3}} \zeta_4 + \frac{62}{3} \frac{\pi}{\sqrt{3}} \text{Ls}_4^{(1)} \left(\frac{2\pi}{3}\right) - \frac{31}{2} \frac{\text{Ls}_5^{(2)}\left(\frac{2\pi}{3}\right)}{\sqrt{3}},$$

$$I_{1,3} = 6S_2 \zeta_2 \ln 3 - 22S_2 \zeta_3 + 27 \frac{\pi}{\sqrt{3}} S_2^2 - \frac{9}{2} S_2 \ln^3 3 - 56\zeta_2 \frac{\text{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}} + 9 \frac{\text{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}} \ln^2 3 - \frac{208}{27} \frac{\text{Ls}_4\left(\frac{\pi}{3}\right)}{\sqrt{3}} \ln 3 - 12 \frac{\text{Ls}_4\left(\frac{2\pi}{3}\right)}{\sqrt{3}} \ln 3 + \frac{398}{27} \frac{\text{Ls}_5\left(\frac{\pi}{3}\right)}{\sqrt{3}} + 6 \frac{\text{Ls}_5\left(\frac{2\pi}{3}\right)}{\sqrt{3}} + \frac{47}{9} \frac{\pi}{\sqrt{3}} \zeta_2 \ln^2 3 + \frac{740}{27} \frac{\pi}{\sqrt{3}} \zeta_3 \ln 3 + \frac{10847}{108} \frac{\pi}{\sqrt{3}} \zeta_4 + 26 \frac{\pi}{\sqrt{3}} \text{Ls}_4^{(1)} \left(\frac{2\pi}{3}\right) - \frac{39}{2} \frac{\text{Ls}_5^{(2)}\left(\frac{2\pi}{3}\right)}{\sqrt{3}},$$

$$I_{0,4} = \frac{2}{3} \frac{\pi}{\sqrt{3}} \ln^4 3 - 24S_2 \ln^3 3 + 48 \frac{\text{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}} \ln^2 3 - 64 \frac{\text{Ls}_4\left(\frac{2\pi}{3}\right)}{\sqrt{3}} \ln 3 + 228 \frac{\pi}{\sqrt{3}} \zeta_4.$$
(B.2)

References

- [1] D. J. Broadhurst, Z. Phys.C 47 (1990) 115.
- [2] D. J. Broadhurst, Z. Phys. C **54** (1992) 599.
- [3] J. van der Bij and M. Veltman, Nucl. Phys. B 231 (1984) 205;
 C. Ford, I. Jack and D. R. T. Jones, Nucl. Phys. B 387 (1992) 373.
- [4] A. I. Davydychev and J. B. Tausk, Nucl. Phys. B **397** (1993) 123.
- [5] A. I. Davydychev and J. B. Tausk, Phys. Rev. D 53 (1996) 7381.
- [6] D. J. Broadhurst, Eur. Phys. J. C 8 (1999) 311.
- [7] J. Fleischer, M. Yu. Kalmykov and A. V. Kotikov, Phys. Lett. B 462 (1999) 169 (hep-ph/9905249).
- [8] E. E. Boos and A. I. Davydychev, Teor. Mat. Fiz. 89 (1991) 56 [Theor. Math. Phys. 89 (1991) 1052];
 - F. A. Berends, M. Buza, M. Böhm and R. Scharf, Z. Phys.C **63** (1994) 227.
- [9] D. J. Broadhurst, J. Fleischer and O. V. Tarasov, Z. Phys. C 60 (1993) 287.
- [10] F. A. Berends and J. B. Tausk, Nucl. Phys. B 421 (1994) 456;
 S. Bauberger, F. A. Berends, M. Böhm and M. Buza, Nucl. Phys. B 434 (1995) 383.
- [11] A. Ghinculov and J. J. van der Bij, Nucl. Phys. B **436** (1995) 30;
 - A. Ghinculov, Phys. Lett. B **385** (1996) 279;
 - A. Ghinculov and Y.-P. Yao, Nucl. Phys. B **516** (1998) 385.
- [12] D. Kreimer, Phys. Lett. B **273** (1991), 277; ibid B **292** (1992), 341;
 - A. Czarnecki, U. Kilian and D. Kreimer, Nucl. Phys. B 433 1995) 259;
 - S. Bauberger, and M. Böhm, Nucl. Phys. B **445** (1995) 25;
 - J. Fleischer et al., Eur. Phys. J. C 2 (1998) 747.
- [13] J. Fleischer and O. V. Tarasov, Z. Phys. C **64** (1994) 413;O. V. Tarasov, Nucl. Phys. B **480** (1996) 397.
- [14] S. Groote, J. G. Körner and A. A. Pivovarov, Phys. Lett. B 443 (1998) 269; Nucl. Phys. B 542 (1999) 515.
- [15] H. R. P. Ferguson and D. H. Bailey, RNR Technical Report, RNR-91-032;
 - H. R. P. Ferguson, D. H. Bailey and S. Arno, NASA Technical Report, NAS-96-005;
 - D. H. Bailey and D. J. Broadhurst, math.na/9905048.

- [16] F. V. Tkachov, Phys. Lett. B 100 (1981) 65;K. G. Chetyrkin and F. V. Tkachov, Nucl. Phys. B 192 (1981) 159.
- [17] D. J. Broadhurst, hep-th/9604128.
- [18] D. Kreimer, Phys. Lett. B **354** (1995) 117; Eur. Phys. J. C **2** (1998) 757.
 - D. J. Broadhurst and D. Kreimer, Int. J. Mod. Phys. C 6 (1995) 519; Phys. Lett. B 393 (1997) 403; Phys. Lett. B 426 (1998) 339;
 - D. J. Broadhurst, R. Delbourgo and D. Kreimer, Phys. Lett. B 366 (1996) 421;
 - D. J. Broadhurst, J. A. Gracey and D. Kreimer, Z. Phys. C 75 (1997) 559.
- [19] J. Fleischer, A. V. Kotikov and O. L. Veretin, Nucl. Phys. B **547** (1999) 343.
- [20] J. A. M. Vermaseren, Int. J. Mod. Phys. A **14** (1999) 2037;
 - J. Blümlein and S. Kurth, Phys. Rev. D **60** (1999) 014018;
 - E. Remiddi and J. A. M. Vermaseren, hep-ph/9905237.
- [21] A. V. Kotikov, Phys. Lett. B **254** (1991) 158; B **259** (1991) 314.
- [22] A. I. Davydychev and R. Delbourgo, J. Math. Phys. 39 (1998) 4299;
 D. J. Broadhurst, hep-th/9806174.
- [23] A. I. Davydychev, hep-th/9908032.
- [24] G. 't Hooft and M. Veltman, Nucl. Phys. B 44 (1972) 189.
- [25] S. Groote, J. G. Körner and A. A. Pivovarov, Phys. Rev. D 60 (1999) 061701.
- [26] L. Lewin, Polylogarithms and associated functions (North-Holland, Amsterdam, 1981).
- [27] N. I. Ussyukina and A. I. Davydychev, Phys. Lett. B 332 (1994) 159.
- [28] A. I. Davydychev, Mainz preprint MZ-TH/99-44 (hep-ph/9910224).
- [29] O V. Tarasov, Nucl. Phys. B **502** (1997) 455.
- [30] F. A. Berends, A. I. Davydychev and N. I. Ussyukina, Phys. Lett. B 426 (1998) 95;
 A. I. Davydychev and V. A. Smirnov, Nucl. Phys. B 554 (1999) 391.
- [31] J. Fleischer, V. A. Smirnov and O. V. Tarasov, Z. Phys. C 74 (1997) 379;
 J. Fleischer, M. Yu. Kalmykov and O. L. Veretin, Phys. Lett. B 427 (1998) 141.
- [32] L. V. Avdeev, J. Fleischer, M. Yu. Kalmykov and M. N. Tentyukov, Nucl. Inst. Meth. A 389 (1997) 343; Comp. Phys. Commun. 107 (1997) 155.
- [33] J. Fleischer and M. Yu. Kalmykov, hep-ph/9907431.

- [34] Leo V. Avdeev, Comput. Phys. Commun. **98** (1996) 15.
- [35] K. G. Chetyrkin and M. Steinhauser, hep-ph/9907509.
- [36] L. Avdeev, J. Fleischer, S. Mikhailov and O. Tarasov, Phys. Lett. B 336 (1994) 560, Erratum-ibid. B 349 (1995) 597.
- [37] J. Fleischer and O. Tarasov, Nucl. Phys. B. (Proc. Suppl.) **37** (1994) 115 (hep-ph/9407235).
- [38] N. Gray, D. J. Broadhurst, W. Grafe and K. Schilcher, Z. Phys. C 48 (1990) 673;
 - D. J. Broadhurst, N. Gray and K. Schilcher, Z. Phys. C 52 (1991) 111;
 - L.V. Avdeev and M. Yu. Kalmykov, Nucl. Phys. B **502** (1997) 419;
 - J. Fleischer, F. Jegerlehner, O. V. Tarasov and O. L. Veretin, Nucl. Phys. B **539** (1999) 671.
- [39] S. Laporta and E. Remiddi, Phys. Lett. B **379** (1996) 283.