

Boundary remnant of Yangian symmetry and the structure of rational reflection matrices

G. W. Delius, N. J. MacKay, B. J. Short

*Department of Mathematics,
University of York,
York YO10 5DD, U.K.¹*

Abstract

For the classical principal chiral model with boundary, we give the subset of the Yangian charges which remains conserved under certain integrable boundary conditions, and extract them from the monodromy matrix. Quantized versions of these charges are used to deduce the structure of rational solutions of the reflection equation, analogous to the ‘tensor product graph’ for solutions of the Yang-Baxter equation. We give a variety of such solutions, including some for reflection from non-trivial boundary states, for the $SU(N)$ case, and confirm these by constructing them by fusion from the basic solutions.

1 The principal chiral model with boundary

1.1 Classical boundary conditions and conserved charges

In a recent paper [1] (to which the reader is referred for more detail and references), two of us explored the classical integrability of the principal chiral model (PCM) with boundary, and the corresponding quantum boundary S -matrices. The model is defined by the action

$$\mathcal{L} = \frac{1}{2} \text{Tr} (\partial_\mu g^{-1} \partial^\mu g) , \quad (1.1)$$

where the field $g(x^\mu)$ takes values in a compact Lie group G , and is defined in 1+1D Minkowski spacetime with $-\infty < x \leq 0$.

¹emails: gwd2, nm15, bjs108 @york.ac.uk

We found two classes of classical boundary condition which preserve the conservation and involution of the charges necessary for integrability. Here we discuss only the ‘chiral’ condition,

$$g(0) \in k_L H k_R^{-1}, \quad (1.2)$$

where $k_{L,R}$ are arbitrary group elements and H is a maximal Lie subgroup of G such that G/H is a symmetric space. For simplicity we also set $k_L = k_R = e$, the identity element. The global $G_L \times G_R$ symmetry of the original bulk model, given by $g \mapsto UgV^{-1}$ and generated by the conserved currents

$$j_\mu^L = \partial_\mu g g^{-1}, \quad j_\mu^R = -g^{-1} \partial_\mu g, \quad (1.3)$$

is thus broken to $H \times H$. For the currents (L or R), and writing the Lie algebras of G and H as \mathfrak{g} and \mathfrak{h} respectively, we have $j_0(0) \in \mathfrak{h}$, while the boundary equation-of-motion then requires $j_1(0) \in \mathfrak{k}$, where $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{k}$. Alternatively, since H is the subgroup of G fixed under an involution σ of \mathfrak{g} , we can write $j_0 = \sigma(j_0)$, and $j_1 = -\sigma(j_1)$, at $x = 0$.

In the bulk model, the $G \times G$ symmetry sits inside a larger $Y(\mathfrak{g}) \times Y(\mathfrak{g})$ symmetry, where $Y(\mathfrak{g})$ is the Yangian algebra. This is generated by charges (where we use the conventions of [2])

$$Q^{(0)a} = \int j_0^a dx \quad (1.4)$$

$$Q^{(1)a} = \int j_1^a dx - \frac{1}{2} f^a_{bc} \int j_0^b(x) \int^x j_0^c(y) dy dx \quad (1.5)$$

using j^L and j^R respectively, decomposed into $j = j^a t_a$ where the t^a are generators of \mathfrak{g} with $[t_a, t_b] = f_{ab}^c t_c$. The integrals are over all space, $(-\infty, \infty)$ for the bulk model. But on the half-line $(-\infty, 0]$, these charges are no longer generally conserved. However, there are two important sets of charges which do remain conserved. Writing \mathfrak{h} -indices as i, j, k, \dots and \mathfrak{k} -indices as p, q, r, \dots , and noting that the only non-zero structure constants are f^i_{jk} and f^i_{pq} (and cycles thereof), these are

$$Q^{(0)i} \quad (1.6)$$

$$\text{and } \tilde{Q}^{(1)p} \equiv Q^{(1)p} + \frac{1}{2} f^p_{qi} Q^{(0)i} Q^{(0)q}. \quad (1.7)$$

The first set generates H and was noted in [1]. To check their conservation, we note that

$$\frac{d}{dt} Q^{(0)i} = j_1^i(0) = 0$$

$$\text{and } \frac{d}{dt} \tilde{Q}^{(1)p} = \frac{d}{dt} Q^{(1)p} + \frac{1}{2} f^p_{qi} Q^{(0)i} j_1^q(0) = 0$$

$$\text{since } \frac{d}{dt} Q^{(1)p} = \frac{1}{2} f^p_{iq} Q^{(0)i} j_1^q(0).$$

It is difficult to prove rigorous quantum results for the PCM, but we shall assume that these charges remain conserved after quantization, in the form

$$Q^{(0)i} \tag{1.8}$$

$$\text{and } \tilde{Q}^{(1)p} \equiv Q^{(1)p} + \frac{1}{4}[C_2^{\mathbf{h}}, Q^{(0)p}]. \tag{1.9}$$

Here we have set $\hbar = 1$ for convenience, and $C_2^{\mathbf{h}} \equiv \gamma_{ij} Q^{(0)i} Q^{(0)j}$ is the quadratic Casimir operator of \mathfrak{g} restricted to \mathbf{h} , with $\gamma_{ij} = f_{ia}{}^b f_{jb}{}^a$.

1.2 Conserved charges from the monodromy matrix

It is straightforward to construct these charges as coefficients of a spectral parameter in a monodromy matrix. The conservation and curvature-freedom of the bulk PCM currents can be expressed through a Lax pair, $[\partial_0 - L_0, \partial_1 - L_1] = 0$ where

$$L_1 = \frac{1}{1-u^2} (j_1 - u j_0), \quad L_0 = \frac{1}{1-u^2} (j_0 - u j_1).$$

The Yangian charges then appear in the monodromy matrix

$$T_{-\infty}^{\infty}(u) \equiv \mathbf{P} \exp \left(\int_{-\infty}^{\infty} L_1 dx \right) = \exp \left(\frac{1}{u} Q^{(0)a} t_a - \frac{1}{u^2} Q^{(1)a} t_a + \dots \right).$$

We can regard our model on the half-line, with boundary condition $j_0 = \sigma(j_0)$ and $j_1 = -\sigma(j_1)$ at $x = 0$, as a restriction of the bulk model with $j_0(x) = \sigma(j_0(-x))$ and $j_1(x) = -\sigma(j_1(-x))$. We then have

$$T_{-\infty}^{\infty}(u) = \sigma \left((T_{-\infty}^0)^{-1}(-u) \right) T_{-\infty}^0(u),$$

which is conserved because $L_0(u) = \sigma(L_0(-u))$. Expanding this gives

$$\begin{aligned} & \exp \left(\frac{1}{u} Q^{(0)a} \sigma(t_a) + \frac{1}{u^2} Q^{(1)a} \sigma(t_a) + \dots \right) \exp \left(\frac{1}{u} Q^{(0)a} t_a - \frac{1}{u^2} Q^{(1)a} t_a + \dots \right) \\ &= 1 + \frac{1}{u} Q^{(0)a} (t_a + \sigma(t_a)) - \frac{1}{u^2} \left\{ Q^{(1)a} (t_a - \sigma(t_a)) - Q^{(0)b} Q^{(0)c} \left(\sigma(t_b) t_c + \frac{1}{2} \sigma(t_b) \sigma(t_c) + \frac{1}{2} t_b t_c \right) \right\} + \dots \\ &= 1 + \frac{2}{u} Q^{(0)i} t_i + \frac{1}{2} \left(\frac{2}{u} Q^{(0)i} t_i \right)^2 - \frac{2}{u^2} \left(Q^{(1)p} t_p - \frac{1}{4} Q^{(0)b} Q^{(0)c} [\sigma(t_b), t_c] \right) + \dots \\ &= \exp \left(\frac{2}{u} Q^{(0)i} t_i - \frac{2}{u^2} \tilde{Q}^{(1)p} t_p + \dots \right), \end{aligned}$$

as required.

2 Rational reflection matrices

2.1 Reflection from the boundary ground state

Recall the structure of the rational solutions of the bulk Yang-Baxter equation (YBE), the ‘ R -matrices’. The R -matrix acting on $U \otimes V$, where U and V are irreducible representations of $Y(\mathfrak{g})$, decomposes into the sum of projectors onto the \mathfrak{g} -irreducible component representations of $U \otimes V$. The coefficients of these projectors can then, in simple cases – where U and V are \mathfrak{g} -irreducible and $U \otimes V$ has no multiplicities –, be deduced from a ‘tensor product graph’ [4] which describes how the rest of $Y(\mathfrak{g})$ relates these components.

We proceed similarly now for the boundary Yang-Baxter or ‘reflection’ equation on $U \otimes V$,

$$\begin{aligned} \check{R}_{VU}(\theta - \phi)(I \otimes K_U(\theta))\check{R}_{UV}(\theta + \phi)(I \otimes K_V(\phi)) = \\ (I \otimes K_V(\phi))\check{R}_{VU}(\theta + \phi)(I \otimes K_U(\theta))\check{R}_{UV}(\theta - \phi), \end{aligned} \quad (2.1)$$

where $\check{R}_{UV} : U \otimes V \rightarrow V \otimes U$ is a solution of the bulk YBE, $K_U : U \rightarrow U$ is the reflection matrix and the variables θ and ϕ are the rapidities of the particles incident on the boundary. Implicit in K ’s acting only on the bulk multiplet U or V is that the boundary has no structure of its own – *i.e.* it is in a singlet state. Another possibility, also discussed below, is that $K_U : U \rightarrow \bar{U}$, in which case the reflection equation becomes

$$\begin{aligned} \check{R}_{\bar{V}\bar{U}}(\theta - \phi)(I \otimes K_U(\theta))\check{R}_{U\bar{V}}(\theta + \phi)(I \otimes K_V(\phi)) = \\ (I \otimes K_V(\phi))\check{R}_{\bar{V}\bar{U}}(\theta + \phi)(I \otimes K_U(\theta))\check{R}_{U\bar{V}}(\theta - \phi). \end{aligned} \quad (2.2)$$

In this paper, in contrast to [1], we describe only the matrix structure of (the individual L and R factors of) K , and do not concern ourselves with the scalar prefactors and crossing/unitarity conditions necessary to construct a valid boundary S -matrix [5].

Let us specialize, as in the bulk case, to U and V which are \mathfrak{g} -irreducible. Conservation of the $Q^{(0)i}$ requires that

$$K_U(\theta)Q^{(0)i} = Q^{(0)i}K_U(\theta)$$

(in which by $Q^{(0)i}$ we mean its appropriate representation) and thus that $K_U(\theta)$ act as the identity on \mathfrak{h} -irreducible components of U . So we have

$$K_U(\theta) = \sum_{W_{\mathfrak{h}} \subset U_{\mathfrak{g}}} \tau_W(\theta)P_W,$$

where the sum is over \mathfrak{h} -representations W into which the \mathfrak{g} -representation U branches, and P_W is the projector onto W . Thus the conjugating case, $K : U \rightarrow \bar{U}$, is only admissible when U and \bar{U} branch to the same \mathfrak{h} -representations.

To deduce relations among the τ_W we use conservation of the $\tilde{Q}^{(1)p}$, recalling² that, on a \mathfrak{g} -irreducible multiplet of rapidity θ , the action of $Q^{(1)}$ is given by the evaluation representation,

$$Q^{(1)a} = \theta \frac{c_A}{2i\pi} Q^{(0)a},$$

where c_A is the value of the quadratic Casimir of \mathfrak{g} in the adjoint representation. So

$$K_U(\theta) \left(\theta \frac{c_A}{2i\pi} Q^{(0)p} + \frac{1}{4} [C_2^{\mathfrak{h}}, Q^{(0)p}] \right) = \left(-\theta \frac{c_A}{2i\pi} Q^{(0)p} + \frac{1}{4} [C_2^{\mathfrak{h}}, Q^{(0)p}] \right) K_U(\theta),$$

and for $W_1, W_2 \subset U_{\mathfrak{g}}$ such that the reduced matrix element $\langle W_1 || Q^{(0)p} || W_2 \rangle \neq 0$ we have

$$\frac{\tau_{W_2}(\theta)}{\tau_{W_1}(\theta)} = [\Delta_{12}], \quad \text{where}^3 [A] \equiv \frac{\frac{i\pi A}{2c_A} + \theta}{\frac{i\pi A}{2c_A} - \theta} \quad (2.3)$$

and $\Delta_{12} = C_2^{\mathfrak{h}}(W_1) - C_2^{\mathfrak{h}}(W_2)$.

To find the W_1, W_2 for which $\langle W_1 || Q^{(0)p} || W_2 \rangle \neq 0$ we note that the $Q^{(0)p}$ (that is, the generators of \mathfrak{k}) form a representation Z of \mathfrak{h} . The Wigner-Eckart theorem then requires $W_1 \subset Z \otimes W_2$. This is strictly a necessary rather than a sufficient condition, but we shall assume its sufficiency in what follows.⁴

We can therefore describe the structure of $K_U(\theta)$ by using a ‘branching graph’, in which the W_i are the nodes, linked by an edge, directed from W_i to W_j and labelled by positive Δ_{ij} , when $W_i \subset Z \otimes W_j$.

2.2 Example : $G = SU(N)$

We shall examine how this works in the case of $G = SU(N)$, for which $c_A = 2N^2$. One can go through the list of all G/H similarly. Case 1 below corresponds to the standard, non-conjugating reflection equation (2.1), and cases 2 and 3 to the conjugating equation (2.2).

1. $H = S(U(M) \times U(N - M))$

We denote representations of $SU(M) \times SU(N - M)$ by (X, Y) where X is an $SU(M)$ - and Y an $SU(N - M)$ -representation, here written as a Young tableau. The singlet representation is written 1. Note that $Z = (\bar{\square}, \square) \oplus (\square, \bar{\square})$ (where $\bar{\square}$ denotes the conjugate of

²Note that our conventions differ from that of [3]: the relative sign of θ is due to our use of the opposite coproduct.

³This definition differs slightly from that used in [1].

⁴This distinction was made in the bulk case as that between the ‘tensor product graph’ and the ‘extended tensor product graph’ [4].

\square), so that the branching graphs are

$$\begin{aligned} U_{\mathbf{g}} = \square : & \quad (1, \square) \xrightarrow{N-2M} (\square, 1) \\ U_{\mathbf{g}} = \boxplus : & \quad (1, \boxplus) \xrightarrow{N-2M-2} (\square, \square) \xrightarrow{N-2M+2} (\boxplus, 1) \\ U_{\mathbf{g}} = \boxplus\boxplus : & \quad \left(1, \boxplus\boxplus\right) \xrightarrow{N-2M-4} (\square, \boxplus) \xrightarrow{N-2M} (\boxplus, \square) \xrightarrow{N-2M+4} (\boxplus\boxplus, 1) \end{aligned}$$

and so on (in agreement with the restriction to $SU(N)$ of the results of [7]).

The general result, for U the r th-rank antisymmetric tensor ($r \leq [N/2]$), and with (p, q) denoting the p th rank $SU(M)$ and q th rank $SU(N - M)$ antisymmetric tensor, is easily read-off from the graph

$$(0, r) \xrightarrow{N-2M-2(r-1)} (1, r-1) \dots \xrightarrow{N-2M-2(r-1)+4(p-1)} (p, r-p) \xrightarrow{N-2M-2(r-1)+4p} \dots (r-1, 1) \xrightarrow{N-2M+2(r-1)} (r, 0),$$

and is

$$K_U(\theta) = \sum_{p=0}^r \prod_{q=0}^p [N - 2M - 2(r-1) + 4(q-1)] P_{(p, r-p)}. \quad (2.4)$$

2. $H = Sp(N)$ (N even)

$U_{\mathbf{g}} = \square$, the vector representation, branches to the single \square of $Sp(N)$, and the reflection matrix is therefore constant. Here⁵ $Z = \boxplus$, and for higher representations we have

$$\begin{aligned} U_{\mathbf{g}} = \boxplus : & \quad \boxplus \xrightarrow{N} 1 \\ U_{\mathbf{g}} = \boxplus\boxplus : & \quad \boxplus\boxplus \xrightarrow{N-2} \square \\ U_{\mathbf{g}} = \boxplus\boxplus\boxplus : & \quad \boxplus\boxplus\boxplus \xrightarrow{N-4} \boxplus \xrightarrow{N} 1 \end{aligned}$$

and in general for the r th rank fundamental tensor of $SU(N)$, and denoting by (p) the p th rank antisymmetric tensor of $Sp(N)$,

$$(r) \longrightarrow (r-2) \dots \xrightarrow{N-2(r-2p)} (r-2p) \xrightarrow{N+4-2(r-2p)} \dots \begin{cases} (2) \xrightarrow{N} (0) & r \text{ even} \\ (3) \xrightarrow{N-2} (1) & r \text{ odd} \end{cases}$$

⁵We continue to use Young tableau notation for representations of $SO(N)$ and $Sp(N)$, but it should be understood that traces have been removed from symmetric tableaux for $SO(N)$, and symplectic traces from antisymmetric tableaux for $Sp(N)$.

3. $H = SO(N)$

Here $Z = \square$ and each reflection matrix is constant, since the r th rank fundamental anti-symmetric tensor representation of $SU(N)$ branches to the same, irreducible representation of $SO(N)$.

2.3 Reflection from a boundary bound state

When a reflection matrix is used to construct a boundary S -matrix, it may have a pole at one of the labels of the branching graph. We then expect a multiplet of boundary states to exist which transforms in an \mathbf{h} -representation corresponding to a subgraph. The results of the previous section can be extended to accommodate such non-trivial boundary states.

Suppose we wish to calculate the reflection matrix $K_U^{[V]}(\theta)$ of bulk multiplet $U_{\mathbf{g}}$ off boundary multiplet $V_{\mathbf{h}}$. The $Q^{(0)i}$ have trivial coproduct, and their conservation enforces the decomposition of $K_U^{[V]}(\theta)$ into projectors P_W onto the \mathbf{h} -irreducible components $W_{\mathbf{h}}$ of $U_{\mathbf{g}} \otimes V_{\mathbf{h}}$,

$$K_U^{[V]}(\theta) = \sum_{W_{\mathbf{h}} \subset U_{\mathbf{g}} \otimes V_{\mathbf{h}}} \tau_W(\theta) P_W.$$

To deduce the τ_W , we need to compute the action of $\tilde{Q}^{(1)p}$ on the spaces W . Using the Yangian coproduct

$$\Delta(Q^{(1)a}) = Q^{(1)a} \otimes 1 + 1 \otimes Q^{(1)a} + \frac{1}{2} f^a_{bc} Q^{(0)b} \otimes Q^{(0)c},$$

we have

$$\begin{aligned} \Delta(\tilde{Q}^{(1)p}) &= \Delta\left(Q^{(1)p} + \frac{1}{4}[C_2^{\mathbf{h}}, Q^{(0)p}]\right) \\ &= Q^{(1)p} \otimes 1 + 1 \otimes Q^{(1)p} + \frac{1}{4}[C_2^{\mathbf{h}}, Q^{(0)p}] \otimes 1 + 1 \otimes \frac{1}{4}[C_2^{\mathbf{h}}, Q^{(0)p}] \\ &\quad + \frac{1}{2} f^p_{iq} Q^{(0)i} \otimes Q^{(0)q} + \frac{1}{2} f^p_{qi} Q^{(0)q} \otimes Q^{(0)i} + \frac{1}{2} [\gamma_{ij} Q^{(0)i} \otimes Q^{(0)j}, Q^{(0)p} \otimes 1 + 1 \otimes Q^{(0)p}] \\ &= \tilde{Q}^{(1)p} \otimes 1 + 1 \otimes \tilde{Q}^{(1)p} + [\gamma_{ij} Q^{(0)i} \otimes Q^{(0)j}, Q^{(0)p} \otimes 1] \\ &= \tilde{Q}^{(1)p} \otimes 1 + 1 \otimes \tilde{Q}^{(1)p} + \frac{1}{2} [\Delta(C_2^{\mathbf{h}}) - C_2^{\mathbf{h}} \otimes 1 - 1 \otimes C_2^{\mathbf{h}}, Q^{(0)p} \otimes 1]. \end{aligned}$$

Thus when U is \mathbf{g} -irreducible and V is \mathbf{h} -irreducible, we have

$$\begin{aligned} &K_U^{[V]}(\theta) \left(\theta \frac{c_A}{2i\pi} Q^{(0)p} \otimes 1 + \frac{1}{4}[C_2^{\mathbf{h}} \otimes 1, Q^{(0)p} \otimes 1] + \frac{1}{2} [\Delta(C_2^{\mathbf{h}}) - C_2^{\mathbf{h}} \otimes 1 - 1 \otimes C_2^{\mathbf{h}}, Q^{(0)p} \otimes 1] \right) \\ &= \left(-\theta \frac{c_A}{2i\pi} Q^{(0)p} \otimes 1 + \frac{1}{4}[C_2^{\mathbf{h}} \otimes 1, Q^{(0)p} \otimes 1] + \frac{1}{2} [\Delta(C_2^{\mathbf{h}}) - C_2^{\mathbf{h}} \otimes 1 - 1 \otimes C_2^{\mathbf{h}}, Q^{(0)p} \otimes 1] \right) K_U^{[V]}(\theta). \end{aligned}$$

So a graph is generated exactly as for the boundary ground state, but now with

$$\frac{\tau_{W_2}(\theta)}{\tau_{W_1}(\theta)} = [\Delta_{12}] .$$

Here

$$\Delta_{12} = 2 (C_2^{\mathbf{h}}(W_1) - C_2^{\mathbf{h}}(W_2)) - (C_2^{\mathbf{h}}(\widetilde{W}_1) - C_2^{\mathbf{h}}(\widetilde{W}_2)) ,$$

where \widetilde{W}_i is the \mathbf{h} -component of $U_{\mathbf{g}}$ from which W_i descends. Note that this subsumes the results of section 2.1: if the boundary is in the ground state, $W_i = \widetilde{W}_i$, and we reproduce (2.3).

2.4 Example : $G = SU(N)$, $H = S(U(M) \times U(N - M))$

We denote by $K_p^{[q]}$ the reflection of the bulk p th antisymmetric tensor off the boundary state in the q th antisymmetric tensor of $SU(M)$. Then

$$K_1^{[1]} : \quad \{(\square, 1) \oplus (1, \square)\} \otimes (\square, 1) \\ (\square, 1) \xleftarrow{N-2M-4} (\square, \square) \xrightarrow{N-2M+4} (\square, 1)$$

$$K_1^{[2]} : \quad \{(\square, 1) \oplus (1, \square)\} \otimes (\square, 1) \\ (\square, 1) \xleftarrow{N-2M-4} (\square, \square) \xrightarrow{N-2M+8} (\square, 1)$$

$$K_2^{[1]} : \quad \{(\square, 1) \oplus (\square, \square) \oplus (1, \square)\} \otimes (\square, 1) \\ (\square, \square) \xrightarrow{N-2M-6} (\square, \square) \\ \downarrow_{N-2M+2} \qquad \qquad \downarrow_{N-2M+2} \\ (\square, \square) \xrightarrow{N-2M-6} (\square, 1) \\ \downarrow_{N-2M+6} \\ (\square, 1)$$

and so on; the result is easily generalized.

3 Fusion of reflection matrices

For the low rank examples that we have given in the previous section it is possible to construct the K -matrices explicitly by fusion [6, 7]. In this section we explain how this is done.

3.1 $G = SU(N)$, $H = S(U(M) \times U(N - M))$

We start from the vector reflection matrix $K_1^{[0]}(\theta)$. For $H = S(U(M) \times U(N - M))$ the matrix part of this is


$$P_1 + [N - 2M] P_2 \quad \text{where} \quad P_1 = \frac{1}{2}(I - E), \quad P_2 = \frac{1}{2}(I + E).$$

Here E is an $N \times N$ matrix, determined by the boundary conditions [1], which satisfies $E^2 = I_N$ (the $N \times N$ identity matrix), so that the above are orthogonal projectors. (Recall [1] that the admissible E locally parametrize G/H , although choosing $k_L = k_R = 1$ makes the specific choice $E = \text{diag}(I_M, -I_{N-M})$.) They project onto $(1, \square)$ and $(\square, 1)$ respectively, so that the vector particle reflection matrix is as given in the previous section.

We calculate the second rank reflection matrix $K_2^{[0]}$ by fusing together two vector multiplets. The vector multiplet bulk S -matrix has a simple pole at $\theta = \frac{2i\pi}{c_A}$ whose residue is the projector onto the second-rank antisymmetric representation of $SU(N)$. This is interpreted according to the bootstrap principle as another particle multiplet, for which the reflection matrix is

$$K_2^{[0]}(\theta) = \check{R}_{11}(2i\pi/N) \left(I \otimes K_1^{[0]}(\theta + i\pi/N) \right) \check{R}_{11}(2\theta) \left(I \otimes K_1^{[0]}(\theta - i\pi/N) \right), \quad (3.1)$$

or, diagrammatically,



We now introduce the diagrammatic notation $I = \text{---}$ and $E = \text{---}\circ\text{---}$, so that, for example,

the permutation operator on two vectors is written \times . (The reader is referred to [1] for further details.) Expressing the above diagram in this notation, we obtain

$$K_2^{[0]}(\theta) \propto \left(\overline{\quad} - \times \right) \left(\overline{\quad} + c \left(\theta + \frac{i\pi}{N} \right) \overline{\quad} \right) \left(\overline{\quad} - \frac{N\theta}{i\pi} \times \right) \left(\overline{\quad} + c \left(\theta - \frac{i\pi}{N} \right) \overline{\quad} \right),$$

where $c = \frac{2N}{i\pi(N-2M)}$. Expanding this, up to an overall scalar factor we obtain

$$K_2^{[0]} \propto P_2^A (P_1 + [N - 2M - 2] (P_2 + [N - 2M + 2] P_3))$$

where $P_2^A = \frac{1}{2} \left(\overline{\quad} - \times \right)$ is the projector onto the second rank antisymmetric representation of $SU(N)$ and $P_{1,2,3}$ are the orthogonal projectors

$$P_1 = \frac{1}{4} \left(\overline{\quad} - \overline{\quad} \right) \left(\overline{\quad} - \overline{\quad} \right), \quad P_2 = \frac{1}{2} \left(\overline{\quad} - \overline{\quad} \right) \quad \text{and} \quad P_3 = \frac{1}{4} \left(\overline{\quad} + \overline{\quad} \right) \left(\overline{\quad} + \overline{\quad} \right).$$

$P_2^A P_{1,2,3}$ project onto the irreducible representations $(1, \boxplus)$, (\boxminus, \boxminus) and $(\boxplus, 1)$ respectively, in agreement with the previous section. Note that these explicit expressions for the projectors make it clear how the \mathfrak{h} -representations are embedded into the parent $SU(N)$ representation \boxplus . A similar but more complicated calculation gives the matrix part of $K_3^{[0]}(\theta)$ in the same way. We also have an inductive construction of the matrix part of $K_n^{[0]}$ which reproduces (2.4).

3.2 $G = SU(N)$, $H = Sp(N)$ and $H = SO(N)$

For these cases (in which the ‘conjugated’ reflection equation (2.2) applies) the matrix part E of $K_1^{[0]}$ is constant, and is symmetric in the $SO(N)$ and antisymmetric in the $Sp(N)$ case. We have again performed the fusion calculations of $K_2^{[0]}$ (see also [7]) and $K_3^{[0]}$, and reproduced the results of section two.

Recall that for $H = SO(N)$ all the K -matrices are constant, whereas for $H = Sp(N)$, due to the non-trivial branching rule, extra structure appears. In the fusion calculation, this distinction becomes apparent when P_2^A is contracted with E : for the symmetric, $SO(N)$ case, such terms vanish, whereas for the antisymmetric, $Sp(N)$ case, they give non-trivial contributions.

3.3 Scattering off an excited boundary


We return now to the Grassmannian case $H = S(U(M) \times U(N - M))$ and consider boundary scattering off excited boundary states. It was noted in [1] that the vector particle

boundary scattering matrix $K_1^{[0]}$ has a simple pole at $\theta = \frac{(N-2M)i\pi}{2N} = \theta_0$ corresponding to the formation of a boundary bound state, transforming in $(\square, 1)$.

We determine the reflection matrix $K_1^{[1]}(\theta)$ of the vector multiplet off this state as follows. The bootstrap principle, applied now to the *boundary* bound state, gives

$$K_1^{[1]}(\theta) = (I \otimes K_1^{[0]}(\theta_0)) \check{R}_{11}(\theta + \theta_0) (I \otimes K_1^{[0]}(\theta)) \check{R}_{11}(\theta - \theta_0), \quad (3.2)$$

or, diagrammatically,



This confirms the results of section 2.4, with explicit expressions for the projectors as follows: for $K_1^{[1]}$ we have

$$\begin{aligned} P_{(\square, \square)} &= \frac{1}{4} \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} - \overline{\circ} \right) \\ P_{(\square, 1)} &= \frac{1}{8} \left(\overline{\square} + \overline{\times} \right) \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} + \overline{\circ} \right) \\ P_{(\square, 1)} &= \frac{1}{8} \left(\overline{\square} - \overline{\times} \right) \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} + \overline{\circ} \right). \end{aligned}$$

Proceeding similarly for $K_1^{[2]}$ we obtain

$$\begin{aligned} P_{(\square, \square)} &= \frac{1}{16} \left(\overline{\square} - \overline{\times} \right) \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} - \overline{\circ} \right) \\ P_{(\square, 1)} &= \frac{1}{48} \left(\overline{\square} - \overline{\times} \right) \left(2\overline{\square} + \overline{\times} + \overline{\times} \right) \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} + \overline{\circ} \right) \\ P_{(\square, 1)} &= \frac{1}{48} \left(\overline{\square} - \overline{\times} \right) \left(\overline{\square} - \overline{\times} - \overline{\times} \right) \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} + \overline{\circ} \right), \end{aligned}$$

and for $K_2^{[1]}$

$$\begin{aligned} P_{(\square, \square)} &= \frac{1}{16} \left(\overline{\square} - \overline{\times} \right) \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} - \overline{\circ} \right) \left(\overline{\square} - \overline{\circ} \right) \\ P_{(\square, \square)} &= \frac{1}{16} \left(\overline{\square} - \overline{\times} \right) \left(\overline{\square} + \overline{\times} \right) \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} - \overline{\circ} \right) \left(\overline{\square} + \overline{\circ} \right) \\ P_{(\square, \square)} &= \frac{1}{32} \left(\overline{\square} - \overline{\times} \right) \left(\overline{\square} - \overline{\times} \right) \left(\overline{\square} + \overline{\circ} \right) \left(\overline{\square} - \overline{\circ} \right) \left(\overline{\square} + \overline{\circ} \right) \end{aligned}$$

$$\begin{aligned}
P_{(\mathbb{A},1)} &= \frac{1}{48} \left(\begin{array}{c} \equiv \\ \equiv \\ \equiv \end{array} - \begin{array}{c} \times \\ \times \\ \times \end{array} \right) \left(2 \begin{array}{c} \equiv \\ \equiv \\ \equiv \end{array} + \begin{array}{c} \overline{\times} \\ \overline{\times} \\ \overline{\times} \end{array} + \begin{array}{c} \times \\ \times \\ \times \end{array} \right) \left(\begin{array}{c} \equiv \\ \equiv \\ \equiv \end{array} + \begin{array}{c} \overline{\circ} \\ \overline{\circ} \\ \overline{\circ} \end{array} \right) \left(\begin{array}{c} \equiv \\ \equiv \\ \equiv \end{array} + \begin{array}{c} \overline{\circ} \\ \overline{\circ} \\ \overline{\circ} \end{array} \right) \left(\begin{array}{c} \equiv \\ \equiv \\ \equiv \end{array} + \begin{array}{c} \overline{\circ} \\ \overline{\circ} \\ \overline{\circ} \end{array} \right) \\
P_{(\mathbb{B},1)} &= \frac{1}{48} \left(\begin{array}{c} \equiv \\ \equiv \\ \equiv \end{array} - \begin{array}{c} \times \\ \times \\ \times \end{array} \right) \left(\begin{array}{c} \equiv \\ \equiv \\ \equiv \end{array} - \begin{array}{c} \overline{\times} \\ \overline{\times} \\ \overline{\times} \end{array} - \begin{array}{c} \times \\ \times \\ \times \end{array} \right) \left(\begin{array}{c} \equiv \\ \equiv \\ \equiv \end{array} + \begin{array}{c} \overline{\circ} \\ \overline{\circ} \\ \overline{\circ} \end{array} \right) \left(\begin{array}{c} \equiv \\ \equiv \\ \equiv \end{array} + \begin{array}{c} \overline{\circ} \\ \overline{\circ} \\ \overline{\circ} \end{array} \right) \left(\begin{array}{c} \equiv \\ \equiv \\ \equiv \end{array} + \begin{array}{c} \overline{\circ} \\ \overline{\circ} \\ \overline{\circ} \end{array} \right).
\end{aligned}$$

4 Concluding remarks

The natural next step is to put all these results together to find the full set of boundary S -matrices in the PCM, and thereby the complete spectrum of boundary bound states and their interactions. Fusion calculations rapidly become intractable with increasing rank, but with these in combination with the graphical methods developed here we hope to be able to progress towards the completion of this programme.

There also remains much to be discovered at the classical level – for example, about what happens when a Wess-Zumino term is added to the PCM action, to make contact with work on D-branes in group manifolds. From the mathematical point of view it remains to integrate our results with those on representations of twisted Yangians [8]. It also remains to understand how these results apply in Gross-Neveu models, and their relationship with the results of [9].

An immediate prospect, and work in progress, is to apply similar ideas to the trigonometric case, where the underlying model is affine Toda field theory. We have constructed the remnant of the quantum affine algebra symmetry generated by non-local charges in [10].

Acknowledgments

We thank Evgueni Sklyanin for discussions, and the UK EPSRC for BJS's studentship and GWD's advanced fellowship. NJM thanks the Nuffield foundation for a grant.

References

- [1] N. MacKay and B. Short, *Boundary scattering, symmetric spaces and the principal chiral model on the half-line*, [hep-th/0104212](#)
- [2] J. M. Evans, M. Hassan, N. J. MacKay and A. Mountain, *Local conserved charges in principal chiral models*, *Nucl. Phys.* **B561** (1999) 385, [hep-th/9902008](#)
- [3] D. Bernard, *Hidden Yangians in 2D massive current algebras*, *Commun. Math. Phys.* **137** (1991) 191

- [4] N. J. MacKay, *Rational R-matrices in irreducible representations*, J. Phys. **A24** (1991), 4017
M. D. Gould, Y.-Z. Zhang and A. Bracken, *From representations of the braid group to solutions of the Yang-Baxter equation*, Nucl. Phys. **B354** (1991), 625
G. W. Delius, M. D. Gould, Y.-Z. Zhang, *On the construction of trigonometric solutions of the Yang-Baxter equation*, Nucl. Phys. **B432** (1994), 377; hep-th/9405030
- [5] S. Ghoshal and A. Zamolodchikov, *Boundary S-matrix and boundary state in 2D integrable quantum field theory*, Int. J. Mod. Phys. **A9** (1994), 3841, hep-th/9306002
- [6] L. Mezincescu and R. Nepomechie, *Fusion procedure for open chains*, J. Phys. **A25** (1992), 2533
- [7] N. J. MacKay, *Fusion of $SO(N)$ reflection matrices*, Phys. Lett. **B357** (1995) 89, hep-th/9505124
- [8] A. Molev, M. Nazarov and G. Ol'shanskii, *Yangians and classical Lie algebras*, Russ. Math. Surveys **51** (1996) 205
A. Molev, *Finite-dimensional irreducible representations of twisted Yangians*, J. Math. Phys. **39** (1998), 5559, q-alg/9711022
- [9] M. Mintchev, E. Ragoucy and P. Sorba, *Spontaneous symmetry breaking in the $gl(N)$ -NLS hierarchy on the half-line*, hep-th/0104079
- [10] G. W. Delius and N. J. MacKay, *Non-local charges and quantum group symmetry in sine-Gordon and affine Toda field theories with an integrable boundary*, in preparation.