Nambu–Goto Strings from SU(N) Born-Infeld model

Stefano Ansoldi^a , Carlos Castro^b , E. I. Guendelman^c , Euro Spallucci^a

^aDipartimento di Fisica Teorica, Università di Trieste and INFN, Sezione di Trieste

^bCenter for Theoretical Studies of Physical Systems, Clark Atlanta University, Atlanta, GA.30314

^cPhysics Department, Ben Gurion University, Beer Sheva, Israel

Abstract

The spectrum of quenched Yang-Mills theory in the large–N limit displays strings and higher dimensional extended objects. The effective dynamics of string-like excitations is encoded into area preserving Schild action. In this letter, we bridge the gap between SU(N) gauge models and fully reparametrization invariant Nambu–Goto string models by introducing an extra matrix degree of freedom in the Yang-Mills action. In the large–N limit this matrix variable becomes the world-sheet auxiliary field allowing a smooth transition between the Schild and Nambu–Goto strings. The new improved matrix model we propose here can be extended to p-branes provided we enlarge the dimensionality of the target spacetime.

In the large–N limit SU(N) Yang-Mills gauge theories display string-like excitations [1]. The effective dynamics of these one-dimensional objects is described by a Schild action which is invariant under area preserving reparametrizations only. This result allowed to establish a relationship between $SU(\infty)$ and symplectic transformations: $\sigma^m \to \sigma'^m = \sigma'^m(\sigma)$, $|\frac{\partial \sigma'}{\partial \sigma}| = 1$, but not between $SU(\infty)$ and the group of general reparametrization encoded into the Nambu–Goto action.

In this letter we are going to show how to recover the reparametrization invariant Nambu–Goto action form the large–N limit of a SU(N) gauge invariant action.

¹ e-mail address: ansoldi@trieste.infn.it

 $^{^2\,}$ e-mail address: castro@ctsps.cau.edu

³ e-mail address:guendel@bgumail.bgu.ac.il

⁴ e-mail address: spallucci@trieste.infn.it

As a starting point, let us consider an action smoothly interpolating between the (area preserving) Schild action and the fully (reparametrization invariant) Nambu–Goto action can be written by introducing an auxiliary world– sheet field $\Phi(\sigma)$ [2], :

$$I\left[\Phi, X\right] \equiv \frac{\mu_0}{2} \int_{\Sigma} d^2 \sigma \left[\frac{\det(\gamma_{mn})}{\Phi(\sigma)} + \Phi(\sigma)\right]$$
(1)

where, $\gamma_{mn} \equiv \eta_{\mu\nu} \partial_m X^{\mu} \partial_n X^{\nu}$ is the *induced metric* on the string Euclidean world-sheet $x^{\mu} = X^{\mu}(\sigma)$, $\operatorname{sign}(\gamma_{mn}) = (+, +)$; finally, $\mu_0 \equiv 1/2\pi\alpha'$ is the string tension. We assigned the following dimension (in natural units) to the various quantities in (1):

$$[\sigma^m] = \text{length}, \quad [X^\mu] = \text{length}, \quad [\Phi] = 1$$
 (2)

For later convenience, we recall the relation between $det(\gamma_{mn})$ and the world manifold Poisson Bracket:

$$\det(\gamma_{mn}) = \{ X^{\mu} , X^{\nu} \}^2$$
(3)

$$\{X^{\mu}, X^{\nu}\}_{\rm PB} \equiv \epsilon^{mn} \,\partial_m \,X^{\mu} \,\partial_n \,X^{\nu} \tag{4}$$

The action (1) is reparametrization invariant provided the auxiliary field $\Phi(\sigma)$ transforms as a world-sheet scalar density:

$$\Phi(\sigma') = \left| \frac{\partial \sigma}{\partial \sigma'} \right| \Phi(\sigma)$$
(5)

Thus, by implementing reparametrization invariance $\Phi(\sigma)$ can be transformed to unity and the Schild action can be recovered as a "gauge fixed" form of (1):

$$I\left[\Phi=1,X\right] = \frac{\mu_0}{2} \int_{\Sigma} d^2 \sigma' \left[det(\gamma_{mn}) + 1\right] = \text{Schild} + \text{const.}$$
(6)

where, the numerical constant is proportional to the area of the integration domain Σ .

On the other hand, by solving $\Phi(\sigma)$ in terms of X from (1) one recovers, "on-shell", the Nambu–Goto action

$$\frac{\delta I}{\delta \Phi} = 0 \to \phi = \sqrt{\det(\gamma_{mn})} \to I = \mu_0 \int_{\Sigma} d^2 \sigma \sqrt{-\det(\gamma_{mn})}$$
(7)

The inverse equivalence relation can be proven by starting from the Schild action [3]

$$I_S \equiv \mu_0 \int_{\Xi} d^2 \varphi \det \left[\gamma_{ab}(\varphi) \right]$$
(8)

and "*lifting*" the original world-sheet coordinates φ^m to the role of dynamical variables by mean of reparametrization $\varphi^m \to \sigma^m = \sigma^m(\varphi)$ [4]:

$$I_{rep} \equiv \mu_0 \int_{\Sigma} d^2 \sigma \, \Phi^{-1} \det \left[\gamma_{ab}(\sigma) \right] \,, \qquad \Phi^{-1} \equiv \epsilon_{ij} \, \epsilon^{mn} \, \partial_m \, \phi^i \, \partial_n \, \phi^j \qquad (9)$$

By variation of I_{rep} with respect to φ^i one gets the field equation

$$\epsilon_{ij} \,\epsilon^{mn} \partial_n \,\phi^j \partial_m \,\left(\frac{\det\left[\gamma_{ab}(\sigma)\right]}{\Phi^2}\right) = 0 \tag{10}$$

Thus,

$$\frac{\det\left[\gamma_{ab}(\sigma)\right]}{\Phi^2} = \text{const.} \equiv \frac{1}{4\mu_0} \tag{11}$$

and the Nambu–Goto action (7) is ained again.

This second option introduces a scalar doublet ϕ^i , i = 1, 2 and expresses the scalar density Φ as a "composite" object, rather than a fundamental one, or as a second "integration measure" [5]. In any case the final result is unchanged. The action (1) is a special case of the general two-parameter family of p-brane actions [2]

$$I_{n}^{p} \equiv \frac{\mu_{0}^{(p+1)/2}}{n} \int_{\Sigma} d^{p+1} \sigma \, e(\,\sigma\,) \, \left[\frac{(\det \gamma_{mn})^{n/2}}{e(\,\sigma\,)^{n}} + n - 1 \right]$$
(12)

where, in our notation $e(\sigma)$ has been replaced by $\Phi(\sigma)$ and $\det \gamma_{mn}$ is now the square of the Nambu-Poisson bracket $\{X^{\mu_1}, \ldots, X^{\mu_{p+1}}\}_{NPB}$.

For n = 2 and p = 1 we obtain (1), while for n = 1 the auxiliary field decouples and we get the Nambu–Goto action.

In a naive, but not too much, way of thinking, one could trace back the correspondence between Yang–Mills gauge fields and Schild strings to the common quadratic form of both actions in the respective sets of "field strengths":

$$\frac{1}{4} \{ X^{\mu}, X^{\nu} \}_{\mathrm{PB}} \{ X_{\mu}, X_{\nu} \}_{\mathrm{PB}} \longleftrightarrow \frac{1}{4} \operatorname{Tr} \mathbf{F}_{\mu\nu} \mathbf{F}^{\mu\nu}$$
(13)

Pushing this formal analogy a little forward, and taking into account a possible gauge field type formulation of string dynamics [6], one would expect to find a similar relation between the Nambu–Goto action and a non-Abelian Born–Infeld type action

$$\sqrt{\{X^{\mu}, X^{\nu}\}_{\mathrm{PB}}\{X_{\mu}, X_{\nu}\}_{\mathrm{PB}}} \longleftarrow (?) \longrightarrow \sqrt{\mathrm{Tr}\mathbf{F}_{\mu\nu}\,\mathbf{F}^{\mu\nu}} \qquad (14)$$

While being suggestive, relation (14) suffers from various problems not present in (13), e.g. the very definition of the non-Abelian version of the Born-Infeld action is ambiguous [7]. As the relation (13) can be obtained through several, non-trivial steps, including "quenching", large–N expansion, Wigner-Weyl-Moyal quantization, it is the purpose of this communication to investigate how this approach can be, eventually, extended to the square root type gauge action in (14).

The non-perturbative aspects of the Yang-Mills models are better described by transforming the original gauge field theory into a *Matrix Quantum Mechanics*. Such a transition is realized through dimensional reduction and quenching. The technical steps which allows to "get rid of" the internal, non-Abelian, indices i, j and replace the spacetime coordinates x^{μ} with two continuous coordinates (σ^0 , σ^1), are described in some detail elsewhere [9], and will not be repeated here. For the reader convenience we shall give only sketch the main steps. The general procedure can be summarized as follows.

Take the large–N limit, i.e. let the row and column labels i, j to range over arbitrarily large values. Thus, $SU(N) \rightarrow U(N)$ and the group of spacetime translations fits into the diagonal part of $U(\infty)$. By neglecting off-diagonal components, spacetime dependent dynamical variables can be shifted to the origin by means of a translation operator $\mathbf{U}(x)$: since the translation group is Abelian one can choose the matrix $\mathbf{U}(x)$ to be a plane wave diagonal matrix [8]

$$\mathbf{U}_{ab}(x) = \delta_{ab} \exp\left(iq^a{}_{\mu}x^{\mu}\right) \,, \tag{15}$$

where $q^a{}_{\mu}$ are the eigenvalues of the four-momentum \mathbf{q}_{μ} . Then

$$\mathbf{A}_{\mu}(x) = \exp\left(-i\mathbf{q}_{\mu}x^{\mu}\right)\mathbf{A}_{\mu}(0)\exp\left(i\mathbf{q}_{\mu}x^{\mu}\right) \equiv \mathbf{U}^{\dagger}(x)\mathbf{A}_{\mu}^{(0)}\mathbf{U}(x)$$

and in view of the equality

$$\mathbf{D}_{\mu}\mathbf{A}_{\nu} = i\mathbf{U}^{\dagger}(x)\left[\mathbf{q}_{\mu} + \mathbf{A}_{\mu}^{(0)}, \mathbf{A}_{\nu}\right]\mathbf{U}(x),$$

which when antisymmetrized yields

$$\mathbf{D}_{[\mu}\mathbf{A}_{\nu]} = i\mathbf{U}^{\dagger}(x)\left[\mathbf{q}_{\mu} + \mathbf{A}_{\mu}^{(0)}, \mathbf{q}_{\nu} + \mathbf{A}_{\nu}^{(0)}\right]\mathbf{U}(x) \equiv i\mathbf{U}^{\dagger}(x)\left[\mathbf{A}_{\mu}^{(q)}, \mathbf{A}_{\nu}^{(q)}\right]\mathbf{U}(x),$$

we can see that the translation is compatible with the covariant differentiation, so that

$$\mathbf{F}_{\mu\nu}(x) = \exp\left(-i\mathbf{q}_{\mu}x^{\mu}\right)\mathbf{F}_{\mu\nu}(0)\exp\left(i\mathbf{q}_{\mu}x^{\mu}\right) \equiv \mathbf{U}^{\dagger}(x)\mathbf{F}_{\mu\nu}^{(0)}\mathbf{U}(x) \ .$$

Quenching Approximation amounts to take into account the contributions of the slow modes, described by the eigenvalues of the momentum matrix \mathbf{q} and "integrate out" the non-diagonal fast modes. The final result is to turn the original gauge theory is transformed into a quantum mechanical model where the physical degrees of freedom are carried by large coordinate independent matrices. Finally, the spacetime volume integration is regularized by enclosing the system in a "box" of four-volume V

$$\int d^4x \quad \longrightarrow \quad V$$

In a Yang-Mills framework one can relate V to the QCD scale, i.e. $V = (2\pi/\Lambda_{\rm QCD})^4$ [11]. Here, we shall determined V by matching the large-N limit of our matrix model with the Nambu–Goto action.

The resulting quenched action is

$$S_{\rm YM/\Phi}^{\rm (q)} = \frac{NV}{4g_0^2} \text{Tr}\left(\left[\mathbf{A}_{\mu}^{\rm (q)}, \mathbf{A}_{\nu}^{\rm (q)}\right]^2\right) \ . \tag{16}$$

The large–N quantum properties of the matrix model (16) can be effectively investigated by means of the Wigner-Weyl-Moyal correspondence between matrices and functions, i.e. Symbols, defined over a noncommutative phase space. The resulting theory is a deformation of an ordinary field theory, where the ordinary product between functions is replaced by a noncommutative *-product. The deformation parameter, measuring the amount of non-commutativity, results to be 1/N, and the classical limit corresponds exactly to the large–N limit. The final result is a string action of the Schild type, which is invariant under area-preserving reparametrization of the worldsheet.

More recently, we have also shown that bag-like objects fit the large-N spectrum of Yang-Mills type theories as well, both in four [9] and higher dimensions.

Here, we would like to explore a different route leading in a straightforward way to a Nambu–Goto string action. Having discussed the role of the auxiliary field Φ in bridging the gap between Schild and Nambu–Goto actions at the classical level, we propose the following improved gauge matrix model

$$S_{\rm YM/\Phi}^{\rm (q)} = \frac{N V}{4g_0^2} \text{Tr} \left[\Phi^{-1} \left(\left[\mathbf{A}_{\mu}^{\rm (q)}, \mathbf{A}_{\nu}^{\rm (q)} \right]^2 \right) \right] + \frac{1}{4} \,\text{Tr} \Phi \,\,. \tag{17}$$

where, $\boldsymbol{\Phi}$ is a $N \times N$ diagonal matrix. In a different framework, a similar matrix variable have been introduced to build a consistent path integral for the matrix version of type IIB superstring model [10].

We see, as things stand up to now there seems to be a qualitative difference between equation (17) and equation (9). This is the absence of the analogous to the last term in eq. (17), the one that contains only $Tr(\Phi)$, which is absent in eq.(9). Nevertheless the corresponding demand that the "measure" Φ be a "total derivative" can be implemented in simple ways also in the matrix model. The first way to implement this is as follows: let us take (17) with the first term in the right hand side only, but let us also say that the variation with respect to Φ has to be performed taking into account that Φ is in some sense the anologous of a "total derivative". But then, what is a total derivative in the matrix formulation? One property of a total derivative is that its integral is is fixed from the boundaries, which are not varied. Let us say for simplicity that we take the integral to be zero (fixing it to another constant will not change anything). In this case we must proceed as follows:

- (1) Take the action as in (17), but only consider the first term in the right hand side.
- (2) Consider the variation of such action, but with the constraint that Φ is a total derivative, which means that $Tr(\Phi) = 0$.
- (3) To do this in practice we add to the action defined in (1) $c Tr(\Phi)$, where c is an undetermined lagrange multiplier which implements condition (2). The resulting theory will have then an arbitrary string tension if we continue from (2). The constant c, i.e. the undetermined lagrange multiplier playing the role of the constant of integration in (11).

Yet another way of proceeding is to explicitly construct a composite measure in the matrix formulation (instead of using a property a total derivative must satisfy and then implement this). The procedure in this case would be:

- i) the world sheet coordinates are replaced by matrices ξ^m ;
- ii) the derivatives are replaced by commutators between the matrix coordinates and the matrix momentum q_a .

The end result is a matrix $\Phi : \Phi = \epsilon^{ab} \epsilon_{mn} [q_a, \xi^m] [q_b, \xi^n]$ which should be used in (17), except that now the second term in (17) and/or a possible lagrange multiplier is now not necessary (the constraint $\text{Tr}(\Phi) = 0$ is now an identity). Now, going back to the action (17), if Φ is taken to be the $N \times N$ *Identity* matrix, then action (17) takes on the form of the Yang-Mills action. On the other hand, if we let Φ to be diagonal, $\Phi_{ab} \equiv \phi_{(a)} \delta_{ab}$ (no summation over the index *a*), and determine its eigenvalues by varying (17), we find

$$\phi_{(a)} = \sqrt{\frac{NV}{g_0^2}} \mathbf{F^{(0)}}_{cd\,\mu\nu} \mathbf{F}^{(0)\,cd\,\mu\nu} \equiv \sqrt{\frac{NV}{g_0^2}} \operatorname{Tr} \mathbf{F^{(0)}}_{\mu\nu} \mathbf{F}^{(0)\,\mu\nu}$$
(18)

All the eigenvalues are degenerate and quadratic in the Yang-Mills field strength. By inserting (18) into (17) one finds:

$$S_{\rm BI-YM}^{\rm (q)} = \sqrt{\frac{NV}{4g_0^2} \mathbf{F}_{cd\,\mu\nu}^{(0)} \mathbf{F}^{(0)cd\,\mu\nu}} \equiv \sqrt{\frac{NV}{4g_0^2} \text{Tr} \mathbf{F}_{\mu\nu}^{(0)} \mathbf{F}^{(0)\mu\nu}}$$
(19)

The action (19) is the "square root" form of a non-Abelian Born-Infeld type action, where the trace is the standard one. The ambiguity in the definition of the trace over internal indices is removed, in our model, by choosing Φ to be diagonal. With hindsight, we know that in the large-N limit the trace over internal SU(N) indices will turn into an integration over world manifold coordinates. Thus, it is compelling to "move out" the trace from the square root, in order to obtain a Nambu–Goto type action integral. What we are going to describe is a procedure such that in the large-N limit:

$$\sqrt{\mathbf{Tr}\left(\ldots\right)} \longrightarrow \int_{\Sigma} d^2 \sigma \sqrt{\ldots}$$
(20)

and the non-commuting Yang–Mills matrices are replaced by commuting string coordinates.

The string world-sheet is the target spacetime image $x^{\mu} = X^{\mu}(\sigma^0, \sigma^1)$ of the world manifold $\Sigma : -\infty \leq \sigma^0 \leq +\infty, 0 \leq \sigma^1 \leq L, X^{\mu}$ belonging to the algebra \mathcal{A} of C^{∞} , of functions over Σ . Thus, to realize our program we must *deform* \mathcal{A} to a *non-commutative "starred" algebra* by introducing a *-product. The general rule is to define the new product between two functions as (for a recent review see [12]):

$$f * g = f g + \hbar P_{\hbar}(f, g) , \qquad (21)$$

where $P_{\hbar}(f, g)$ is a bilinear map $P_{\hbar} : \mathcal{A} \times \mathcal{A} \to \mathcal{A}$. \hbar is the *deformation* parameter, which, in our case, is defined as $\hbar \equiv 2\pi/N$. The Moyal product is defined as the deformed *-product

$$f(\sigma) * g(\sigma) \equiv \exp\left[i\frac{\hbar}{2}\omega^{mn}\frac{\partial^2}{\partial\sigma^m\partial\xi^n}\right]f(\sigma)g(\xi)\Big|_{\xi=\sigma},\qquad(22)$$

where ω^{mn} is a non-degenerate, antisymmetric matrix, which can be locally written as

$$\omega^{mn} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} . \tag{23}$$

The Moyal product (22) takes a simple looking form in Fourier space

$$F(\sigma) * G(\sigma) = \int \frac{d^2\xi}{(2\pi)} \exp\left(i\frac{\hbar}{2}\omega_{mn}\sigma^m\xi^n\right) F\left(\frac{\sigma}{2} + \xi\right) G\left(\frac{\sigma}{2} - \xi\right) , \quad (24)$$

where F and G are the Fourier transform of f and g. Let us consider the Heisenberg algebra

$$[\mathbf{K}, \mathbf{P}] = i\,\hbar\;; \tag{25}$$

Weyl suggested, many years ago, how an operator $\mathbf{O}_F(\mathbf{K}, \mathbf{P})$ can be written as a sum of algebra elements as

$$\mathbf{O}_F = \frac{1}{(2\pi)} \int dp \, dk \, F\left(p, k\right) \exp\left(ip \, \mathbf{K} + ik \mathbf{P}\right) \,. \tag{26}$$

The Weyl map (26) can be inverted to associate functions, or more exactly *symbols*, to operators

$$F(q,k) = \int \frac{d\xi}{(2\pi)} \exp\left(-ik\xi\right) \left\langle q + \hbar \frac{\xi}{2} \middle| \mathbf{O}_F(\mathbf{K},\mathbf{P}) \middle| q - \hbar \frac{\xi}{2} \right\rangle; \quad (27)$$

moreover it translates the commutator between two operators \boldsymbol{U} , \boldsymbol{V} into the *Moyal Bracket* between their corresponding symbols $\mathcal{U}(\sigma)$, $\mathcal{V}(\sigma)$

$$\frac{1}{i\hbar} \left[\boldsymbol{U}, \boldsymbol{V} \right] \longleftrightarrow \left\{ \boldsymbol{\mathcal{U}}, \boldsymbol{\mathcal{V}} \right\}_{\mathrm{MB}} \equiv \frac{1}{i\hbar} \left(\boldsymbol{\mathcal{U}} \ast \boldsymbol{\mathcal{V}} - \boldsymbol{\mathcal{V}} \ast \boldsymbol{\mathcal{U}} \right)$$

and the quantum mechanical trace into an integral over Fourier space. A concise but pedagogical introduction to the deformed differential calculus and its application to the theory of integrable system can be found in [13]. We are now ready to formulate the alleged relationship between the quenched model (19) and string model: the symbol of the matrix $\mathbf{A}^{(q)}_{\mu}$ is proportional to the string coordinates $X^{\mu}(\sigma^0, \sigma^1)$. Going through the steps discussed above the action $S^{(q)}_{\text{YM/}\Phi}$ transforms into its symbol $W^{(q)}_{\text{YM/}\Phi}$:

$$S_{\rm YM/\Phi}^{\rm (q)} \to W_{\rm YM/\Phi}^{\rm (q)} = \frac{NV}{8\pi g_0^2} \int_{\Sigma} d^2\sigma \left[\frac{1}{\Phi(\sigma)} * \{ A_{\mu}(\sigma) , \mathbf{A}_{\nu}(\sigma) \}_{\rm MB}^2 + \frac{g_0^2}{NV} \Phi(\sigma) \right] . (28)$$

and we rescale the Yang–Mills charge and field $^5\,$ as

 $[\]frac{1}{5}$ For the sake of clarity, let us summarize the canonical dimensions in natural units

$$\frac{N}{g_0^2} \longrightarrow \frac{1}{g^2}, \qquad \text{Tr} \longmapsto \frac{1}{2\pi\sqrt{V}} \int_{\Sigma} d^2\sigma$$
 (29)

$$A^{\mu} \longmapsto V^{-1/2} X^{\mu} , \qquad \mathbf{F}^{(0)\mu\nu} \longmapsto V^{-1/2} \left\{ X^{\mu} , X^{\nu} \right\}_{\mathrm{MB}} . \tag{30}$$

Finally, if $N\gg 1$ the Moyal bracket can be approximated by the Poisson bracket

$$\{X^{\mu}, X^{\nu}\}_{\mathrm{MB}} \longmapsto \{X^{\mu}, X^{\nu}\}_{\mathrm{PB}}$$

and (28) takes the form

$$W_{\rm YM/\Phi}^{\rm (q)} \to S_{\rm NG} = \frac{1}{8\pi g^2 \sqrt{V}} \int_{\Sigma} d^2\sigma \left[\frac{1}{\Phi(\sigma)} \left\{ X_{\mu}(\sigma) , X_{\nu}(\sigma) \right\}_{\rm PB}^2 + g^2 \Phi(\sigma) \right] (31)$$

which is (1) provided we identify

$$\mu_0 \longrightarrow \equiv \frac{1}{4\pi \, g \sqrt{V}} \tag{32}$$

According with the initial discussion we can establish the following, large N, correspondence:

$$S_{\rm BI-YM}^{\rm (q)} \approx S_{\rm YM/\Phi}^{\rm (q)} \to \frac{1}{4\pi \, g \, \sqrt{V}} \int_{\Sigma} d^2 \sigma \, \sqrt{-det(\gamma_{mn})} \tag{33}$$

As a concluding remark, it may be worth mentioning that the approach discussed above can be extended in a straightforward way to a more general non-Abelian Born-Infeld action including topological terms. Instead of starting from (17), one can consider

$$S_{\rm YM/\Phi}^{\rm (q)} = \frac{M^4 V}{2} \operatorname{Tr} \left\{ \mathbf{\Phi}^{-1} \left[\mathbf{I} + \frac{N}{4g_0^2 M^4} \mathbf{F}_{\mu\nu} \mathbf{F}_{\mu\nu} + \left(\frac{N}{4g_0^2 M^4} \right)^2 \left(\epsilon^{\lambda\mu\nu\rho} \mathbf{F}_{\lambda\mu} \mathbf{F}_{\nu\rho} \right)^2 \right] - \mathbf{I} \right\} + \frac{M^4 V}{2} \operatorname{Tr} \mathbf{\Phi} .$$
(34)

where, **I** is the $N \times N$ identity matrix and M a new mass scale. The coefficient in front of the topological density has been assigned in analogy to the Abelian of various quantities:

$$[A_{\mu}{}^{a}(x)] \equiv [\mathbf{A}_{\mu}^{(q)}] = (\text{length})^{-1} , \qquad [F_{\mu\nu}{}^{a}(x)] \equiv [\mathbf{F}_{\mu\nu}{}^{(q)}] = (\text{length})^{-2} [g_{0}] \equiv [g] = (\text{length})^{0} = 1 , \qquad [V] = (\text{length})^{4} , \qquad [\mu_{0}] = (\text{length})^{-2} .$$

case, where no trace ambiguity occurs and the square root form is derived by expanding the determinant of the matrix $M_{\mu\nu} \equiv \delta_{\mu\nu} + \text{const.} \times F_{\mu\nu}$. The topological density contribution trivially vanishes in the large-N limit:

$$\epsilon^{\lambda\mu\nu\rho} \mathbf{F}_{\lambda\mu} \mathbf{F}_{\nu\rho} \longrightarrow \epsilon^{\lambda\mu\nu\rho} \{ X_{\lambda}(\sigma) , X_{\mu}(\sigma) \}_{\mathrm{PB}} \{ X_{\nu}(\sigma) , X_{\rho}(\sigma) \}_{\mathrm{PB}} \equiv 0$$
(35)

A non-vanishing contribution from the topological term can only be obtained if strings are replaced by higher dimensional extended objects [9]. Thus, as far as strings are concerned, one finds

$$S_{\rm YM/\Phi}^{\rm (q)} \to S_{\rm BI}^{\rm (q)} = \frac{M^4 \sqrt{V}}{2\pi} \int_{\Sigma} d^2 \sigma \left[\sqrt{\left(1 + \frac{1}{4g^2 M^4 V} \det[\gamma_{mn}] \right)} - 1 \right] (36)$$

Recent results about UV/IR interplay, in the framework of noncommutative Yang-Mills theories [15], suggest to investigate the behavior of the action (36) by changing the size of the quantization volume V. In the *small vol*ume limit, i.e. $V \ll 1/4g^2 M^4$, the main contribution to $S_{\rm BI}^{(q)}$ comes from the det[γ_{mn}] and the Nambu–Goto action is recovered again. The quantization volume drops out and the string tension, i.e. $\mu_0 = M^2/4\pi g$, is determined by the only relevant mass scale M.

In the opposite, *large volume*, limit, i.e. $V >> 1/4g^2 M^4$, the second term in the square root is small with respect to 1 and the first non-vanishing contribution of the Taylor expansion is the Schild action. In this regime the relevant energy scale is set $V^{-1/4}$ and the corresponding string tension is $\mu_0 = 1/32\pi g^2 \sqrt{V}$.

From a different point of view, Fairlie has recently pointed out some intriguing analogy between the Born-Infeld and Nambu–Goto actions [14]. Our results supports this connection. We believe we bridged the gap between four dimensional gauge theories and fully reparametrization invariant string models. We found new, non-trivial, relationship between a class of generalized SU(N)models and Born-Infeld/Nambu–Goto strings, provided a new matrix degree of freedom, $\mathbf{\Phi}$, is introduced. In the original SU(N) model $\mathbf{\Phi}$ connects the Yang-Mills phase, at $\mathbf{\Phi} = \mathbf{I}$, with the, non-linear, Born-Infeld phase, where the eigenvalues $\phi_{(a)}$ are given by equation (18). In the large–N limit the order parameter becomes a world-sheet auxiliary field linearizing a square root type Born-Infeld string model. The small, respectively, large volume limits of this model correspond to Nambu–Goto and Schild actions.

References

I. Bars Phys. Lett. **245B** 35 (1990)
 E.G. Floratos, J. Illiopulos and G. Tiktopoulos Phys. Lett. **B 217** 285 (1989)

D. Fairlie and C.K. Zachos Phys. Lett. **B224** 101 (1989)

- [2] I. Oda Chaos, Sol. & Fract. 10, n.2/3, 483 (1999)
 S. Ansoldi, C. Castro, E. Spallucci Class. Quant. Grav. L97 (2001)
- [3] A. Schild Phys. Rev. D16 1722 (1977)
- [4] S. Ansoldi, A. Aurilia, E. Spallucci Phys. Rev. D53 870 (1996)
- [5] E. I. Guendelman Class. Quant. Grav. 17 3673 (2000)
 E. I. Guendelman Phys. Rev. D 63 046006 (2001)
- [6] A. Aurilia, A. Smailagic, E. Spallucci Phys. Rev. D47 2536 (1993)
- [7] T. Hagiwara, J. Phys. A14, 3059 (1981)
 A. A. Tseytlin Nucl. Phys. B501 41 (1997)
 J. H. Park Phys. Lett. B 458 471 (1999)
- [8] Y. Kitazawa, S.R. Wadia Phys. Lett. **120B** 377 (1983)
- [9] S. Ansoldi , C. Castro, E. Spallucci Phys. Lett. B504 174 (2001)
 S. Ansoldi , C. Castro, E. Spallucci Class. Quant. Grav. 18 L17 (2001)
- [10] A. Fayyazuddin, Y. Makeenko, P. Olesen, D.J. Smith, K. Zarembo Nucl. Phys. B499 159 (1997)
 for a review see Y. Makeenko Three Introductory Lectures in Helsinki on Matrix Models of Superstrings, hep-th/9704075 (1997)
- [11] S. Ansoldi, C. Castro, E.Spallucci Class. Quantum Grav. 18 2865 (2001)
- [12] Harvey J A Komaba Lectures on Noncommutative Solitons and D-Branes, hepth/0102076 (2001)
- [13] I. A. B. Strachan J. Geom. and Phys. 21 255 (1997)
- [14] D. B. Fairlie Phys. Lett. **B456**, 141 (1999)
- [15] R. J. Szabo Quantum Field Theory on Noncommutative Spaces, hep-th/0109162 (2001)