

On the β -Function and Conformal Anomaly of Noncommutative QED with Adjoint Matter Fields

Néda Sadooghi ^{†,a} and Mojtaba Mohammadi ^{‡,b}

^{†,‡} *Department of Physics, Sharif University of Technology*

P.O. Box 11365-9161, Tehran-Iran

and

[†] *Institute for Studies in Theoretical Physics and Mathematics (IPM)*

School of Physics, P.O. Box 19395-5531, Tehran-Iran

Abstract

In the first part of this work, a perturbative analysis up to one-loop order is carried out to determine the one-loop β -function of noncommutative $U(1)$ gauge theory with matter fields in the adjoint representation. In the second part, the conformal anomaly of the same theory is calculated using the Fujikawa's path integral method. The value of the one-loop β -function calculated in both methods coincides. As it turns out, noncommutative QED with matter fields in the adjoint representation is asymptotically free for the number of flavor degrees of freedom $N_f < 3$.

PACS No.: 11.15.Bt, 11.10.Gh, 11.25.Db

Keywords: Noncommutative Field Theory, Conformal Anomaly, β -function

^aElectronic address: sadooghi@theory.ipm.ac.ir

^bElectronic address: m.mohammadi@mehr.sharif.edu

1 Introduction

In the past few years, the noncommutative gauge theories have attracted much attention and are studied seriously by many authors [1]. They are interesting due to their realization in the String Theory. In the decoupling limit, noncommutative gauge theories can occur in the world volume of D-branes in the presence of the constant background $B_{\mu\nu}$ -field [2]. But, noncommutative gauge theories can also be regarded as nontrivial deformation of the ordinary commutative gauge theories. This deformation is simply realized by replacing the ordinary commutative product of functions by the noncommutative \star -product:

$$f(x) \star g(x) \equiv e^{\frac{i\theta_{\mu\nu}}{2} \frac{\partial}{\partial \xi_\mu} \frac{\partial}{\partial \zeta_\nu}} f(x + \xi) g(x + \zeta) \Big|_{\xi=\zeta=0},$$

where $\theta_{\mu\nu}$ is a real, and antisymmetric tensor, reflecting the noncommutativity of the space-time coordinates.

The most natural candidate for such noncommutative gauge theories is the noncommutative QED, where the coupling of the matter fields to the gauge fields can be given in the fundamental, antifundamental or adjoint representations.

Different aspects of noncommutative $U(1)$ gauge theory with *fundamental* matters are studied intensively in the literature, and various results are achieved [for a review, see Ref. [1] and the references therein]. A perturbative analysis of the one-loop β -function of this theory shows, that for the number of flavor degrees of freedom $N_f < 6$, the NC- $U(1)$ gauge theory is asymptotically free [3]. Asymptotically free theories appear also in the ordinary commutative non-Abelian gauge theories. Commutative $SU(3)$, *e.g.*, exhibits asymptotic freedom for $N_f \leq 16$.

Recently, the same one-loop β -function of noncommutative QED with *fundamental* matter fields is also calculated by studying the relation between the conformal anomaly of noncommutative QED and the β -function of the theory [see Ref. [4]]. As is known, noncommutative gauge theories include the noncommutativity parameter θ of dimension [length]². It is therefore expected, that the Weyl symmetry of noncommutative gauge theories is broken at classical level, even in the massless limit. But, as in the ordinary commutative case, the Weyl symmetry of noncommutative $U(1)$ is also broken by quantum corrections. This is defined as the conformal anomaly of noncommutative QED. These quantum corrections can be calculated using, *e.g.*, the Fujikawa's path integral method [5], which is also used to calculate the global and gauge anomalies of noncommutative $U(1)$ and $U(N)$ gauge theories [6]. As it is shown in Ref. [4], the value of the one-loop β -function of noncommutative QED with fundamental matters calculated from the conformal anomaly coincides with the result of perturbative analysis from Ref. [3].

What concerns the noncommutative QED with *adjoint* matter field coupling, there are good reasons to study these theories too. As it is argued by Seiberg in Ref. [7], noncommutative Yang-Mills theory appears as a manifestation of a matrix model and the underlying model forces the coupling of the matter to the gauge fields to be in the adjoint representation of noncommutative algebra. Noncommutative QED with matter fields in the adjoint representation is first studied in Ref. [8], where the one-loop vacuum polarization tensor of the theory was calculated in the framework of perturbation theory. It is shown, that in noncommutative gauge theories with matter fields in the adjoint representations, nonplanar Feynman diagrams appear which exhibit the important UV/IR mixing phenomena of noncommutative Field Theory [9]. The study of noncommutative gauge theories with adjoint matter fields is also interesting from the point of view of noncommutative standard model and a possible θ -dependent correction to the neutrino mass [10, 11].

In this letter, the one-loop β -function of NC- $U(1)$ gauge theory with adjoint matter fields will be calculated by making use of two different methods. We will first introduce the noncommutative QED with adjoint matter fields and its corresponding Feynman rules in Sect. 2. Then, in Sect. 3, a perturbative analysis up to one-loop order will be performed to calculate the one-loop β -function of the theory. In Sect. 4, the conformal anomaly of noncommutative QED with adjoint matter fields will be calculated using the Fujikawa's path integral method [5]. As in the fundamental case, the value of the one-loop β -function coincides in both methods. We will show that, noncommutative QED with adjoint matter fields is asymptotically free for $N_f < 3$. Sect. 5, is devoted to discussions.

2 Noncommutative QED with Adjoint Matter Fields

The action of noncommutative QED with matter fields in the adjoint representation is given by:

$$S[\psi, \bar{\psi}; A_\mu] = S_{gauge}[A_\mu] + S_{matter}[\psi, \bar{\psi}] + S_{ghost}[c, \bar{c}], \quad (2.1)$$

with the gauge part

$$S_{gauge}[A_\mu] = -\frac{1}{4} \int d^D x F_{\mu\nu}(x) \star F^{\mu\nu}(x). \quad (2.2)$$

The field strength tensor is defined by:

$$F_{\mu\nu}(x) = \partial_\mu A_\nu(x) - \partial_\nu A_\mu(x) - ig[A_\mu(x), A_\nu(x)]_\star. \quad (2.3)$$

The matter field part of the action is given by

$$S_{matter}[\psi, \bar{\psi}] = \int d^D x \left[i\bar{\psi}(x)\gamma^\mu \star D_\mu\psi(x) - m\bar{\psi}(x) \star \psi(x) \right], \quad (2.4)$$

with the covariant derivative

$$D_\mu \psi(x) = \partial_\mu \psi(x) - ig[A_\mu(x), \psi(x)]_\star. \quad (2.5)$$

Finally the ghost part of the action reads

$$S_{ghost} = \int d^D x \bar{c}(x) \star (-\partial_\mu D^\mu) c(x), \quad (2.6)$$

where $c(x)$ are the ghost fields in the adjoint representation. The covariant derivative in Eq. (2.6) is defined by the derivative from Eq. (2.5). The Feynman rules of noncommutative QED with adjoint matter fields can be calculated from the above actions and read [see also Ref. [8]]:

i) *Fermion Propagator:*

$$\begin{array}{c} \bullet \text{-----} \bullet \\ p \end{array} \quad S(p) = \frac{i}{\not{p} - m + i\varepsilon}. \quad (2.7)$$

ii) *Photon Propagator in the Feynman Gauge*

$$\begin{array}{c} \bullet \text{-----} \bullet \\ \mu \quad k \quad \nu \end{array} \quad D_{\mu\nu}(k) = \frac{-ig_{\mu\nu}}{k^2}. \quad (2.8)$$

iii) *Ghost Propagator*

$$\begin{array}{c} \bullet \text{-----} \bullet \\ p \end{array} \quad G(p) = \frac{i}{p^2 + i\varepsilon}. \quad (2.9)$$

iv) *The Vertex of Two Fermions and One Gauge Field*

$$\begin{array}{c} \begin{array}{c} \text{wavy line } k_1, \mu \\ \swarrow \quad \searrow \\ \text{fermion } p_1 \quad \text{fermion } p_2 \end{array} \end{array} \quad V_\mu(p_1, p_2; k_1) = (2\pi)^4 \delta^4(p_1 + p_2 + k_1) 2g\gamma_\mu \sin(p_1 \times p_2). \quad (2.10)$$


v) *The Vertex of Two Ghost Fields and One Gauge Field*

$$\begin{array}{c} \begin{array}{c} \text{wavy line } k_1, \mu \\ \swarrow \quad \searrow \\ \text{ghost } p_1 \quad \text{ghost } p_2 \end{array} \end{array} \quad G_\mu(p_1, p_2; k_1) = (2\pi)^4 \delta^4(p_1 + p_2 + k_1) 2g(p_2)_\mu \sin(p_1 \times p_2). \quad (2.11)$$

vi) *Three-Photon Vertex*

$$\begin{array}{c} \begin{array}{c} \text{wavy line } p_1, \mu_1 \\ \swarrow \quad \searrow \\ \text{wavy line } p_2, \mu_2 \quad \text{wavy line } p_3, \mu_3 \end{array} \end{array} \quad W_{\mu_1\mu_2\mu_3}(p_1, p_2, p_3) = -(2\pi)^4 \delta^4(p_1 + p_2 + p_3) 2g \sin(p_1 \times p_2) \\ \times \left[g_{\mu_1\mu_2}(p_1 - p_2)_{\mu_3} + g_{\mu_1\mu_3}(p_3 - p_1)_{\mu_2} + g_{\mu_3\mu_2}(p_2 - p_3)_{\mu_1} \right]. \quad (2.12)$$

vii) *Four-Photon Vertex*



$$\begin{aligned}
 W_{\mu_1\mu_2\mu_3\mu_4}(p_1, p_2, p_3, p_4) = & -4ig^2 (2\pi)^4 \delta^4(p_1 + p_2 + p_3) \\
 & \times \left[(g^{\mu_1\mu_3} g^{\mu_2\mu_4} - g^{\mu_1\mu_4} g^{\mu_2\mu_3}) \sin(p_1 \times p_2) \sin(p_3 \times p_4) \right. \\
 & + (g^{\mu_1\mu_4} g^{\mu_2\mu_3} - g^{\mu_1\mu_2} g^{\mu_3\mu_4}) \sin(p_3 \times p_1) \sin(p_2 \times p_4) \\
 & \left. + (g^{\mu_1\mu_2} g^{\mu_3\mu_4} - g^{\mu_1\mu_3} g^{\mu_2\mu_4}) \sin(p_1 \times p_4) \sin(p_2 \times p_3) \right]. \quad (2.13)
 \end{aligned}$$

Here we have used the notation: $p_1 \times p_2 \equiv \frac{1}{2}\theta_{\mu\nu}p_1^\mu p_2^\nu$.

3 Perturbative Calculation of the One-Loop β -Function

In this section, we present the result of the perturbative calculation of the one-loop β -function of noncommutative U(1) gauge theory with adjoint matter field coupling. But, let us briefly review the result of the one-loop β -function of noncommutative U(1) with matter fields in the fundamental representation, which is first presented in Ref. [3].

i) One Loop β -Function of NC-U(1) with Matter Fields in the Fundamental Representation

The one-loop β -function receives contributions from one-loop fermion self-energy, vacuum polarization tensor and vertex function. In Figs. [1]-[3], all one-loop diagrams contributing to these functions are presented. To calculate the one-loop β -function, we would like to consider only the contributions from the planar Feynman integrals corresponding to the fermion and photon self-energy and vertex function. The nonplanar integrals are assumed to be finite for finite noncommutativity parameter θ and do not contribute to the β -function of the theory [3]. The β -function of the theory is therefore not altered by the UV/IR mixing phenomena [9].

As in the commutative QED, the β -function of the theory is given by

$$\beta(g) = \mu \frac{\partial}{\partial \mu} g(\mu). \quad (3.1)$$

In terms of the bare coupling constant g_0 and the renormalization constants $Z_i, i = 1, 2, 3$, the renormalized coupling constant $g(\mu)$ is defined by:

$$g_0 = \mu^{-\epsilon/2} g(\mu) Z_1 Z_2^{-1} Z_3^{-1/2}, \quad (3.2)$$

with $\epsilon = 4 - D$. In one-loop order, the renormalization constant Z_1 receives contribution from one-loop vertex function, whereas one-loop fermion and photon self-energy diagrams contribute to Z_2 and

Z_3 , respectively. After regulating the planar part of the corresponding one-loop Feynman integrals to these diagrams, using *e.g.* dimensional regularization, the one-loop contribution to $Z_i, i = 1, 2, 3$ read

$$\begin{aligned} Z_1^{Fund.} &= 1 - \frac{g^2}{16\pi^2\epsilon}, \\ Z_2^{Fund.} &= 1 - \frac{g^2}{8\pi^2\epsilon}, \\ Z_3^{Fund.} &= 1 + \frac{g^2}{16\pi^2\epsilon} \left(\frac{20}{3} - \frac{8N_f}{3} \right). \end{aligned} \quad (3.3)$$

Now using Eqs. (3.2) and (3.3), the one-loop β -function of noncommutative QED with the matter fields in the fundamental representation can be given by:

$$\beta(g) \Big|_{1-loop}^{Fund.} = -\frac{g^3}{16\pi^2} \left(\frac{22}{3} - \frac{4N_f}{3} \right), \quad (3.4)$$

where N_f is the number of the flavor degrees of freedom. For $N_f < 6$ the noncommutative QED with fundamental matters is asymptotically free.

ii) One Loop β -Function of NC-U(1) with Matter Fields in the Adjoint Representation

Let us now consider the noncommutative QED with adjoint matter fields coupling. Using the propagators and vertices of noncommutative QED from Eqs. (2.7)-(2.13), the Feynman integral corresponding to the one-loop fermion self energy from Fig. 1 can be given by

$$-i\Sigma(p) = -4g^2 \int \frac{d^D k}{(2\pi)^D} \frac{\gamma^\mu (\not{k} + m) \gamma_\mu}{(k^2 - m^2)(p - k)^2} \sin^2\left(\frac{p \times k}{2}\right). \quad (3.5)$$

Using now the relation $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$, the above integral can be separated into a planar and a nonplanar part. The planar part reads:

$$-i\Sigma_{planar} = -2g^2 \int \frac{d^D k}{(2\pi)^D} \frac{\gamma^\mu (\not{k} + m) \gamma_\mu}{(k^2 - m^2)(p - k)^2}. \quad (3.6)$$

It is exactly twice the value of the one-loop fermion self energy from commutative QED. The wave function renormalization constant Z_2 is then given by:

$$Z_2^{Adj.} = 1 - \frac{g^2}{4\pi^2} \frac{1}{(4 - D)}. \quad (3.7)$$

The one-loop photon self energy receives contributions from four diagrams (2.a)-(2.d) from Fig. 2. It turns out that diagrams (2.a)-(2.c) have both planar and nonplanar parts, whereas diagram (2.d) has only a finite nonplanar part. We will consider only the planar parts of the corresponding Feynman integrals, because they are the only relevant parts to calculate the one-loop β -function of the theory. The planar parts of the diagrams (2.a)-(2.c) are given by:

$$i\Pi_{a-planar}^{\mu\nu} = -2g^2 \int \frac{d^D k}{(2\pi)^D} \frac{\text{Tr}(\gamma^\mu (\not{k} - \not{p} + m) \gamma^\nu (\not{k} + m))}{(k^2 - m^2)(k - p)^2},$$

$$\begin{aligned}
i\Pi_{b\text{-planar}}^{\mu\nu} &= -2g^2 \int \frac{d^D k}{(2\pi)^D} \frac{k^\mu k^\nu - k^\mu p^\nu}{k^2(k-p)^2}, \\
i\Pi_{c\text{-planar}}^{\mu\nu} &= +2g^2 \int \frac{d^D k}{(2\pi)^D} \frac{(D-6)p^\mu p^\nu - (2D-3)(k^\mu p^\nu + k^\nu p^\mu - 2k^\mu k^\nu) + g^{\mu\nu}(5p^2 - 2p \cdot k + 2k^2)}{k^2(k-p)^2}.
\end{aligned} \tag{3.8}$$

Using dimensional regularization, it is possible to isolate the UV-divergent part of the above integrals, which is given by

$$i \left[\Pi_{a\text{-planar}}^{\mu\nu} + \Pi_{b\text{-planar}}^{\mu\nu} + \Pi_{c\text{-planar}}^{\mu\nu} \right]_{\text{div.}} = \frac{ig^2}{16\pi^2} (p^\mu p^\nu - g^{\mu\nu} p^2) \left(\frac{-20}{3} + \frac{16N_f}{3} \right) \frac{1}{(4-D)}. \tag{3.9}$$

The wave function renormalization constant Z_3 can then be obtained by making use of standard field theoretical definitions and reads:

$$Z_3^{\text{Adj.}} = 1 + \frac{g^2}{16\pi^2} \left(\frac{20}{3} - \frac{16N_f}{3} \right) \frac{1}{(4-D)}. \tag{3.10}$$

As next, let us consider the one-loop correction to the vertex function, which receives contribution from two diagrams (3.a) and (3.b) from Fig. 3. The corresponding Feynman integrals to diagram (3.a) is given by:

$$\begin{aligned}
2g \sin\left(\frac{p \times p'}{2}\right) \Gamma_a^\mu(p, p', q = p' - p, m) = \\
+ 8ig^3 \int \frac{d^D k}{(2\pi)^D} \frac{\gamma^\alpha(\not{k} + m)\gamma^\mu(\not{k} + \not{q} + m)\gamma_\alpha}{(k^2 - m^2)((k+q)^2 - m^2)(p-k)^2} \sin\left(\frac{p \times k}{2}\right) \sin\left(\frac{(q+k) \times p'}{2}\right) \sin\left(\frac{k \times q}{2}\right),
\end{aligned} \tag{3.11}$$

and the one-loop Feynman integral corresponding to diagram (3.b) reads

$$\begin{aligned}
2g \sin\left(\frac{p \times p'}{2}\right) \Gamma_b^\mu(p, p', q = p' - p, m) = \\
- 8ig^3 \int \frac{d^D k}{(2\pi)^D} \frac{\gamma_\alpha(\not{k} + m)\gamma_\eta[(p - 2p' + k)^\alpha g^{\mu\eta} + (p + p' - 2k)^\mu g^{\beta\eta} + (k + p' - 2p)^\eta g^{\mu\beta}]}{(k^2 - m^2)(p-k)^2(p'-k)^2} \\
\times \sin\left(\frac{p \times k}{2}\right) \sin\left(\frac{k \times p'}{2}\right) \sin\left(\frac{(k-p') \times q}{2}\right).
\end{aligned} \tag{3.12}$$

Using again some standard trigonometrical relations, the planar and the nonplanar parts of these integrals can be separated. The planar parts are given by

$$\Gamma_{a\text{-planar}}^\mu(p, p', q = p' - p, m) = g^2 \int \frac{d^D k}{(2\pi)^D} \frac{\gamma^\alpha(\not{k} + m)\gamma^\mu(\not{k} + \not{q} + m)\gamma_\alpha}{(k^2 - m^2)((k+q)^2 - m^2)(p-k)^2}, \tag{3.13}$$

for diagram (3.a) and

$$\begin{aligned}
\Gamma_{b\text{-planar}}^\mu(p, p', q = p' - p, m) = \\
ig^2 \int \frac{d^D k}{(2\pi)^D} \frac{\gamma_\alpha(\not{k} + m)\gamma_\eta[(p - 2p' + k)^\alpha g^{\mu\eta} + (p + p' - 2k)^\mu g^{\beta\eta} + (k + p' - 2p)^\eta g^{\mu\beta}]}{(k^2 - m^2)(p-k)^2(p'-k)^2},
\end{aligned} \tag{3.14}$$

for diagram (3.b). Standard dimensional regularization is used, as next, to obtain the divergent part of these two Feynman integrals, which is given by

$$\left[\Gamma_{a-planar}^\mu + \Gamma_{b-planar}^\mu \right]_{\text{div.}} = -\frac{8g^2\gamma^\mu}{16\pi^2} \frac{1}{(4-D)}. \quad (3.15)$$

The scaling factor Z_1 is therefore given by

$$Z_1^{Adj.} = 1 - \frac{8g^2}{16\pi^2} \frac{1}{(4-D)}. \quad (3.16)$$

Using now Eqs. (3.7), (3.10) and (3.16) and the relations (3.2) as well as (3.1), we obtain the one-loop β -function of noncommutative QED with matter fields in the adjoint representation

$$\beta(g) \Big|_{1-loop}^{Adj.} = -\frac{g^3}{16\pi^2} \left(\frac{22}{3} - \frac{8N_f}{3} \right). \quad (3.17)$$

As it turns out, the noncommutative QED with adjoint matters is asymptotically free for $N_f < 3$. In the next section, the conformal anomaly of the noncommutative U(1) gauge theory with adjoint matters will be calculated explicitly.

4 The Conformal Anomaly and the One-Loop β -Function

In this section, the conformal anomaly of noncommutative QED with matter fields in the adjoint representation will be calculated using the Fujikawa's path integral method [5]. We will first introduce the background field method to obtain the relevant path integral for the effective action. Using this path integral, we will then calculate the contributions of the matter, ghost and gauge fields to the conformal anomaly, separately. The conformal anomaly of noncommutative QED with adjoint matter field coupling will be calculated eventually. Using the conformal anomaly, it is then possible to obtain the one-loop β -function of the theory. Its value coincides with the result which we have obtained in the previous section in the framework of perturbative expansion [see Eq. (3.17)].

4.1 The Background Field Method and the One-Loop Effective Action

Let us first redefine the gauge field A_μ by $A_\mu \rightarrow \frac{1}{g}A_\mu$. In the background field method, the gauge field must be separated into two parts:

$$A_\mu(x) = B_\mu(x) + a_\mu(x), \quad (4.1)$$

with B_μ as the classical part and a_μ the quantum fluctuating part. In terms of a_μ , the field strength tensor from Eq. (2.3) is then given by:

$$F_{\mu\nu} = F_{\mu\nu}[B] + D_\mu[B]a_\nu - D_\nu[B]a_\mu - i[a_\mu, a_\nu]_\star, \quad (4.2)$$

where $F_{\mu\nu}[B]$ and $D_\mu[B]$ are defined by

$$\begin{aligned} F_{\mu\nu}[B] &\equiv \partial_\mu B_\nu - \partial_\nu B_\mu - i[B_\mu, B_\nu]_\star, \\ D_\mu[B]a_\nu &\equiv \partial_\mu a_\nu - i[B_\mu, a_\nu]_\star. \end{aligned} \quad (4.3)$$

Using these definitions, the pure gauge part of the action is given by

$$\begin{aligned} S_{gauge}[a_\mu, B_\mu] &= -\frac{1}{4g^2} \int d^4x \left\{ F_{\mu\nu}[B] \star F^{\mu\nu}[B] + 4ia^\mu \star [F_{\mu\nu}[B], a^\nu]_\star \right. \\ &\quad \left. - 2a_\mu \star D_\nu[B]D^\nu[B]a^\mu - 2D_\nu[B]a^\nu \star D_\mu[B]a^\mu + O(a_\mu^3, a_\mu^4) \right\}. \end{aligned} \quad (4.4)$$

Like in the commutative gauge theory, the linear terms in a_μ vanish by making use of the classical equation of motion for B_μ . The cubic and quadratic terms in a_μ can also be neglected for small quantum fluctuations a_μ . For constant B_μ , this action is invariant under the following infinitesimal transformation of $a_\mu(x)$

$$\delta a_\mu(x) = D_\mu[B]\alpha(x) - i[a_\mu(x), \alpha(x)]_\star, \quad (4.5)$$

where $\alpha(x)$ is an arbitrary function of x . Here, like in the commutative non-Abelian gauge theory, the gauge must be fixed. The gauge fixing and the Fadeev-Popov parts of the action can then be given by:

$$S_{GF+FP}[a_\mu, c, \bar{c}] = -\frac{1}{2g^2} \int d^4x \left\{ D_\mu[B]a^\mu \star D_\nu[B]a^\nu - i\bar{c} \star D^\mu[B] (D_\mu[B]c - i[a_\mu, c]_\star) \right\}. \quad (4.6)$$

In terms of the background field B_μ , the matter field part of the action reads

$$S_{matter}[\psi, \bar{\psi}] = \int d^4x \bar{\psi}(x) \star (i\not{D}[B] - m)\psi(x), \quad (4.7)$$

with

$$D_\mu[B]\psi(x) = \partial_\mu\psi(x) - i[B_\mu(x), \psi(x)]_\star. \quad (4.8)$$

The resulting total action, is invariant under the following infinitesimal transformations:

$$\begin{aligned} \delta B_\mu &= \partial_\mu\alpha(x) - i[B_\mu(x), \alpha(x)]_\star, & \delta a_\mu &= i[a_\mu(x), \alpha(x)]_\star \\ \delta c &= i[c(x), \alpha(x)]_\star, & \delta \bar{c} &= i[\bar{c}(x), \alpha(x)]_\star, \\ \delta \psi &= -ig\alpha(x) \star \psi(x), & \delta \bar{\psi} &= ig\bar{\psi}(x) \star \alpha(x). \end{aligned} \quad (4.9)$$

The effective action of the noncommutative QED with matter fields in the adjoint representation is therefore given by:

$$\exp(-W[B]) = \int D\psi D\bar{\psi} Da_\mu Dc D\bar{c} \exp(S_{gauge}^{gauge}[a_\mu] + S_{quad}^{ghost}[c, \bar{c}] + S_{matter}[\psi, \bar{\psi}]), \quad (4.10)$$

with

$$S_{quad}^{gauge}[a_\mu] = \frac{1}{2g^2} \int d^4x (a_\mu(x) \star D_\nu[B] D^\nu[B] a^\mu(x) - 2ia_\mu(x) \star [F^{\mu\nu}[B], a_\nu(x)]_\star), \quad (4.11)$$

and

$$S_{quad}^{ghost}[c, \bar{c}] = \frac{i}{4g^2} \int d^4x \bar{c}(x) \star D_\mu[B] D^\mu[B] c(x). \quad (4.12)$$

Further, $S_{matter}[\psi, \bar{\psi}]$ is given in Eq. (4.7).

4.2 The Conformal Anomaly of NC-QED

As it is mentioned in Sect. 1, even in the massless limit, the scale invariance of the noncommutative QED with matter fields in the adjoint representation is broken explicitly due to the noncommutativity parameter θ . The theory is at the same time anomalous under the global conformal transformations of matter fields ψ and $\bar{\psi}$

$$\psi(x) \rightarrow \psi'(x) = \exp\left(-\frac{1}{2}\alpha\right)\psi(x), \quad \text{and} \quad \bar{\psi}(x) \rightarrow \bar{\psi}'(x) = \exp\left(-\frac{1}{2}\alpha\right)\bar{\psi}(x), \quad (4.13)$$

gauge fields a_μ as well as ghost fields $c(x)$

$$\begin{aligned} a_\mu(x) \rightarrow a'_\mu(x) &= e^{-\alpha} a_\mu(x), \\ c(x) \rightarrow c'(x) &= e^{-2\alpha} c(x), \quad \text{and} \quad \bar{c} \rightarrow \bar{c}'(x) = \bar{c}(x). \end{aligned} \quad (4.14)$$

In this section, we intend to calculate the conformal anomaly of noncommutative QED with adjoint matters arising from these global transformations. The contributions of matter fields, gauge and ghost fields to the conformal anomaly will be calculated separately. We then calculate the corresponding one-loop β -function of the theory and will compare our result with the result from one-loop perturbative calculation in the previous section.

The Contribution of the Matter Fields

Let us assume that the Dirac operator $\mathcal{D}[B]$ has a complete set of eigenfunctions $\varphi_n(x)$ satisfying the eigenvalue equation:

$$\mathcal{D}[B]\varphi_n(x) = \lambda_n\varphi_n(x), \quad (4.15)$$

with λ_n the eigenvalues corresponding to the eigenfunctions $\{\varphi_n(x)\}$. The orthonormality and completeness relations for the set $\{\varphi_n(x)\}$ are defined formally by the noncommutative \star -product and read

$$\int d^4x \varphi_m^\dagger(x) \star \varphi_n(x) = \delta_{nm}, \quad \sum_n \varphi_n^\dagger(x) \star \varphi_n(y) = \delta^4(x - y). \quad (4.16)$$

The matter fields can therefore be written as a sum of these eigenfunctions

$$\psi(x) = \sum_n a_n \varphi_n(x), \quad \bar{\psi}(x) = \sum_m \varphi_m^\dagger(x) \bar{b}_m, \quad (4.17)$$

with the coefficients a_n and \bar{b}_m as Grassmannian numbers. Let us now consider the matter field part of the effective action

$$\int D\psi D\bar{\psi} D a_\mu Dc D\bar{c} e^{S_{matter}[\psi, \bar{\psi}]}, \quad (4.18)$$

with $S_{matter}[\psi, \bar{\psi}]$ given in Eq. (4.7). Under infinitesimal conformal transformations of matter fields $\psi \rightarrow \psi' = (1 - \frac{\alpha}{2})\psi$ and $\bar{\psi} \rightarrow \bar{\psi}' = (1 - \frac{\alpha}{2})\bar{\psi}$ the fermionic measure of the above action transforms as:

$$D\psi D\bar{\psi} \rightarrow D\psi' D\bar{\psi}' = J_\psi[\alpha] D\psi D\bar{\psi}, \quad (4.19)$$

with the Jacobian

$$J_\psi[\alpha] = \det \left[\delta_{nm} - \frac{1}{2} \alpha \int d^4x \varphi_m^\dagger(x) \star \varphi_n(x) \right]^{-2} = \exp \left(\alpha \sum_n \int d^4x \varphi_n^\dagger(x) \star \varphi_n(x) \right). \quad (4.20)$$

Here, we have used the relations $\det C = \exp[\text{Tr} \ln C]$ and $\ln(1 + \alpha) = \alpha + \mathcal{O}(\alpha^2)$ for $\alpha \ll 1$. The expression on the r.h.s. of Eq. (4.20) must be regulated using a Gaussian damping factor. We arrive at

$$J_\psi[\alpha] = \lim_{\epsilon \rightarrow 0} \exp \left(\alpha \sum_n \int d^4x \exp(-\epsilon \lambda_n^2) \varphi_n^\dagger(x) \star \varphi_n(x) \right). \quad (4.21)$$

Now using the eigenvalue equation (4.15) we get

$$\lim_{\epsilon \rightarrow 0} \exp \left(\alpha \sum_n \int d^4x \exp_\star(-\epsilon \mathcal{D}^2) \varphi_n^\dagger(x) \star \varphi_n(x) \right) \equiv \exp \left(\alpha \int d^4x A_\psi(x) \right), \quad (4.22)$$

where A_ψ is defined to be the contribution of matter fields to the global conformal anomaly. After Fourier transforming the eigenfunctions φ_n 's and using the completeness relation of their Fourier transformed, we obtain:

$$\int d^4x A_\psi(x) = \lim_{\epsilon \rightarrow 0} \int d^4x \int \frac{d^4k}{(2\pi)^4} e^{-ikx} \star \text{Tr}[\exp_\star(-\epsilon \mathcal{D}^2)] \star e^{ikx}. \quad (4.23)$$

As next let us evaluate the exponent \mathcal{D}^2 . Using the definition of the covariant derivative from Eq. (4.8) in the adjoint representation, we have:

$$[D_\mu[B], D_\nu[B]]\chi(x) = -i \int \frac{d^4k}{(2\pi)^4} F_{\mu\nu}(x; k) \tilde{\chi}(k) e^{ikx}. \quad (4.24)$$

Here $\chi(x)$ is an arbitrary function of x and

$$F_{\mu\nu}(x; k) \equiv -2i \int \frac{d^4p}{(2\pi)^4} \tilde{F}_{\mu\nu}(p) e^{ipx} \sin\left(\frac{p \times k}{2}\right). \quad (4.25)$$

Choosing now $\chi(x) = [D_\alpha, D_\beta]\varphi(x)$ we obtain

$$\begin{aligned} & [D_\mu[B], D_\nu[B]] [D_\alpha[B], D_\beta[B]] \varphi(x) = \\ & = -4 \int \frac{d^4 p}{(2\pi)^4} \frac{d^4 q}{(2\pi)^4} \tilde{F}_{\mu\nu}(p) \tilde{F}_{\alpha\beta}(q) e^{i(p+q)x} \sin^2\left(\frac{p \times q}{2}\right) \varphi(x). \end{aligned} \quad (4.26)$$

Using further the relation $2\sin^2 x = (1 - \cos 2x)$ to separate the planar from the nonplanar part of the above expression, we arrive at

$$[D_\mu[B], D_\nu[B]] [D_\alpha[B], D_\beta[B]] \Big|_{\text{planar}} = -2F_{\mu\nu}(x)F_{\alpha\beta}(x), \quad (4.27)$$

which leads automatically to

$$[D_\mu, D_\nu] \Big|_{\text{planar}} = -i\sqrt{2}F_{\mu\nu}. \quad (4.28)$$

Now let us go back to the equation (4.23) to evaluate $\mathcal{D}^2 = D_\mu D_\nu \gamma_\mu \gamma_\nu$. After a rescaling of the momentum $k_\mu \rightarrow \frac{k_\mu}{\sqrt{\epsilon}}$ with ϵ the cutoff appearing in the damping factor from Eq. (4.23), and using the relations $\gamma_\mu \gamma_\nu = -\delta_{\mu\nu} + \sigma_{\mu\nu}$ with $\sigma_{\mu\nu} = \frac{1}{2}[\gamma_\mu, \gamma_\nu]$ and $e^{-ikx} \star \partial_\mu e^{ikx} = \partial_\mu + ik_\mu$, we arrive at

$$\begin{aligned} & \int d^4 x A_\psi(x) \Big|_{\text{planar}}^{\text{Adj.}} = \\ & = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon^2} \int d^4 x \int \frac{d^4 k}{(2\pi)^4} e^{k_\mu k^\mu} \text{Tr} \left[\exp_\star \left(-2i\sqrt{\epsilon} k^\mu D_\mu - \epsilon D_\mu D^\mu - \frac{i\sqrt{2}}{2} \epsilon \sigma_{\mu\nu} F^{\mu\nu}(x) \right) \right]. \end{aligned} \quad (4.29)$$

As next, the trace appearing on the r.h.s. of the above equation must be expanded in the orders of ϵ . Due to the factor $\frac{1}{\epsilon^2}$ appearing before the integral, the expansion must be taken up to the fourth order in ϵ . All higher orders in ϵ vanish in the limit $\epsilon \rightarrow 0$. Making use of the relations $\text{Tr}(\gamma^\mu \gamma^\nu) = 4\delta^{\mu\nu}$, $\text{Tr}(\sigma^{\mu\nu}) = 0$, and $\text{Tr}(\sigma^{\mu\nu} \sigma^{\alpha\beta}) = 4(\delta^{\mu\beta} \delta^{\alpha\nu} - \delta^{\mu\alpha} \delta^{\beta\nu})$ and the integrals

$$\int \frac{d^4 k}{(2\pi)^4} e^{k^\rho k_\rho} [1, k^\mu k^\nu, k^\mu k^\nu k^\alpha k^\beta] = \frac{1}{(4\pi)^2} [1, \frac{-\delta^{\mu\nu}}{2}, \frac{1}{4}(\delta^{\mu\alpha} \delta^{\beta\nu} + \delta^{\mu\beta} \delta^{\alpha\nu} + \delta^{\mu\nu} \delta^{\beta\alpha})], \quad (4.30)$$

the contribution of the adjoint matter fields to the global anomaly of noncommutative QED with respect to the scale transformations of matter fields from Eq. (4.13) is given by

$$\int d^4 x A_\psi(x) \Big|_{\text{planar}}^{\text{Adj.}} = \frac{1}{12\pi^2} \int d^4 x F^{\mu\nu}(x) \star F_{\mu\nu}(x). \quad (4.31)$$

The Contribution of the Ghost and Gauge Fields

As next, the contributions of the gauge and ghost fields to the conformal anomaly are to be calculated. After Eq. (4.14) the infinitesimal global scale transformation of gauge and ghost fields are given by:

$$a_\mu(x) \rightarrow a'_\mu(x) = (1 - \alpha)a_\mu(x), \quad \text{and} \quad c(x) \rightarrow c'(x) = (1 - 2\alpha)c(x), \quad (4.32)$$

respectively.

Let us now consider the eigenvalue equations of the operators $D_\nu[B]D^\nu[B] - 2i[F^{\mu\nu}[B], \cdot]_\star$ from the gauge part of the effective action S_{quad}^{gauge} [see Eq. (4.11)] and $D_\nu[B]D^\nu[B]$ from the ghost part of the effective action S_{quad}^{ghost} [see Eq. (4.12)]

$$\begin{aligned} D_\nu[B]D^\nu[B]V_n^\mu(x) - 2i[F^{\mu\nu}[B](x), V_{n,\nu}(x)]_\star &= \xi_n V_n^\mu(x), \\ D_\nu[B]D^\nu[B]S_n(x) &= \zeta_n S_n(x), \end{aligned} \quad (4.33)$$

where ξ_n and ζ_n are the eigenvalues corresponding to the eigenfunctions $\{V_n^\mu(x)\}$ as well as $\{S_n(x)\}$. Both sets of eigenfunctions build a complete and orthonormal basis of the Hilbert space and can therefore be used to describe the gauge and the ghost fields

$$a_\mu(x) = \sum_n a_n V_{n,\mu}(x), \quad c(x) = \sum_n b_n S_n(x), \quad (4.34)$$

where a_n are c-number, whereas b_n are Grassmannian numbers. As next, let us consider the gauge and ghost part of the effective action, which are given by

$$\int D\psi D\bar{\psi} Da_\mu Dc D\bar{c} e^{S_{quad}^{gauge}[a_\mu]}, \quad \text{and} \quad \int D\psi D\bar{\psi} Da_\mu Dc D\bar{c} e^{S_{quad}^{ghost}[c,\bar{c}]}, \quad (4.35)$$

with $S_{quad}^{gauge}[a_\mu]$ and $S_{quad}^{ghost}[c,\bar{c}]$ from Eq. (4.11) and (4.12), respectively. Under the infinitesimal scale transformation, the measures of the above integrals transform as:

$$Da'_\mu = J_a[\alpha]Da_\mu, \quad \text{and} \quad Dc' D\bar{c}' = J_c[\alpha]Dc D\bar{c}, \quad (4.36)$$

with

$$\begin{aligned} J_a[\alpha] &= \det[\delta_{nm} + \alpha \int d^4x V_{m,\mu}(x) \star V_n^\mu(x)]^{-1}, \\ J_c[\alpha] &= \det[\delta_{nm} - 2\alpha \int d^4x S_m(x) \star S_n(x)]^{-1}. \end{aligned} \quad (4.37)$$

As it turns out both Jacobians J_a and J_c are divergent and have to be regulated using an ordinary exponential damping factor. The contributions of the gauge and ghost fields to the conformal anomaly, A_a and A_c , are then defined by

$$\begin{aligned} \int d^4x A_a(x) &\equiv \lim_{\epsilon \rightarrow 0} \left(- \sum_n \int d^4x V_{n,\mu}(x) \star e^{-\epsilon \xi_n} V_n^\mu(x) \right), \\ \int d^4x A_c(x) &\equiv \lim_{\epsilon \rightarrow 0} \left(2 \sum_n \int d^4x e^{-\epsilon \zeta_n} S_n(x) \star S_n(x) \right). \end{aligned} \quad (4.38)$$

As next, the eigenvalue equations (4.33) must be used to replace the eigenvalues ξ_n and ζ_n by their corresponding operators. After a rescaling $k_\mu \rightarrow \frac{k_\mu}{\sqrt{\epsilon}}$ with ϵ as the same cutoff appearing in the damping

factors and using the same algebraic relations leading from Eq. (4.22) to Eq. (4.29) in the fermionic case, we arrive at:

$$\begin{aligned} \int d^4x A_a(x) &= \\ &= \lim_{\epsilon \rightarrow 0} \frac{-1}{\epsilon^2} \int d^4x \int \frac{d^4k}{(2\pi)^4} e^{k^\rho k_\rho} \text{Tr} \left[\exp \left(-2i\sqrt{\epsilon} k^\alpha D_\alpha[B] \delta^{\mu\nu} - \epsilon D^2[B] + 2i\epsilon F^{\mu\nu}(x; k) \right) \right], \end{aligned} \quad (4.39)$$

for the contribution of the gauge fields and

$$\int d^4x A_c(x) = \lim_{\epsilon \rightarrow 0} \frac{2}{\epsilon^2} \int d^4x \int \frac{d^4k}{(2\pi)^4} e^{k^\rho k_\rho} \exp \left(-2i\sqrt{\epsilon} k^\mu D_\mu - \epsilon D_\mu[B] D^\mu[B] \right), \quad (4.40)$$

for the contribution of the ghost fields. The function $F_{\mu\nu}(x; k)$ on the r.h.s. of the Eq. (4.39) is defined in the Eq. (4.25). Going through the same algebraic manipulations leading from Eq. (4.29) to Eq. (4.31) in the fermionic case, we arrive at the contributions of the gauge and ghost fields to the conformal anomaly:

$$\begin{aligned} \int d^4x A_a(x) \Big|_{planar} &= -\frac{5}{24\pi^2} \int d^4x F_{\mu\nu}(x) \star F^{\mu\nu}(x), \\ \int d^4x A_c(x) \Big|_{planar} &= -\frac{1}{48\pi^2} \int d^4x F^{\mu\nu}(x) \star F_{\mu\nu}(x). \end{aligned} \quad (4.41)$$

Now, adding these contributions to the contribution of the fermionic fields from Eq. (4.31), the total conformal anomaly of the noncommutative QED with matter fields in the adjoint representation is then given by

$$\begin{aligned} \int d^4x A(x) \Big|_{planar}^{Adj.} &= N_f \int d^4x A_\psi(x) \Big|_{planar} + \int d^4x A_a(x) \Big|_{planar} + \int d^4x A_c(x) \Big|_{planar} \\ &= \frac{1}{(4\pi)^2} \left(\frac{4}{3} N_f - \frac{11}{3} \right) \int d^4x F^{\mu\nu}(x) \star F_{\mu\nu}(x). \end{aligned} \quad (4.42)$$

Here, N_f is the number of flavor degrees of freedom. In Ref. [4] the conformal anomaly of noncommutative QED with matter fields in the fundamental representation was calculated using the same path integral method. It was given by

$$\int d^4x A(x) \Big|_{planar}^{Fund.} = \frac{1}{(4\pi)^2} \left(\frac{2}{3} N_f - \frac{11}{3} \right) \int d^4x F^{\mu\nu}(x) \star F_{\mu\nu}(x). \quad (4.43)$$

Comparing to the result from Eq. (4.42), only the contribution to the matter fields to the scale anomaly differs from the corresponding contribution to the anomaly of the noncommutative $U(1)$ gauge theory with adjoint matter fields.

4.3 One-loop β -Function of Noncommutative QED with Adjoint Matters

As in the ordinary commutative case, in noncommutative QED there is a relation between the scale anomaly and the β -function of the theory which is given by

$$\int d^4x A(x) = \frac{\beta(g)}{2g^3} \int d^4x F^{\mu\nu}(x) \star F_{\mu\nu}(x). \quad (4.44)$$

The function $A(x)$ on the l.h.s. of this equation is the scale anomaly including the contributions of matter, gauge and ghost fields. This is an exact relation and is satisfied nonperturbatively for all orders of perturbative expansion.

To calculate the one-loop β -function of the theory using Eq. (4.44), it is enough to consider the conformal anomaly arising from the quadratic terms in a_μ and neglecting all higher order terms. Here, a_μ describes the quantum fluctuations of the gauge fields from the constant background field B_μ .

Using now the result from Eq. (4.42) and the relation (4.44), the one-loop β -function of noncommutative QED with adjoint matters can be calculated and reads:

$$\beta(g) \Big|_{1-loop}^{Adj.} = -\frac{g^3}{(4\pi)^2} \left(\frac{22}{3} - \frac{8}{3} N_f \right). \quad (4.45)$$

This result coincides with the one-loop β -function from Eq. (3.17), which was found by performing an explicit perturbative calculation up to one-loop order. In Ref. [4], the one-loop β -function of noncommutative QED with fundamental matter fields was calculated using the same relation (4.44) between the anomaly and the β -function of the theory. It was given by the same perturbative result from Eq. (3.4).

5 Conclusion

In this paper, the one-loop β -function of noncommutative QED with adjoint matter fields is calculated using two different methods. First, a perturbative analysis up to one-loop order is performed and the value of the one-loop β -function is determined. To obtain the renormalization constants $Z_i, i = 1, 2, 3$, only the planar parts of the one-loop fermion- and photon self energy, as well as vertex function are evaluated using dimensional regularization. The nonplanar part of the corresponding one-loop Feynman integrals are assumed to be finite for finite noncommutativity parameter θ . The value of the one-loop β -function is therefore not altered by the UV/IR mixing phenomena of noncommutative Field Theory. Since the β -function of the noncommutative QED does not include any noncommutativity parameter θ , it is not possible to compare its value with the β -function of the ordinary commutative QED. The one-loop β -function of noncommutative QED is more comparable to the β -function of a commutative non-Abelian gauge theory, like QCD. We have found that noncommutative QED with matter fields in the adjoint representation is asymptotically free for $N_f < 3$. As is known from Ref. [3], noncommutative QED with fundamental matter fields is asymptotically free for $N_f < 6$.

The one-loop β -function of the theory is then determined by calculating the one-loop contributions to the conformal anomaly of the noncommutative QED with adjoint matter field coupling. It can be shown that classically, even in the massless limit, the scale invariance of noncommutative Field

Theories is explicitly broken. This is due to the noncommutativity of the space-time coordinates. In this paper, only the quantum corrections to the conformal anomaly are calculated up to one-loop order using the Fujikawa's path integral method. Considering only the planar contributions to the conformal anomaly, we have shown that the value of the one-loop β -function coincides in both methods.

6 Acknowledgment

The authors thanks F. Ardalan, and H. Arfaei for useful discussions.

References

- [1] M. R. Douglas, and N. A. Nekrasov, *Noncommutative Field Theory*, Rev. Mod. Phys. **73** (2002) 977, hep-th/0106048.
I. Ya. Aref'eva, D. M. Belov, A. A. Giriyavets, A. S. Koshelev, and P. B. Medvedev, *Noncommutative Field Theories and (Super)String Field Theories*, hep-th/0111208.
- [2] M. R. Douglas, and C. Hull, *D-branes and the Noncommutative Torus*, JHEP **9802** (1998) 008, hep-th/9711165.
F. Ardalan, H. Arfaei, and M.M. Sheikh-Jabbari, *Mixed Branes and M(atrrix) Theory on Noncommutative Torus*, hep-th/9803067; *Noncommutative Geometry From Strings and Branes*, JHEP **9902** (1999) 016, hep-th/9810072; *Dirac Quantization of Open Strings and Noncommutativity in Branes*, Nucl. Phys. **B576** (2000) 578, hep-th/9906161.
Y.-K. E. Cheung, and M. Krogh, *Noncommutative Geometry from 0-Branes in a Background B Field*, Nucl. Phys. **B528** (1998) 185.
C.-S. Chu, and P.-M. Ho, *Noncommutative Open String and D-brane*, Nucl. Phys. **B550** (1999) 151, hep-th/9812219; *Constrained Quantization of Open String in Background B Field and Noncommutative D-brane*, Nucl. Phys. **B568** (2000) 447, hep-th/9906192.
V. Shomerus, *D-branes and Deformation Quantization*, JHEP **9906** (1999) 030, hep-th/9903205.
N. Seiberg, and E. Witten, *String Theory and Noncommutative Geometry*, JHEP **9909** (1999) 032, hep-th/9908142.
- [3] M. Hayakawa, *Perturbative Analysis on Infrared and Ultraviolet Aspects of Noncommutative QED on R^4* , hep-th/9912167; *Perturbative Analysis on Infrared Aspects of Noncommutative QED on R^4* , Phys. Lett. **B478** (2000) 394, hep-th/9912094.

- [4] T. Nakajima, *Conformal Anomalies in Noncommutative Gauge Theories*, hep-th/0108158.
- [5] K. Fujikawa, *Path Integral for Gauge Theories with Fermions*, Phys. Rev. **D21** (1980) 2884;
Nondiagrammatic Calculation of the One-Loop β Function in QCD, Phys. Rev. **D48** (1993) 3922.
- [6] F. Ardalan, and N. Sadooghi, *Axial Anomaly of Noncommutative QED on R^4* , Int. J. Mod. Phys. **A16** (2001) 3151; *Anomaly and Nonplanar Diagrams of Noncommutative Gauge Theories*, Int. J. Mod. Phys. **A17** (2002) 123.
- J. M. Garcia-Bondia, and C. P. Martin, Phys. Lett. **B479** (2000) 321.
- L. Bonara, M. Schnabl, and A. Tomasiello, Phys. Lett. **B485** (2000) 311.
- E. F. Moreno, and F. A. Shaposnik, JHEP **0003** (2000) 032.
- [7] N. Seiberg, JHEP **0009** (2000) 003.
- [8] A. Matusis, L. Susskind, and N. Toumbas, *The IR/UV Connection in the Noncommutative Gauge Theories*, JHEP **0012** (2000) 002, hep-th/0002075.
- [9] S. Minwalla, M. Van. Raamsdonk, and N. Seiberg, *Noncommutative Perturbative Dynamics*, JHEP **0002** (2000) 020, hep-th/9912072.
- [10] M. Chaichian, P. Presnajder, M. M. Sheikh-Jabbari, and A. Tureanu, *Noncommutative Standard Model: Model Building*, hep-th/0107055.
- [11] X. Calmet, B. Jurco, P. Schupp, J. Wess, and M. Wohlgenannt, *The Standard Model on Noncommutative space-time*, Eur. Phys. J. **C23** (2002) 363.

7 Figures




Figure 1: One-loop contribution to the fermion self energy.




Figure 2: One-loop contributions to the photon self energy.




Figure 3: One-loop contributions to the vertex function.