

# Holographic Charge Excitation on Horizontal Boundary

M.Hotta

*Department of Physics, Tohoku University,  
Sendai 980-8578, Japan*

## Abstract

We argue that states with nontrivial horizontal charges of BTZ black hole can be excited by ordinary falling matter including Hawking radiation. The matter effect does not break the integrability condition of the charges on the horizon. Thus we are able to trace the processes in which the matter imprints the information on the horizon by use of the charged states. It is naturally expected that in the thermal equilibrium with the Hawking radiation the black hole wanders ergodically through different horizontal states due to thermal fluctuation of incoming matter. This fact strengthens plausibility of the basic part of Carlip's idea [1]. We also discuss some aspects of the quantum horizontal symmetry and conjecture how the precise black hole entropy will be given from our point of view.

# 1 Introduction

The black hole entropy problem [2] has been attacked for long years by many people and the effort yields significant progresses. Especially, the state counting on BPS branes has been achieved in the string theory and the precise entropy form of the corresponding black branes,  $\mathcal{A}/4G$ , is obtained, using the stability of the BPS states against the strong coupling of the gravitational interaction [3]. It is quite an interesting result because the analysis really proves that the origin of the black hole entropy is entirely statistical. Even though such a crucial success has been performed, it should be stressed that there still a lot of questions remain. For example, we do not know where the states contributing to the entropy live in the macroscopical black-hole spacetimes. In addition, we must explain explicitly how the no-hair theorem is reconciled with the hairy black hole with large entropy. In the string argument the BPS states are counted just in the weak coupling region, that is, in the flat spacetime. Hence no real curved spacetime with horizon appears in the step and we cannot find any resolution in the argument to answer the above questions.

From the view point that the black hole horizon is essential to resolve the questions, Carlip [1] proposed a quite stimulating scenario. He pointed out a possibility that a part of gauge freedom of the general covariance becomes dynamical on the horizon and generates physical states contributing to the entropy. The idea itself does not depend on the spacetime dimensions and the gravity theories (general relativity, dilatonic gravity and so on) as long as the theories possess the general covariance in the action. In his idea the reconciliation between the no-hair theorem and the tremendous amount of the entropy becomes fairly evident. The point is that the charges mentioned in the no-hair theorem, mass  $M$ , angular momentum  $J$  and gauge charges  $Q$ , characterize just the geometrical property of the black-hole spacetimes. In fact the charges appear in geometrical quantities like curvature in any coordinate system. Hence the change of the charges means generation of different geometry which cannot be yielded by any coordinate transformation. On the other hand the horizontal charges are proposed to be generators of a part of general coordinate transformation. Therefore the transformation generated by the horizontal charges does not make a different geometry at all. The

situation may be easily understood by recalling that momentum  $P$  of a black hole is a significant physical charge, but translation generated by  $P$  does not change the geometry itself. Clearly two quantum states with different  $P$  values are orthogonal to each other, thus the two states are physically different and, for example, will contribute independently to the partition function and the entropy of a black-hole many-body system. The similar mechanism will happen near horizon of a black hole and the horizontal charges are expected to generate many different states without deformation of the geometry, respecting the no-hair theorem.

As a path to reach the universal realization of his basic idea in arbitrary dimensions, a lot of people including himself [4] tried to find a Virasoro symmetry with classically nonvanishing central charge on the horizon in order to get the entropy via Cardy's CFT formula [5]. Unfortunately, various difficulties prevent the scenario from succeeding so far[6]. For example, some models need to pick up a circle  $S^1$  without any convincing reason in higher-dimensional geometries, some have ill-defined boundary charge or no classical central extension for Cardy's formula and some suffer from ghost states with negative norm. Hence fully-satisfactory resolution based on the Virasoro-algebra scenario has not been achieved yet to explain the entropy.

In order to reach the realization of essential part of the Carlip's idea, another way may be still open. It has been pointed out [7] that the Schwarzschild black hole admits local time translation and angular diffeomorphism on the horizon to be a well-defined asymptotic symmetry and that nontrivial representation with nonvanishing canonical charge can be really constructed. In four dimensions the charges are labeled by the spherical harmonic indices  $(lm)$  like  $Q_{lm}$ . The symmetry appears naturally in arbitrary dimensions (higher than two dimensions) and does not request any selection of a circle  $S^1$  near the horizon. Though the algebra does not have any classical central extension, the regular symmetry on the horizon is the first example, as far as we know, which possesses a nonsinglet representation even in the classical theory. The number of the charges, which is classically infinite, looks so large as to incorporate huge degeneracy of the black hole states into the representational space. Available values of the charges distinguish a tremendous number of different physical states and might generate the exact black hole entropy.

In this paper we prove that falling matter into the horizon does not break the integrability condition of the charges in the perturbation level. Conse-

quently it is argued that the states with the nontrivial charges of BTZ black hole are actually excited by ordinary falling matter including Hawking radiation. From this observation it is naturally expected that, in the thermal equilibrium with the Hawking radiation, the black hole wanders ergodically through different horizontal states due to thermal fluctuation of incoming matter. This fact strengthens plausibility of the basic part of the Carlip's idea. In the final section we discuss some aspects of the quantum symmetry and conjecture how the precise black hole entropy will be given from our point of view.

## 2 Horizontal Charge and Boundary Condition

Let us consider the three-dimensional Einstein gravity with negative cosmological constant. The action reads

$$S = \frac{1}{16\pi G} \int d^3x \sqrt{-g} (R + 2\mu^2), \quad (1)$$

where the constant  $\mu (> 0)$  fixes the cosmological constant scale. In the system it is well known that the black hole solution exists, so called the BTZ black hole. The metric is given as follows.

$$ds^2 = -A(r)dt'^2 + \frac{dr^2}{A(r)} + r^2 \left( d\phi' - \frac{J}{2r^2} dt' \right)^2, \quad (2)$$

$$A(r) = \mu^2 r^2 - M + \frac{J^2}{4r^2} = \frac{\mu^2}{r^2} (r^2 - r_+^2)(r^2 - r_-^2), \quad (3)$$

where  $0 \leq \phi' \leq 2\pi$ ,  $r_+ \geq r_-$ ,  $M$  and  $J$  are mass and angular momentum of the black hole normalized in the planck unit and  $r_+$  is the radius of the horizon. Introducing a constant  $\epsilon > 0$ , which will be set the planck scale later, we consider the following regular transformation.

$$t = 4\pi T_H \epsilon t', \quad (4)$$

$$\rho = 4\pi T_H \epsilon \left[ - \int \frac{dr}{A(r)} + \frac{1}{2\pi T_H} \ln(4\pi T_H \epsilon) \right], \quad (5)$$

$$\phi = \phi' - 2\pi T_H \epsilon \frac{J}{r_+^2} t', \quad (6)$$

where  $0 \leq \phi \leq 2\pi$  and  $T_H = \mu^2 \frac{r_+^2 - r_-^2}{2\pi r_+}$  is the Hawking temperature. Using the transformation the BTZ metric is reexpressed near horizon ( $\rho \sim \infty$ ) as

$$d\bar{s}^2 = \Delta(-dt^2 + d\rho^2) + (r_+^2 + O(\Delta))d\phi^2 + O(\Delta^2). \quad (7)$$

Here  $\Delta$  has been defined as

$$\Delta = e^{-\frac{\rho}{\epsilon}} \sim 0. \quad (8)$$

Then we propose asymptotic metrics to the BTZ background in eqn(7) in the same spirit of [7] as follows.

$$\begin{bmatrix} g_{tt} & g_{t\rho} & g_{t\phi} \\ g_{\rho t} & g_{\rho\rho} & g_{\rho\phi} \\ g_{\phi t} & g_{\phi\rho} & g_{\phi\phi} \end{bmatrix} = \begin{bmatrix} -\Delta + O(\Delta^2) & O(\Delta^2) & O(\Delta) \\ O(\Delta^2) & \Delta + O(\Delta^2) & O(\Delta) \\ O(\Delta) & O(\Delta) & O(1) \end{bmatrix}, \quad (9)$$

where  $O(\Delta^k)$  implies that

$$\left| \lim_{\rho \rightarrow \infty} \frac{O(\Delta^k)}{\Delta^k} \right| = |f(t, \phi)| < \infty. \quad (10)$$

Note that the boundary congruence defined by

$$\rho = \infty, \quad (11)$$

$$\phi = \text{const.} \quad (12)$$

is always null in regular spacetimes with the metric in eqn(9). This implies that the boundary is just analogous concept to the Rindler horizon in the flat spacetime, that is, any physical matter across the boundary cannot come back to the outside. Taking account of the gravitational backreaction, falling matter from the thermal bath may temporarily shift the physical position of the boundary at  $\rho = \infty$  apart from the global event horizon. (Here the shift should be measured by appropriate geometrical length  $d\tilde{l}$  between the

congruence and the horizon without gauge ambiguity.) However the deviation is not expected to become so large because the black hole swallows both positive energy flux (associated with on-shell incoming matter) and negative energy flux (associated with quanta pair-created with positive-energy particles of the Hawking radiation). The average magnitudes of the both fluxes take the same value because of the thermal equilibrium. Hence the average shift of the boundary position will be suppressed.

It is rather straightforward to prove that infinitesimal coordinate transformations which preserve the asymptotic condition in eqn(9) take the following form.

$$\xi^t = U(\phi) + O(\Delta), \quad (13)$$

$$\xi^\rho = O(\Delta), \quad (14)$$

$$\xi^\phi = V(\phi) + O(\Delta), \quad (15)$$

where  $U(\phi)$  and  $V(\phi)$  are arbitrary periodic functions of  $\phi$ . The algebra of the generators is now easily written down. It is convenient to introduce mode expansions for  $U$  and  $V$ :

$$U(\phi) = \sum_m U_m e^{im\phi}, \quad (16)$$

$$V(\phi) = \sum_m V_m e^{im\phi}. \quad (17)$$

Then the generators are defined as

$$\mathcal{Q}_m^{(T)} = e^{im\phi} \partial_t, \quad (18)$$

$$\mathcal{Q}_m^{(\Phi)} = e^{im\phi} \partial_\phi, \quad (19)$$

and the algebra is explicitly calculated as follows.

$$[\mathcal{Q}_m^{(T)}, \mathcal{Q}_{m'}^{(T)}] = 0, \quad (20)$$

$$[\mathcal{Q}_m^{(\Phi)}, \mathcal{Q}_{m'}^{(T)}] = im' \hat{Q}_{m+m'}^{(T)}, \quad (21)$$

$$[\mathcal{Q}_m^{(\Phi)}, \mathcal{Q}_{m'}^{(\Phi)}] = i(m' - m) \hat{Q}_{m+m'}^{(\Phi)}. \quad (22)$$

Stress that this is just a classical  $U(1)$  current algebra.

It must be checked next that canonical charges of the transformations in eqns(13)  $\sim$  (15) are well-defined or integrable. Decomposing the asymptotic metric into the ADM form as

$$ds^2 = -N^2 dt^2 + h_{ij}(dx^i + N^i dt)(dx^j + N^j dt), \quad (23)$$

the canonical theory of the system can be constructed. Introducing the surface deformation vector:

$$\hat{\xi}^t = N\xi^t, \quad (24)$$

$$\hat{\xi}^i = \xi^i + N^i\xi^t, \quad (25)$$

the deviation of the canonical charge takes ordinary form as

$$\begin{aligned} \delta Q_H[\xi] = \oint_{\Delta=0} d\phi \left[ \frac{\sqrt{h}}{16\pi G} (h^{ac}h^{b\rho} - h^{ab}h^{c\rho}) (\hat{\xi}^t \delta h_{ab|c} - \hat{\xi}^t_{|c} \delta h_{ab}) \right. \\ \left. + 2\hat{\xi}^a \delta \Pi_a^\rho - \hat{\xi}^\rho \Pi^{ab} \delta h_{ab} \right]. \end{aligned} \quad (26)$$

Using the asymptotic condition in eqn(9), it is shown that  $\int \delta Q_H[\xi]$  really exists and is analytically integrable for the transformations in eqns(13)  $\sim$  (15). This results in

$$Q_H[\xi] = \int_0^{2\pi} d\phi \left[ \frac{\sqrt{h_{\phi\phi}}}{16\pi G\epsilon} \xi^t + 2\Pi_\phi^\rho \xi^\phi \right]_{\rho=\infty}. \quad (27)$$

The charges in eqn(27) are divided into the following two parts. The local time translation on the horizon is associated with the charges:

$$Q_m^{(T)} = \int_0^{2\pi} d\phi e^{im\phi} \left[ \frac{\sqrt{h_{\phi\phi}}}{16\pi G\epsilon} \right]_{\rho=\infty}. \quad (28)$$

The angular diffeomorphism on the horizon is associated with the charges:

$$Q_m^{(\Phi)} = \int_0^{2\pi} d\phi e^{im\phi} \left[ 2\Pi_\phi^\rho \right]_{\rho=\infty}. \quad (29)$$

The finite transformation corresponding to the infinitesimal one in eqns (13)  $\sim$  (15) is given as

$$t' = t + T(\phi) + O(\Delta), \quad (30)$$

$$\rho' = \rho + O(\Delta), \quad (31)$$

$$\phi' = \Phi(\phi) + \Delta\Phi_1(t, \phi) + O(\Delta^2), \quad (32)$$

where  $T(\phi + 2\pi) = T(\phi)$ ,  $\Phi(\phi + 2\pi) = \Phi(\phi) + 2\pi$ ,  $\Phi_1(\phi + 2\pi) = \Phi_1(\phi)$  and  $d\Phi/d\phi > 0$ . Let us consider a BTZ black hole metric with  $J = 0$  and  $M = (\mu r_+)^2$ :

$$ds^2 = e^{-\rho'/\epsilon}(-dt'^2 + d\rho'^2) + r_+^2 d\phi'^2 + \dots \quad (33)$$

The asymptotic transformation induces the following excited metric from the metric.

$$\begin{aligned} ds^2 = & \Delta(-dt^2 + d\rho^2) \\ & -2\Delta[\dot{T}(\phi) - r_+^2 \dot{\Phi}(\phi) \partial_t \Phi_1(t, \phi)] dt d\phi \\ & + r_+^2 \dot{\Phi}(\phi)^2 d\phi^2 - 2\Delta \frac{r_+^2}{\epsilon^2} \dot{\Phi}(\phi) \Phi_1(t, \phi) d\rho d\phi + \dots, \end{aligned} \quad (34)$$

where  $\dot{T} = \partial_\phi T$  and  $\dot{\Phi} = \partial_\phi \Phi$ . The charges of the excited metric in eqn(34) are explicitly evaluated as

$$Q_m^{(T)} = \frac{r_+^2}{16\pi G\epsilon} \int_0^{2\pi} \frac{d\Phi}{d\phi} e^{im\phi} d\phi, \quad (35)$$

$$Q_m^{(\Phi)} = \frac{r_+^2}{16\pi G\epsilon} \int_0^{2\pi} \frac{dT}{d\phi} \frac{d\Phi}{d\phi} e^{im\phi} d\phi. \quad (36)$$

Consequently it is clear that they certainly take nonzero values in general, that is, the representation is not singlet even in the classical theory.

In the next section we will discuss falling matter effect. So let us change the variables as follows.

$$u = t - \rho, \quad (37)$$

$$\rho = \rho, \quad (38)$$

$$\phi = \phi. \quad (39)$$

The asymptotic condition in eqn(9) is now rewritten as

$$\begin{bmatrix} g_{uu} & g_{u\rho} & g_{u\phi} \\ g_{\rho u} & g_{\rho\rho} & g_{\rho\phi} \\ g_{\phi u} & g_{\phi\rho} & g_{\phi\phi} \end{bmatrix} = \begin{bmatrix} -\Delta + O(\Delta^2) & -\Delta + O(\Delta^2) & O(\Delta) \\ -\Delta + O(\Delta^2) & O(\Delta^2) & O(\Delta) \\ O(\Delta) & O(\Delta) & O(1) \end{bmatrix}, \quad (40)$$

where  $O(\Delta^k)$  means that

$$\left| \lim_{\rho \rightarrow \infty} \frac{O(\Delta^k)}{\Delta^k} \right| = |F^{(k)}(u, \phi)| < \infty. \quad (41)$$



For the condition in eqn(40), the asymptotic symmetry is reexpressed as

$$\xi^u = U(\phi) + O(\Delta), \quad (42)$$

$$\xi^\rho = O(\Delta), \quad (43)$$

$$\xi^\phi = V(\phi) + O(\Delta). \quad (44)$$

Because the ADM decomposition is not available in the new coordinates, we check the integrability of the symmetric charges via the covariant phase-space formulation [8]. The deviation of the charge is now defined as

$$\delta\mathcal{H}[\xi] = \delta \int_{\partial\mathcal{C}} \frac{1}{2} \epsilon_{\beta\alpha\mu} Q^{\beta\alpha} + \int_{\partial\mathcal{C}} \epsilon_{\beta\alpha\mu} \xi^\beta \Theta^\alpha, \quad (45)$$

where

$$Q^{\beta\alpha} = \frac{1}{16\pi G} [\nabla^\alpha \xi^\beta - \nabla^\beta \xi^\alpha] \quad (46)$$

and

$$\Theta^\alpha = \frac{1}{16\pi G} [g_{\mu\nu} \nabla^\alpha \delta g^{\mu\nu} - \nabla_\nu \delta g^{\nu\alpha}]. \quad (47)$$

The first term of r.h.s in eqn(45) is clearly integrable, however, integrability of the second term is not trivial. We prove using eqn(40) that the Wald-Zoupas integrability condition [9] strictly holds for the transformations in eqns(42)  $\sim$  (44) and show that the second term in eqn(45) is integrable. Actually the form of the local time translation charge, for example, is given as

$$Q_m^{(T)} = \frac{1}{16\pi G\epsilon} \oint_{\rho=\infty} d\phi e^{im\phi} [1 - 2\epsilon\partial_u] \sqrt{g_{\phi\phi}}. \quad (48)$$

In the static case the second term in eqn(48) vanishes and eqn(48) coincides precisely with eqn(28) in the canonical theory, as it should be.

### 3 Excitations with the Horizontal Charge

In this section we argue that falling matter does not break the asymptotic boundary condition in eqn(40) at least in the perturbative level. The horizontal states with nontrivial charges are actually excited by the matter.

Let us consider the following spinless BTZ black hole solution with horizon radius  $r_+$  as the background for the perturbation.

$$d\bar{s}^2 = -\frac{1}{\sinh^2(\mu\rho)} \left( du^2 + 2dud\rho \right) + r_+^2 \coth^2(\mu\rho) d\phi^2. \quad (49)$$

In the background the action of free massive scalar field reads

$$S_{matter} = \int d^3x \sqrt{-\bar{g}} \left[ -\frac{1}{2} (\bar{\nabla}\varphi)^2 - \frac{M^2}{2} \varphi^2 \right], \quad (50)$$

where  $M$  is the mass of the field. Solving the equation of motion of the field, the general in-coming-wave solution can be explicitly written down as

$$\begin{aligned} \varphi = & \sum_{L=-\infty}^{\infty} \int_0^{\infty} dE A_L^{in}(E) [2 \sinh(\mu\rho)]^{-i\frac{E}{\mu}} [\coth(\mu\rho)]^{-i\frac{L}{\mu r_+}} e^{iE(u+\rho)+iL\phi} \\ & \times F\left(a(E, L), b(E, L), c(E, L); -\sinh^{-2}(\mu\rho)\right) \\ & + c.c., \end{aligned} \quad (51)$$

where

$$a = \frac{1}{2} \left( 1 + \sqrt{1 + \frac{M^2}{\mu^2}} \right) + \frac{i}{2} \left( \frac{E}{\mu} - \frac{L}{\mu r_+} \right), \quad (52)$$

$$b = \frac{1}{2} \left( 1 - \sqrt{1 + \frac{M^2}{\mu^2}} \right) + \frac{i}{2} \left( \frac{E}{\mu} - \frac{L}{\mu r_+} \right), \quad (53)$$

$$c = 1 + i\frac{E}{\mu}. \quad (54)$$

Near the horizon ( $\rho \sim \infty$ ), the solution behaves as

$$\varphi = \sum_L \int_0^{\infty} dE \left( A_L^{in}(E) e^{iEu+iL\phi} + A_L^{in}(E)^* e^{-iEu-iL\phi} \right) + O\left(e^{-\frac{\rho}{\mu}}\right). \quad (55)$$

Note that the quantum effect can be taken into account simply by replacing the coefficients  $A_L^{in}(E)$  and  $A_L^{in}(E)^*$  with operators  $\hat{A}_L^{in}(E)$  and  $\hat{A}_L^{in}(E)^\dagger$

which satisfy

$$[\hat{A}_L^{in}(E), \hat{A}_{L'}^{in}(E')] = 0, \quad (56)$$

$$[\hat{A}_L^{in}(E), \hat{A}_{L'}^{in}(E')^\dagger] = \frac{1}{8\pi^2 r_+ E} \delta_{LL'} \delta(E - E'). \quad (57)$$

In order to solve the Einstein equation, let us introduce the Chern-Simons variables. The dual variable of the spin connection  $\omega_\mu^{ab}$  is defined as

$$\omega_\mu^c = \frac{1}{2} \epsilon^{cab} \omega_{\mu ab}. \quad (58)$$

Using  $\omega_\mu^a$  and the triad variable  $e_\mu^a$ , let us define the following two Chern-Simons gauge fields.

$$A_\mu^a = \omega_\mu^a + \mu e_\mu^a, \quad (59)$$

$$\bar{A}_\mu^a = \omega_\mu^a - \mu e_\mu^a. \quad (60)$$

Then the Einstein equation can be rewritten as

$$\epsilon^{\mu\nu\lambda} [\partial_\nu A_{\lambda c} - \partial_\lambda A_{\nu c} + \epsilon_{abc} A_\nu^a A_\lambda^b] = 16\pi G e T_c^\mu, \quad (61)$$

$$\epsilon^{\mu\nu\lambda} [\partial_\nu \bar{A}_{\lambda c} - \partial_\lambda \bar{A}_{\nu c} + \epsilon_{abc} \bar{A}_\nu^a \bar{A}_\lambda^b] = 16\pi G e T_c^\mu, \quad (62)$$

where  $T_c^\mu$  is the stress tensor of the matter. Now taking variation of eqns(61) and (62), we get the first-order perturbative equation of motion. Here let us choose the gauge fixing as

$$\delta A_u^c = \delta \bar{A}_u^c = 0. \quad (63)$$

In the original variables  $\omega$  and  $e$ , the conditions are rewritten as

$$\delta e_u^a = 0, \quad (64)$$

$$\delta \omega_u^{ab} = 0. \quad (65)$$

Note that these six conditions fix both general covariance( with three degrees of freedom) and local Lorentz symmetry (with other three degrees of freedom). Thus in principle they can be expressed only in terms of metric variables  $g_{\mu\nu}$ . However the analytic achievement is too complicated except a single condition  $\delta g_{uu} = 0$ . Fortunately it is found that we do not need

detailed expression of the gauge fixing in the metric variables. The perturbative equation corresponding to eqns (61) and (62) can be integrated out formally just by using the expressions in eqn(63). The result is summarized as follows.

$$\delta \mathbf{A}_\rho = \int_{-\infty}^u \exp[(u - u')K(\rho)] \mathbf{J}_\phi(u', \rho, \phi) du', \quad (66)$$

$$\delta \mathbf{A}_\phi = - \int_{-\infty}^u \exp[(u - u')K(\rho)] \mathbf{J}_\rho(u', \rho, \phi) du', \quad (67)$$

$$\delta \bar{\mathbf{A}}_\rho = \int_{-\infty}^u \exp[(u - u')\bar{K}(\rho)] \mathbf{J}_\phi(u', \rho, \phi) du', \quad (68)$$

$$\delta \bar{\mathbf{A}}_\phi = - \int_{-\infty}^u \exp[(u - u')\bar{K}(\rho)] \mathbf{J}_\rho(u', \rho, \phi) du', \quad (69)$$

where the matrices  $K$  and  $\bar{K}$  are defined as

$$K = \begin{bmatrix} 0 & \beta & 0 \\ \beta & 0 & -\alpha \\ 0 & \alpha & 0 \end{bmatrix}, \quad \bar{K} = \begin{bmatrix} 0 & \beta & 0 \\ \beta & 0 & \alpha \\ 0 & -\alpha & 0 \end{bmatrix}, \quad (70)$$

and the functions  $\alpha$  and  $\beta$  are

$$\alpha = \frac{\mu}{\sinh(\mu\rho)}, \quad (71)$$

$$\beta = \mu \coth(\mu\rho). \quad (72)$$

The source terms of the matter is expressed as

$$\mathbf{J}_\rho = 8\pi G e \begin{bmatrix} \delta T^{\rho 0} \\ \delta T^{\rho 1} \\ \delta T^{\rho 2} \end{bmatrix}, \quad (73)$$

$$\mathbf{J}_\phi = 8\pi G e \begin{bmatrix} \delta T^{\phi 0} \\ \delta T^{\phi 1} \\ \delta T^{\phi 2} \end{bmatrix}, \quad (74)$$

where

$$\delta T_{\alpha\beta} = \nabla_\alpha \varphi \nabla_\beta \varphi - \frac{1}{2} g_{\alpha\beta} [(\nabla\varphi)^2 - M^2 \varphi^2]. \quad (75)$$

Substituting the general solution in eqn(51) into eqn(75) and taking the limit of  $\rho \rightarrow \infty$ , it is proven that the asymptotic metric condition in eqn(40) still

holds because the perturbative correction induced by the matter field is given in general as

$$\begin{bmatrix} \delta g_{uu} & \delta g_{u\rho} & \delta g_{u\phi} \\ \delta g_{\rho u} & \delta g_{\rho\rho} & \delta g_{\rho\phi} \\ \delta g_{\phi u} & \delta g_{\phi\rho} & \delta g_{\phi\phi} \end{bmatrix} = \begin{bmatrix} 0 & O(\Delta^2) & O(\Delta) \\ O(\Delta^2) & O(\Delta^2) & O(\Delta) \\ O(\Delta) & O(\Delta) & O(1) \end{bmatrix}, \quad (76)$$

where the coordinate variables  $(u, \rho)$  have been rescaled as

$$u' = 2\mu\epsilon u, \quad (77)$$

$$\rho' = 2\mu\epsilon \left( \rho + \frac{1}{\mu} \ln(2\mu\epsilon) \right). \quad (78)$$

Here the first component in eqn(76),  $\delta g_{uu} = 0$ , comes just from our gauge choice. Besides this there exist two other nontrivial constraints of our gauge fixing among the metric components in eqn(76). However the point is that whatever gauge constraints appears additionally in the metric the existence condition of the charge is satisfied as long as eqn(76) holds. Therefore we did not need to translate our gauge choice into the metric language.

Initially the horizontal charges  $Q_m^{(T)}$  vanish for the background metric in eqn(49) except  $m = 0$ . After the matter falls into the black hole, they get the following corrections, which are generally nonzero.

$$\delta Q_m^{(T)} = r_+ \oint d\phi e^{im\phi} \left[ \int_{-\infty}^u (\partial_u \varphi(u', \infty, \phi))^2 du' \right]. \quad (79)$$

Therefore it is concluded that the matter certainly excites the states with  $Q_{m \neq 0}^{(T)}$  nonvanishing.

Here it may be worthwhile to stress that even if one may remove the charge at the late time by an asymptotic transformation in eqns(42)  $\sim$  (44):

$$\delta Q_m^{(T)'}(u = \infty) = 0, \quad (80)$$

the initial nonzero deviations appear due to the transformation itself:

$$\delta Q_m^{(T)'}(u = -\infty) \neq 0. \quad (81)$$

Hence not the absolute value but the difference of the charges generated through the process has physical significance.

Next, as a simple example, let us consider a shock-wave solution of the matter field near the horizon:

$$(\partial_u \varphi)^2 = \kappa(\phi)\delta(u) + O(\Delta), \quad (82)$$

where  $\kappa(\phi)$  is arbitrary periodic positive function of  $\phi$ . Passing across the boundary, the wave imprints the nonzero value on the charges  $Q_m^{(T)}$  as

$$\delta Q_m^{(T)}(u > 0) = r_+ \oint e^{im\phi} \kappa(\phi) d\phi. \quad (83)$$

Interestingly note that if one observes the charges  $\delta Q_m^{(T)}$  after the matter falling, the functional form of  $\kappa(\phi)$  is reproduced completely as follows.

$$\kappa(\phi) = \frac{1}{2\pi r_+} \sum_m \delta Q_m^{(T)} e^{-im\phi}. \quad (84)$$

Thus the horizontal charge plays a role of recorder on the horizon.

For the quantum field the charges may be obtained by replacing the classical stress tensor into expectation value of the stress operator in a vacuum state:

$$\delta Q_m^{(T)} = r_+ \oint d\phi e^{im\phi} \left[ \int_{-\infty}^u \langle \hat{T}_{--}(u', \infty, \phi) \rangle du' \right], \quad (85)$$

where, due to the quantum effect, the flux  $\langle \hat{T}_{--} \rangle$  can take negative values, losing the mass of the black hole. It is easy to see that the quantum radiation is also able to excite the horizontal states, similarly to the classical field.

If the black hole completely evaporates swallowing negative quantum flux, the information registered in the horizontal charges will be released into the space via the black hole radiation because of the conservation law of the charges. (Note that the black hole radiation does not always look isotropic in the coordinate system in which the asymptotic condition in eqn(40) is satisfied at each time.)

## 4 Discussion about Quantum Hair States

We analyzed so far the classical gravity theory and showed the existence of the nontrivial horizontal charges and its excited states. In this section we try to make a hand-waving but quite suggestive argument of the quantum gravitational states.

As shown in eqns(20)  $\sim$  (22), the classical algebra is just U(1) current algebra in a circle. Hence it tempts us to suppose that its quantum version with central extension:

$$[\hat{Q}_m^{(T)}, \hat{Q}_{m'}^{(T)}] = \frac{k}{2}m\delta_{m+m'}, \quad (86)$$

$$[\hat{Q}_m^{(\Phi)}, \hat{Q}_{m'}^{(T)}] = -m'\hat{Q}_{m+m'}^{(T)}, \quad (87)$$

$$[\hat{Q}_m^{(\Phi)}, \hat{Q}_{m'}^{(\Phi)}] = (m - m')\hat{Q}_{m+m'}^{(\Phi)} + \frac{c}{12}(m^3 - m)\delta_{m+m',0} \quad (88)$$

gives some inspiration of the quantum treatment of the horizontal state counting. Though there may be some q-deformation of the algebra in the real quantum system, let us investigate to what extent the above algebra leads us to the black hole entropy.

Note that the reflection symmetry,  $\phi \rightarrow -\phi$ , prefers representations with  $k = 0$  and  $c = 0$  for the black hole system. If  $k \neq 0$ , define

$$a_m = \sqrt{\frac{2}{|k|m}} Q_{\text{sgn}(k) \times m}^{(T)} \quad (89)$$

for positive  $m$ . Then the following relation holds.

$$[a_m, a_{m'}^\dagger] = \delta_{mm'}. \quad (90)$$

In order to make unitary representation, we must impose that

$$a_m|0\rangle = 0 \quad (91)$$

for *only* positive  $m$ . Hence the prescription prevents the symmetry,  $m \rightarrow -m$ , from being respected. Similarly, if  $c > 0$ , the vacuum state in the unitary representation is defined as

$$\hat{Q}_m^{(\Phi)}|0\rangle = 0 \quad (92)$$

with  $m \geq -1$ . Therefore the reflectional symmetry is not realized in the treatment. Thus the representation with  $k = c = 0$  seems favored from the viewpoint of the angular reflection.

In the case with  $k = c = 0$ :

$$[\hat{Q}_m^{(T)}, \hat{Q}_{m'}^{(T)}] = 0, \quad (93)$$

$$[\hat{Q}_m^{(\Phi)}, \hat{Q}_{m'}^{(T)}] = -m' \hat{Q}_{m+m'}^{(T)}, \quad (94)$$

$$[\hat{Q}_m^{(\Phi)}, \hat{Q}_{m'}^{(\Phi)}] = (m - m') \hat{Q}_{m+m'}^{(\Phi)}, \quad (95)$$

the structure of the algebra is quite similar to that of the Poincaré algebra, which is constructed by the momentum  $P_a$  and the Lorentz generator  $J_{ab}$ . In fact, if  $\hat{Q}_m^{(T)}$  are regarded as  $P_a$  and  $\hat{Q}_m^{(\Phi)}$  as  $J_{ab}$ , the skeleton structure of algebras in eqns (93)  $\sim$  (95) reads

$$[P, P] = 0, \quad (96)$$

$$[J, P] = P, \quad (97)$$

$$[J, J] = J. \quad (98)$$

This precisely coincides with that of the Poincaré algebra. Consequently the irreducible unitary representation may be constructed by use of the Wigner's little-group argument [10].

Let us think a vector made of the charges of the background in eqn(7),  $(\bar{Q}_0^{(T)}, \bar{Q}_1^{(T)}, \bar{Q}_{-1}^{(T)}, \dots)$  with

$$\bar{Q}_m^{(T)} = \frac{r_+^2}{8G\epsilon} \delta_{m0}, \quad (99)$$

as a reference vector for the little group of our symmetry. Then the little group is defined as a subgroup generated by a part of  $\hat{Q}_m^{(\Phi)}$ . Under the little-group transformation the reference vector  $(\bar{Q}_m^{(T)})$  must remain unchanged:

$$\delta_L \bar{Q}_m^{(T)} = 0. \quad (100)$$

After some easy manipulation, it is proven that the little group is generated only by  $\hat{Q}_0^{(\Phi)}$ . Thus the little group is the rigid rotation of the horizon and just has  $O(2)$  group structure.

Because the operators  $\hat{Q}_m^{(T)}$  commute with each other, we can diagonalize simultaneously all the operators. In order to construct unitary representation, we assume that  $\hat{Q}_m^{(T)}$  are hermitian. Along the Wigner's argument, let a state  $|\mathbf{0}; s\rangle$  denote an eigenstate of the operator  $\hat{Q}_m^{(T)}$  with its eigenvalue  $\bar{Q}_m^{(T)}$ :

$$\hat{Q}_m^{(T)} |\mathbf{0}; s\rangle = \bar{Q}_m^{(T)} |\mathbf{0}; s\rangle. \quad (101)$$



Here the additional index  $s$  is prepared for possible intrinsic spin freedom which is transformed by the little group  $O(2)$ .

Next we define the act of the little group on the states. Because the little group does not change the reference vector ( $\bar{Q}_m^{(T)}$ ), the subspace spanned by  $\{|\mathbf{0}, s \rangle\}$  must be a representation of the little group:

$$\hat{Q}_0^{(\Phi)}|\mathbf{0}, s \rangle = \sum_{s'} J_{ss'}|\mathbf{0}, s' \rangle, \quad (102)$$

where  $[J_{ss'}]$  is arbitrary matrix representation of  $O(2)$  group. If one impose irreducibility on the representation, it must be singlet, that is,  $[J_{ss'}]$  is a real number which  $s$  denotes:

$$\hat{Q}_0^{(\Phi)}|\mathbf{0}, s \rangle = s|\mathbf{0}, s \rangle. \quad (103)$$

Note that in this algebra, subtraction of constant from  $\hat{Q}_0^{(\Phi)}$  is always admitted. Consequently we can set  $s = 0$  without loss of generality. Therefore the index  $s$  will be suppressed later.

By virtue of eqn(94), the finite transformation operator of the angular diffeomorphism  $\hat{L}(\Lambda)$  is introduced as

$$\hat{L}(\Lambda)^{-1}\hat{Q}_m^{(T)}\hat{L}(\Lambda) = \sum_{m'} \Lambda_{mm'}\hat{Q}_{m'}^{(T)}, \quad (104)$$

where  $\Lambda$  is arbitrary element of the fundamental representation of the group and, in the infinitesimal case:  $\Lambda_{mm'} = \delta_{mm'} + m\varepsilon_{m'-m}$ , the operator  $\hat{L}(\Lambda)$  is reduced into the infinitesimal one:

$$\hat{L}(\Lambda) = 1 + \sum_m \varepsilon_m \hat{Q}_m^{(\Phi)}. \quad (105)$$

Using eqn(104), it is shown that the state  $\hat{L}(\Lambda)|\mathbf{0} \rangle$  is an eigenstate with a new eigenvalue of  $\hat{Q}_m^{(T)}$ :

$$\hat{Q}_m^{(T)}\hat{L}(\Lambda)|\mathbf{0} \rangle = \sum_{m'} \Lambda_{mm'}\bar{Q}_{m'}^{(T)}\hat{L}(\Lambda)|\mathbf{0} \rangle. \quad (106)$$

Let us define here coefficients  $f_m$  as

$$f_m = \frac{8G\epsilon}{r_+^2} \sum_{m'} \Lambda_{mm'}\bar{Q}_{m'}^{(T)}. \quad (107)$$

The coefficients  $f_m$  are related with the original function  $\Phi(\phi)$  of the eigenvalues in eqn (35) as follows.

$$\frac{d\Phi}{d\phi} = \sum_m f_m e^{-im\phi}. \quad (108)$$

Note that  $f_0 = 1$  for arbitrary eigenstates of  $\hat{Q}_0^{(T)}$ . Just as in the Poincaré algebra, we can consistently introduce positive norm for properly rescaled eigenvectors:

$$|f_m \rangle \propto \hat{L}(\Lambda) |\mathbf{0} \rangle, \quad (109)$$

$$\hat{Q}_m^{(T)} |f_m \rangle = \frac{r_+^2}{8G\epsilon} f_m |f_m \rangle, \quad (110)$$

that is, the following relations can be set up.

$$\begin{aligned} \langle f_m | f_{m'} \rangle &= \prod_{m>0} \delta(\text{Re} f_m - \text{Re} f_{m'}) \\ &\times \prod_{m>0} \delta(\text{Im} f_m - \text{Im} f_{m'}). \end{aligned} \quad (111)$$

Gathering all possible eigenstates generated by  $\hat{L}(\Lambda)$ , we make a vector space spanned by  $\{|f_m \rangle\}$ . By construction it is clearly a unitary irreducible representation of the algebra.

Now let us try to discuss the number of the horizontal states roughly. Because the eigenstates  $|f_m \rangle$  with different eigenvalues are orthogonal to each other, they must be treated as physically different states. Thus we should know the number of possible values of  $f_m$  at each  $m$ , denoted by  $N_m$ . Note that the magnitude of the eigenvalues  $|f_m|$  must not be larger than one:

$$|f_m| \leq 1. \quad (112)$$

This is easily noticed from the following manipulation:

$$\begin{aligned} 1 &= \frac{1}{2\pi} \int_0^{2\pi} \frac{d\Phi}{d\phi} \left| e^{im\phi} \right| d\phi = \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{d\Phi}{d\phi} e^{im\phi} \right| d\phi \\ &\geq \left| \frac{1}{2\pi} \int_0^{2\pi} \frac{d\Phi}{d\phi} e^{im\phi} d\phi \right| = |f_m|, \end{aligned} \quad (113)$$

where we have used  $\oint d\Phi = 2\pi$  and monotonically increasingness of  $\Phi(\phi)$ .

Though the eigenvalues run only in the bounded regions  $|f_m| \leq 1$  as seen above, they are essentially continuous and shows a band structure in the representation. The continuous spectrum makes the state counting confusing. As in the free gas system, the number of states might be naively defined by the form of  $V^{(T)} d^2 f_m / (2\pi\hbar)^2$  where  $V^{(T)}$  is ‘‘volume’’ of the space canonically conjugate to the  $f_m$  space. However the principle of determination of the value of  $V^{(T)}$  has not been established and seems rather ambiguous so far.

Fortunately, the bounded regions are independent of  $m$ . Therefore it may be sufficient that we just assign a proper integer  $N$  to the number of the values of each  $f_m$ . Therefore the number of the states of the black hole  $N_{BH}$  may be roughly expressed as

$$N_{BH} \sim \prod_{m=1} N_m \sim \prod_{m=1} N. \quad (114)$$

However, it is noticed soon that the estimation is divergent because the subscript  $m$  can take any large integer. This means that the quantum treatment discussed here is too naive to count the black hole states precisely and needs some (q-)modification. The estimation in eqn(114) seems to suggest that the fully quantized gravity theory supplies some cutoff  $m_{max}$ , analogously to that of noncommutative spheres, and improves the estimation as

$$N_{BH} \sim \prod_{m=1}^{m_{max}} N = N^{m_{max}}. \quad (115)$$

Consequently the entropy will be written down as

$$S_{BH} = \ln N_{BH} \sim \ln N^{m_{max}}. \quad (116)$$

Because the cutoff  $m_{max}$  is naturally expected of order of the horizon circumference divided by the Planck length  $\epsilon_{pl}$ :

$$m_{max} \sim \frac{2\pi r_+}{\epsilon_{pl}}, \quad (117)$$

by taking the gravitational constant as

$$G = \frac{\epsilon_{pl}}{4 \ln N}, \quad (118)$$

the correct black hole entropy will be certainly reproduced:

$$S_{BH} \sim \frac{\mathcal{A}}{4G}, \quad (119)$$

where  $\mathcal{A}$  is the circumference of the horizon:

$$\mathcal{A} = 2\pi r_+. \quad (120)$$

The last step of the entropy estimation must be said quite crude. However, the same prescription also works in higher dimensional black holes and reproduces the correct entropy form,  $\mathcal{A}/4G$ . Therefore it may be possible that the analysis indicates some features of the exact quantum black hole physics, that is, the horizon might become a “quantum boundary” which is constructed from area-unit elements of order of the planck scale on the horizon. To establish the exact estimation of the entropy, more information is clearly needed about the quantum gravity theory and expected to be gained in the future work.

## Acknowledgement

I would like to thank J.Koga for helpful discussions.

## References

- [1] S.Carlip, Phys.Rev.**D51**,632,(1995).
- [2] J.D.Bekenstein, Phys.Rev.**D7**,2333,(1973).  
S.W.Hawking, S.W.Hawking, Commun.Math.Phys.**43**,199,(1975).
- [3] A.Strominger and C.Vafa, Phys.Lett.**B379**,99,(1996).
- [4] S.Carlip, Phys.Rev.Lett.**82**,2828,(1999).
- [5] S.N.Solodukhin, Phys.Lett.**B454**,213,(1999).  
F.-L.Lin and Y.-S.Wu, Phys.Lett.**B453**,222,(1999).  
D.Navarro, J.Navarro-Salas, P.Navarro, Nucl.Phys.**B580**,311,(2000).

- R.Brustein, Phys.Rev.Lett.**86**,576,(2001).  
 S.Das, A.Chosh and P.Mitra, Phys.Rev.**D63**,024023,(2001).  
 J.Jing and M.-L. Yan, Phys.Rev.**D64**,064015,(2001).
- [6] M.-I.Park and J.Ho, Phys.Rev.Lett.**83**,5595,(1999).  
 S.Carlip, Phys.Rev.Lett.**83**,5596,(1999).  
 M.-I. Park and J.H.Yee, Phys.Rev.**D61**,088501,(2000).  
 O.Dreyer, A.Ghosh and J.Wisniewski,  
 Class.Quant.Grav.**18**,1929,(2001).  
 J.Koga, Phys.Rev.**D64**,124012,(2001).
- [7] M.Hotta, K.Sasaki and T.Sasaki, Class.Quant.Grav.**18**,1823,(2001).
- [8] J.Lee and R.M.Wald, J.Math.Phys.**31**,725,(1990).  
 V.Iyer and R.M.Wald, Phys.Rev.**D50**,846,(1994);  
 Phys.Rev.**D52**,4430,(1995).
- [9] R.M.Wald and A.Zoupas, Phys.Rev.**D61**,084027,(2000).
- [10] E.P.Wigner, Ann.Math.**40**,149,(1939).