

# Quasinormal Modes of Near Extremal Black Branes

ANDREI O. STARINETS

*Institute for Nuclear Theory, University of Washington, Seattle, WA 98195*

## Abstract

We find quasinormal modes of near extremal black branes by solving a singular boundary value problem for the Heun equation. The corresponding eigenvalues determine the dispersion law for the collective excitations in the dual strongly coupled  $\mathcal{N} = 4$  supersymmetric Yang-Mills theory at finite temperature.

*In memory of Aleksandr Sergeevich Vshivtsev*

July 2002

---

starina@phys.washington.edu

## 1. Introduction

The gauge theory/gravity correspondence [1,2,3] provides useful insights into the properties of a strongly coupled supersymmetric Yang-Mills (SYM) theory at non-zero temperature. A well-known example is the finite-temperature  $\mathcal{N} = 4$   $SU(N)$  SYM theory in  $4d$  which in the large  $N$ , large 't Hooft coupling limit is dual to the gravitational background of  $N$  parallel near extremal three-branes, with temperature related to the parameter of non extremality. For this theory, a number of quantities such as the free energy [4,5], the Wilson loop [6,7], the shear viscosity [8,9], the  $R$ -charge diffusion constant [9], and the Chern-Simons diffusion rate [10] have been computed using the methods of gauge theory/gravity duality. Since supersymmetry is broken at finite temperature and thus no non-renormalization theorem is expected to hold, quantities computed in the strong coupling regime using a gravity dual differ from their analogues obtained at weak coupling via perturbation theory. The results of a gravity calculation are then regarded as a prediction for the SYM theory, assuming that the AdS/CFT correspondence is valid at finite temperature<sup>1</sup>.

Dynamical properties of a thermal gauge theory are encoded in its Green's functions. In the context of AdS/CFT, Minkowski space Green's functions can be computed from gravity using the recipe given in [10]. Unfortunately, for a non-extremal background, only approximate expressions for the correlators are usually obtained. For example, the retarded propagator of the gauge invariant local operator  $F^2$  (dual to the dilaton) defined by

$$G^R(\omega, \mathbf{k}) = -i \int dt d^3x e^{-i\omega t + i\mathbf{k}\mathbf{x}} \theta(t) \langle [F^2(x), F^2(0)] \rangle \quad (1.1)$$

can be explicitly computed only at zero or very high (with respect to the absolute value of momentum) temperature. At zero temperature, the retarded propagator (1.1) has a branch cut singularity for  $|\omega| > |\mathbf{k}|$ ,

$$G^R(\omega, \mathbf{k}) = \frac{N^2(-\omega^2 + \mathbf{k}^2)^2}{64\pi^2} (\ln |-\omega^2 + \mathbf{k}^2| - i\pi\theta(\omega^2 - \mathbf{k}^2) \text{sgn } \omega) . \quad (1.2)$$

In the high temperature limit  $\omega/T \ll 1$ ,  $|\mathbf{k}|/T \ll 1$ , the propagator is analytic in the complex  $\omega$ -plane,

$$G^R(\omega, \mathbf{k}) = -\frac{N^2 T^2}{16} (i 2\pi T \omega + \mathbf{k}^2) . \quad (1.3)$$

---

<sup>1</sup> A non-trivial check of the validity of the correspondence at finite temperature was made recently in [9].

However, for generic values of  $\omega$  and  $\mathbf{k}$ , we expect  $G^R(\omega, \mathbf{k})$  to have poles corresponding to the spectrum of collective excitations of the SYM plasma.

One can compare the situation to the simpler case of the  $2d$  CFT dual to the Bañados-Teitelboim-Zanelli (BTZ) black hole background. There, the retarded Green's functions can be computed exactly. For illustration, consider the case of the conformal dimension  $\Delta = 2$ . Then

$$G_{2d}^R(\omega, k) = \frac{\omega^2 - k^2}{4\pi^2} \left[ \psi \left( 1 - i \frac{\omega - k}{4\pi T} \right) + \psi \left( 1 - i \frac{\omega + k}{4\pi T} \right) \right], \quad (1.4)$$

where we have put  $T_L = T_R$  and ignored the constant prefactor for simplicity. The high temperature limit of (1.4) is an analytic function of  $\omega$ . In general, however,  $G_{2d}^R(\omega, k)$  has infinitely many poles located at

$$\omega_n = \pm k - i 4\pi T (n + 1), \quad n = 0, 1, \dots \quad (1.5)$$

When  $T \rightarrow 0$ , the poles merge forming branch cuts. One can use the asymptotic expansion  $\psi(z) \sim \log z - 1/2z + \dots$  (valid for  $|\arg z| < \pi/2$ ) to find the zero-temperature limit of (1.4) (ignoring the contact terms) :

$$G_{2d}^R \sim \frac{\omega^2 - k^2}{4\pi^2} \log |\omega^2 - k^2| - i \frac{\omega^2 - k^2}{4\pi} \theta(\omega^2 - k^2) \operatorname{sgn} \omega. \quad (1.6)$$

For spacelike momenta,  $|\omega| < k$ , the imaginary part of  $G_{2d}^R$  is exponentially suppressed.

Even though the retarded Green's function in  $4d$  cannot be found explicitly, the location of its singularities and thus the “dispersion law” of thermal excitations at strong coupling can be determined precisely. As shown in [10], this amounts to finding the quasinormal frequencies of dilaton's fluctuation in the dual near-extremal black brane background as functions of the spatial momentum<sup>2</sup>. Recently, there has been considerable interest (inspired by the AdS/CFT duality conjecture) in studying quasinormal modes of Schwarzschild- AdS black holes<sup>3</sup> [12,13,14,15,16,17,18]. Here, we consider the non-compact case directly. This corresponds to finding quasinormal frequencies for an infinitely large AdS black hole.

---

<sup>2</sup> Note that the poles (1.5) of  $G_{2d}^R(\omega, k)$  coincide with the quasinormal frequencies of BTZ black hole [11].

<sup>3</sup> References to the early works on quasinormal modes in asymptotically AdS spacetime as well as to works considering bulk dimension other than five can be found in [10], [12].

Computing the scalar quasinormal frequencies in the black brane background is equivalent to solving the two-parameter singular spectral problem for the equation with four regular singularities (Heun equation). In this paper, we solve this problem by using the elegant method based on Pincherle’s theorem. The eigenvalue equation is written in terms of continued fractions and solved numerically. Similar approach had been used by G. Jaffé [19] in 1933 to find the spectrum of the hydrogen molecule ion. Later, it has been applied to the problem of quasinormal modes in asymptotically flat spacetimes by E.W. Leaver [20].

The paper is organized as follows. In the next section, we formulate the problem of finding the quasinormal modes of near-extremal black three-brane in terms of the singular boundary value problem for Heun equation. In section 3 this boundary value problem is solved by analyzing the associated linear difference equation and applying Pincherle’s theorem. The results (quasinormal frequencies  $\omega_n$  and the “dispersion law”  $\omega(\mathbf{k})$ ) are presented in section 4. The discussion follows in section 5.

## 2. Quasinormal modes and the boundary value problem for Heun equation

The metric corresponding to the collection of  $N$  parallel non-extremal three-branes in the near-horizon limit is given by

$$ds^2 = \frac{r^2}{R^2} (-f dt^2 + d\mathbf{x}^2) + \frac{R^2}{r^2} (f^{-1} dr^2 + r^2 d\Omega_5^2), \quad (2.1)$$

where  $f(r) = 1 - r_0^4/r^4$ . According to the gauge theory/gravity correspondence, this background with the parameter of non-extremality  $r_0$  is dual to the  $\mathcal{N} = 4$   $SU(N)$  SYM at finite temperature  $T = r_0/\pi R^2$  in the limit  $N \rightarrow \infty$ ,  $g_{YM}^2 N \rightarrow \infty$ .

Using the new coordinate  $z = 1 - r_0^2/r^2$  and the Fourier decomposition

$$\phi(z, t, \mathbf{x}) = \int \frac{d^4 k}{(2\pi)^4} e^{-i\omega t + i\mathbf{k}\cdot\mathbf{x}} \phi_k(z), \quad (2.2)$$

the equation for the minimally coupled massless scalar in the background (2.1) reads

$$\phi_k'' + \frac{1 + (1 - z)^2}{z(1 - z)(2 - z)} \phi_k' + \frac{\lambda^2}{4z^2(1 - z)(2 - z)^2} \phi_k - \frac{q^2}{4z(1 - z)(2 - z)} \phi_k = 0, \quad (2.3)$$

where  $\lambda = \omega/\pi T$ ,  $q = |\vec{k}|/\pi T$ . Eq. (2.3) has four regular singularities at  $z = 0, 1, 2, \infty$ , the corresponding pairs of characteristic exponents being respectively  $\{-i\lambda/4, i\lambda/4\}$ ;  $\{0, 2\}$ ;  $\{-\lambda/4, \lambda/4\}$ ;  $\{0, 0\}$ .

Quasinormal modes are defined as solutions of Eq.(2.3) obeying the “incoming wave” boundary condition at the horizon  $z = 0$  and the vanishing Dirichlet boundary condition at spatial infinity  $z = 1$ . The first condition singles out the exponent  $\nu_0^{(1)} = -i\lambda/4$  at  $z = 0$ .

The most straightforward way to find quasinormal modes would be to construct a local series solution  $\varphi_{loc}(z, \lambda)$  to Eq.(2.3) with the exponent  $\nu_0^{(1)}$  near the origin, prove its convergence at  $z = 1$ , and determine the eigenfrequencies  $\lambda_n(q)$  by solving the equation  $\varphi_{loc}(1, \lambda) = 0$  numerically. This approach works quite well for the low-level eigenfrequencies, and in fact it has been successfully used in a number of publications on quasinormal modes in asymptotically *AdS* space. Here we would like to solve the above eigenvalue problem in a somewhat different way which, in our opinion, is more appealing both analytically and numerically.

By making a transformation of the dependent variable

$$\phi(z) = z^{-\frac{i\lambda}{4}} (z - 2)^{-\frac{\lambda}{4}} y(z), \quad (2.4)$$

Eq.(2.3) can be reduced to the standard form of the Heun equation [21,22],

$$y'' + \left[ \frac{\gamma}{z} + \frac{\delta}{z-1} + \frac{\epsilon}{z-2} \right] y' + \frac{\alpha\beta z - Q}{z(z-1)(z-2)} y = 0, \quad (2.5)$$

where  $\alpha, \beta, \gamma, \epsilon$  depend on  $\lambda$ ,

$$\alpha = \beta = -\frac{\lambda(1+i)}{4}, \quad \gamma = 1 - \frac{i\lambda}{2}, \quad \delta = -1, \quad \epsilon = 1 - \lambda/2, \quad (2.6)$$

and  $Q$  is the so called “accessory parameter” given in our case by

$$Q = \frac{q^2}{4} - \frac{\lambda(1-i)}{4} - \frac{\lambda^2(2-i)}{8}. \quad (2.7)$$

We would like to determine values of  $\lambda$  and  $q$  for which Eq. (2.5) on the interval  $[0, 1]$  has solutions obeying the boundary conditions  $y(0) = 1, y(1) = 0$ .

Before turning to the solution, let us remark that Eq. (2.3) with  $\lambda = 0$  has received some attention previously in connection with the so called “glueball mass spectrum” in *QCD*<sub>3</sub> [23], [24],[25], [26]. Approximate analytic expressions for the “glueball” masses squared  $M_n^2 = -q_n^2$  were found either via the “brute force” WKB calculation [26] or by appealing to the unpublished results of the RIMS group [23] (both approaches give satisfactory agreement with the results obtained by numerical integration). Introducing a non-zero  $\lambda$  and using the “incoming wave” boundary condition makes the problem considerably more complicated, as is evident from Eqs. (2.5) and (2.6) .

### 3. Solving the boundary value problem

#### 3.1. Local solutions

The Frobenius set of local solutions near each of the singularities can be easily constructed.

At  $z = 0$ , the local series solution corresponding to the index  $\nu = 0$  and normalized to 1 is given by

$$y_0(z) = \sum_{n=0}^{\infty} a_n(\lambda, k) z^n, \quad (3.1)$$

where  $a_0 = 1$ ,  $a_1 = \frac{Q}{2\gamma}$ , and the coefficients  $a_n$  with  $n \geq 2$  obey the three-term recursion relation

$$a_{n+2} + A_n(\lambda) a_{n+1} + B_n(\lambda) a_n = 0, \quad (3.2)$$

where

$$A_n(\lambda) = -\frac{(n+1)(2\delta + \epsilon + 3(n+\gamma)) + Q}{2(n+2)(n+1+\gamma)}, \quad (3.3)$$

$$B_n(\lambda) = \frac{(n+\alpha)(n+\beta)}{2(n+2)(n+1+\gamma)}. \quad (3.4)$$

The series (3.1) is absolutely convergent for  $|z| < 1$  and, in general, is divergent for  $|z| > 1$ . The condition for convergence at  $|z| = 1$  involves parameters of the equation and will be investigated below.

At  $z = 1$ , the difference of the exponents has an integer value, and we expect the local solution there to contain logarithms. Indeed, the set of local solutions is given by

$$y_1(z) = (1-z)^2 \left( 1 + b_1^{(1)}(1-z) + b_2^{(1)}(1-z)^2 + \dots \right), \quad (3.5)$$

$$y_2(z) = 1 + b_1^{(2)}(1-z) + h y_1(z) \log(1-z) + b_2^{(2)}(1-z)^2 + \dots, \quad (3.6)$$

where  $b_1^{(1)} = (q^2 - \lambda^2 + 3\lambda(1-i))/12$ ,  $b_1^{(2)} = (-q^2 + \lambda^2 + \lambda(1-i))/4$ ,  $h = -(q^2 - \lambda^2)^2/32$ , and the coefficients  $b_n^{(1,2)}$ ,  $n \geq 2$ , can be found recursively from the relation similar to the one in (3.2). Solving the differential equation Eq. (2.5) essentially means finding the connection between the sets of local solutions.

#### 3.2. The connection problem

From the general theory of linear differential equations, it follows that three solutions  $y_0(z)$ ,  $y_1(z)$ ,  $y_2(z)$  are connected on  $[0, 1]$  by the linear relation

$$y_0(z) = \mathcal{A}(\lambda, q) y_1(z) + \mathcal{B}(\lambda, q) y_2(z), \quad (3.7)$$

where  $\mathcal{A}$ ,  $\mathcal{B}$  are independent of  $z$ . The vanishing Dirichlet boundary condition for quasinormal modes at  $z = 1$  implies  $\mathcal{B} = 0$ , which is the equation for eigenfrequencies  $\lambda_n$ . If  $\lambda = \lambda_n$ ,  $y_0(z)$  is proportional to  $y_1(z)$ , and thus it is simultaneously a Frobenius solution about  $z = 0$  (with exponent 0) and a Frobenius solution about  $z = 1$  (with exponent 2). Such a solution is called a Heun function [27]. We learn that quasinormal modes of black branes are Heun functions. Unfortunately, the connection problem for the Heun equation remains unsolved, and explicit expressions for the coefficients  $\mathcal{A}$ ,  $\mathcal{B}$  are unavailable, with the exception of some special cases<sup>4</sup>. There is, however, an indirect way of determining for which values of  $\lambda$  and  $q$  the connection coefficient  $\mathcal{B}$  vanishes. This is achieved through an analysis of convergence.

### 3.3. The analysis of convergence

The convergence of the series (3.1) can be analyzed by studying the large  $n$  asymptotic behavior of the linear difference equation (3.2). One finds<sup>5</sup> that Eq. (3.2) possesses two linearly independent asymptotic solutions of the form

$$a_n^{(1)} \sim 2^{-n} n^{-1-\frac{\lambda}{2}} \sum_{s=0}^{\infty} \frac{c_s^{(1)}}{n^s}, \quad (3.8)$$

$$a_n^{(2)} \sim n^{-3} \sum_{s=0}^{\infty} \frac{c_s^{(2)}}{n^s}, \quad (3.9)$$

where the coefficients  $c_s^{(1,2)}$  can be found recursively using the asymptotic expansion of  $A(n)$ ,  $B(n)$ . In particular, we have  $c_1^{(1)} = (q^2 - \lambda^2 + \lambda(1+i) + 2\lambda^2(1+i))/4$ ,  $c_1^{(2)} = (12 - q^2 + \lambda^2 - 3(1+i)\lambda)/4$ . Equation (3.8) is called a *minimal* solution to Eq. (3.2), and Eq. (3.9) represents a *dominant* one. This distinction reflects the property

$$\lim_{n \rightarrow \infty} \frac{a_n^{(1)}}{a_n^{(2)}} = 0, \quad (3.10)$$

---

<sup>4</sup> Schäfke and Schmidt [28] give the connection coefficients in terms of the  $n \rightarrow \infty$  limit of  $a_n$  obeying Eq. (3.2). Although the asymptotic behavior of  $a_n$  near  $n = \infty$  can be studied by means of local analysis, it only determines  $a_n$  up to a  $\lambda$ -dependent coefficient which is essentially the very quantity we are looking for.

<sup>5</sup> Necessary information about second order linear difference equations can be found, for example, in the excellent review paper by Wong and Li [29].

and will be useful later on. Using Eqs. (3.8) , (3.9) , we obtain

$$\left| \frac{a_{n+1}^{(1)}}{a_n^{(1)}} \right| = \frac{1}{2} \left[ 1 - \frac{2 + \operatorname{Re} \lambda}{2n} + O\left(\frac{1}{n^2}\right) \right], \quad (3.11)$$

$$\left| \frac{a_{n+1}^{(2)}}{a_n^{(2)}} \right| = 1 - \frac{3}{n} + O\left(\frac{1}{n^2}\right). \quad (3.12)$$

Equations (3.11) , (3.12) imply<sup>6</sup> that generically the series (3.1) converges absolutely for  $|z| \leq 1$  for *any* value of  $\lambda$ . This proves that the equation

$$\sum_{n=0}^{\infty} a_n(\lambda, k) = 0 \quad (3.13)$$

can indeed be used to compute the quasinormal frequencies numerically. What is more interesting, however, is that in some cases the radius of convergence increases. It happens when the minimal solution (3.8) exists. In that case the series (3.1) converges absolutely for  $|z| < 2$  and thus represents an analytic function (the Heun function) in that region.

Thus, even though the connection coefficient  $\mathcal{B}(\lambda, q)$  in Eq. (3.7) remains unknown, finding zeros of  $\mathcal{B}(\lambda, q)$  is equivalent to finding a condition under which the minimal solution to Eq. (3.2) exists. Such a condition is conveniently supplied by Pincherle's theorem [31] which states that the minimal solution exists if and only if the continued fraction

$$-\frac{B_0(\lambda)}{A_0(\lambda)-} \frac{B_1(\lambda)}{A_1(\lambda)-} \frac{B_2(\lambda)}{A_2(\lambda)-} \dots \quad (3.14)$$

converges. Moreover, in case of convergence one has

$$\frac{a_{n+1}}{a_n} = -\frac{B_n(\lambda)}{A_n(\lambda)-} \frac{B_{n+1}(\lambda)}{A_{n+1}(\lambda)-} \frac{B_{n+2}(\lambda)}{A_{n+2}(\lambda)-} \dots. \quad (3.15)$$

The right hand side of Eq. (3.15) is generated by a simple algorithm. Define  $r_n = a_{n+1}/a_n$ . Then Eq.(3.2) can be written as

$$r_n = -\frac{B_n(\lambda)}{A_n(\lambda) + r_{n+1}}. \quad (3.16)$$

Applying (3.16) repeatedly, one gets Eq. (3.15) . Now, setting  $n = 0$  in Eq. (3.15) we have

$$\frac{Q}{2 - i\lambda} = -\frac{B_0(\lambda)}{A_0(\lambda)-} \frac{B_1(\lambda)}{A_1(\lambda)-} \frac{B_2(\lambda)}{A_2(\lambda)-} \dots. \quad (3.17)$$

---

<sup>6</sup> See [30] , §2.37.



This is the transcendental eigenvalue equation which determines the quasinormal frequencies. Equation (3.17) can be solved numerically with great efficiency using a variety of methods (see [32], which also includes a discussion of the error analysis). Here we use a nonlinear backward recursion which amounts to breaking the continued fraction (3.16) by setting  $r_n$  to zero<sup>7</sup> for some large  $n_*$  and computing backward to get  $r_0$ . Stability can be checked by choosing a larger value of  $n_*$  and repeating the calculation.

#### 4. Quasinormal frequencies

For  $q = 0$ , the lowest 15 quasinormal frequencies  $\lambda_n$  obtained by solving Eq. (3.17) numerically are listed in Table 1 and shown in Fig. 1. The results suggest that the number of frequencies is infinite, and that their large  $n$  asymptotic behavior is given by the simple formula

$$\lambda_n^\pm = \lambda_0^\pm \pm 2n(1 \mp i), \quad (4.1)$$

where  $\lambda_0^\pm \approx \pm 1.2139 - 0.7775i$ . In terms of the original variable  $\omega$ , Eq. (4.1) is written as

$$\omega_n^\pm = \omega_0^\pm \pm 2\pi T n(1 \mp i), \quad (4.2)$$

where the coefficient in front of the parentheses is the bosonic Matsubara frequency and  $\omega_0^\pm = \pi T \lambda_0^\pm$ .

The lowest eigenfrequency in Table 1 can be compared with the result of Horowitz and Hubeny [12] for a large  $5d$  Schwarzschild- AdS black hole. Normalizing the  $r_+ = 100$  entry in Table 1 of [12] appropriately ( $\lambda = \omega R^2/r_+$ ,  $R = 1$ ), we have  $\lambda^{(100)} \approx 3.119627 - 2.746655i$  which is fairly close to our result  $\lambda^{(\infty)} \approx 3.119452 - 2.746667i$ .

The dependence of the lowest five quasinormal frequencies on  $q$  for  $q \in [0, 100]$  is shown in Figs. 2-4. All five branches stay above the line  $\text{Re } \lambda = q$  ( $\text{Re } \omega = |\mathbf{k}|$ ) slowly approaching it in the low-temperature limit  $q \rightarrow \infty$ . The imaginary part of the branches tends to zero in the same limit. Figures 2-4 bear resemblance to the dispersion law of thermal excitations in a weakly coupled Yang-Mills plasma<sup>8</sup>.


---

<sup>7</sup> To improve the convergence of continued fractions, one can instead set  $r_n$  to its asymptotic value at  $n_*$ ,  $r_n = 1/2 - (2 + \lambda)/4n_* + \dots$ , as suggested by Nollert [33].

<sup>8</sup> Note, however, that in this paper we consider correlators of the gauge-invariant operators rather than the gluon propagator.


$n$	$\text{Re } \lambda_n$	$\text{Im } \lambda_n$
1	$\pm 3.119452$	$-2.746676$
2	$\pm 5.169521$	$-4.763570$
3	$\pm 7.187931$	$-6.769565$
4	$\pm 9.197199$	$-8.772481$
5	$\pm 11.202676$	$-10.774162$
6	$\pm 13.206247$	$-12.775239$
7	$\pm 15.208736$	$-14.775979$
8	$\pm 17.210558$	$-16.776515$
9	$\pm 19.211943$	$-18.776919$
10	$\pm 21.213025$	$-20.777232$
11	$\pm 23.213896$	$-22.777489$
12	$\pm 25.213896$	$-24.777489$
13	$\pm 27.213896$	$-26.777489$
14	$\pm 29.213896$	$-28.777489$
15	$\pm 31.213896$	$-30.777489$

Table 1: The lowest quasinormal frequencies  $\lambda_n$  for  $q = 0$ .




**Fig. 1:** The lowest 15 quasinormal frequencies in the complex  $\lambda$ -plane for  $q = 0$ .

All quasinormal frequencies have negative imaginary parts. This fact can be proven rigorously using an argument based on the “energy-type” integral [34] or by writing Eq. (2.3) in the Regge-Wheeler form and essentially repeating the proofs for AdS black holes given recently in [12], [14]. It reflects the stability of the near-extremal metric against



**Fig. 2:**  $\text{Re } \lambda$  vs  $q$  (with the interval  $\Delta q = 0.5$ ) for the lowest five quasinormal frequencies. The zeros approach the line  $\text{Re } \lambda = q$  as  $q \rightarrow \infty$ .




**Fig. 3:**  $\text{Re } \lambda$  vs  $q$  (with the interval  $\Delta q = 1$ ) for the lowest five quasinormal frequencies.

a scalar perturbation. On the field theory side, the negative sign corresponds to the damping of the plasma excitations.

The computed frequencies appear to be symmetric with respect to the imaginary  $\lambda$  axis, i.e., the eigenvalue equation (3.17) seems to give complex-conjugate pairs of solutions in terms of the variable  $i\lambda$ . This symmetry does not seem to be explicit in Eq. (3.17)<sup>9</sup>.

---

<sup>9</sup> The symmetry, however, is obvious in the original form of the equation (2.3). I thank A. V. Shchepetilov for pointing this out to me.



**Fig. 4:**  $-\text{Im } \lambda$  vs  $q$  (with the interval  $\Delta q = 1$ ) for the lowest five quasinormal frequencies.

## 5. Discussion of results

The results are compatible with expectations outlined in the Introduction. The retarded Green’s function has infinitely many poles in the complex  $\omega$ -plane whose location depends on the spatial momentum. There is a “mass gap”: for small enough values of  $\lambda$  the propagator  $G^R(\omega, \mathbf{k})$  is analytic in agreement with Eq. (1.3). In the low-temperature limit  $\lambda \rightarrow \infty$ ,  $q \rightarrow \infty$  singularities merge, forming branch cuts of the zero-temperature propagator <sup>10</sup> (1.2) as in the BTZ case.

The distribution of quasinormal frequencies in the complex  $\omega$  plane for an asymptotically AdS background appears to follow a much simpler pattern than the one corresponding to the asymptotically flat case (compare Fig. 1 in [20] ). In particular, Chandrasekhar’s “algebraically special” solution is absent: there are no frequencies with  $\text{Re } \lambda = 0$ . Obviously, all statements about the behaviour of higher order modes are conjectural. It would be very desirable to confirm the asymptotic formula (4.1) analytically, possibly by using the complex WKB method. We remark, however, that the analogous problem remains unsolved even in the much studied case of a Schwarzschild black hole in flat space.

Spectra of excitations in more realistic theories can be similarly studied provided their gravitational duals are known explicitly. It would also be interesting to determine the poles of the current and energy-momentum tensor correlators at finite temperature,

---

<sup>10</sup> The zero-temperature limit (1.2) of  $G^R(\omega, \mathbf{k})$  can be obtained from the Heun equation using the Langer-Olver asymptotic expansion (see [10]).

whose hydrodynamic limit has recently been computed in [9] .

Finally, studying gravitational quasinormal modes may be important in investigating the stability of the non extremal black brane background.

### **Acknowledgments**

It is a pleasure to thank P.K. Kovtun, D.T. Son and L.G. Yaffe for discussions. I would also like to thank A. Nuñez-Nikitin and A. V. Shchepetilov for valuable comments on the manuscript. This work is supported in part by DOE grant No. DOE-ER-41132.

## References

- [1] J. M. Maldacena, “The large  $N$  limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2**, 231 (1998) [*Int. J. Theor. Phys.* **38**, 1113 (1999)] [arXiv:hep-th/9711200].
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Phys. Lett. B* **428**, 105 (1998) [arXiv:hep-th/9802109].
- [3] E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2**, 253 (1998) [arXiv:hep-th/9802150].
- [4] S. S. Gubser, I. R. Klebanov and A. W. Peet, “Entropy and Temperature of Black 3-Branes,” *Phys. Rev. D* **54**, 3915 (1996) [arXiv:hep-th/9602135].
- [5] S. S. Gubser, I. R. Klebanov and A. A. Tseytlin, “Coupling constant dependence in the thermodynamics of  $N = 4$  supersymmetric Yang-Mills theory,” *Nucl. Phys. B* **534**, 202 (1998) [arXiv:hep-th/9805156].
- [6] S. J. Rey, S. Theisen and J. T. Yee, “Wilson-Polyakov loop at finite temperature in large  $N$  gauge theory and anti-de Sitter supergravity,” *Nucl. Phys. B* **527**, 171 (1998) [arXiv:hep-th/9803135].
- [7] A. Brandhuber, N. Itzhaki, J. Sonnenschein and S. Yankielowicz, “Wilson loops in the large  $N$  limit at finite temperature,” *Phys. Lett. B* **434**, 36 (1998) [arXiv:hep-th/9803137].
- [8] G. Policastro, D. T. Son and A. O. Starinets, “The shear viscosity of strongly coupled  $N = 4$  supersymmetric Yang-Mills plasma,” *Phys. Rev. Lett.* **87**, 081601 (2001) [arXiv:hep-th/0104066].
- [9] G. Policastro, D. T. Son and A. O. Starinets, “From AdS/CFT correspondence to hydrodynamics I,II,” arXiv:hep-th/0205052, hep-th/0210220.
- [10] D. T. Son and A. O. Starinets, “Minkowski-space correlators in AdS/CFT correspondence: Recipe and applications,” *JHEP* **0209**, 042 (2002) [arXiv:hep-th/0205051].
- [11] D. Birmingham, I. Sachs and S. N. Solodukhin, “Conformal field theory interpretation of black hole quasi-normal modes,” *Phys. Rev. Lett.* **88**, 151301 (2002) [arXiv:hep-th/0112055].
- [12] G. T. Horowitz and V. E. Hubeny, “Quasinormal modes of AdS black holes and the approach to thermal equilibrium,” *Phys. Rev. D* **62**, 024027 (2000) [arXiv:hep-th/9909056].
- [13] B. Wang, C. Y. Lin and E. Abdalla, *Phys. Lett. B* **481**, 79 (2000) [arXiv:hep-th/0003295].
- [14] V. Cardoso and J. P. Lemos, “Quasi-normal modes of Schwarzschild anti-de Sitter black holes: Electromagnetic and gravitational perturbations,” *Phys. Rev. D* **64**, 084017 (2001) [arXiv:gr-qc/0105103].
- [15] W. T. Kim and J. J. Oh, *Phys. Lett. B* **514**, 155 (2001) [arXiv:hep-th/0105112].

- [16] V. Cardoso and J. P. Lemos, “Quasi-normal modes of toroidal, cylindrical and planar black holes in anti-de Sitter spacetimes,” *Class. Quant. Grav.* **18**, 5257 (2001) [arXiv:gr-qc/0107098].
- [17] I. G. Moss and J. P. Norman, *Class. Quant. Grav.* **19**, 2323 (2002) [arXiv:gr-qc/0201016].
- [18] R. A. Konoplya, “On quasinormal modes of small Schwarzschild-anti-de-Sitter black hole,” arXiv:hep-th/0205142.
- [19] G. Jaffé, “Zur Theorie des Wasserstoffmolekülions”, *Z. Phys.* **87**, 535 (1933).
- [20] E.W. Leaver, “An analytic representation for the quasi-normal modes of Kerr black holes”, *Proc. R. Soc. Lond. A* **402** 285 (1985).
- [21] K. Heun, “Zur Theorie der Riemann’s sehen Functionen zweiter Ordnung mit vier Verzweigungspunkten”, *Math. Ann.* **33** (1889) 61.
- [22] *Heun’s differential equation*, ed. A. Ronveaux, F. Arscott, Oxford University Press, Oxford (1995).
- [23] C. Csaki, H. Ooguri, Y. Oz and J. Terning, “Glueball mass spectrum from supergravity,” *JHEP* **9901**, 017 (1999) [arXiv:hep-th/9806021].
- [24] R. de Mello Koch, A. Jevicki, M. Mihailescu and J. P. Nunes, “Evaluation of glueball masses from supergravity,” *Phys. Rev. D* **58**, 105009 (1998) [arXiv:hep-th/9806125].
- [25] M. Zyskin, *Phys. Lett. B* **439**, 373 (1998) [arXiv:hep-th/9806128].
- [26] J. A. Minahan, *JHEP* **9901**, 020 (1999) [arXiv:hep-th/9811156].
- [27] *Higher transcendental functions, vol. III*, ed. A. Erdelyi, McGraw Hill, New York (1955).
- [28] R. Schäfke and D. Schmidt, “The connection problem for general linear ordinary differential equations at two regular singular points with applications to the theory of special functions”, *SIAM J. Math. Anal.* **11** (1980) 848.
- [29] R. Wong and H. Li, “Asymptotic expansions for second order linear difference equations”, *J. Comput. Appl. Math.* **41** (1992) 65.
- [30] E.T. Whittaker and G.N. Watson, *A course of modern analysis*, Cambridge University Press, 1927.
- [31] S. Pincherle, “Delle funzioni ipergeometriche e de varie questione ad esse attinenti”, *Giorn. mat. Battaglini* **32** (1894) 209.
- [32] *Theory and Application of Special Functions*, Ed. R. Askey, Academic Press, New York (1975).
- [33] H.-P. Nollert, “Quasinormal modes of Schwarzschild black holes: The determination of quasinormal frequencies with very large imaginary parts”, *Phys. Rev. D* **47** (1993) 5253.
- [34] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, Oxford University Press, New York, 1983.