

Notes on Five-dimensional Kerr Black Holes

Makoto Sakaguchi* and Yukinori Yasui*

* *Osaka City University Advanced Mathematical Institute (OCAMI)*

* *Department of Mathematics and Physics, Graduate School of Science,
Osaka City University*

Sumiyoshi, Osaka 558-8585, JAPAN

Abstract

The geometry of five-dimensional Kerr black holes is discussed based on geodesics and Riemannian curvatures. Kerr-Star space, Star-Kerr space and Kruskal space are naturally introduced by using special null geodesics. We show that the geodesics of AdS Kerr black hole are integrable, which generalizes the result of Frolov and Stojkovic.

*msakaguc@sci.osaka-cu.ac.jp

*yasui@sci.osaka-cu.ac.jp

1 Introduction

Black holes have attracted renewed interests in the recent developments of string theory and Riemannian geometry, such as the AdS/CFT correspondence [1] and a certain relation between compact Einstein manifolds and black holes [2] [3] [4] [5]. These developments motivate us to study the geometry of black holes especially in four-, five- and seven-dimensions among higher dimensional black holes.

This paper is devoted to a first step to investigate the global geometry of five-dimensional Kerr black holes constructed by Myers and Perry [6], and Hawking et al [7]. For four-dimensional Kerr black holes, a valuable textbook [8] written by B. O'Neill from mathematical point of view has been published. In the textbook, the differential geometry based on special null geodesics (principal null geodesics) were fully analyzed. Following the textbook we try to generalize the analysis to five-dimensional black holes. In this paper we do not stick to mathematical completeness, but develop some key points of the geometry; integrability of geodesics and curvature property. These points were studied in the previous works [9] and [10], however, our method is different from them. In addition, AdS black holes have not been discussed in the textbook.

This paper is organized as follows. In the next section, we examine the curvature property of five-dimensional black holes. We show that a linear map on two-forms constructed from Riemannian curvature is diagonalizable, and derive the eigenvalues and the degeneracy. In section 3, we examine the integrability of geodesics based on the Euler-Lagrange equations. The section 4 is devoted to the global analysis of five-dimensional Kerr black holes. In the last section, we generalize the analysis to the case with cosmological constant.

2 5-dimensional Kerr Black Holes

Let us write the Ricci flat metric for the 5-dimensional Kerr black hole [6] in the Boyer-Lindquist coordinates $(t, \phi, \psi, r, \theta)$;

$$g_{BH} = -dt^2 + \frac{\rho^2}{\Delta_r} dr^2 + \rho^2 g_{S^3} + (a^2 - b^2)(\sin^2 \theta d\phi)^2 + (b^2 - a^2)(\cos^2 \theta d\psi)^2 + \frac{2m}{\rho^2} (dt - a \sin^2 \theta d\phi - b \cos^2 \theta d\psi)^2, \quad (2.1)$$

where

$$\rho^2 = r^2 + a^2 \cos^2 \theta + b^2 \sin^2 \theta, \quad (2.2)$$

$$\Delta_r = \frac{(r^2 + a^2)(r^2 + b^2)}{r^2} - 2m . \quad (2.3)$$

The non-negative parameters a , b and m correspond to angular momenta and mass, respectively. The metric g_{S^3} is given by

$$g_{S^3} = d\theta^2 + \sin^2 \theta d\phi^2 + \cos^2 \theta d\psi^2 \quad (2.4)$$

with the ranges $0 \leq \theta \leq \pi/2$ and $0 \leq \phi, \psi \leq 2\pi$. If we introduce an orthonormal coordinate (x, y, z, w) ;

$$x = \sin \theta \cos \phi , \quad y = \sin \theta \sin \phi , \quad z = \cos \theta \cos \psi , \quad w = \cos \theta \sin \psi , \quad (2.5)$$

then

$$g_{S^3} = dx^2 + dy^2 + dz^2 + dw^2 , \quad (2.6)$$

which represents the standard metric on a three-sphere S^3 . Also, the one-forms $\sin^2 \theta d\phi$ and $\cos^2 \theta d\psi$ in the metric (2.4) are written as

$$\sin^2 \theta d\phi = xdy - ydx , \quad \cos^2 \theta d\psi = zdw - wdz , \quad (2.7)$$

and hence these are well-defined as one-forms on S^3 . Thus g_{BH} can be regarded as a metric on the Lorentzian space

$$M = \mathbb{R}_+^2 \times S^3 - H , \quad (2.8)$$

where $\mathbb{R}_+^2 = \{(t, r) \mid t \in \mathbb{R}, r \in \mathbb{R}^+\}$. The horizon $H \simeq \mathbb{R} \times S^3$ defined by $\Delta_r = 0$ is removed from the space since $\rho^2/\Delta_r \rightarrow \infty$ on H . The equation $\Delta_r = 0$ has two positive real roots $r = r_{\pm}$,

$$r_{\pm}^2 = \frac{1}{2} \left(2m - a^2 - b^2 \pm \sqrt{(2m - (a - b)^2)(2m - (a + b)^2)} \right) \quad (2.9)$$

in the region $m > (a + b)^2/2$. We have three open sets I, II and III (called Boyer-Lindquist blocks) of M :

$$\text{I} : r_+ < r , \quad \text{II} : r_- < r < r_+ , \quad \text{III} : 0 < r < r_- . \quad (2.10)$$

It should be noticed that three-dimensional submanifolds $M|_{\theta=0}$ and $M|_{\theta=\pi/2}$ are time-like totally geodesic, while two-dimensional submanifold $M|_{t, \phi, \psi = \text{const}}$ is space-like totally

geodesic. This can be seen from calculations of the Christoffel symbol; $\Gamma_{\mu\nu}^\theta \propto \sin 2\theta$ and $\Gamma_{\mu\nu}^i = 0$ ($i = t, \phi, \psi$), where μ and ν run the tangential indices to the submanifolds.

Let us calculate the Riemannian curvature of the black hole metric. It is convenient to use the following orthonormal frame $\{e_a\}$ ($a = 1, 2, \dots, 5$) [10]:

$$e_1 = \frac{1}{r^2 \sqrt{\Delta_r \varepsilon} \rho} \left(\Sigma_a^2 \Sigma_b^2 \frac{\partial}{\partial t} + a \Sigma_b^2 \frac{\partial}{\partial \phi} + b \Sigma_a^2 \frac{\partial}{\partial \psi} \right), \quad (2.11)$$

$$e_2 = \frac{1}{r \rho \sin \theta} \left[\Sigma_b \left(\frac{\partial}{\partial \phi} + a \sin^2 \theta \frac{\partial}{\partial t} \right) + \frac{b \Sigma_a}{\Sigma_a \Sigma_b + r \rho} \left(a \sin^2 \theta \frac{\partial}{\partial \psi} - b \cos^2 \theta \frac{\partial}{\partial \phi} \right) \right], \quad (2.12)$$

$$e_3 = \frac{1}{r \rho \cos \theta} \left[\Sigma_a \left(\frac{\partial}{\partial \psi} + b \cos^2 \theta \frac{\partial}{\partial t} \right) - \frac{a \Sigma_b}{\Sigma_a \Sigma_b + r \rho} \left(a \sin^2 \theta \frac{\partial}{\partial \psi} - b \cos^2 \theta \frac{\partial}{\partial \phi} \right) \right], \quad (2.13)$$

$$e_4 = \frac{\sqrt{\Delta_r \varepsilon}}{\rho} \frac{\partial}{\partial r}, \quad (2.14)$$

$$e_5 = \frac{1}{\rho} \frac{\partial}{\partial \theta}, \quad (2.15)$$

where $\Sigma_a = \sqrt{r^2 + a^2}$ and $\Sigma_b = \sqrt{r^2 + b^2}$. The inner product is given by $\langle e_a, e_b \rangle_{g_{BH}} = \eta_{ab}$ with $\eta_{ab} = \text{diag}(-\varepsilon, +1, +1, \varepsilon, +1)$. The time-like vector is e_1 in I \cup III, while e_4 in II, so that we choose $\varepsilon = 1$ in I \cup III and $\varepsilon = -1$ in II.

The curvature R_{cd}^{ab} may be regarded as a linear map (curvature transformation) on two-forms $\Lambda^2(M)$. Then, the matrix $R : \Lambda^2(M) \rightarrow \Lambda^2(M)$ is of the form $R_{cd}^{ab} = \frac{2mr^2}{\rho^6} \hat{R}_{cd}^{ab}$. Explicitly, non-zero components of \hat{R}_{cd}^{ab} are given as

$(a, b) \setminus (c, d)$	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(2, 3)
(1, 2)	$-1 + \mathbf{I}^2 - \mathbf{J}^2$	$-2\mathbf{IJ}$			
(1, 3)	$-2\mathbf{IJ}$	$-1 - \mathbf{I}^2 - \mathbf{J}^2$			
(1, 4)			$3 - \mathbf{I}^2 - \mathbf{J}^2$		
(1, 5)				$-1 + \mathbf{I}^2 + \mathbf{J}^2$	
(2, 3)					$1 + \mathbf{I}^2 + \mathbf{J}^2$
(2, 4)				$-2\varepsilon\mathbf{I}$	
(2, 5)			$-4\mathbf{I}$		
(3, 4)				$2\varepsilon\mathbf{J}$	
(3, 5)			$4\mathbf{J}$		
(4, 5)	$-2\varepsilon\mathbf{I}$	$2\varepsilon\mathbf{J}$			

$(a, b) \setminus (c, d)$	$(2, 4)$	$(2, 5)$	$(3, 4)$	$(3, 5)$	$(4, 5)$
$(1, 2)$					$2\varepsilon \mathbf{I}$
$(1, 3)$					$-2\varepsilon \mathbf{J}$
$(1, 4)$		$4\mathbf{I}$		$-4\mathbf{J}$	
$(1, 5)$	$2\varepsilon \mathbf{I}$		$-2\varepsilon \mathbf{J}$		
$(2, 3)$					
$(2, 4)$	$-1 + \mathbf{I}^2 - \mathbf{J}^2$		$-2\mathbf{IJ}$		
$(2, 5)$		$1 - 3\mathbf{I}^2 + \mathbf{J}^2$		$4\mathbf{IJ}$	
$(3, 4)$	$-2\mathbf{IJ}$		$-1 - \mathbf{I}^2 + \mathbf{J}^2$		
$(3, 5)$		$4\mathbf{IJ}$		$1 + \mathbf{I}^2 - 3\mathbf{J}^2$	
$(4, 5)$					$-1 + \mathbf{I}^2 + \mathbf{J}^2$

(2.16)

where

$$\mathbf{I} = \frac{a \cos \theta}{r} \frac{\Sigma_a r + \Sigma_b \rho}{\Sigma_a \Sigma_b + r \rho}, \quad \mathbf{J} = \frac{b \sin \theta}{r} \frac{\Sigma_b r + \Sigma_a \rho}{\Sigma_a \Sigma_b + r \rho}. \quad (2.17)$$

We find that the 10×10 matrix R_{cd}^{ab} is diagonalizable and has the following eigenvalues

$$2 : \lambda_1 = \frac{2mr^2}{\rho^6} (1 + \mathbf{I}^2 + \mathbf{J}^2) = \frac{2m}{\rho^4}, \quad (2.18)$$

$$2 : \lambda_2 = -\frac{2mr^2}{\rho^6} (1 + \mathbf{I}^2 + \mathbf{J}^2) = -\frac{2m}{\rho^4}, \quad (2.19)$$

$$\begin{aligned} 2 : \lambda_3 &= \frac{2mr^2}{\rho^6} (-1 + \mathbf{I}^2 + \mathbf{J}^2 + 2\sqrt{-\mathbf{I}^2 - \mathbf{J}^2}) \\ &= \frac{2m}{\rho^6} \left(\rho^2 - 2r^2 + 2ir\sqrt{\rho^2 - r^2} \right), \end{aligned} \quad (2.20)$$

$$\begin{aligned} 2 : \lambda_4 &= \frac{2mr^2}{\rho^6} (-1 + \mathbf{I}^2 + \mathbf{J}^2 - 2\sqrt{-\mathbf{I}^2 - \mathbf{J}^2}) \\ &= \frac{2m}{\rho^6} \left(\rho^2 - 2r^2 - 2ir\sqrt{\rho^2 - r^2} \right), \end{aligned} \quad (2.21)$$

$$\begin{aligned} 1 : \lambda_5 &= \frac{2mr^2}{\rho^6} (2 - 2\mathbf{I}^2 - 2\mathbf{J}^2 + \sqrt{1 - 14\mathbf{I}^2 - 14\mathbf{J}^2 + \mathbf{I}^4 + 2\mathbf{I}^2\mathbf{J}^2 + \mathbf{J}^4}) \\ &= \frac{2m}{\rho^6} \left(-2\rho^2 + 4r^2 + \sqrt{(r^2 - (7 + 4\sqrt{3})(\rho^2 - r^2))(r^2 - (7 - 4\sqrt{3})(\rho^2 - r^2))} \right), \end{aligned} \quad (2.22)$$

$$\begin{aligned} 1 : \lambda_6 &= \frac{2mr^2}{\rho^6} (2 - 2\mathbf{I}^2 - 2\mathbf{J}^2 - \sqrt{1 - 14\mathbf{I}^2 - 14\mathbf{J}^2 + \mathbf{I}^4 + 2\mathbf{I}^2\mathbf{J}^2 + \mathbf{J}^4}) \\ &= \frac{2m}{\rho^6} \left(-2\rho^2 + 4r^2 - \sqrt{(r^2 - (7 + 4\sqrt{3})(\rho^2 - r^2))(r^2 - (7 - 4\sqrt{3})(\rho^2 - r^2))} \right), \end{aligned}$$

$$(2.23)$$

where the number in the left hand side represents the degeneracy.

The curvature invariants $\text{tr}(R^n)$ ($n = 1, 2, 3, \dots$) can be simply written as

$$\text{tr}(R^n) = 2 \sum_{i=1}^4 (\lambda_i)^n + (\lambda_5)^n + (\lambda_6)^n = \left(\frac{2m}{\rho^6} \right)^n P_n(r^2, \theta) , \quad (2.24)$$

where P_n is an n -polynomial with respect to r^2 . For example, we have

$$P_1 = 0 , \quad (2.25)$$

$$P_2 = 6r^4(3\mathbf{I}^2 + 3\mathbf{J}^2 - 2)(\mathbf{I}^2 + \mathbf{J}^2 - 3) , \quad (2.26)$$

$$P_3 = -24r^6(\mathbf{I}^2 + \mathbf{J}^2 - 1) ((\mathbf{I}^2 + \mathbf{J}^2)^2 - 6\mathbf{I}^2 - 6\mathbf{J}^2 + 1) . \quad (2.27)$$

The invariants are finite for $ab \neq 0$. However, when we extend the square of the radial coordinate r^2 to a negative region, a singularity appears at $\rho^2 = 0$, i.e. $r^2 = -a^2 \cos^2 \theta - b^2 \sin^2 \theta$.

For the two-dimensional subspace Π_{ab} spanned by $\{e_a, e_b\}$, the sectional curvature is defined by $K(\Pi_{ab}) = \eta_{bb} R_{abab}$. We find from (2.16) and (2.18)-(2.21) the following relation between the sectional curvature and the eigenvalue with degeneracy 2:

$$K(\Pi_{15}) = K(\Pi_{45}) = -\text{Re}\lambda_3 = -\text{Re}\lambda_4 , \quad (2.28)$$

$$K(\Pi_{23}) = \lambda_1 = -\lambda_2 . \quad (2.29)$$

3 Geodesics

In this section, we examine geodesics and show the integrability of them. The integrability of geodesics has been shown in [9] by using Hamilton-Jacobi formulation, while we work in the Euler-Lagrange formulation.

The geodesic equations are given by the Euler-Lagrange equations. From the metric (2.1) we obtain a Lagrangian

$$\mathcal{L} = \frac{1}{2} g_{\alpha\beta} \frac{dx^\alpha}{ds} \frac{dx^\beta}{ds} , \quad (\alpha, \beta = t, \phi, \psi, r, \theta) , \quad (3.1)$$

where

$$g_{tt} = -1 + \frac{2m}{\rho^2} , \quad (3.2)$$

$$g_{t\phi} = -\frac{2am \sin^2 \theta}{\rho^2} , \quad (3.3)$$

$$g_{t\psi} = -\frac{2bm \cos^2 \theta}{\rho^2} , \quad (3.4)$$

$$g_{\phi\psi} = \frac{2abm \cos^2 \theta \sin^2 \theta}{\rho^2} , \quad (3.5)$$

$$g_{\phi\phi} = \sin^2 \theta \left(r^2 + a^2 + \frac{2ma^2 \sin^2 \theta}{\rho^2} \right) , \quad (3.6)$$

$$g_{\psi\psi} = \cos^2 \theta \left(r^2 + b^2 + \frac{2mb^2 \cos^2 \theta}{\rho^2} \right) , \quad (3.7)$$

$$g_{rr} = \frac{\rho^2}{\Delta_r} , \quad (3.8)$$

$$g_{\theta\theta} = \rho^2 . \quad (3.9)$$

The metric has three Killing vector fields $\{\frac{\partial}{\partial t}, \frac{\partial}{\partial \phi}, \frac{\partial}{\partial \psi}\}$, and hence there are three conservations L_t , L_ϕ and L_ψ . In addition, we have a “total energy” Q corresponding to the Hamiltonian. Let us write a geodesic as $\gamma(s) = (t(s), \phi(s), \psi(s), r(s), \theta(s))$. Then the conservations above are given by

$$L_t = \langle \gamma', \frac{\partial}{\partial t} \rangle , \quad L_\phi = \langle \gamma', \frac{\partial}{\partial \phi} \rangle , \quad L_\psi = \langle \gamma', \frac{\partial}{\partial \psi} \rangle , \quad Q = \langle \gamma', \gamma' \rangle , \quad (3.10)$$

where $\gamma' = \frac{d\gamma}{ds}$, and $\langle \cdot, \cdot \rangle$ represents an inner product with respect to the black hole metric.

In addition, there exists another conservation generalizing the Carter constant in the four-dimensional Kerr black hole [11].

Theorem 1. (see also [9])

There exists a constant K for a geodesic γ satisfying

$$\rho^4 \left(\frac{dr}{ds} \right)^2 + R(r) = 0 , \quad (3.11)$$

$$\rho^4 \left(\frac{d\theta}{ds} \right)^2 + \Theta(\theta) = 0 , \quad (3.12)$$

where

$$\Theta = (a^2 \sin^2 \theta + b^2 \cos^2 \theta) L_t^2 + \frac{L_\phi^2}{\sin^2 \theta} + \frac{L_\psi^2}{\cos^2 \theta} - (a^2 \cos^2 \theta + b^2 \sin^2 \theta) Q - K , \quad (3.13)$$

$$R = \frac{R_{-2}}{r^2} + R_0 + r^2 R_2 + r^4 R_4 . \quad (3.14)$$

The coefficients are explicitly given by

$$R_4 = -Q - L_t^2 , \quad (3.15)$$

$$R_2 = -2(a^2 + b^2)L_t^2 + Q(a^2 + b^2 - 2m) + K , \quad (3.16)$$

$$R_0 = -(a^4 + b^4 + 3a^2b^2)L_t^2 - (a^2 - b^2)(L_\phi^2 - L_\psi^2) - 4mL_t(aL_\phi + bL_\psi) - a^2b^2Q + (a^2 + b^2 - 2m)K , \quad (3.17)$$

$$R_{-2} = -a^2b^2(a^2 + b^2 + 2m)L_t^2 - b^2(a^2 - b^2 + 2m)L_\phi^2 - a^2(b^2 - a^2 + 2m)L_\psi^2 - 4abm(L_t(bL_\phi + aL_\psi) + L_\phi L_\psi) + Ka^2b^2 . \quad (3.18)$$

Proof

For simplicity we prove this theorem in a special case; the totally geodesic submanifold $N \equiv M|_{t,\phi,\psi=\text{const.}}$. The metric restricted to N is given by $g_N = \frac{\rho^2}{\Delta_r}dr^2 + \rho^2d\theta^2$. The Euler-Lagrange equations are

$$\frac{d}{ds} \left(\frac{\rho^2}{\Delta_r} \frac{dr}{ds} \right) = \frac{1}{2} \left[\frac{\partial}{\partial r} \left(\frac{\rho^2}{\Delta_r} \right) \left(\frac{dr}{ds} \right)^2 + \frac{\partial \rho^2}{\partial r} \left(\frac{d\theta}{ds} \right)^2 \right] , \quad (3.19)$$

$$\frac{d}{ds} \left(\rho^2 \frac{d\theta}{ds} \right) = \frac{1}{2} \left[\frac{\partial}{\partial \theta} \left(\frac{\rho^2}{\Delta_r} \right) \left(\frac{dr}{ds} \right)^2 + \frac{\partial \rho^2}{\partial \theta} \left(\frac{d\theta}{ds} \right)^2 \right] . \quad (3.20)$$

From (3.20), we have

$$\frac{d}{ds} \left(\rho^2 \frac{d\theta}{ds} \right) = -\frac{1}{2}(a^2 - b^2) \sin 2\theta \left(\frac{1}{\Delta_r} \left(\frac{dr}{ds} \right)^2 + \left(\frac{d\theta}{ds} \right)^2 \right) . \quad (3.21)$$

By using the conservation

$$Q = \langle \gamma', \gamma' \rangle_{g_N} = \rho^2 \left(\frac{1}{\Delta_r} \left(\frac{dr}{ds} \right)^2 + \left(\frac{d\theta}{ds} \right)^2 \right) , \quad (3.22)$$

the equation (3.21) can be transformed to

$$\frac{d}{ds} \left[\left(\rho^2 \frac{d\theta}{ds} \right)^2 - \frac{1}{2}Q(a^2 - b^2) \cos 2\theta \right] = 0 , \quad (3.23)$$

which yields

$$\rho^4 \left(\frac{d\theta}{ds} \right)^2 = \frac{1}{2}Q(a^2 - b^2) \cos 2\theta + K \quad (3.24)$$

where K is an integration constant. It is easy to show that

$$\rho^4 \left(\frac{dr}{ds} \right)^2 = \Delta_r \left(Q(r^2 + \frac{a^2 + b^2}{2}) - K \right) . \quad (3.25)$$

The equations (3.24) and (3.25) give the first-order geodesic equations on N , which are identical to the theorem specialized to the case $L_t = L_\phi = L_\psi = 0$ with a trivial constant shift of K . Using similar arguments we can prove the theorem in the general case, although the calculation is complicated. \square

4 Maximal Extension of Kerr spacetime

Let us consider special geodesics $\gamma_{\text{in}}^{\text{out}}(s)$ with constants (where θ is an arbitrary constant)

$$\begin{aligned} Q &= 0, \quad L_t = -1, \quad L_\phi = a \sin^2 \theta, \quad L_\psi = b \cos^2 \theta, \\ K &= 2(a^2 \sin^2 \theta + b^2 \cos^2 \theta) \end{aligned} \quad (4.26)$$

for which $\Theta = 0$ and $R(r) = -\rho^4$:

$$\gamma'_{\text{in}}^{\text{out}} = \pm \frac{\partial}{\partial r} + \frac{1}{r^2 \Delta_r} V(r), \quad (4.27)$$

where

$$V = (r^2 + a^2)(r^2 + b^2) \frac{\partial}{\partial t} + a(r^2 + b^2) \frac{\partial}{\partial \phi} + b(r^2 + a^2) \frac{\partial}{\partial \psi}. \quad (4.28)$$

For the limit $r \rightarrow \infty$, $\gamma'_{\text{in}}^{\text{out}} \rightarrow \pm \frac{\partial}{\partial r} + \frac{\partial}{\partial t}$, which represents an outgoing (or ingoing, respectively) null geodesic in the Boyer-Lindquist block I. The direction of the geodesic $\gamma_{\text{in}}^{\text{out}}$ coincides with $e_1 \pm e_4$,

$$\pm \frac{\partial}{\partial r} + \frac{1}{r \Delta_r} V = \frac{\rho}{\sqrt{\Delta_r \varepsilon}} (e_1 \pm e_4). \quad (4.29)$$

It should be noticed that the ingoing null geodesic γ_{in} coincides with the Kerr-Schild null geodesic given in [6].

We introduce a new coordinate (Kerr-Star coordinate) defined by

$$t^* = t + T(r), \quad (4.30)$$

$$\phi^* = \phi + A(r), \quad (4.31)$$

$$\psi^* = \psi + B(r), \quad (4.32)$$

together with $r^* = r$ and $\theta^* = \theta$, where

$$T(r) = \int \frac{(r^2 + a^2)(r^2 + b^2)}{r^2 \Delta_r} dr, \quad (4.33)$$

$$A(r) = \int \frac{a(r^2 + b^2)}{r^2 \Delta_r} dr, \quad (4.34)$$

$$B(r) = \int \frac{b(r^2 + a^2)}{r^2 \Delta_r} dr. \quad (4.35)$$

Then, the null geodesics are written as [†]

$$\gamma'_{\text{in}} = -\frac{\partial}{\partial r^*}, \quad (4.36)$$

$$\gamma'_{\text{out}} = \frac{\partial}{\partial r^*} + \frac{2}{r^2 \Delta_r} V. \quad (4.37)$$

[†]Note that $\frac{\partial}{\partial r^*} = \frac{\partial}{\partial r} - \frac{1}{r^2 \Delta_r} V$, although $r^* = r$. We have also $\frac{\partial}{\partial t^*} = \frac{\partial}{\partial t}$, $\frac{\partial}{\partial \phi^*} = \frac{\partial}{\partial \phi}$, $\frac{\partial}{\partial \psi^*} = \frac{\partial}{\partial \psi}$.




Figure 1: *Kerr-Star space* M^* . Horizontal lines crossing the horizons represent ingoing null geodesics, while vertical lines outgoing null geodesics. Ingoing null geodesics run from $r = \infty$ to $r = 0$. Outgoing null geodesics in the Boyer-Lindquist block I(III) run from $r = r_+$ ($r = 0$) to $r = \infty$ ($r = r_-$); the radial coordinate increases since $\Delta_r > 0$. In the block II, outgoing null geodesics run from $r = r_+$ to $r = r_-$; the radial coordinate decreases since $\Delta_r < 0$.

The metric in the Kerr-Star coordinate is given by

$$g_{BH} = \sum_{i,j=t,\phi,\psi} g_{ij} dx^{*i} dx^{*j} + g_{\theta\theta} d\theta^2 + 2dt^* dr - 2a \sin^2 \theta dr d\phi^* - 2b \cos^2 \theta dr d\psi^*, \quad (4.38)$$

where the components g_{ij} and $g_{\theta\theta}$ are same form as ones written by the Boyer-Lindquist coordinate (see (2.1) or (3.2)-(3.9)).

The dangerous term at the horizon, $g_{rr} \propto 1/\Delta_r$, is absent, and so g_{BH} is extended to the metric g_{BH}^* on the space $M^* = \mathbb{R}_+^2 \times S^3$ (see figure 1). The outgoing null geodesic γ_{out} is not defined on H . However, multiplying the factor $r^2 \Delta_r / 2$ with the right hand side in (4.37), we have a vector field

$$X^* = \frac{r^2 \Delta_r}{2} \frac{\partial}{\partial r^*} + V, \quad (4.39)$$

which is defined on M^* including H . Thus, one may understand the outgoing null geodesic γ_{out} as an integral curve of X^* . On the surface $r = \text{constant}$, the determinant of $g_{BH}^*|_{r=\text{const.}}$ is calculated as $-\rho^2 r^2 \Delta_r \sin^2 \theta \cos^2 \theta$, so that the restricted metric is degenerate if $\Delta_r = 0$, i.e. the horizons $H_{\pm} = H|_{r=r_{\pm}}$ are null hypersurfaces. At H_{\pm} , X^* in (4.39) reduces to the Killing vector fields

$$V_{\pm} = V|_{r=r_{\pm}} = (r_{\pm}^2 + a^2)(r_{\pm}^2 + b^2) \frac{\partial}{\partial t} + a(r_{\pm}^2 + b^2) \frac{\partial}{\partial \phi} + b(r_{\pm}^2 + a^2) \frac{\partial}{\partial \psi}. \quad (4.40)$$

The vector fields V_{\pm} are tangential to H_{\pm} and also perpendicular to H_{\pm} . The integral curves of V_{\pm} generate the horizons, which become totally geodesic null hypersurfaces in M^* .




Figure 2: *Star-Kerr space* *M . Horizontal lines crossing the horizons represent outgoing null geodesics, while vertical lines ingoing null geodesics. Star-Kerr space is opposite to the Kerr-Star space in these respects.

If we introduce a coordinate (Star-Kerr coordinate),

$${}^*t = t - T(r) , \quad {}^*\phi = \phi - A(r) , \quad {}^*\psi = \psi - B(r) , \quad {}^*r = r , \quad {}^*\theta = \theta \quad (4.41)$$

instead of the Kerr-Star coordinate (T , A and B are the same functions as (4.33), (4.34) and (4.35)), the null geodesics are given by

$$\gamma'_{\text{in}} = -\frac{\partial}{\partial {}^*r} + \frac{2}{r^2 \Delta_r} V(r) , \quad (4.42)$$

$$\gamma'_{\text{out}} = \frac{\partial}{\partial {}^*r} . \quad (4.43)$$

Then, we obtain a metric

$$g_{BH} = \sum_{i,j=t,\phi,\psi} g_{ij} d{}^*x^i d{}^*x^j + g_{\theta\theta} d\theta^2 - 2d{}^*t dr + 2a \sin^2 \theta dr d{}^*\phi + 2b \cos^2 \theta dr d{}^*\psi , \quad (4.44)$$

which differs from (4.38) only in the last three terms. The Star-Kerr space ${}^*M = \mathbb{R}_+^2 \times S^3$ is related to the Kerr-Star space M^* by the following isometric mapping $f : {}^*M \rightarrow M^*$; ${}^*t = -t^*$, ${}^*\phi = -\phi^*$, ${}^*\psi = -\psi^*$ together with ${}^*r = r^*$ and ${}^*\theta = \theta^*$ (see figure 2).

We may write the black hole metric in terms of the Kruskal coordinates $(u^\pm, v^\pm, \phi^\pm, \psi^\pm, \theta)$. These coordinates are defined on the regions $D(r_\pm)$ (see figures 4,3 and 5); the relation between Boyer-Lindquist and Kruskal coordinates are given by

$$|u^\pm| = \exp[\kappa_\pm(T(r) - t)] , \quad (4.45)$$

$$|v^\pm| = \exp[\kappa_\pm(T(r) + t)] , \quad (4.46)$$

$$\phi^\pm = \phi - \frac{at}{r_\pm^2 + a^2} , \quad (4.47)$$

$$\psi^\pm = \psi - \frac{bt}{r_\pm^2 + b^2} , \quad (4.48)$$




Figure 3: *Kerr-Star' space and Star-Kerr' space.* Kerr-Star' space (1) (Star-Kerr' space (2)) is defined by reversing the directions of ingoing and outgoing null geodesics of Kerr-Star space (Star-Kerr space, respectively).




Figure 4: Kruskal spaces $D(r_{\pm})$.

Kruskal spaces $D(r_{\pm})$ are building blocks of the maximal extension (see figure 5). Dot \bullet at the center represents the crossing three-sphere $S^3(r_{\pm})$.




Figure 5: *Maximal extension.* Each row corresponds to a Kerr-Star(ℓ) space while each column a Star-Kerr(ℓ) space. The Boyer-Lindquist block III (or III') is extended to a negative region of the square of radial coordinate r^2 [6], which is depicted in gray. The wavy lines in III represent curvature singularities defined by $r^2 = -a^2 \sin^2 \theta - b^2 \cos^2 \theta$.

where the constants

$$\kappa_{\pm} = \frac{r_{\pm}(r_{\pm}^2 - r_{\mp}^2)}{(r_{\pm}^2 + a^2)(r_{\pm}^2 + b^2)} \quad (4.49)$$

are the surface gravities of the horizons H_{\pm} . The integral (4.33) is calculated as

$$T(r) = r + \frac{1}{2\kappa_+} \log \left| \frac{r - r_+}{r + r_+} \right| + \frac{1}{2\kappa_-} \log \left| \frac{r - r_-}{r + r_-} \right|. \quad (4.50)$$

Then the radial coordinate $r = r(u^{\pm}v^{\pm})$ is given implicitly by

$$u^{\pm}v^{\pm} = \frac{1}{G_{\pm}(r)} \left(\frac{r - r_{\pm}}{r + r_{\pm}} \right), \quad (4.51)$$

where

$$G_{\pm}(r) = \pm \left| \frac{r - r_{\mp}}{r + r_{\mp}} \right|^{r_{\mp}/r_{\pm}} \exp(-2\kappa_{\pm}r) \quad (4.52)$$

are non-zero analytic functions on $D(r_{\pm})$. Note that two hypersurfaces $u^{\pm} = 0$ and $v^{\pm} = 0$ express the horizons, which cross at the region $D(r_{\pm})|_{u_{\pm}=v_{\pm}=0} \simeq S^3(r_{\pm})$ (crossing three-sphere). The Killing vector fields V_{\pm} on the horizons become

$$V_{\pm} = \kappa_{\pm}(r_{\pm}^2 + a^2)(r_{\pm}^2 + b^2) \left(-u^{\pm} \frac{\partial}{\partial u^{\pm}} + v^{\pm} \frac{\partial}{\partial v^{\pm}} \right) \quad (4.53)$$

in the Kruskal coordinate. It follows that the crossing spheres are fixed sets of V_{\pm} .

On $D(r_+)$ the metric takes the form,

$$\begin{aligned} g_{BH} = & \frac{G_+^2}{4\kappa_+^2} \frac{(r + r_+)^3(r^2 - r_-^2)}{(r - r_+)r^2\rho^2} \left(\frac{r^4\rho^4}{(r^2 + a^2)^2(r^2 + b^2)^2} - \frac{r_+^4\rho_+^4}{(r_+^2 + a^2)^2(r_+^2 + b^2)^2} \right) \\ & \times (u^{+2}dv^{+2} + v^{+2}du^{+2}) \\ & + \frac{G_+}{2\kappa_+^2} \frac{(r + r_+)^2(r^2 - r_-^2)}{r^2\rho^2} \left(\frac{r^4\rho^4}{(r^2 + a^2)^2(r^2 + b^2)^2} + \frac{r_+^4\rho_+^4}{(r_+^2 + a^2)^2(r_+^2 + b^2)^2} \right) du^+dv^+ \\ & + \frac{G_+^2}{4\kappa_+^2} \frac{(r + r_+)^4(a^2b^2\rho_+^4 + a^2(r_+^2 + b^2)^2r^2\sin^2\theta + b^2(r_+^2 + a^2)^2r^2\cos^2\theta)}{r^2\rho^2(r_+^2 + a^2)^2(r_+^2 + b^2)^2} \\ & \times (u^+dv^+ - v^+du^+)^2 \\ & + \frac{aG_+}{\kappa_+} \frac{(r + r_+)^2(r^2 + a^2 + \rho_+^2)}{\rho^2(r_+^2 + a^2)} \sin^2\theta d\phi^+(u^+dv^+ - v^+du^+) \\ & + \frac{bG_+}{\kappa_+} \frac{(r + r_+)^2(r^2 + b^2 + \rho_+^2)}{\rho^2(r_+^2 + b^2)} \cos^2\theta d\psi^+(u^+dv^+ - v^+du^+) \\ & + g_{S^3(r_+)}, \end{aligned} \quad (4.54)$$

where $\rho_+^2 = r_+^2 + a^2\cos^2\theta + b^2\sin^2\theta$ and the last term

$$g_{S^3(r_+)} = \sum_{i,j=\phi^+, \psi^+} g_{ij} dx^i dx^j + \rho^2 d\theta^2, \quad (4.55)$$

(g_{ij} are the same functions as (3.5)-(3.7)) gives a metric on the crossing sphere $S^3(r_+)$. In this coordinate, $S^3(r_+)$ becomes a totally geodesic submanifold of $D(r_+)$. The metric on $D(r_-)$ is given by replacing plus indices with minus indices. If we put $a = b = 0$ (Schwarzschild metric), for which we have $r_+ = 1/\kappa_+ = \sqrt{2m}$ and $r_- = 0$. then the metric reduces to

$$g_{BH} = 2m \left(1 + \frac{\sqrt{2m}}{r}\right)^2 \exp\left(-\sqrt{\frac{2}{m}}r\right) du^+ dv^+ + r^2(d\theta^2 + \sin^2\theta d\phi^2 + \cos^2\theta d\psi^2) \quad (4.56)$$

with

$$u^+ v^+ = \left(\frac{r - \sqrt{2m}}{r + \sqrt{2m}}\right) \exp\left(\sqrt{\frac{2}{m}}r\right). \quad (4.57)$$

5 Five-dimensional AdS Kerr black holes

The analysis persists in the case with a cosmological constant. The Kerr AdS metric in the Boyer-Lindquist coordinate is given as [7]

$$\begin{aligned} g_5^{BH} = & -\frac{\Delta_r}{\rho^2} \left[dt - \frac{a \sin^2 \theta}{\Xi_a} d\phi - \frac{b \cos^2 \theta}{\Xi_b} d\psi \right]^2 + \frac{\rho^2}{\Delta_r} dr^2 + \frac{\rho^2}{\Delta_\theta} d\theta^2 \\ & + \frac{\Delta_\theta \sin^2 \theta}{\rho^2} \left(a dt - \frac{r^2 + a^2}{\Xi_a} d\phi \right)^2 + \frac{\Delta_\theta \cos^2 \theta}{\rho^2} \left(b dt - \frac{r^2 + b^2}{\Xi_b} d\psi \right)^2 \\ & + \frac{1 + r^2 \ell^2}{r^2 \rho^2} \left[ab dt - \frac{b(r^2 + a^2) \sin^2 \theta}{\Xi_a} d\phi - \frac{a(r^2 + b^2) \cos^2 \theta}{\Xi_b} d\psi \right]^2, \end{aligned} \quad (5.1)$$

where

$$\begin{aligned} \rho^2 = & r^2 + a^2 \cos^2 \theta + b^2 \sin^2 \theta, \quad \Delta_r = \frac{1}{r^2} (r^2 + a^2)(r^2 + b^2)(1 + r^2 \ell^2) - 2m, \\ \Delta_\theta = & 1 - a^2 \ell^2 \cos^2 \theta - b^2 \ell^2 \sin^2 \theta, \quad \Xi_a = 1 - a^2 \ell^2, \quad \Xi_b = 1 - b^2 \ell^2. \end{aligned} \quad (5.2)$$

We find that the theorem 1 is generalized as follows.

Theorem 2.

There exists a constant K' for a geodesic γ satisfying

$$\rho^4 \left(\frac{d\theta}{ds} \right)^2 + \Theta_\ell(\theta) = 0, \quad (5.3)$$

$$\rho^4 \left(\frac{dr}{ds} \right)^2 + R_\ell(r) = 0, \quad (5.4)$$

where

$$\begin{aligned}\Theta_\ell = & (a^2 \sin^2 \theta + b^2 \cos^2 \theta - a^2 b^2 \ell^2) L_t^2 + \frac{\Xi_a^2 (1 - b^2 \ell^2 \sin^2 \theta)}{\sin^2 \theta} L_\phi^2 + \frac{\Xi_b^2 (1 - a^2 \ell^2 \cos^2 \theta)}{\cos^2 \theta} L_\psi^2 \\ & - 2\ell^2 (a^2 \sin^2 \theta + b^2 \cos^2 \theta - a^2 b^2 \ell^2) L_t (a L_\phi + b L_\psi) - 2ab\ell^2 \Xi_a \Xi_b L_\phi L_\psi \\ & - \Delta_\theta (a^2 \cos^2 \theta + b^2 \sin^2 \theta) Q - \Delta_\theta K' ,\end{aligned}\quad (5.5)$$

$$R_\ell = R_{-2} r^{-2} + R_0 + R_2 r^2 + R_4 r^4 + R_6 r^6 . \quad (5.6)$$

The coefficients are explicitly given by

$$R_6 = -Q\ell^2 , \quad (5.7)$$

$$R_4 = -L_t^2 + 2\ell^2 L_t (a L_\phi + b L_\psi) - (1 + a^2 \ell^2 + b^2 \ell^2) Q + \ell^2 K' , \quad (5.8)$$

$$\begin{aligned}R_2 = & (-2(a^2 + b^2) + a^2 b^2 \ell^2) L_t (L_t - 2\ell^2 (a L_\phi + b L_\psi)) + 2ab\ell^2 \Xi_a \Xi_b L_\phi L_\psi \\ & + b^2 \ell^2 \Xi_a^2 L_\phi^2 + a^2 \ell^2 \Xi_b^2 L_\psi^2 - (a^2 + b^2 + a^2 b^2 \ell^2 - 2m) Q + (1 + a^2 \ell^2 + b^2 \ell^2) K' \end{aligned} \quad (5.9)$$

$$\begin{aligned}R_0 = & (-a^4 \Xi_b - b^4 \Xi_a - 3a^2 b^2) L_t^2 + \Xi_a^2 (b^2 - a^2 + a^2 b^2 \ell^2 + b^4 \ell^2) L_\phi^2 \\ & + \Xi_b^2 (a^2 - b^2 + a^2 b^2 \ell^2 + a^4 \ell^2) L_\psi^2 - 2ab\ell^2 \Xi_a \Xi_b (a^2 + b^2) L_\phi L_\psi \\ & + 2(a^4 \ell^2 \Xi_b + b^4 \ell^4 \Xi_a + 3a^2 b^2 \ell^2 - 2m) L_t (a L_\phi + b L_\psi) \\ & - a^2 b^2 Q + (a^2 + b^2 + a^2 b^2 \ell^2 - 2m) K' ,\end{aligned}\quad (5.10)$$

$$\begin{aligned}R_{-2} = & -a^2 b^2 (a^2 + b^2 - a^2 b^2 \ell^2 + 2m) L_t^2 - b^2 \Xi_a^2 (a^2 \Xi_b^2 - b^2 + 2m) L_\phi^2 \\ & - a^2 \Xi_b^2 (b^2 \Xi_a^2 - a^2 + 2m) L_\psi^2 + 2ab^2 (a^2 \ell^2 (a^2 \Xi_b + b^2) + 2m \Xi_a) L_t L_\phi + a^2 b^2 K' \\ & + 2a^2 b (b^2 \ell^2 (b^2 \Xi_a + a^2) + 2m \Xi_b) L_t L_\psi + 2ab \Xi_a \Xi_b (a^2 b^2 \ell^2 - 2m) L_\phi L_\psi\end{aligned}\quad (5.11)$$

with

$$Q = \langle \gamma', \gamma' \rangle , \quad L_i = \langle \gamma', \frac{\partial}{\partial x^i} \rangle , \quad x^i = (t, \phi, \psi). \quad (5.12)$$

These equations reduce to (3.13) and (3.14) when we set $\ell = 0$.

The special geodesics in (4.27) are generalized to

$$\gamma'_{\text{in}}^{\text{out}} = \pm \frac{\partial}{\partial r} + \frac{1}{r^2 \Delta_r} V(r) \quad (5.13)$$

with

$$V(r) = (r^2 + a^2)(r^2 + b^2) \frac{\partial}{\partial t} + a \Xi_a (r^2 + b^2) \frac{\partial}{\partial \phi} + b \Xi_b (r^2 + a^2) \frac{\partial}{\partial \psi} . \quad (5.14)$$

The constants are

$$\begin{aligned}Q = & 0 , \quad L_t = -1 , \quad L_\phi = \frac{a \sin^2 \theta}{\Xi_a} , \quad L_\psi = \frac{b \cos^2 \theta}{\Xi_b} , \\ K = & \frac{2}{\Xi_a \Xi_b} (a^2 \sin^2 \theta + b^2 \cos^2 \theta - a^2 b^2 \ell^2) ,\end{aligned}\quad (5.15)$$

for which we have $\Theta_\ell = 0$ and $R_\ell = -\rho^4$.

In the same way as the Ricci flat case, Kerr-Star coordinate and Star-Kerr coordinate are introduced:

$${}^*t^* = t \pm T(r) , \quad {}^*\phi^* = \phi \pm A(r) , \quad {}^*\psi^* = \psi \pm B(r) , \quad (5.16)$$

where

$$T(r) = \int \frac{(r^2 + a^2)(r^2 + b^2)}{r^2 \Delta_r} dr , \quad (5.17)$$

$$A(r) = \int \frac{a \Xi_a (r^2 + b^2)}{r^2 \Delta_r} dr , \quad (5.18)$$

$$B(r) = \int \frac{b \Xi_b (r^2 + a^2)}{r^2 \Delta_r} dr . \quad (5.19)$$

Then, the metric is given by

$$g_{BH} = \sum_{i,j=t,\phi,\psi} g_{ij} d^*x^{*i} d^*x^{*j} + g_{\theta\theta} d\theta^2 \mp 2d^*t^* dr \pm \frac{2a \sin^2 \theta}{\Xi_a} dr d^*\phi^* \pm \frac{2b \cos^2 \theta}{\Xi_b} dr d^*\psi^* , \quad (5.20)$$

where

$$g_{tt} = -1 + \frac{2m}{\rho^2} - \ell^2 (r^2 + a^2 \sin^2 \theta + b^2 \cos^2 \theta) , \quad (5.21)$$

$$g_{t\phi} = -\frac{2ma \sin^2 \theta}{\rho^2 \Xi_a} + \ell^2 \frac{a \sin^2 \theta}{\Xi_a} (r^2 + a^2) , \quad (5.22)$$

$$g_{t\psi} = -\frac{2mb \cos^2 \theta}{\rho^2 \Xi_b} + \ell^2 \frac{b \cos^2 \theta}{\Xi_b} (r^2 + b^2) , \quad (5.23)$$

$$g_{\phi\phi} = \sin^2 \theta \left[\frac{r^2 + a^2}{\Xi_a} + \frac{2 \sin^2 \theta a^2 m}{\rho^2 \Xi_a^2} \right] , \quad (5.24)$$

$$g_{\psi\psi} = \cos^2 \theta \left[\frac{r^2 + b^2}{\Xi_b} + \frac{2 \cos^2 \theta b^2 m}{\rho^2 \Xi_b^2} \right] , \quad (5.25)$$

$$g_{\phi\psi} = \frac{2abm \cos^2 \theta \sin^2 \theta}{\rho^2 \Xi_a \Xi_b} . \quad (5.26)$$

In this coordinate system, the special geodesics (5.13) reduce to (4.42) and (4.43), or (4.36) and (4.37), with V in (5.14) and Δ_r in (5.2).

Note added:

While this paper was in preparation we received [12] which overlaps with the result on the integrability of geodesics on the five-dimensional AdS Kerr black hole. However their result has been obtained using a different approach.

Acknowledgements

The authors thank Yoshitake Hashimoto, Hideki Ishihara and Ken-ichi Nakao for useful discussions. This paper is supported by the 21 COE program “Constitution of wide-angle mathematical basis focused on knots”. Research of Y.Y. is supported in part by the Grant-in Aid for scientific Research (No. 14540073 and No. 14540275) from Japan Ministry of Education. The preliminary version of this work was presented by Y.Y. in “Quantum Cohomology and Mirror Symmetry Day” held at Tokyo Metropolitan University (21 January, 2005).

References

- [1] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2** (1998) 231 [*Int. J. Theor. Phys.* **38** (1999) 1113] [arXiv:hep-th/9711200]; S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Phys. Lett. B* **428** (1998) 105 [arXiv:hep-th/9802109]; E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2** (1998) 253 [arXiv:hep-th/9802150].
- [2] D. Page, “A Compact Rotating Gravitational Instanton,” *Phys. Lett. B* **79** (1978) 235.
- [3] Y. Hashimoto, M. Sakaguchi and Y. Yasui, “New infinite series of Einstein metrics on sphere bundles from AdS black holes,” arXiv:hep-th/0402199, *Commun. Math. Phys.* in press.
- [4] G. W. Gibbons, H. Lu, D. N. Page and C. N. Pope, “The general Kerr-de Sitter metrics in all dimensions,” *J. Geom. Phys.* **53** (2005) 49 [arXiv:hep-th/0404008]; “Rotating black holes in higher dimensions with a cosmological constant,” *Phys. Rev. Lett.* **93** (2004) 171102 [arXiv:hep-th/0409155].
- [5] Y. Hashimoto, M. Sakaguchi and Y. Yasui, “Sasaki-Einstein twist of Kerr-AdS black holes,” arXiv:hep-th/0407114, *Phys. Lett. B* **600** (2004) 270.
- [6] R. C. Myers and M. J. Perry, “Black Holes In Higher Dimensional Space-Times,” *Annals Phys.* **172** (1986) 304.

- [7] S. W. Hawking, C. J. Hunter and M. M. Taylor-Robinson, “Rotation and the AdS/CFT correspondence,” *Phys. Rev. D* **59** (1999) 064005 [arXiv:hep-th/9811056].
- [8] B. O’Neill, “The Geometry of Kerr Black Holes,” A K Peters, Ltd., 1995.
- [9] V. P. Frolov and D. Stojkovic, “Particle and light motion in a space-time of a five-dimensional rotating black hole,” *Phys. Rev. D* **68** (2003) 064011 [arXiv:gr-qc/0301016].
- [10] P. J. De Smet, “The Petrov type of the five-dimensional Myers-Perry metric,” *Gen. Rel. Grav.* **36** (2004) 1501 [arXiv:gr-qc/0312021].
- [11] B. Carter, “Global Structure Of The Kerr Family Of Gravitational Fields,” *Phys. Rev.* **174** (1968) 1559.
- [12] H. K. Kunduri and J. Lucietti, “Integrability and the Kerr-(A)dS black hole in five dimensions,” arXiv:hep-th/0502124.