#### BAXTERIZATION, DYNAMICAL SYSTEMS, AND THE SYMMETRIES OF INTEGRABILITY

#### C.-M. Viallet $^{\rm 1}$

**Abstract**. We resolve the 'baxterization' problem with the help of the automorphism group of the Yang-Baxter (resp. star-triangle, tetrahedron, ...) equations. This infinite group of symmetries is realized as a non-linear (birational) Coxeter group acting on matrices, and exists as such, *beyond the narrow context of strict integrability*. It yields among other things an unexpected elliptic parametrization of the non-integrable sixteen-vertex model. It provides us with a class of discrete dynamical systems, and we address some related problems, such as characterizing the complexity of iterations.

To appear in the proceedings of the the Third Baltic Rim Student Seminar HELSINKI (September 1993)

work supported by CNRS

<sup>&</sup>lt;sup>1</sup> Laboratoire de Physique Théorique et des Hautes Energies, Centre National de la Recherche Scientifique, Université de Paris 6- Paris 7, Tour 16, 1<sup>er</sup> étage, boîte 126. 4 Place Jussieu/ F-75252 PARIS Cedex 05 / FRANCE / e-mail: viallet@lpthe.jussieu.fr

## 1 Introduction

The results presented here originate in a long-standing collaboration of M. Bellon, J.M. Maillard and the author [1, 2, 3, 4, 5, 6, 7], augmented by further elaborations (as [8, 9]).

The Yang-Baxter equations and their variations (star-triangle equations, tetrahedron equations,...) are nowadays considered as a characteristic feature of integrability of mechanical systems (classical hamiltonian systems, as well as quantum and statistical systems) [10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20].... This field has expanded very much for the last twenty years. We shall not dwell here on its innumerable developments, and relegate general considerations to the end.

We shall start from the Yang-Baxter equations and view them as an algebraic system of equations on matrix entries.

We first recall (section 2) what the two most common forms of the equations for vertex models are.

In section 3 we introduce certain groups generated by involutions (Coxeter groups) [21], together with some realizations in terms of birational transformations of projective spaces. These realizations are generically denoted  $\Gamma$  in the sequel.

In section 4 we build up, from the groups  $\Gamma$ , an infinite group of automorphisms (denoted  $\mathcal{A}$ ) of the Yang-Baxter equations. One should keep in mind that the equations we analyze form an *overdetermined* system, making the existence of a large group of symmetry quite remarkable.

We next show, for the paradigmatic example of the Baxter symmetric eightvertex model and with both a picture and algebraic results, how our automorphism group reconstructs the well known elliptic curves of solutions. This is a first solution to the 'baxterization' problem [22].

In section 6 we describe more groups of matrix transformations. These are very similar to the ones of section 3, but are adapted to the star-triangle relations, tetrahedron equations, and higher dimensional generalizations.

In section 7 we complete the solution to the baxterization problem, emphasizing the notions of  $\Gamma$ -covariant versus  $\Gamma$ -invariant quantities.

We next produce (section 8) an algorithm to calculate the covariant polynomials as well as the algebraic invariants.

One of the features of  $\Gamma$  is that it is defined outside of the space of solutions of the Yang-Baxter equations, and we may look at its action *beyond integrability*. It becomes an ordinary discrete dynamical system on the space of parameters, possibly non-conservative. In section 9, we describe this action for the general 16-vertex model, i.e. the most general two-state vertex model on a two-dimensional square lattice, which is known not to be integrable. The outcome is quite surprising: the orbits of  $\Gamma$  stay within 1-dimensional subvarieties of the 15-dimensional space of parameters. This reveals the actuality of an amazingly large number of algebraically independent invariants of  $\Gamma$ , and yields an elliptic parametrization of the non-integrable model, without reference to the Yang-Baxter equations.

In section 11 we turn to the notion of complexity of the iterations and show how it fits with the existence of algebraic invariants.

In the last section (12), we exemplify some of our results on a definite system, related to a three dimensional vertex model.

# 2 The Yang-Baxter equations (vertex models)

Many presentations of the Yang-Baxter equations appear in the literature, dividing into two classes:

The first class contains a parameter, also called spectral parameter [18], and reads, for vertex models:

$$\sum_{\alpha_{1},\alpha_{2},\alpha_{3}} R^{i_{1}i_{2}}_{\alpha_{1}\alpha_{2}}(\lambda_{1},\lambda_{2}) R^{\alpha_{1}i_{3}}_{j_{1}\alpha_{3}}(\lambda_{1},\lambda_{3}) R^{\alpha_{2}\alpha_{3}}_{j_{2}j_{3}}(\lambda_{2},\lambda_{3})$$
(1)  
=  $\sum_{\beta_{1},\beta_{2},\beta_{3}} R^{i_{2}i_{3}}_{\beta_{2}\beta_{3}}(\lambda_{2},\lambda_{3}) R^{i_{1}\beta_{3}}_{\beta_{1}j_{3}}(\lambda_{1},\lambda_{3}) R^{\beta_{1}\beta_{2}}_{j_{1}j_{2}}(\lambda_{1},\lambda_{2}).$ 

In this system of equations, R is a matrix of size  $q^2 \times q^2$ , whose entries are functions of the parameters  $\lambda_i$ , usually through the difference  $\lambda_i - \lambda_j$  (at least when  $\lambda$  labels a point on an elliptic curve).

The parameter first appeared as parametrizing the solution. It was later understood as a "spectral parameter" in the context of the quantum inverse scattering method. The origin of the parameter is one of the issues we wish to address.

A given family of models is obtained when one specifies the size q and the relations between the entries of R. Typical relations are equalities between different entries or vanishing of some others (see for example section 5).

Since the entries of R are allowed to be different for the three copies of R entering (1), one may rewrite (1) with three explicitly different matrices A, B, and C of the same size  $q^2 \times q^2$ .

$$\sum_{\alpha_1,\alpha_2,\alpha_3} A^{i_1i_2}_{\alpha_1\alpha_2} B^{\alpha_1i_3}_{j_1\alpha_3} C^{\alpha_2\alpha_3}_{j_2j_3} = \sum_{\beta_1,\beta_2,\beta_3} C^{i_2i_3}_{\beta_2\beta_3} B^{i_1\beta_3}_{\beta_1j_3} A^{\beta_1\beta_2}_{j_1j_2} \tag{2}$$

or shortly

 $\epsilon$ 

$$A_{12}B_{13}C_{23} = C_{23}B_{13}A_{12} \tag{3}$$

with now usual notations. The latter form has the interest of not referring to any explicit parametrization.

The second class contains no parameter, and is sometimes called "constant" Yang-Baxter equation. It reads:

$$R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12} \tag{4}$$

that is to say like (3) but with A = B = C = R.

To go from equation (1) to equation (4) requires essentially an adequate choice of  $\lambda_1, \lambda_2, \lambda_3$ , namely  $\lambda_1 = \lambda_2 = \lambda_3$ . The reverse move, i.e. to recover the parameter dependence from a constant solution was given the name of 'baxter-ization' [22].

The *use* of the spectral parameter is clear in the context of the Bethe ansatz, since it serves building up the generating functional for commuting conserved quantities, but we shall not be concerned with this aspect. Its *origin* will be explained in the next sections.

## **3** Some operations on matrices: the groups $\Gamma$

We describe here some elementary operations on matrices of various sizes. The matrices we consider are defined up to an overall multiplicative factor. The space of parameters is thus some projective space  $\mathbf{P}_n$  with the entries of the matrix as homogeneous coordinates.

Let us start with the matrix R of a two-dimensional q-state vertex model on a square lattice. R has the structure of a tensor product and the indices of Rare pairs of indices:

$$R_{kl}^{ij} \qquad i, j, k, l = 1..q \tag{5}$$

We may define the inverse I up to a factor (well defined in the projective space)

$$\sum_{\alpha\beta} (IR)^{ij}_{\alpha\beta} R^{\alpha\beta}_{kl} = \mu \, \delta^i_k \delta^j_l \qquad i, j, k, l = 1..q \tag{6}$$

with  $\mu$  an arbitrary multiplicative factor, the transposition t:

$$(tR)_{kl}^{ij} = R_{ij}^{kl} \qquad i, j, k, l = 1..q$$
 (7)

as well as two partial transpositions  $t_l$  (index l for left) and  $t_r$  respectively by:

$$(t_l R)_{kl}^{ij} = R_{il}^{kj} \qquad (t_r R)_{kl}^{ij} = R_{kj}^{il} \qquad i, j, k, l = 1..q$$
(8)

Of course

$$t = t_l t_r = t_r t_l, \qquad I^2 = t^2 = t_l^2 = t_r^2 = 1, \quad \text{and} \quad I t = t I$$
 (9)

However

$$t_l I \neq I t_l \quad \text{and} \ t_r \ I \neq I \ t_r$$
 (10)

The two partial transpositions do not commute with the inversion while their product t does. The transformation  $t_l I$  and  $t_r I$  are of infinite order. Notice of course that they may have finite orbits when acting on certain non-generic matrices.

The group  $\Gamma$  generated by  $t_l, t_r$  and I essentially consists of the iterates of  $t_l I$  and its inverse  $It_l$  (or  $t_r I$  and  $It_r$ ) up to multiplication by an element of

its center. It is generated by involutions realized as rational (possibly linear) transformations of some projective space  $\mathbf{P}_n$ .

We shall describe in section 6 more instances of groups  $\Gamma$  acting as rational transformations of matrices, and generated by involutions.

# 4 The automorphisms of the Yang-Baxter equations

Suppose we have a solution (A, B, C) of (3):

$$A_{12}B_{13}C_{23} = C_{23}B_{13}A_{12} \tag{11}$$

We may take the partial transpose  $t_1$  of (11) along space 1 and get:

$$(t_1 B_{13}) (t_1 A_{12}) C_{23} = C_{23} (t_1 A_{12}) (t_1 B_{13})$$

$$(12)$$

Taking the partial transpose  $t_2$  of (12) yields

$$(t_1B_{13}) (t_2C_{23}) (t_1t_2A_{12}) = (t_1t_2A_{12}) (t_2C_{23}) (t_1B_{13})$$

$$(13)$$

that is to say:

$$(t_l B)_{13} (t_l C)_{23} (tA)_{12} = (tA)_{12} (t_l C)_{23} (t_l B)_{13}$$
(14)

Multiplying both sides of (14) to the left and the right by the inverse ItA of tA gives:

$$(tIA)_{12} (t_l B)_{13} (t_l C)_{23} = (t_l C)_{23} (t_l B)_{13} (tIA)_{12}$$
(15)

which shows that the transformation

$$\mathcal{K}_A: (A, B, C) \longrightarrow (tIA, t_l B, t_l C)$$
 (16)

takes a solution of (3) into a solution of (3).

We could similarly construct  $\mathcal{K}_B$  (resp.  $\mathcal{K}_C$ ) which inverts and transposes B (resp. C) and acts by partial transpositions on A and C (resp. A and B).

The automorphisms  $\mathcal{K}_A, \mathcal{K}_B, \mathcal{K}_C$  are involutive and generate an infinite group of automorphisms  $\mathcal{A}$  of the Yang-Baxter equations. It is possible to show that  $\mathcal{A}$  is isomorphic to the Weyl group of an affine Lie algebra of type  $A_2^{(1)}$  [3, 4].

The group  $\Gamma$  appears when we look at the action of  $\mathcal{A}$  on one of the individual copies of R entering equation (1). However, the action of  $\Gamma$  is defined without reference to (1), and we will now concentrate on it, and somehow forget about (1).

## 5 Baxterization of the Baxter model

The exemplary two-dimensional integrable vertex model is the Baxter model [23, 24, 12], with matrix of Boltzmann weights:

$$R = \begin{pmatrix} a & 0 & 0 & d \\ 0 & b & c & 0 \\ 0 & c & b & 0 \\ d & 0 & 0 & a \end{pmatrix}$$
(17)

where one sees the vanishing conditions and equality relations between the entries. It is straightforward to see that the form (17) is stable by all the generators of  $\Gamma$ . We say that the pattern (17), i.e. the collection of vanishing conditions and equalities between entries is admissible (see [1, 2]).

The transformations  $t_l$  and  $t_r$  read:

$$a \to a, \quad b \to b, \quad c \to d, \quad d \to c,$$

and I reads

$$a \to \frac{a}{a^2 - d^2}, \quad b \to \frac{b}{b^2 - c^2}, \quad c \to \frac{-c}{b^2 - c^2}, \quad d \to \frac{-d}{a^2 - d^2}.$$

One may look at the action of  $\Gamma$  on R. Take one starting point in the space  $\mathbf{P}_3$  of parameters, iterate R on it and just look at the orbit. One gets the following picture, showing a three-dimensional perspective of the orbit, in inhomogeneous coordinates u = b/a, v = c/a, w = d/a, with starting point (\*):



#### Figure 1: 'Baxterization' of the Baxter model: a perspective view

One sees from this figure that the orbit lies inside a curve. The action of  $\Gamma$  must have at least two algebraically independent invariants. Indeed there are two such invariants:

$$\Delta_1 = \frac{a^2 + b^2 - c^2 - d^2}{ab + cd}, \qquad \Delta_2 = \frac{ab - cd}{ab + cd}.$$
 (18)

The curves  $\Delta_1 = constant$ ,  $\Delta_2 = constant$ , are precisely the elliptic curves which appear in the solution of (1), but we have obtained them with the form (17) as only input, and in particular without having to solve (1). The resolution of (1) becomes straightforward once the adequate parametrization of the above curves is used. The orbit we get, although discrete, tends to be dense inside one dimensional curves. The phenomenon is similar to the iteration of a rotation (obtained as the composition of two axial symmetries) with irrational rotation number.

Warning: It is in no way the existence of these curves which ensures the compatibility of the Yang-Baxter equations!

# 6 Other transformation groups

There are many other equations like (1). They also have automorphism groups generated by (bi)rational transformations. This means we may construct many different realizations  $\Gamma$ . Here are a few more.

#### 6.1 From spin model with interactions along the edges

As was shown in [3], the star-triangle equation also has an infinite group of automorphisms. The matrices under consideration are ordinary  $q \times q$  matrices m, of entries  $m_{ij}$ . The generators of  $\Gamma$  are the matrix inverse I as previously

$$\sum_{\alpha} (Im)_{i\alpha} \ m_{\alpha j} = \mu \ \delta_{ij} \tag{19}$$

with  $\mu$  an arbitrary multiplicative constant, and the element-by-element inverse (Hadamard inverse):

$$U: \qquad m_{ij} \longrightarrow 1/m_{ij} \tag{20}$$

I and J are two non-commuting rational involutions. The product  $\varphi = IJ$  is a birational transformation of infinite order.

#### 6.2 From vertex models in three (and more) dimensions

The matrices under consideration in this case again have multi-indices. The matrices are of size  $q^3 \times q^3$ , with entries of the form:

$$R_{lmn}^{ijk} \tag{21}$$

By examining the symmetries [4, 5] of the tetrahedron equations [25, 26, 27]:

$$R_{123}R_{145}R_{246}R_{356} = R_{356}R_{246}R_{145}R_{123}.$$
(22)

with the usual notations, one is lead to the group  $\Gamma$  generated by the inverse Iand the *three* partial transpositions  $t_q$ ,  $t_m$  and  $t_d$  with

$$(t_g R)^{i_g i_m i_d}_{j_g j_m j_d} = R^{j_g i_m i_d}_{i_g j_m j_d}, (23)$$

and similar definitions for  $t_m$  and  $t_d$ .

One could as well consider multi-index matrices of size  $q^d \times q^d$  written in the form  $M_{j_1j_2...j_d}^{i_1i_2...i_d}$ . Section 4 deals with d = 2. The tetrahedron equation corresponds to d = 3, and so on. There exist d different partial transpositions  $t_1, t_2, \ldots, t_d$  with obvious definitions. Combined with the inverse I, they generate an *infinite* group  $\Gamma$ . We will return to the question of the size of these groups in sections 8 and 11.

#### 6.3 More groups

One may further enrich the representations by imposing constraints on the entries of the matrices, provided the transformations are compatible with these constraints (see [1] for the notion of admissible patterns). This yields realizations on projective spaces of lower dimensions.

If the dimension n of the projective space is 1, the only birational transformations are homographies. If n = 2, the group, called Cremona group [28], is already much larger. Its elements may however be written as products of linear transformations and the basic blow-up J. The interesting subgroup made out of polynomial and polynomially invertible transformations of the plane has a more elementary structure [29, 30]. When n is larger than 2, the structure of the group of birational transformations is much more complex.

Let us stress that, at the level of the realization, there may exist additional relations between the generators, possibly making it finite.

## 7 Baxterization

It appears clearly in sections 4 and 5 that, for the Baxter model, the existence of the parameter comes from the existence of sufficiently many algebraic invariants of  $\Gamma$  to confine its orbits, and therefore the ones of  $\mathcal{A}$ , to curves. The general situation is more subtle, but it always amounts to the following: find the  $\Gamma$ -covariant varieties passing through a constant solution, and/or look for solutions only on covariant varieties.

Different situations may appear, and we illustrate them by specific examples:

– the simplest one was encountered in section 5, recalling that the only input information was the form of the matrix (17). The parameter is just the uniformizing parameter for the elliptic curve which pops out of the action of  $\Gamma$ on any of its generic points. In this case, the action of  $\Gamma$  has INVARIANTS to imprison the orbit, and the entire space of parameters is foliated by invariant subvarieties (curves for the Baxter model). Each generic spectral curve has an infinite group of automorphisms, and they consequently all have genus 0 or 1. The solution of (3) is obtained in the following way: choose any point A in the space of parameters. Through A passes one curve of the foliation. Choose B anywhere on this curve. The last matrix C is completely determined then. What is special about the Baxter model is that there is no restriction on the first choice (A), within the space  $\mathbf{P}_3$  of parameters. This is far from being the general occurrence.

– a more subtle situation is encountered for the chiral Potts model (as well as for the free fermion condition [31, 32, 33]). Integrability appears on a algebraic variety of which all points have a periodic orbit under  $\Gamma$ . This variety is automatically globally invariant by  $\Gamma$ , but there may very well be NO INVARI-ANT of the action of  $\Gamma$ . Besides, the resolution of the Yang-Baxter equations may lead to consider further algebraic conditions. Solutions are to be found on a possibly isolated subvariety of the space of parameters. In this situation, the spectral curve may have genus higher than 0 or 1, since it has only a finite number of automorphisms [34, 35, 36].

- a similar situation is encountered in the case of the Jaeger-Higman-Sims model [37, 38]. The parameter space of the model is  $\mathbf{P}_2$ . The matrix of Boltzmann weights belongs to a 3-dimensional abelian algebra of matrices of size  $100 \times 100$ , with three generators  $\mathbf{1}, A, \mathcal{J} - \mathbf{1} - A$  where  $\mathbf{1}$  is the unit matrix, A the adjacency matrix of the Higman-Sims graph, and  $\mathcal{J}$  the matrix with all entries equal to 1. Two products exist in this algebra, respectively the usual matrix product and the element by element product. The symmetry group is generated by the inverses I and J, corresponding to these two products, as in section (6.1), equations (19, 20). The two inverses are related by a collineation C (linear map of  $\mathbf{P}_2$ ):

$$I = C^{-1}JC$$

In terms of homogeneous coordinates  $[x_0, x_1, x_2]$ :

$$U: [x_0, x_1, x_2] \to [x_1 x_2, x_0 x_2, x_1 x_2]$$
$$C = \begin{bmatrix} 1 & 22 & 77 \\ 1 & -8 & 7 \\ 1 & 2 & -3 \end{bmatrix}$$

We can find three  $\Gamma$ -invariant subvarieties of  $\mathbf{P}_2$ : one line D, and two hyperbolae H1, H2:

$$D: \quad x_2 - x_1 = 0$$
  
H1: 
$$x_2^2 + 3 x_1 x_2 - 3 x_0 x_2 - x_0 x_1 = 0$$
  
H2: 
$$x_1 x_2 + 2 x_1^2 - 2 x_0 x_2 - x_0 x_1 = 0$$

Hyperbola H1 is the one of reference [39]. The two hyperbolae contain an infinite number of singular points, and we know this prevents the existence of an invariant [9], but not of covariant polynomials, and a fortiori not of covariant ideals.

## 8 Covariants and invariants

It is fortunately possible to go further in the analysis of the invariant subvarieties [9].

We start from a group G generated by by  $\nu$  involutions  $I_1, I_2, \ldots, I_k$ ,  $(k = 1 \ldots \nu)$ , verifying no relations other than the involution property. The group G is infinite and there are two essentially different situations.

If  $\nu = 2$ , the group is the infinite dihedral group  $\mathbf{Z}_2 \times \mathbf{Z}$ , and all elements may uniquely be written  $I_1^{\alpha}(I_1I_2)^q$ , with  $\alpha = 0, 1$  and  $q \in \mathbf{Z}$ . The number of elements of given length l is 2.

If  $\nu \geq 3$ , the number of elements of length l grows exponentially with l, and the group is in a sense bigger (still countable).

As an example, for the groups described in section 6.2, the number  $\nu$  of generators depends on the dimension d of the lattice: it is just  $2^{d-1}$  so that if d = 2, G is generated by two involutions and if  $d \ge 3$ , G is generated by more than three involutions.

We then construct various realizations  $\Gamma$  of G by explicit transformations of some projective space. They are obtained by specifying the realization of the generators. We use the same notation  $I_k$  for the generators of G and their representatives in  $\Gamma$ . The realizations  $\Gamma$  of G may be written as polynomial transformations in terms of the homogeneous coordinates.

Each involution  $I_k$  defines a characteristic polynomial  $\phi_k$  of degree  $d_k^2 - 1$  in the following manner. The  $I_k$  being involutions,  $I_k^2$  appears as the multiplication by a degree  $d_k^2 - 1$  polynomial  $\phi_k(x_0, \ldots, x_n)$ .

A  $\Gamma$ -covariant polynomial P verifies:

$$P(\gamma(x)) = a(\gamma, x)P(x) \qquad \forall \gamma \in \Gamma, \forall x \in \mathbf{P}_n$$
(24)

The coefficient  $a(\gamma, x)$  has to fulfill the cocycle condition:

$$a(\gamma_1\gamma_2, x) = a(\gamma_1, \gamma_2 x)a(\gamma_2, x) \tag{25}$$

Indeed relation (24) demands

$$P(\gamma_1\gamma_2 x) = a(\gamma_1\gamma_2, x)P(x)$$
  
=  $a(\gamma_1, \gamma_2, x)P(\gamma_2 x)$   
=  $a(\gamma_1, \gamma_2, x)a(\gamma_2, x)P(x)$ 

The cocycle *a* will be completely determined by the values of  $a(I_k, x), k = 1...\nu$ . These values may be found easily: when applied to  $\gamma_1 = \gamma_2 = I_k$ , condition (25) shows that  $a(I_k, x)$  has to divide a suitable power of  $\phi_k$ .

Finding an invariant  $\Delta = P/Q$  implies finding two polynomials P and Q of the same degree, which transform the same way (are covariant with the same cocycle) under all the generators, i.e:

$$P(I_k(x)) = a(I_k, x) \cdot P(x) \quad \text{and} \quad Q(I_k(x)) = a(I_k, x) \cdot Q(x) \quad (26)$$

Once the cocycle a is chosen, solving (26) becomes a handable linear problem, of which the compatibility can be further studied [9]. We have proved that the proliferation of singularities impeaches the existence of any invariants, but the converse is not true. One of the outcomes is that, in this general setting, the existence of invariants is exceptional.

Finding all invariant subvarieties is more tricky. What we have obtained so far is invariant subvarieties of codimension 1, or subvarieties determined by invariants of  $\Gamma$ . Smaller subvarieties have more than one equation, and (24) has to be replaced by a matrix relation:

$$\Pi(\gamma(x)) = A(\gamma, x)\Pi(x) \qquad \forall \gamma \in \Gamma, \forall x \in \mathbf{P}_n$$
(27)

where  $\Pi$  is a vector constructed from the (non canonical) list of equations, and A a matrix. In other words we demand that the ideal defining the subvariety be invariant by  $\Gamma$ , keeping in mind that the subvariety we look for does not have to be a complete intersection.

Let us describe an example, coming from the hard hexagon model [40, 41]. Consider the two involutions of  $\mathbf{P}_4$  given in terms of homogeneous coordinates  $[x_0, x_1, x_2, x_3, x_4]$ :

$$\begin{array}{rcl} i_1 & \longrightarrow & \left[ x_5 \, x_1 \, x_4, x_4 \, \left( x_0 \, x_5 - x_2^2 \right), -x_2 \, x_1 \, x_4, x_1 \, \left( x_0 \, x_5 - x_2^2 \right), x_0 \, x_1 \, x_4 \right] \\ i_2 & \longrightarrow & \left[ x_4 \, x_2 \, x_5, -x_1 \, x_2 \, x_5, x_5 \, \left( x_0 \, x_4 - x_1^2 \right), x_0 \, x_2 \, x_5, x_2 \, \left( x_0 \, x_4 - x_1^2 \right) \right] \end{array}$$

The two inverses are related by a linear transformation, i.e  $i_2 = \tau i_1 \tau$  with

$$\tau : [x_0, x_1, x_2, x_4, x_5] \longrightarrow [x_0, x_2, x_1, x_5, x_4]$$
(28)

and  $\phi_1 = i_1^2 = x_1^2 x_4^2 (x_0 x_5 - x_2^2)^2$ . If

$$\begin{aligned} \Pi_1 &= x_0^2 x_2^2 x_4 + x_0^2 x_1^2 x_5 - x_0^3 x_4 x_5 - x_4^2 x_5 x_2^2 - x_4 x_5^2 x_1^2 + x_0 x_4^2 x_5^2 - x_1^2 x_2^2 x_0 \\ \Pi_2 &= x_0^2 - x_4 x_5 - \lambda x_1 x_2 \end{aligned}$$

with  $\lambda$  a free parameter, then we do have relation (27) with a matrix

$$A_{1} = \begin{bmatrix} x_{1}^{3}x_{4}^{3} (x_{0} x_{5} - x_{2}^{2})^{2} & 0 \\ -x_{4} & -x_{4}^{2} (x_{0} x_{5} - x_{2}^{2}) \end{bmatrix}$$

and

$$A_1(i_1(x))A_1(x) = \begin{pmatrix} \phi_1^5(x) & 0\\ 0 & \phi_1^2(x) \end{pmatrix}$$

Both  $\Pi_1$  and  $\Pi_2$  are unchanged by  $\tau$ ,  $\Pi_1$  is covariant, and consequently the variety  $\Pi_1 = 0$  is invariant by  $\Gamma$ . In contrast,  $\Pi_2$  is not covariant and only the intersection of  $\Pi_2 = 0$  with  $\Pi_1 = 0$  is invariant by  $\Gamma$ .

#### 9 The sixteen-vertex model

We abandon here the strict context of integrability, and turn to problems of discrete dynamical systems, defined in the space of parameters of the models.

The field of discrete dynamical system is a well developed one. The notion of integrable map goes back, in the context of hamiltonian dynamics, to Poincaré, who founded the subject ([42, 43], see also [44, 45]). The last 30 years have seen the subject expand a lot, specifically with the advent of computer calculations. In order to gain simplicity, one was lead long ago to studying iterations of polynomial and polynomially invertible transformations of the plane [46]), eventually renouncing to hamiltonian structures. Analytic maps have also been fruitfully analyzed, especially in one complex dimension [47, 48]. More recently, some remarkable multi-parameter families of maps, eventually integrable, have been constructed from soliton equations [49, 50, 51, 52, 53, 54].

We want to analyze the behaviour of iterations of a typical infinite order element of our groups  $\Gamma$  such as the product of two generating involutions. The first example we will examine is of great interest for statistical mechanics on the lattice. It comes from the general 2-state vertex model on the square lattice in dimension 2, i.e. the sixteen-vertex model. Take R to be the general  $2 \times 2$ matrix. This matrix up to a multiplicative constant represents an element of the 15-dimensional projective space  $\mathbf{P}_{15}$ . A typical orbit of the adequate  $\Gamma$ , when projected to a 2-plane of coordinates, looks like:



Figure 2: One generic orbit of  $\Gamma$  for the 16-vertex model

This figure shows how efficient the graphical method can be. The situation is

very favorable since the orbit is confined to a low dimensional variety (a curve). What can be proved [6], following the ideas of section (8), is the existence of a collection of invariants, of algebraic rank 14. More precisely, there are 17 invariants with 3 relations. As in Figure 1, the discrete orbits are generically dense in a curve, and this helps making the 'graphical detector of invariants' so useful.

#### 10 More pictures

One should not believe that  $\Gamma$ -orbits are always curves. Let us introduce a class of transformations in  $\mathbf{P}_2$  which contains in particular the transformations (24). Consider  $\Gamma$  generated by I and J, given in homogeneous coordinates  $[x_0, x_1, x_2]$  by:

$$J: [x_0, x_1, x_2] \to [x_1 x_2, x_0 x_2, x_1 x_2]$$
(29)

$$I = C^{-1}JC \tag{30}$$

with C a projective linear transformation of  $\mathbf{P}_2$ . This class of realizations, parametrized by the collineation matrix C already contains all kinds of behaviours. Moreover, many nice examples obtained from matrix inversions do implement relation (30).

#### **10.1** Non-symmetric $Z_7$

Suppose *m* is the following cyclic  $7 \times 7$  matrix:

$$m = \begin{pmatrix} x & y & y & z & y & z & z \\ z & x & y & y & z & y & z \\ z & z & x & y & y & z & y \\ y & z & z & x & y & y & z \\ z & y & z & z & x & y & y \\ y & z & y & z & z & x & y \\ y & y & z & y & z & z & x \end{pmatrix}$$
(31)

The matrix inverse verifies (30) with<sup>2</sup>

$$C_{Z_7} = \begin{bmatrix} 2 & 6 & 6\\ 2 & -1 - i\sqrt{7} & -1 + i\sqrt{7}\\ 2 & -1 + i\sqrt{7} & -1 - i\sqrt{7} \end{bmatrix}$$
(32)

The iteration of  $\varphi = IJ$ , with I the matrix inversion and J the element by element inversion, yields the following picture, in the inhomogeneous coordinates u = y/x, v = z/x, if we represent a number of orbits at the same time:

<sup>&</sup>lt;sup>2</sup>This form is a particular value of an eigenmatrix of conference digraph [55]



Figure 3: A collection of chaotic orbits

Figure 3 shows there should not be any invariant. Indeed, as proved in [9], there cannot be any because  $\Gamma$  has an infinite number of singular points. The first singular points are visible on the figure. Their location is very easy to compute.

#### 10.2A finite diagram model

Take

$$C = \begin{bmatrix} 2 & 0 & 2\\ 1 & 1 & -1\\ -1 & 1 & 1 \end{bmatrix}$$
(33)

This has been introduced in [9] and the denomination 'finite diagram model' refers to the finite number of singular points of  $\Gamma$ . Although such a transformation was not directly obtained from a matrix inverse, it (or similar ones) appears in properly defined reductions of the inversion of matrices of larger sizes [56]. We have found no invariant for this model. The graphical method is quite ineffectual at generic points. It proves nevertheless useful in the vicinity of a fixed point.



- • 9 • • • • • •

Figure 4 represents a collection of orbits not far away from a fixed point of  $(IJ)^2$ . The fixed point  $[\alpha^2, \alpha, -1], \alpha = (1 + \sqrt{5})/2, (x_1/x_0 \simeq .618, x_2/x_0 \simeq -.382)$  is an elliptic fixed point at which the eigenvalues of the linear differential are the roots of  $s^2 - 17/16 \ s + 1$ . We see that the orbits explode when one gets away from the fixed point. This is exactly the situation encountered in [57], see [58].

# 11 The complexity of iterations

More insight into the behaviour of the iterations can be gained by looking at their complexity, in a way we will define.

The first grasp at the complexity of  $\Gamma$  is through its size (section 8), but we shall not comment on this here.

We may satisfy ourselves with examining the simple case  $\nu = 2$ , where the group is essentially two copies of **Z**, and describe in the next section a specific example (section 12) with  $\nu = 3$ .

The simplest situation is the one where  $\Gamma$  acts by transformations of the 2-dimensional projective space  $\mathbf{P}_2$ . We have given in section 10 a class of such a realization, and the study of 'the dynamics of singularities' as explored in [9] proves extremely useful.

It is nevertheless something else we have in mind here: suppose  $\varphi = IJ$  is of degree w. Its iterates  $\varphi^k$  do not have to be of degree  $w^k$ . Indeed there is a simple mechanism for the lowering of the degree d(k) of  $\varphi^k$ , for one has to factorize out

common factors from the expressions of the homogeneous coordinates of  $\varphi^{(k)}$ . We may analyze the sequence d(k), as a function of the order of the iteration k. This measure of the complexity of the iterations actually coincides with the one introduced by Arnold [59]; see [60] for the particular case of bi-polynomial transformations of the plane. It brings a variety of behaviours, lying between the generic exponential growth and periodicity, with the particular instance of polynomial (or polynomially bounded) growth. In a given system of coordinates the notion of degree is well defined. Of course, changes of coordinates may change the degree, but allowed birational changes will preserve the nature of the growth.

We have calculated the degree of the successive iterates of  $\varphi$  in a number of cases, the simplest ones being given by the class (29,30).

If  $C = C_{Z_7}$  given by (32), the first terms are:

$$1, \quad 4, \quad 12, \quad 33, \quad 88, \quad 232, \quad 609, \quad \dots \quad (34)$$

This may be seen as the first terms of the sequence:

$$d(k) = \sum_{i=0}^{k} f_{2i}$$

with  $f_i$  the Fibonacci sequence:

$$1, \quad 2, \quad 3, \quad 5, \quad 8, \quad 13, \quad 21, \quad 34, \quad \dots$$

This behaviour is found for any *generic* elements of the four parameter family [61] of collineations,

$$\begin{bmatrix} 1 & p & q-p-1 \\ 1 & s & -s-1 \\ 1 & r & -r-1 \end{bmatrix}$$
(35)

Notice that the Jaeger-Higman-Sims model belongs to this family (compare collineations (24) and (35)), and has the same sequence (34).

For the 'finite diagram model' with C given by (33) the degree reaches its maximal value:

$$d(k) = 4^k$$

For

$$C(w) = \begin{bmatrix} 1 & w - 1 & w \\ 1 & -1 & 0 \\ 1 & 0 & -1 \end{bmatrix},$$
(36)

 $\Gamma$  is known to have an invariant, and the orbits lie within elliptic curves of equation

$$P_w^2(x_0, x_1, x_2)Q_w(x_0, x_1, x_2) - \lambda(x_1 + x_2)^4(x_0 - x_2)^2(x_0 - x_1) = 0$$

where

$$P_w = (1 - w)(x_2^2 - x_0 x_1) + (w - 3)x_2(x_0 - x_1),$$
$$Q_w = (1 - w^2)(x_1^3 - x_0 x_2^2) + (w^2 - 4w - 1)x_1^2(x_0 - x_2) + 2(w - 1)^2x_1x_2(x_0 - x_1))$$

and the sequence of degrees starts with:

1, 4, 10, 20, 34, 53, 75, 102, 132, 167, 206, 249, 295, 347, 402, 461,  $\dots$ 

that is to say the second derivative d(k+1) - d(k-1) - 2d(k) is periodic of period 12. Notice that (36) has the form (35). It is just not generic.

As a last example, in the Baxter model, and also for the 16-vertex model, the iteration of  $\varphi = t_g I$  yields the sequence of degrees:

 $1, \quad 3, \quad 9, \quad 19, \quad 33, \quad 51, \quad 73, \quad 99, \quad 129, \quad 163, \quad 201, \quad \dots \quad (37)$ 

from which one infers

$$d(k) = 2k^2 + 1$$

We have examined many more examples, and they all lead to the conjecture [9]: chaotic behaviour implies exponential growth, as regular behaviour implies polynomial bounds.

The proof of this conjecture should follow from considerations on addition on elliptic curves [62] at least for realizations in  $\mathbf{P}_2$ , the main idea being that  $\varphi$ is a constant shift on some curve.

Another approach goes through the study of the factorization properties: the reduction of the degree d(k) from exponential to polynomially bounded comes from a sufficiently regular factorization process. An analysis of this process was successfully undertaken in [63, 56] for various realizations  $\Gamma$  and confirms up to now our conjecture. We shall give one more example in the next section.

# 12 An example

In [7] we introduced a restriction of the general 2-state model on the cubic lattice in three dimensions, by imposing the following relations on the entries of the R-matrix:

$$R_{j_1j_2j_3}^{i_1i_2i_3} = R_{-j_1,-j_2,-j_3}^{-i_1,-i_2,-i_3}$$
(38)

$$R_{j_1j_2j_3}^{i_1i_2i_3} = 0 \quad \text{if} \quad i_1i_2i_3j_1j_2j_3 = -1 \tag{39}$$

R is the direct product of two times the same  $4 \times 4$  submatrix [5]. It is further possible to impose that this  $4 \times 4$  matrix is symmetric, since the product  $t = t_l t_m t_r$  acts as its transposition.

$$R_{j_1j_2j_3}^{i_1i_2i_3} = R_{i_1i_2i_3}^{j_1j_2j_3} \tag{40}$$

We use the following notations for the 10 homogeneous entries of this  $4 \times 4$  submatrix:

$$\begin{pmatrix} a & d_1 & d_2 & d_3 \\ d_1 & b_1 & c_3 & c_2 \\ d_2 & c_3 & b_2 & c_1 \\ d_3 & c_2 & c_1 & b_3 \end{pmatrix}.$$
(41)

The generators of  $\Gamma$  are the inversion of R, which acts as a matrix inverse on (41), and the permutations of entries  $t_l$ ,  $t_m$ , and  $t_r$  being respectively:

$t_l$ :	$c_3 \leftrightarrow d_3,$	$c_2 \leftrightarrow d_2$
$t_m$ :	$c_1 \leftrightarrow d_1,$	$c_3 \leftrightarrow d_3$
$t_r$ :	$c_2 \leftrightarrow d_2,$	$c_1 \leftrightarrow d_1$

A numerical iteration gives the following picture of an orbit, projected on a 2-plane of coordinates.



Figure 5: A 'two-dimensional' orbit

This is another example of the efficiency of the graphical detector: we see that the orbit is confined to a 2-dimensional surface. This may be easily proven. If one defines

$$p_3 = ab_3 + b_1b_2 - c_3^2 - d_3^2, \qquad q_3 = c_1d_1 - c_2d_2, \tag{42}$$

together with the polynomial  $p_2$  and  $p_1$  obtained by permutation of the indices 1,2, and 3, and (with a corrected misprint from [7]):

$$\begin{array}{rcl} r_3 &=& ab_3-b_1b_2+c_3^2-d_3^2,\\ s_3 &=& (a+b_3)c_3-d_1d_2-c_1c_2,\\ t_3 &=& (b_1+b_2)d_3-d_1c_2-c_1d_2. \end{array}$$

then

- The polynomials  $p_i, q_i, i = 1, 2, 3$  form a five dimensional space, and any ratio of these polynomials is invariant by the *whole group*  $\Gamma$ . This means that the orbit of  $\Gamma$  is confined to a 5-dimensional subspace of the original 9-dimensional parameter space.
- If we consider the subgroup generated by I and  $t_r$ , then  $r_3, s_3, t_3$  are three more covariant polynomials, and they furnish three more invariants, proving that the orbit is confined to 2 dimensions.

As far as the complexity is concerned, the non-chaotic image fits with the sequence of degrees, which is the same as (37). The analysis of the degrees can be performed easily: Let M be a generic matrix of the form (41), and let  $M_0 = t_r M$  be the starting point of the iteration. We iterate  $t_r I$  defined polynomially in terms of the entries, and keep track of the common factors (factorized out from the result of the action of K). Let  $M_{k+1} = t_r I(M_k)$ ,  $f_k$  the extracted factor, and  $\Delta_k = det(M_k)$ . One gets  $d_0 = det(t_r M)$ ,  $f_0 = 1$ ,  $d_1 = det(t_r I t_r M)$ ,  $f_2 = 1$ . Then a factorization of the determinants appears regularly, defining in the course an additional sequence  $\delta_k$  such that:

$$\Delta_2 = \Delta_0^3 \delta_2 \quad \Delta_3 = \Delta_1^3 \delta_3 \quad \Delta_4 = \delta_2^3 \delta_4 \quad \Delta_5 = \delta_3^2 \delta_5 \quad \dots \quad \Delta_n = \delta_{n-2}^3 \delta_n \\ \dots \qquad f_3 = \Delta_0^2 \quad f_4 = \Delta_1^2 \quad f_5 = \Delta_2^2 / \Delta_0^6 = \delta_2^2 \quad \dots \quad f_n = \delta_{n-3}^2$$

Denoting by  $u_k, v_k, g_k, x_k$  the degrees of respectively  $\Delta_k, \delta_k, f_k$  and of the matrix elements of  $M_k$  in terms of the entries of  $M_0$ , this implies

$$u_{k} = 3 v_{k-2} + v_{k}$$
  

$$g_{k} = 2 v_{k-3}$$
  

$$u_{k} = 4 x_{k}$$
  

$$x_{k} = 3 x_{k-1} - g_{k}$$

resolved by

$$x_{k+3} - 3 x_{k+2} + 3 x_{k+1} - x_k = 0$$
 and  $d(k) = 2k^2 + 1$ 

This is an instance of the general result obtained in [56] (formula (3.19), for q = 4), giving a good flavour of the method.

## 13 Conclusion and perspectives

Inversion relations have proved to be a powerful tool in the study of statistical mechanical models, leading in particular to exact relations on the partition functions, even for non-integrable or higher dimensional models [64, 65, 66].

The description often appeals to a diagrammatic representation [67], which we have not used here.

The Yang-Baxter equations in their various forms have brought an enormous amount of exact results in the field. They have unified branches of physics and mathematics such as ice models [68, 69] and solitonic wave equations [70, 71], gave new insights into knot polynomials [72, 73, 74, 75], and produced successful offsprings like quantum groups [76, 77, 78].

What [3, 4] contain is a concrete link between discrete dynamical systems and quantum integrability, producing a large number of interesting dynamical systems. We have been able to use a little part of the paraphernalia available from the field of dynamical systems –the simplest being numerical calculations–. One outcome is that the rational nature of the transformations captures a good proportion of the algebro-geometric content of the Yang-Baxter equations [33, 6].

One issue for future work is the resolution of the tetrahedron equations, and the finding of truly three-dimensional integrability. Indeed there is a competition then between the higher overdetermination of the systems of equations to solve, and the increasing size of the corresponding group  $\Gamma$ . This antagonism might be resolved either by a relative triviality of the solutions, which are disguised twodimensional solutions, either by a low complexity of the actual realization of  $\Gamma$ , and this leaves room for interesting solutions. The most natural to conjecture is that  $\Gamma$  has a *finite realizations*, or that its trajectories lie on abelian varieties [79].

The remarkable elliptic parametrization [6] of the 16-vertex model should lead to interesting properties of the corresponding transfer matrices, through controlled functional relations on the exact partition function and/or via a construction of physically relevant spaces of states. This applies as well to three-dimensional models [7], and prompts us to a return to basics, i.e. Bethe Ansatz [80].

Acknowledgment. I would like to thank F. Jaeger for a very useful discussion, and J. Avan, M. Bellon, J.-M. Maillard, G. Rollet, M. Talon for continued stimulating exchanges.

# References

- M.P. Bellon, J-M. Maillard, and C-M. Viallet, *Integrable Coxeter Groups*. Physics Letters A 159 (1991), pp. 221–232.
- [2] M.P. Bellon, J-M. Maillard, and C-M. Viallet, *Higher dimensional map*pings. Physics Letters A 159 (1991), pp. 233–244.

- [3] M.P. Bellon, J-M. Maillard, and C-M. Viallet, Infinite Discrete Symmetry Group for the Yang-Baxter Equations: Spin models. Physics Letters A 157 (1991), pp. 343–353.
- M.P. Bellon, J-M. Maillard, and C-M. Viallet, *Infinite Discrete Symmetry Group for the Yang-Baxter Equations: Vertex Models.* Phys. Lett. B 260 (1991), pp. 87–100.
- [5] M.P. Bellon, J-M. Maillard, and C-M. Viallet, *Rational Mappings, Arborescent Iterations, and the Symmetries of Integrability.* Physical Review Letters 67 (1991), pp. 1373–1376.
- [6] M.P. Bellon, J-M. Maillard, and C-M. Viallet, Quasi integrability of the sixteen-vertex model. Phys. Lett. B 281 (1992), pp. 315–319.
- [7] M.P. Bellon, S. Boukraa, J-M. Maillard, and C-M. Viallet, Towards threedimensional Bethe Ansatz. Phys. Lett. B 314 (1993), pp. 79–88.
- [8] M.P. Bellon, J-M. Maillard, G. Rollet, and C-M. Viallet, Deformations of dynamics associated to the chiral Potts model. Int. J. Mod. Phys. B 6 (1992), pp. 3575–3584.
- G. Falqui and C.-M. Viallet, Singularity, complexity, and quasi-integrability of rational mappings. Comm. Math. Phys. 154 (1993), pp. 111–125.
- [10] L. Onsager, Crystal statistics: I. A two-dimensional model with an orderdisorder transition. Phys. Rev 65 (1944), pp. 117–149.
- J.B. McGuire, Studies of exactly solvable one-dimensional N-body problems.
   J. Math. Phys. 5 (1964), pp. 622–636.
- [12] R.J. Baxter. Exactly solved models in statistical mechanics. London Acad. Press, (1981).
- [13] C.N. Yang, Some exact results for the many-body problem in one dimension with repulsive delta-function interaction. Phys. Rev. Lett. 19(23) (1967), pp. 1312–1315.
- [14] L.A. Takhtajan and L.D. Faddeev, The quantum inverse problem and the XYZ Heisenberg model. Russian Math. Surveys 34(5) (1979), pp. 11–68.
- [15] L.D. Faddeev, E.K. Sklyanin, and L.A. Takhtajan, *The quantum inverse problem method.* Theor. Math. Phys **40** (1980), p. 688.
- [16] E.K. Sklyanin, Quantum version of the method of inverse scattering. transl. of Zap. Nauch. Sem. LOMI Steklov 95 (1980), pp. 55–128.
- [17] P.P. Kulish and E.K. Sklyanin. volume 151 of Lecture Notes in Physics, pages 61–119, (1982).

- [18] L.D. Faddeev. Integrable models in 1+1 dimensional quantum field theory. In Les Houches Lectures (1982), Amsterdam, (1984). Elsevier.
- [19] M. Gaudin. La fonction d'onde de Bethe. Collection du C.E.A. Série Scientifique. Masson, Paris, (1983).
- [20] E.K. Sklyanin. Quantum inverse scattering method. selected topics. In Quantum Groups and Quantum Integrable Systems, Singapore, (1992). World Scientific. Proceedings of the Nankai Lectures 1991 (and preprint hep-th-9211111).
- [21] H.S.M. Coxeter and W.O.J. Moser. Generators and relations for discrete groups. Springer Verlag, second edition, (1965).
- [22] V.F.R. Jones, *Baxterization*. Int. J. Mod. Phys. B 4 (1990), pp. 701–713. proc. of 'Yang-Baxter equations, conformal invariance and integrability in statistical mechanics and field theory', Canberra, 1989.
- [23] R.J. Baxter, Eight-vertex model in lattice statistics. Phys. Rev. Lett. 26(14) (1971), pp. 832–833.
- [24] R.J. Baxter, One dimensional anisotropic Heisenberg chain. Phys. Rev. Lett. 26(14) (1971), p. 834.
- [25] A.B. Zamolodchikov, Tetrahedron equations and the relativistic S-matrix of straight-strings in 2+1 dimensions. Comm. Math. Phys. 79 (1981), p. 489.
- [26] M. T. Jaekel and J. M. Maillard, Symmetry relations in exactly soluble models. J. Phys. A15 (1982), pp. 1309–1325.
- [27] J.M. Maillet and F. Nijhoff, Integrability for multidimensional lattice models. Phys. Lett. B 224 (1989), pp. 389–396.
- [28] M. Demazure, Sous-groupes algébriques de rang maximum du groupe de Cremona. Ann. Scient. Ec. Norm. Sup. 4e série t.3 (1971), pp. 507–588.
- [29] H. Jung, Über ganze birationale Transformationen der Ebene. J. Reine Angew. Math. 184 (1942), pp. 161–172.
- [30] S. Friedland and J. Milnor, Dynamical properties of plane polynomial automorphisms. Ergod. Theory Dyn. Systems 9 (1989), pp. 67–99.
- [31] C. Fan and F.Y. Wu, General lattice statistical model of phase transition. Phys. Rev. B 2 (1970), pp. 723–733.
- [32] B.U. Felderhof, Diagonalization of the transfer matrix of the free-fermion model. II. Physica 66 (1973), pp. 279–297.

- [33] I.M. Krichever, Baxter's equations and algebraic geometry. Funct. Anal. and its Appl. 15 (1981), pp. 92–103.
- [34] H. Au-Yang, B.M. Mc Coy, J.H.H. Perk, S. Tang, and M.L. Yan, Commuting transfer matrices in the chiral Potts models: solutions of the startriangle equations with genus ≥ 1. Phys. Lett. A123 (1987), p. 219.
- [35] R.J. Baxter, J.H.H. Perk, and H. Au-Yang, New solutions of the startriangle relations for the chiral Potts model. Phys. Lett. A128 (1988), p. 138.
- [36] D. Hansel and J. M. Maillard, Symmetries of models with genus > 1. Phys. Lett. A 133 (1988), p. 11.
- [37] F. Jaeger, Strongly regular graphs and spin models for the Kauffman polynomial. Geometriæ Ded. 44 (1992), pp. 23–52.
- [38] P. de la Harpe, Spin models for link polynomials, strongly regular graphs and Jaeger's Higman-Sims model. Pacific J. Math. **162**(1) (1994), pp. 57–96.
- [39] V.F.R Jones, On a certain value of the Kauffman polynomial. Comm. Math. Phys. 125 (1989), pp. 459–467.
- [40] R.J. Baxter and P.A. Pearce, Hard hexagons: interfacial tension and correlation length. J. Phys. A(15) (1982), pp. 897–910.
- [41] J. Avan, M. Talon, J.M. Maillard, and C.M. Viallet, New local relations for lattice models. Int. J. Mod. Phys. B 4 (1990), pp. 1895–1912.
- [42] H. Poincaré. Oeuvres. Tomes I-XI. Gauthier-Villars, Paris, (1952).
- [43] H. Poincaré. Les méthodes nouvelles de la mécanique céleste. Gauthier-Villars, Paris, (1892).
- [44] G. D. Birkhoff, Dynamical systems with two degrees of freedom. Trans. Amer. Math. Soc. 18 (1917), pp. 199–300.
- [45] A.S. Wightman, The mechanics of stochasticity in classical dynamical systems. Perspectives in Statistical Physics (1981), pp. 343–363. Reprinted in "Hamiltonian dynamical systems", R.S. MacKay and J.D. Meiss editors, Adam Hilger (1987).
- [46] M. Henon, A two-dimensional mapping with a strange attractor. Comm. Math. Phys. 50 (1976), pp. 69–77.
- [47] C.L. Siegel, Iteration of analytic functions. Ann. Math. 43 (1942), pp. 807–812.

- [48] J.-C. Yoccoz, Conjugaison différentiable des difféomorphismes du cercle dont le nombre de rotation vérifie une condition diophantienne. Ann. Sc. E.N.S. 4eme série t. 17 (1984), pp. 333–359.
- [49] G.R.W. Quispel, J.A.G. Roberts, and C.J. Thompson, *Integrable Mappings and Soliton Equations*. Phys. Lett. A **126** (1988), p. 419.
- [50] G.R.W. Quispel, J.A.G. Roberts, and C.J. Thompson, *Integrable Mappings and Soliton Equations II.* Physica D34 (1989), pp. 183–192.
- [51] J. Moser and A.P. Veselov, Discrete versions of some classical integrable systems and factorization of matrix polynomials. Comm. Math. Phys. 139 (1991), pp. 217–243.
- [52] B. Grammaticos, A. Ramani, and V. Papageorgiou, Do integrable mappings have the Painlevé property? Phys. Rev. Lett. 67 (1991), pp. 1825–1827.
- [53] V.G. Papageorgiou, F.W. Nijhoff, and H.W. Capel, Integrable mappings and nonlinear integrable lattice equations. Phys. Lett. A147 (1990), pp. 106–114.
- [54] O. Ragnisco. Restricted flows of toda hierarchy as integrable maps. In Proceedings of the XIX I.C.G.T.M.P, (1992).
- [55] Takuya Ikuta. Non-existence of spin models corresponding to non symmetric association schemes of class 2 on 4m + 2 vertices with  $m \ge 1$ . preprint.
- [56] S. Boukraa, J-M. Maillard, and G. Rollet. *Integrable mappings and poly*nomial growth. LPTHE preprint 93-26, to appear in Physica A.
- [57] M. Henon, Numerical study of quadratic area preserving maps. Q. J. Appl. Math. 27 (1969), pp. 291–312.
- [58] M.C. Gutzwiller. Chaos in Classical and Quantum Mechanics. Spinger Verlag, New-York, Berlin, Heidelberg, (1991). Interdisciplinary Applied Mathematics.
- [59] V.I. Arnold, Dynamics of complexity of intersections. Bol. Soc. Bras. Mat. 21 (1990), pp. 1–10.
- [60] A.P. Veselov, Growth and Integrability in the Dynamics of Mappings. Comm. Math. Phys. 145 (1992), pp. 181–193.
- [61] A. Neumaier, Duality in coherent configurations. Combinatorica 9(1) (1989), pp. 59–67.
- [62] M. Bellon. Addition on curves and complexity of quasi-integrable rational mappings. in preparation.

- [63] S. Boukraa, J-M. Maillard, and G. Rollet. Determinantal identities on integrable mappings. LPTHE preprint 93-25.
- [64] R.J. Baxter, The Inversion Relation Method for Some Two-dimensional Exactly Solved Models in Lattice Statistics. J. Stat. Phys. 28 (1982), pp. 1–41.
- [65] M. T. Jaekel and J. M. Maillard, Inverse functional relations on the Potts model. J. Phys. A15 (1982), pp. 2241–2257.
- [66] D. Hansel, J.M. Maillard, J. Oitmaa, and M.J. Velgakis, Analytical properties of the anisotropic cubic Ising model. J. Stat. Phys. 48 (1987), pp. 69–80.
- [67] R.J. Baxter, Solvable eight-vertex model on an arbitrary planar lattice. Phil. Trans. R. Soc. London 289 (1978), p. 315.
- [68] E.H. Lieb, Exact solution of the F model of an antiferroelectric. Phys. Rev. Lett. 18(24) (1967), pp. 1046–1048.
- [69] E.H. Lieb, Exact solution of the problem of the entropy of two-dimensional ice. Phys. Rev. Lett. 18(17) (1967), pp. 692–694.
- [70] V.E. Zakharov and L.D. Faddeev, The Korteweg-de Vries equation, a completely integrable hamiltonian system. Funct. Anal. and its Appl. 5 (1971), pp. 280–287.
- [71] L.D. Faddeev and L.A. Takhtajan. Hamiltonian methods in the theory of solitons. Springer Verlag, Heidelberg, (1986).
- [72] L.H. Kauffman, State models and the Jones polynomial. Topology 26 (1987), pp. 395–407.
- [73] V.G. Turaev, The Yang-Baxter equation and invariants of links. Invent. Math. 92 (1988), pp. 527–553.
- [74] V.F.R. Jones, On knots invariants related to some statistical mechanical models. Pacific J. Math. 137 (1989), pp. 311–334.
- [75] F.Y. Wu, Knot theory and statistical mechanics. Rev. Mod. Phys. 64(4) (1992), pp. 1099–1131.
- [76] M. Jimbo, A q-difference analogue of U(G) and the Yang-Baxter equation. Lett. Math. Phys. 10 (1985), p. 63.
- [77] V.G. Drinfel'd. Quantum groups. In Proceedings of the International Congress of Mathematicians. Berkeley, (1986).

- [78] S.L. Woronowicz, Twisted SU(2) Group. An Example of Noncommutative Differential Calculus. Publ. Res. Inst. Math. Sci. 23 (1987), pp. 117–181.
- [79] I.G. Korepanov. Vacuum Curves, Classical Integrable Systems in Discrete Space-Time and Statistical Physics. preprint hep-th 9312197, (1993).
- [80] H. Bethe, Zur Theorie der Metalle. I. Eigenwerte und Eigenfunktionen der linearen Atomkette. Z. Phys. 71 (1931), pp. 205–226.