A MathematicaTM Package for Computing N = 2 Superfield Operator Product Expansions *

S. Krivonos^{\dagger}

JINR – Bogoliubov Laboratory of Theoretical Physics, 141980 Dubna, Moscow Region, Russia. and INFN-Laboratori Nazionali di Frascati,

P.O. Box 13, I-00044 Frascati, Italy.

K. Thielemans[‡]

Theoretical Physics Group Imperial College, London SW7 2BZ, UK.

November 1995

Abstract

We describe a general purpose $Mathematica^{TM}$ package for computing Superfield Operator Product Expansions in meromorphic N = 2superconformal field theory. Given the SOPEs for a set of "basic" superfields, SOPEs of arbitrarily complicated composites can be computed automatically. Normal ordered products are always reduced to a standard form. It is possible to check the Jacobi identities, and to compute Poisson brackets ("classical SOPEs"). We present two explicit examples: a construction of the "small" N = 4 superconformal algebra in terms of N = 2 superfields, and a realisation of the N = 2superconformal algebra in terms of chiral and antichiral fermionic superfields.

^{*} This work was partly carried out in the framework of the project "Gauge theories, applied supersymmetry and quantum gravity", contract SC1-CT92-0789 of the European Economic Community.

 $^{^{\}dagger}\mathrm{Email}$ address : krivonos@thsun1.jinr.dubna.su

[‡]Email address : k.thielemans@ic.ac.uk

1 Introduction

It is well known that Operator Product Expansions (OPEs) form a powerful tool in two-dimensional Conformal Field Theory (CFT). Once the OPEs for the currents are known, it is possible to compute their correlation functions algebraically. One also derives differential equations for the other correlation functions in the theory [1]. In gauge theories, one can compute the anomaly in the BRST nilpotency, find the spectrum, etc. Although the calculations are in principal straightforward. In practice, they can become quite cumbersome to do by hand. In the case of a meromorphic CFT, these calculations can be performed with the *OPEdefs* package [3, 4], implemented in *Mathematica*^{TM 1}, an interactive environment for performing symbolic computations.

In the case of supersymmetric conformal theories, it is often useful to work in a superspace formalism. This drastically reduces the number of fields one needs to consider, and also the amount of algebraic manipulations. For example, a general N = 2 superfield contains 4 components fields (two bosonic and two fermionic), therefore one SOPE for the N = 2 superfield defines 10 OPEs for the components. Moreover, the checking of one Jacobi identity in terms of N = 2 superfields is equivalent to 64 identities in terms of components.

Unfortunately, although the formulas to deal with super OPEs are similar to the ones for ordinary OPEs, they are more complicated and calculations are more error-prone. This paper describes a *Mathematica* package that enables one to perform computations automatically in the case of N = 2 supersymmetry.

Of course, the simplest supersymmetric extension of the conformal field theories possesses N = 1 supersymmetry. But an N = 1 superfield contains only one additional fermionic component with respect to the bosonic case and so it is possible to use the *OPEdefs* package. On the other hand, when dealing with higher supersymmetry $N \ge 3$, most of the superfields are constrained, but they can be nicely represented in terms of N = 2 ones. In the case of N = 2, the constraints (chirality or antichirality) can be easily taken into account in *SOPEN2defs*. This is why we decided to limit ourselves for the time being to the case of N = 2 superfields.

The package is of a similar nature to *OPEdefs*. One needs to declare the superfields which are used, and give their respective SOPEs. Nonlinear algebras can be handled by using a point–splitting definition for composites. It is then possible to check the Jacobi identities for the SOPEs. If one introduced arbitrary constants in the SOPEs, it is possible in this way to find the values for which the algebra is associative, and hence to construct a super *W*–algebra. In this way, the superfield version of $N = 2 W_3^{(2)}$ was constructed in [5]. Once the basic SOPEs are given, it is in principle possible to compute any OPE (or a particular pole) of any composite. Of course, this

¹ Mathematica is a trademark of Wolfram Research Inc. For details, see [2].

is restricted by the available memory and CPU–time. The package is able to compute SOPEs, but also Poisson brackets ("classical" SOPEs). It has been tested by a number of people, already resulting in many publications.

The package can be extended to perform calculations using dummy indices. This is still a limited facility. Not all possible simplifications are performed when tensors appear which have symmetries when interchanging indices. However, for many cases this possibility is quite useful.

This paper is setup as follows. In a first section we explain our conventions for N = 2 superspace. Then we derive the formulas for SOPEs. The next section consists of a user's guide for the package. We end with two explicit examples.

Notation

Input for and output from Mathematica is written in typeset font. Input lines are preceded by "In[n] :=", and corresponding output statements by "Out[n] =", as in Mathematica.

2 N = 2 superspace. Notations and conventions

In this section we will introduce our conventions concerning the N = 2 superfield formalism.

The N = 2 superspace $Z = \{z, \theta, \overline{\theta}\}$ can be described by one real bosonic coordinate z and pair of two conjugate Grassman cordinates $\theta, \overline{\theta}$:

$$\theta^2 = \bar{\theta}^2 = 0, \qquad \theta \bar{\theta} = -\bar{\theta} \theta.$$
 (1)

The integration measure in N = 2 superspace is $dZ \equiv dz d\theta d\theta$ and the standard convention for the integration over $\theta, \bar{\theta}$ is assumed.

To deal with N = 2 superfields $\Phi(Z)$, it is useful to introduce chiral and antichiral spinor derivatives $\mathcal{D}, \overline{\mathcal{D}}$

$$\mathcal{D} = \frac{\partial}{\partial \theta} - \frac{1}{2} \bar{\theta} \frac{\partial}{\partial z}, \quad \overline{\mathcal{D}} = \frac{\partial}{\partial \bar{\theta}} - \frac{1}{2} \theta \frac{\partial}{\partial z}, \tag{2}$$

which obey the following relations

$$\mathcal{D}^2 = \overline{\mathcal{D}}^2 = 0, \quad \left\{ \mathcal{D}, \overline{\mathcal{D}} \right\} = -\partial_z.$$
 (3)

Now we can define chiral F and antichiral \overline{F} superfields which are subjected to the following constraints

$$\mathcal{D}F = \overline{\mathcal{D}}\bar{F} = 0, \tag{4}$$

respectively.

It is also instructive to write down the Taylor expansion for a general superfield $\Phi(Z)$

$$\Phi(Z_1) = \sum_{n \ge 0} \frac{Z_{12}^n}{n!} \partial_2^n \left\{ 1 + \theta_{12} \mathcal{D}_2 + \bar{\theta}_{12} \overline{\mathcal{D}}_2 - \theta_{12} \bar{\theta}_{12} \frac{1}{2} \left[\mathcal{D}_2, \overline{\mathcal{D}}_2 \right] \right\} \Phi(Z_2), \quad (5)$$

where

$$\theta_{12} = \theta_1 - \theta_2, \quad \bar{\theta}_{12} = \bar{\theta}_1 - \bar{\theta}_2, \quad Z_{12} = z_1 - z_2 + \frac{1}{2} \left(\theta_1 \bar{\theta}_2 - \theta_2 \bar{\theta}_1 \right).$$
 (6)

We give the SOPE for the simplest case of N = 2 superconformal algebra (SCA):

$$J(Z_1)J(Z_2) = \frac{c/4}{Z_{12}^2} + \frac{\theta_{12}\bar{\theta}_{12}J(Z_2)}{Z_{12}^2} + \frac{\bar{\theta}_{12}\overline{\mathcal{D}}J(Z_2)}{Z_{12}} - \frac{\theta_{12}\mathcal{D}J(Z_2)}{Z_{12}} + \frac{\theta_{12}\bar{\theta}_{12}\partial J(Z_2)}{Z_{12}}, \quad (7)$$

where J(Z) is a general bosonic superfield [6].

3 Formulas for SOPEs

In principle, the formulas to work with SOPEs follow by expanding everything in components, applying the standard OPE formulas, and reassembling again into superfields. However, this approach is impractical. It is easier (and conceptually clearer) to construct the formulas by using Taylor expansions, contour integration ... in superspace. We will discuss SOPEs here and briefly comment on which changes are needed for the Poisson bracket case.

In fact, it will be useful to introduce a notation valid for any number of supersymmetries N [6]. For a list of numbers $[i_l]$, we will write $\theta^{[i]}$ for $\theta^{i_1}\theta^{i_2}\ldots$ and $\theta^{[N]}$ is used for $\theta^{i_1}\ldots\theta^{i_N}$. The "complement" N-[i] is defined such that $\theta^{[N]} = \theta^{N-[i]}\theta^{[i]}$.

We also introduce projectors \mathcal{P} to extract a component of superfield or a SOPE. \mathcal{P}_{\emptyset} projects on the term without θ 's, $\mathcal{P}_{[1]}$ recovers the term proportional to θ^1 and so on:

$$\mathcal{P}_{[i]} = \int d\theta^{[N]} \ \theta^{N-[i]} \tag{8}$$

We will use the following notation for SOPEs:

$$A(Z_1)B(Z_2) = \sum_{n < =h(A,B)} \frac{[AB]_n(Z_2, \theta_{12}^i)}{(Z_{12})^n} \quad , \tag{9}$$

where h(A, B) is some finite number, and is usually given in terms of the conformal dimensions of A and B as $h(A, B) = [d_A + d_B + N/2]$. Note that

with this definition (9), $[AB]_n$ depends on the θ_{12}^i , but not on z_1 . The term in $[AB]_0$ which has no θ_{12} dependence is called the normal ordered product of A and B:

$$:AB: = \mathcal{P}_{\emptyset}[AB]_0. \tag{10}$$

With the definition (9), we have:

$$\mathcal{P}_{[i]}[AB]_n(Z_2,\theta_{12}^i) = \oint_{C_2} \frac{dz_1}{2\pi i} \int d\theta_1^{[N]} \ \theta_{12}^{N-[i]}(Z_{12})^{n-1} A(Z_1) B(Z_2) \,, \quad (11)$$

where the contour C_2 encircles z_2 . If we replace the SOPE of A and B by the Poisson bracket $\{A(Z_1), B(Z_2)\}_{PB}$, we adopt the above formula as the definition for $[AB]_n$ with n > 0 in the Poisson brackets ("classical" SOPEs).

Formulas

With all these definitions, we can start to derive how we have to compute with SOPEs. First, there are some formulas for computing an SOPE of two superfields when one of them is a derivative. These follow simply by taking derivatives of the SOPE. We give only one example (in N = 2):

$$[\mathcal{D}AB]_n(Z_2,\theta_{12}^i) = \frac{(n-1)}{2}\bar{\theta}_{12}[AB]_{n-1}(Z_2,\theta_{12}^i) + \mathcal{D}_1[AB]_n(Z_2,\theta_{12}^i), \quad (12)$$

where the subindex on \mathcal{D}_1 indicates it acts on coordinates Z_1 . This implies

$$\mathcal{P}_{\theta}[AB]_{n} = \mathcal{P}_{\emptyset}[\mathcal{D}AB]_{n} \,. \tag{13}$$

For a term in the regular part of an OPE, we find:

$$[A B]_{-n} = \frac{1}{n!} [\partial^n A B]_0 .$$
 (14)

The SOPE $B(Z_1)A(Z_2)$ is by analytic continuation equal to $A(Z_2)B(Z_1)$. It remains then to apply a Taylor expansion (5) to the fields in the latter. The result is straightforward. We list only the special case for normal ordering, where drastical simplifications occurs because of the projection \mathcal{P}_{\emptyset} in (10):

$$:BA: = (-1)^{|A||B|} :AB: + (-1)^{|A||B|} \sum_{l \ge 1} \frac{(-1)^l}{l!} \partial^l \mathcal{P}_{\emptyset}[AB]_l$$
(15)

Applying this formula for A = B a fermionic superfield, one gets an expression for : AA: in terms of the poles in the SOPE $A(Z_1)A(Z_2)$.

Associativity of the SOPE means that SOPEs can be computed in any order (inside correlation functions). This can be used to derive the Jacobi identities. As a first step, we select a particular term in a double SOPE by using contour integrals:

$$\mathcal{P}_{[i]}[A \mathcal{P}_{[j]}[BC]_p]_q(Z_3) = \oint_{C_3} \frac{dZ_1}{2\pi i} \theta_{13}^{N-[i]} Z_{13}^{q-1}$$
$$\oint_{C_3} \frac{dZ_2}{2\pi i} \theta_{23}^{N-[j]} Z_{23}^{p-1} A(Z_1) B(Z_2) C(Z_3), \qquad (16)$$

where C_2 denotes a contour which encircles z_2 once anti-clockwise. We can now use a contour deformation argument relating the contour integral in eq. (16) to a contour integral where the integration over Z_2 is performed last. In fact, the contour deformation works only for the θ -independent part of the integral. For the Berezin integral over θ , we just interchange the order of integration (taking signs into account). The resulting integral has two terms: one where the z_1 contour is around z_3 , and one where it is around z_2 . We find:

$$\mathcal{P}_{[i]}[A \mathcal{P}_{[j]}[BC]_p]_q = (-1)^{(|A|+|i|)(B+|j|)} \mathcal{P}_{[j]}[B \mathcal{P}_{[i]}[AC]_q]_p + (-1)^{(|A|+|i|)(|j|)} \sum_{l>0} \binom{q-1}{l-1} \mathcal{P}_{[j]}[\mathcal{P}_{[i]}[AB]_l C]_{p+q-l}.$$
 (17)

This equation has to hold (inside correlators) for consistency of the OPEformalism. It is valid for any integers p, q, *i.e.* also negative numbers. However, in practice we use the Jacobi identities eq. (17) for positive p, q as equations for the singular part of the SOPEs, while for p or q zero, they define how one should calculate with composites. For example, to compute an SOPE with a composite, we insert $p = 0, [j] = \emptyset$ into eq. (17). When $q = 0, [i] = \emptyset$ as well, we find:

$$:A : BC ::= (-1)^{|A||B|} : B : AC :: + : (:AB : -(-1)^{|A||B|} : BA :) C: , (18)$$

where we used eq. (15).

For Poisson brackets, the normal ordered product of two operators is simply replaced by (noncommutative) multiplication. In particular, this means that double contractions (l < q) in eq. (17) for $p = 0, [j] = \emptyset$ drop out. However, the Jacobi identities eq. (17) for positive p, q remain exactly the same.

Improvements

The rules given in this section up to now are sufficient to compute any SOPE, and to reorder any composite into a standard form. For example, to compute an SOPE where the first operator is a composite, we could use the fact that $A(Z_1)B(Z_2) = B(Z_2)A(Z_1)$ and (17). However, it is possible to construct a rule by using contour integrals to do this in one step. We find:

$$[\mathcal{P}_{\emptyset}[AB]_{0} C]_{q} = \sum_{l \ge 0} \frac{1}{l!} [\partial^{l} A \ [BC]_{l+q}]_{0} + (-1)^{|A||B|} \sum_{l \ge 0} \frac{1}{l!} [\partial^{l} B \ [AC]_{l+q}]_{0} + (-1)^{|A||B|} \sum_{l=1}^{q-1} [B \ [AC]_{q-l}]_{l},$$
(19)

where $q \geq 1$ and we used eq. (14). To have the formula in this simple form, we used the convention that any terms with $\theta_{12}, \bar{\theta}_{12}$ move to the left of the OPE bracket. For instance, if $[BC]_m = \theta_{12}D$, then $[A \ [BC]_m]_n] = (-1)^{|A|} \theta_{12} [AD]_n$.

For composites, we find:

$$::AB: C: = :A :BC::+ \sum_{l>0} \frac{1}{l!} :\partial^{l}A \mathcal{P}_{\emptyset}[BC]_{l}:+(-1)^{|A||B|} \sum_{l>0} \frac{1}{l!} :\partial^{l}B \mathcal{P}_{\emptyset}[AC]_{l}: (20)$$

4 User's Guide

This section is intended as a user's guide to the package SOPEN2defs. Explicit examples are given for most operations. SOPEN2defs is based on OPEdefs 3.1 [4] and most commands are exactly the same. SOPEN2defs requires Mathematica 2.0 or later.

As SOPEN2defs is implemented as a Mathematica package, it has to be loaded *before* any of its global symbols are used. Loading the package a second time will clear all previous definitions of operators and OPEs, as well as all stored intermediate results. Assuming that the package is located in the Mathematica-path, e.g. in your current directory², issue:

In[1] := <<SOPEN2defs.m

After loading *SOPEN2defs* into *Mathematica*, help for all the global symbols is provided using the standard help-mechanism, *e.g.* ?OPE.

Now, you need to declare the operators that will be used. If you want to define bosonic (resp. fermionic) operators, use Bosonic (resp. Fermionic). For chiral operators, prefix with Chiral, for antichiral ones, use AChiral. For instance, to define a bosonic field J and an antichiral bosonic field W, the statements are:

In[2] := Bosonic[J]In[3] := AChiralBosonic[W]

For derivatives we use the following notation:

$$\partial_z J \to J', \quad \mathcal{D}J \to \mathrm{DT}[J], \quad \overline{\mathcal{D}}J \to \mathrm{DBT}[J], \quad [\mathcal{D},\overline{\mathcal{D}}]J \to \mathrm{DDB}[J] \quad (21)$$

The order of the declarations fixes also the ordering of operators used by the program:

$$J' < DBT[J] < DDB[J] < DT[J] < J < W$$
(22)

By default, spatial derivatives of an operator are considered "smaller" than the operator itself. This can be reversed using the global option NOOrdering (see below).

Finally, the nonregular OPEs between the basic operators have to be given by listing the operators that occur at the poles, the first operator in

²Use the *Mathematica* command SetDirectory for this.

the list is the one at the highest non-zero pole, the last operator has to be the one at the first order pole. For example, to give J the OPE (7), and W of a conformal dimension 1 primary antichiral superfield with U(1)-charge -2:

Note the operator One which specifies the unit-operator. The symbols t12, tb12 denote $\theta_{12}, \bar{\theta}_{12}$. In an OPE, they should always be multiplied with an operator using **. In particular, one could have t12**One.

Warning: it is important that the operators occuring as arguments of OPE in a *definition* should be given in the order the operators are declared (22), otherwise wrong results will be generated.

A normal ordered product : AB: (10) is entered in the form NO[A,B]. Multiple composites can be entered using only one NO head, e.g. NO[A,B,C]. This input is effectively translated into NO[A, NO[B, C]]. All output is normal ordered with the same convention, *i.e.* from right to left (input can be in any order). Also, the operators in composites will always be ordered according to the standard order (22).

```
In[6] := OPE[J,NO[J,J]]
Out[6] = << 4|| c t12**tb12**One/ 4 ||3|| 0 ||2|| c J/2 +
        tb12**DBT[J] + t12**DT[J] + 2 t12**tb12**N0[J, J]
        ||1|| tb12**DBT[J'] + 2 tb12**N0[J, DBT[J]] +
        t12**DT[J'] - 2 t12**N0[J, DT[J]] +
        2 t12**tb12**N0[J', J] >>
```

This also shows what the output for an OPE looks like. For an OPE with n poles, the output is $<< n || \dots ||n-1|| \dots ||1|| \dots >>$.

Warning: when computing OPEs with composites, or when reordering composites, SOPEN2defs remembers by default some intermediate results. Thus, it is dangerous to change the definition of the basic OPEs after some calculations have been performed. For example, consider a constant a in an OPE. If calculations are performed after assigning a value to a, the intermediate results are stored with this value. Changing a afterwards will give wrong results.

The other globally defined functions available from the package are:

• OPEOperator[operator_, parity_] provides a more general way to declare an operator than Bosonic and Fermionic. The second argu-

ment is the parity of the operator such that $(-1)^{\text{parity}}$ is +1 for a boson, and -1 for a fermion. It can be a symbolic constant. This can be used to declare a *bc*-system of unspecified parity. In such cases, the operator can contain a named pattern:

In[7] := OPEOperator[J[i_],parity[i]]

If one wants to declare more operators, one can group each operator and its parity in a list:

In[8] := OPEOperator[{b[i_],parity[i]},{c[i_],parity[i]}]
See also SetOPEOptions[ParityMethod, _].

- ChiralQ[operator_] tests if operator is chiral. One can use ChiralQ[A] = True.
- AChiralQ[operator_] is the same for antichiral operators.
- OPEPole[n_] [ope_] gets a single pole of an OPE: In[9] := OPEPole[2] [Out[6]]

Out[9] = c J/2 + tb12**DBT[J] + t12**DT[J] + 2 t12**tb12**N0[J, J]

OPEPole[n_] [A_,B_] can also be used to compute only one pole term of an OPE:

In[10] := % - OPEPole[2][J, J]//Expand
Out[10] = 0

OPEPole can also give terms in the regular part of the OPE:

```
In[11] := OPEPole[-1][J, J]
```

Out[11] = NO[J',J] + tb12**NO[DBT[J'],J] + t12**NO[DT[J'],J] - t12**tb12**NO[DDB[J'],J]/2

- MaxPole[ope_] gives the order of the highest pole in the OPE.
- **OPEParity**[A_] returns an even (odd) integer if A is bosonic (fermionic).
- OPESimplify[ope_, Function -> function_] "collects" all terms in ope with the same operator and applies function on the coefficients. The default setting for the option Function is Expand, but this can be changed using SetOptions.

OPESimplify[pole_, Function -> function_] does the same simplifications on sums of operators.

The alternative syntax OPESimplify[ope_, function_] is also allowed.

- OPEMap[function_, ope_] maps function to all poles of ope.
- OPEMapAt[function_, ope_, position_] maps function to a part of ope specified by position (see MapAt). For example, OPEMapAt[Expand, ope, {1}]

maps Expand on the highest order pole of ope.

- GetCoefficients[expr_] returns a list of all coefficients of operators in expr which can be OPEs or poles. Applied on a list, it maps on the elements of the list.
- GetOperators [expr_] returns a list of all operators in expr which can be OPEs or poles. Applied on a list, it maps on the elements of the list.
- OPEJacobi [op1_, op2_, op3_] computes the Jacobi-identities (17) for the singular part of the OPEs of the three arguments. In general, all different orderings should be tried to ensure associativity. OPEJacobi returns a list of which all should be zero up to null fields

to be associative.

OPEJacobi accepts an option Function -> function which it passes to OPESimplify at intermediate steps.

- Delta[i_, j_] is the Kronecker delta symbol δ_{ij} .
- Epsilon[i_,j__] is the antisymmetric symbol (with any number of arguments.
- N20PEToComponents [J_] gives the N = 0 components of a superfield J. For $J = A + \theta B + \overline{\theta}C + \theta\overline{\theta}D$ the components are $\{A, B, C, D\}$. In SOPEN2defs, the components are represented as

$$\{J, \mathcal{D}J, \overline{\mathcal{D}}J, -\frac{1}{2}[\mathcal{D}, \overline{\mathcal{D}}]J\},\$$

where a projection on the θ independent part is understood. N2OPEToComponents [ope_, J1_, J2_] computes the 16 OPEs of the components of J_1 and J_2 . It is a double list where the (m, n)-th element is the N = 0 OPE of the *m*-th component of J_1 with the *n*-th component of J_2 .

- ClearOPESavedValues [] clears all stored intermediate results, but not the definition of the operators and their OPEs. To clear everything, reload the package.
- TeXForm[ope_] gives $T_{E}X$ output for an OPE. The arguments are always Z_1, Z_2 .
- TeXFormTD can be assigned a string which will be used by tt TeXForm for the output of DT, DBT, DDB.
- **OPESave[filename_]** (with **filename** a string between double quotes) saves the intermediate results that *OPEdefs* remembers to file (see the option **OPESaving** below).
- SetOPEOptions is a function to set the global options of the package. The current options are:

- SetOPEOptions[NOOrdering, n_] : if n is negative, order higher spatial derivatives to the left (default), if n is positive, order them to the right.
- SetOPEOptions [ParityMethod, 0|1]: makes it possible to use operators of an unspecified parity. When the second argument is 0 (default), all operators have to be declared to be bosonic or fermionic. When the argument is 1, OPEOperator can be used with a symbolic parity. Note that in this case, powers of -1 are used to compute signs, which is slightly slower than the boolean function which is used by the first method.

This option is not normally needed as the use of **OPEOperator** with a non-integer second argument sets this option automatically.

- SetOPEOptions[OPESaving, boolean_] : if boolean evaluates to True (default), OPEdefs stores the intermediate results when computing OPEs of composites and when reordering composites. This option is useful if Mathematica runs short of memory in a large calculation, or when computing with dummy indices (see section 5.2).
- SetOPEOptions[OPEMethod, method_]: with method set to QuantumOPEs enables normal OPE computations (default setting), while ClassicalOPEs enables Poisson bracket computations. Using this option implicitly calls the function ClearOPESavedValues[].
- SetOPEOptions[EnableDummies, True] : loads the Dummies' package and sets it up.

5 Examples

5.1 The N = 4 superconformal algebra

As a first example of using the SOPEdefs package, let us construct the OPEs of N = 4 SU(2) SCA [7] in terms of N = 2 superfields. The N = 4 SU(2) SCA contains the following component currents: the stress-tensor T(z), affine su(2) currents $J^i(z), i = 1, 2, 3$ and two doublets of fermionic spin 3/2 currents $G_{\alpha}, \bar{G}^{\beta}, \alpha, \beta = 1, 2$ which form respectively the fundamental and conjugated representations of su(2). Due to the spin structure, the simplest way to put these component currents into N = 2 supermultiplets is to start with a general spin 1 N = 2 superfield J(Z) which forms the N = 2 SCA (spin contents is $\{1, 3/2, 3/2, 2\}$ together with two bosonic spin 1 chiral-antichiral superfields $\bar{W}(Z), W(Z), D\bar{W} = \overline{D}W = 0$ (spin content $\{1, 3/2, 1, 3/2\}$).

First of all we need to define all superfields we are dealing with:

In[1] := <<SOPEN2defs.m

In[2] := Bosonic[J]

In[3] :=AChiralBosonic[W] In[4] :=ChiralBosonic[WB] As the next step we define the OPE between all the super currents: In[5] := $OPE[J,J] = MakeOPE[{c/4 One + t12**tb12**J}]$ tb12**DBT[J]-t12**DT[J]+t12**tb12**J'}]; In[6] := $OPE[J,W] = MakeOPE[\{t12**tb12**W,$ -2 W - t12**DT[W]+t12**tb12**W'}]; In[7] :=OPE[J,WB] = MakeOPE[{t12**tb12**WB, 2 WB+tb12**DBT[WB]+t12**tb12**WB'}]; In[8] :=OPE[W,WB] = MakeOPE[{a1 t12**tb12**One, a2 One + t12**tb12**J, a3 J + a4 t12**DT[J]}];

Let us note that due to the first OPE, J(Z) obeys the N = 2 SCA, while the second and third OPEs define W(Z) and $\overline{W}(Z)$ as antichiral-chiral superfields primary spin 1 with respect to J(Z). As concerning the last OPE we introduce the arbitrary coefficients a1 - a4 for all the terms because we want to determine the exact form of this OPE. The possible terms and their structure in this OPE are completely fixed by the antichiral-chiral structure of the W, \overline{W} supercurrents (the coefficient before term $\theta_{12} * *\overline{\theta}_{12} * *J$ can be chosen to be 1 due to the scaling invariance $W \to \alpha W$).

In[10] := sol1 = Solve[GetCoefficients[%]==0]

 $Out[10] = \{\{a1 \rightarrow -c/4, a2 \rightarrow -c/4, a3 \rightarrow 2, a2 \rightarrow 2\}\}$

Now we can check that all other Jacobi identities are satisfied:

In[11] := OPESimplify[OPEJacobi[W,W,WB]/.First[sol1]]//Union $<math>Out[11] = \{0\}$

By this we finished the construction of the N = 4 SU(2) SCA in terms of N = 2 superfields.

5.2 A Miura–like realisation of N = 2

As a second example we will demonstrate how to use the SOPEN2defs package to find a realisation of the N = 2 SCA in terms of M pairs of chiralantichiral fermionic superfields (Miura type realisation). We also show here our technique to deal with summing over repeated indices, which is implemented in a separate package *Dummies*. This package can also be used together with *OPEdefs*, or even independently. We only explain some basic

 $^{^3}$ This statement takes 5 seconds on a Pentium 90 MHz running Mathematica 2.2 for Windows.

commands of *Dummies* here.

The first step is loading the package.

In[1] := <<SOPEN2defs.m

Enable working with dummy indices.

In[2] := SetOPEOptions[EnableDummies, True]

We now define which indices we are going to use. All summation indices will have the form i[1], i[2], etc. That is, they all have Head i.

In[3] := DefineDummy[i]

For convenience, we define an abbreviation for the summation range.

In[4] := dimension[i] = M;

Now we define the superfields we are going to play with. In the case at hand we can define M pairs of chiral-antichiral fermionic superfields F_i, \bar{F}_j through two Mathematica statements:

 $In[5] := ChiralFermionic[F[_]]$

In[6] := AChiralFermionic[FB[_]]

Using indices the SOPEs between superfields F_i and \bar{F}^j can be defined mostly as they are defined in the textbooks:

In[7] := OPE[F[i_],FB[j_]] :=

Delta[i,j] MakeOPE[$\{-1/2 \ t12**tb12**One, \ One\}$] Now we will introduce a composite superfield J(Z) which should satisfy the N = 2 SCA eq. (7):

$$J = a_1 \sum_i : F_i \overline{F}^i : +a_2 \overline{\mathcal{D}} F_1 + a_3 \mathcal{D} \overline{F}^1$$
(23)

where besides the standard term $F_i \overline{F}^i$ we also introduced the N = 2 analog of a Feigin-Fuchs term. In *Mathematica* this becomes:

Note the NewDummies statement and the assignment using ':='. Together, they make sure that everytime J is used, the summation indices will have a new number.

After these definitions we are ready to do real calculations. For example, to check for which values of parameters a_1, a_2, a_3 the superfield J (23) spans N = 2 SCA, let us calculate the SOPE J with J and subtract the known result (7):

where DummySimplify renumbers the dummy indices to get a unique form⁴ For brevity, we omitted a number of terms (all proportional to $a_1 - 1$) in the output and replaced them by an ellipsis. As can be immediately seen, the coefficient a_1 must be equal to 1, while a_2, a_3 are arbitrary. Let us note that our result Out[9] also gives the expression for the central charge of the N = 2 SCA:

$$c = 4M - 8a_2a_3 \tag{24}$$

Thus we reproduced the standard Miura realisation for the N = 2 SCA.

We hope that these simple examples will help users to use the *SOPEN2defs* package in more complicated calculations.

6 How to get it, and the future

If you are interested in *SOPEN2defs*, you can get it by Email from the authors. You'll also find it at http://euclid.tp.ph.ic.ac.uk/~krthie/ or via anonymous ftp at euclid.tp.ph.ic.ac.uk. Please put a reference to this paper in your paper when you use it. Questions, remarks and improvements are welcome. The package will be extended to other numbers of supersymmetries N.

Acknowledgements

K.T. thanks the JINR for the pleasant welcome during the weeks when this work was started. He also acknowledges the Instituut voor Theoretische Fysica from the KU Leuven (Belgium) where the first version of the package was completed. S.K. wishes to thank Imperial College for the hospitality in the final stage of this work. We also thank Lucy Wenham for improving our English.

References

- A.A. Belavin, A.M. Polyakov, A.B. Zamolodchikov, Nucl. Phys. B241 (1984) 333.
- [2] S. Wolfram, Mathematica, A System for Doing Mathematics by Computer, (Addison-Wesley Publishing Company, Inc.) 1991.
- [3] K. Thielemans, Int. J. Mod. Phys. C Vol. 2, No. 3 (1991) 787.
- [4] K. Thielemans, An Algorithmic Approach to Operator Product Expansions, W-algebras and W-strings, PhD thesis KU Leuven, June 1994, hep-th/9506159.

⁴In fact, the result of DummySimplify is not unique when tensors with certain symmetries appear, e.g. $J^{ij} = J^{ji}$.

- [5] E. Ivanov, S. Krivonos, A. Sorin, Mod. Phys. Lett. A10 (1995) 2439.
- [6] K. Schoutens, Nucl. Phys. **B295**[FS21] (1988) 634.
- [7] M. Ademollo, L. Brink, A. D'Adda, R. D'Auria, E. Napolitano, S. Sciutto, E. Del Giudice, P. Di Vecchia, S. Ferrara, F. Gliozzi, R. Musto, R. Pettorino, Nucl. Phys. B111 (1976) 77.