

# A QFT approach to $W_{1+\infty}$

B. N. Bakalov \*

*Differential Equations, Department of Mathematics and Informatics,  
Sofia University, Bld. J. Bourchier 5, BG-1126 Sofia, Bulgaria*

L. S. Georgiev †

*Institute for Nuclear Research and Nuclear Energy,  
Bulgarian Academy of Sciences,  
Tsarigradsko Chaussee 72, BG-1784 Sofia, Bulgaria*

I. T. Todorov ‡§

*Department of Mathematics, Massachusetts Institute of Technology,  
Cambridge, MA 02139-4307, USA*

## Abstract

$W_{1+\infty}$  is defined as an infinite dimensional Lie algebra spanned by the unit operator and the Laurent modes of a series of local quasiprimary chiral fields  $V^l(z)$  of dimension  $l+1$  ( $l = 0, 1, 2, \dots$ ). These fields are neutral with respect to the  $u(1)$  current  $J(z) = V^0(z)$ ; as a result the  $(l+2)$ -fold commutator of  $J$  with  $V^l$  vanishes. We outline a construction of rational conformal field theories with stress energy tensor  $T(z) = V^1(z)$  whose chiral algebras include all  $V^l$ 's. It is pointed out that earlier work on local extensions of the  $u(1)$  current algebra solves the problem of classifying all such theories for Virasoro central charge  $c = 1$ .

---

\*E-mail address: bbakalov@fmi.uni-sofia.bg

†E-mail address: lgeorg@bgearn.acad.bg

‡E-mail address: todorov@math.mit.edu

§On leave from the Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Tsarigradsko Chaussee 72, BG-1784 Sofia, Bulgaria.

# Introduction

The infinite dimensional Lie algebra  $W_{1+\infty}$  has emerged in various contexts. It was first introduced as the non-trivial central extension  $\widehat{\mathcal{D}}$  of the Lie algebra of differential operators on the circle [KP]. The theory of its quasifinite highest weight representations was developed by Kac, Radul et al. [KR, FKRW] (see also [AFMO1, AFMO2], the latter reference containing a review and a bibliography on the subject). Among the many applications we should like to single those to the quantum Hall effect [CTZ1, CTZ2] and to the bispectral problem (for a recent development and further references see [BHY]).

The present paper starts with a quantum field theoretic (QFT) characterization of a larger algebra,  $\widehat{gl}$ , in terms of a basic bilocal field  $V(z_1, z_2)$ .  $W_{1+\infty}$  is spanned by the local quasiprimary fields  $V^l$  entering the expansion of  $V$  around  $z_1 = z_2$ . We outline a program for constructing rational conformal field theory (RCFT) extensions of the chiral  $W_{1+\infty}$  algebra and point out that this program has been carried out in [BMT] and [PT] for Virasoro central charge  $c = 1$  within the study of local extensions of the  $u(1)$  current algebra.

## 1 Chiral algebra generated by a bilocal “dipole field”

We shall begin by describing the QFT counterpart of  $\widehat{gl}$ , the central extension of the Lie algebra of infinite matrices with finitely many non-zero diagonals – see [FKRW], Sec. 2. Our starting point will be a chiral algebra  $\mathcal{A}_c = \mathcal{A}_c(1, -1)$  generated by a bilocal field  $V(z_1, z_2)$  satisfying the commutation relations (CR)

$$\left[ V(z_1, z_2), V(z_3, z_4) \right] = \left\{ V(z_1, z_4) + \frac{c}{z_{14}} \right\} \delta(z_{23}) - \left\{ V(z_3, z_2) + \frac{c}{z_{32}} \right\} \delta(z_{14}) \quad (1.1)$$

where  $z_{ij} = z_i - z_j$  and the complex variable  $\delta$ -function can be defined through its Laurent expansion (cf. [K])

$$\delta(z_{12}) = \frac{1}{z_{12}} + \frac{1}{z_{21}}, \quad \frac{1}{z_{ij}} = \sum_{n=0}^{\infty} \frac{z_j^n}{z_i^{n+1}} \quad (1.2)$$

and is characterized by the properties

$$z_{12}\delta(z_{12}) = 0, \quad \oint_{|z_1|=|z_2|} f(z_1)\delta(z_{12})\frac{dz_1}{2\pi i} = f(z_2). \quad (1.3)$$

The *central charge*  $c$  ( $[c, V(z_1, z_2)] = 0$ ) will be regarded as a real number that labels our chiral algebra (along with the pair  $\{1, -1\}$  of opposite charges to be identified shortly). The inhomogeneous term

$$\Psi(z_1, z_2; z_3, z_4) = \frac{c}{z_{14}}\delta(z_{23}) - \frac{c}{z_{32}}\delta(z_{14})$$

in (1.1) satisfies the 2-cocycle condition

$$\begin{aligned} & \delta(z_{23})\Psi(z_1, z_4; z_5, z_6) - \delta(z_{14})\Psi(z_3, z_2; z_5, z_6) + \delta(z_{25})\Psi(z_3, z_4; z_1, z_6) - \\ & \delta(z_{16})\Psi(z_3, z_4; z_5, z_2) + \delta(z_{36})\Psi(z_5, z_4; z_1, z_2) - \delta(z_{45})\Psi(z_1, z_2; z_3, z_6) = 0 \end{aligned}$$

which guarantees the Jacobi identity for double commutators. In order to display the connection of the above Lie algebra structure with  $\widehat{gl}$  it suffices to insert the double Laurent expansion of  $V$

$$V(z_1, z_2) = \sum_{i, j \in \mathbb{Z}} E_{ij} z_1^i z_2^{-j-1} \quad (1.4)$$

in (1.1) and compute the resulting CR for the Weyl matrices  $E_{ij}$ :

$$\begin{aligned} [E_{ij}, E_{kl}] &= \delta_{jk} E_{il} - \delta_{il} E_{kj} + c \{ \theta(j) - \theta(i) \} \delta_{il} \delta_{jk}, \\ \theta(i) &= 1 \quad \text{for } i > 0, \quad \theta(i) + \theta(-i) = 1. \end{aligned} \quad (1.5)$$

The CR (1.1) have a well defined limit when we set, after a possible differentiation, the arguments of  $V$  equal to each other. The resulting local fields will be viewed as elements of the same chiral algebra  $\mathcal{A}_c$ . The first two of them, the  $u(1)$  current

$$J(z) = V(z, z) = \sum_{n \in \mathbb{Z}} J_n z^{-n-1} \quad (1.6)$$

and the (chiral) stress energy tensor

$$T(z) = \frac{1}{2} \left\{ (\partial_1 - \partial_2) V(z_1, z_2) \right\} \Big|_{z_1=z_2=z} = \sum_{n \in \mathbb{Z}} L_n z^{-n-2} \quad (1.7)$$

$(\partial_i = \frac{\partial}{\partial z_i})$  generate a Lie subalgebra of  $\widehat{gl}$ . Inserting (1.6) into (1.1) we find

$$\left[ V(z_1, z_2), J(z_3) \right] = \left\{ V(z_1, z_2) + \frac{c}{z_{12}} \right\} \left( \delta(z_{23}) - \delta(z_{13}) \right) \quad (1.8)$$

$$\left[ V(z_1, z_2), J_n \right] = \left\{ V(z_1, z_2) + \frac{c}{z_{12}} \right\} (z_2^n - z_1^n). \quad (1.9)$$

These relations justify the  $\{1, -1\}$  charge assignment for  $V$  (and the term “*dipole field*”). In the limit  $z_2 \rightarrow z_1$  we recover the  $u(1)$  current algebra:

$$\left[ J(z_1), J(z_2) \right] = -c\delta'(z_{12}) \iff \left[ J_m, J_n \right] = cm\delta_{m,-n}. \quad (1.10)$$

Similarly, (1.1) and (1.7) imply

$$\begin{aligned} \left[ V(z_1, z_2), T(z_3) \right] &= \left\{ \frac{c}{z_{12}^2} - \partial_1 V(z_1, z_2) \right\} \delta(z_{13}) - \left\{ \frac{c}{z_{12}^2} + \partial_2 V(z_1, z_2) \right\} \delta(z_{23}) \\ &\quad - \frac{1}{2} \left\{ V(z_1, z_2) + \frac{c}{z_{12}} \right\} \left\{ \delta'(z_{13}) + \delta'(z_{23}) \right\}, \end{aligned} \quad (1.11)$$

$$\begin{aligned} \left[ L_n, V(z_1, z_2) \right] &= \left\{ z_1^n \left( D_1 + \frac{n+1}{2} \right) + z_2^n \left( D_2 + \frac{n+1}{2} \right) \right\} V(z_1, z_2) \\ &\quad + cd_n(z_1, z_2), \quad D_i = z_i \partial_i; \end{aligned} \quad (1.12)$$

the cocycle  $d_n(z_1, z_2)$  ( $= -d_n(z_2, z_1)$ ) is homogeneous of degree  $n - 1$  in its arguments:

$$d_n(z_1, z_2) = \frac{1}{2z_{12}} \sum_{i=1}^{n-1} (z_1^i - z_2^i)(z_1^{n-i} - z_2^{n-i}) \quad \text{for } n \geq 2, \quad (1.13)$$

$$d_{-n}(z_1, z_2) = \frac{1}{z_1 z_2} d_n\left(\frac{1}{z_1}, \frac{1}{z_2}\right) \quad (n \geq 2), \quad d_n = 0 \text{ for } n = 0, \pm 1. \quad (1.14)$$

The  $L_n$  generate the Virasoro algebra:

$$\left[ L_n, L_m \right] = (n - m)L_{n+m} + c \frac{n^3 - n}{12} \delta_{n,-m}. \quad (1.15)$$

The vanishing of the cocycle  $d_n$  for  $n = 0, \pm 1$  can be interpreted by saying that  $V$  is a *quasiprimary bilocal field*. We recall that  $V^l(z)$  is a local quasiprimary field of (conformal) dimension  $l + 1$  provided

$$\left[ L_n, V^l(z) \right] = z^n \left( D + (n + 1)(l + 1) \right) V^l(z) \quad \text{for } n = 0, \pm 1. \quad (1.16)$$

( $V^l$  is called *primary* if (1.16) holds for all  $n \in \mathbb{Z}$ .)

The importance of the concept of a quasiprimary field stems from the fact that we shall be dealing, to start with, with the *vacuum representation* of the chiral field algebra in which the lowest weight (*vacuum*) *vector*  $|0\rangle$ , defined by

$$\begin{aligned} E_{ij}|0\rangle &= 0 \quad \text{for } i \leq j \\ \left( \Rightarrow J_n|0\rangle &= 0 \text{ for } n \geq 0, \quad L_n|0\rangle = 0 \text{ for } n \geq -1 \right) \end{aligned} \quad (1.17)$$

is *Möbius* or  $\text{su}(1,1)$  *invariant* (it is annihilated by the *conformal energy operator*  $L_0$  and by  $L_{\pm 1}$ ). Möbius invariance determines local (2- and) 3-point functions of quasiprimary fields (up to a normalization) – see e.g. [FST]. Equations (1.1), (1.4) and (1.17) allow to write down the correlation functions of the basic bilocal field  $V$ . We have, in particular,

$$\langle 0|V(z_1, z_2)V(z_3, z_4)|0\rangle = \frac{c}{z_{14}z_{23}} \quad (1.18)$$

$$\langle 0|V(z_1, z_2)V(z_3, z_4)V(z_5, z_6)|0\rangle = \frac{c}{z_{16}z_{23}z_{45}} - \frac{c}{z_{14}z_{25}z_{36}}. \quad (1.19)$$

A realization of the above algebra for a positive integer  $c$  is provided by the composite field (cf. [BPRSS])

$$V_{\psi^*\psi}(z_1, z_2) = \sum_{i=1}^c :\psi_i^*(z_1)\psi_i(z_2): \quad (1.20)$$

where  $\psi_i$  and their conjugates are free Fermi fields satisfying the canonical anticommutation relations

$$\begin{aligned} \left[ \psi_i(z_1), \psi_j(z_2) \right]_+ &= 0 = \left[ \psi_i^*(z_1), \psi_j^*(z_2) \right]_+ \\ \left[ \psi_i(z_1), \psi_j^*(z_2) \right]_+ &= \delta_{ij}\delta(z_{12}) \end{aligned} \quad (1.21)$$

or, in Fourier modes,

$$\left[ \psi_{i\mu}, \psi_{j\nu}^* \right]_+ = \delta_{\mu,-\nu}\delta_{ij}. \quad (1.22)$$

In the vacuum (Neveu–Schwarz) sector  $\psi^{(*)}(z)$  is expanded in integer powers of  $z$ :

$$\psi^{(*)}(z) = \sum_{n \in \mathbb{Z}} \psi_{-n-\frac{1}{2}}^{(*)} z^n.$$

The *normal product*  $::$  in (1.20) is defined by either subtracting the vacuum expectation value,

$$:\psi^*(z_1)\psi(z_2): = \psi^*(z_1)\psi(z_2) - \frac{1}{z_{12}},$$

or, equivalently, by ordering the Fourier modes:

$$:\psi_\mu^*\psi_\nu: = \begin{cases} \psi_\mu^*\psi_\nu & \text{for } \nu > 0 \\ -\psi_\nu\psi_\mu^* & \text{for } \nu < 0. \end{cases}$$

The main goal of this section is to establish an expansion formula for  $V(z_1, z_2)$  in terms of a basis  $V^l(z)$  of local quasiprimary fields (of conformal dimension  $l + 1$ ) that is used in [CTZ1, CTZ2] for characterizing the  $W_{1+\infty}$  algebra.

**Theorem 1.1** *Under the above assumptions the bilocal field  $V$  admits the following expansion in local quasiprimary fields*

$$V(z_1, z_2) = \sum_{l=0}^{\infty} \frac{(2l+1)!}{(l!)^3} \int_{z_2}^{z_1} \frac{(z_1 - z)^l (z - z_2)^l}{z_{12}^{l+1}} V^l(z) dz \quad (1.23)$$

where  $V^l$  satisfy (1.16) and the orthonormalization condition

$$\langle 0|V^l(z_1)V^{l'}(z_2)|0\rangle = c \frac{(l!)^4}{(2l)!} z_{12}^{-2l-2} \delta_{ll'}. \quad (1.24)$$

As a consequence of (1.24) the 3-point function of  $V$  and  $V^l$  is computed to be

$$\langle 0|V(z_1, z_2)V^l(z_3)|0\rangle = l! \binom{2l}{l}^{-1} \frac{cz_{12}^l}{(z_{13}z_{23})^{l+1}}. \quad (1.25)$$

Conversely,  $V^l$  is computed in terms of  $V$ :

$$V^l(z) = \binom{2l}{l}^{-1} \lim_{z_1, z_2 \rightarrow z} \sum_{k=0}^l \binom{l}{k}^2 \partial_1^k (-\partial_2)^{l-k} V(z_1, z_2) \quad (1.26)$$

$$= \frac{l!}{(2l)!} \lim_{z_1, z_2 \rightarrow z} \partial_1^l (-\partial_2)^l \left\{ z_{12}^l V(z_1, z_2) \right\}. \quad (1.27)$$

Defining the Fourier modes of  $V^l$  by the expansion formula

$$V^l(z) = \sum_{n \in \mathbb{Z}} V_n^l z^{-n-l-1} \quad (1.28)$$

one deduces that  $V_n^l$  satisfy CR of the form

$$\begin{aligned} [V_m^k, V_n^l] &= (lm - kn)V_{m+n}^{l+k-1} + \sum_{\nu=1}^{\lfloor \frac{l+k-1}{2} \rfloor} p_\nu(k, m, l, n) V_{m+n}^{l+k-1-2\nu} \\ &\quad + c \frac{(l!)^4}{(2l)!} \binom{l+m}{m-l-1} \delta_{m,-n} \delta_{kl} \end{aligned} \quad (1.29)$$

where  $p_\nu$  are polynomials in their arguments.

**Proof.** We first check the consistency among (1.23), (1.24) and (1.25). To this end we take the matrix element of both sides of (1.23) between  $V^l(z_3)|0\rangle$  and  $\langle 0|$  using (1.24) and the integral formula

$$\frac{(2l+1)!}{(l!)^2} \int_{z_2}^{z_1} \frac{(z_1 - z)^l (z - z_2)^l}{z_{12}^{l+1} (z - z_3)^{2l+2}} dz = \frac{z_{12}^l}{(z_{13}z_{23})^{l+1}}. \quad (1.30)$$

This expression coincides with the unique up to a normalization Möbius invariant 3-point function. Conversely, using the uniqueness of the solution of the corresponding momenta problem we deduce the expression for the kernel multiplying  $V^l(z)$  in the integrand of (1.23). The numerical coefficients are then computed from the requirement that the expansion (1.23) with  $V^l$  satisfying (1.25) reproduces the 4-point function (1.18). The latter property is a consequence of the identity

$$\frac{1}{1-\eta} = 1 + \sum_{l=1}^{\infty} \binom{2l-2}{l-1}^{-1} \eta^l F(l, l; 2l; \eta) \quad (1.31)$$

where the Gauss hypergeometric function is identified from the integral representation

$$\begin{aligned} F(l, l; 2l; \eta) &= \frac{(2l-1)!}{[(l-1)!]^2} \frac{(z_{13}z_{24})^l}{z_{34}^{2l-1}} \int_{z_4}^{z_3} \frac{(z_3-z)^{l-1}(z-z_4)^{l-1}}{(z_1-z)^l(z_2-z)^l} dz \\ &= \frac{1}{B(l, l)} \int_0^1 \frac{t^{l-1}(1-t)^{l-1}}{(1-\eta t)^l} dt, \quad \left( B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} \right), \end{aligned} \quad (1.32)$$

$t$  and  $\eta$  being the cross-ratios

$$t = \frac{(z-z_4)z_{13}}{(z_1-z)z_{34}}, \quad \eta = \frac{z_{12}z_{34}}{z_{13}z_{24}}. \quad (1.33)$$

The expression (1.26) for  $V^l$  (apart from the normalization factor) can be computed in two ways yielding the same answer. First, it follows from  $L_1$  covariance (being indeed a special case of eqs. (3.68–71) of [FST]). Indeed, setting

$$V^l(z) = \lim_{z_{1,2} \rightarrow z} \mathcal{D}_l(\partial_1, \partial_2) V(z_1, z_2), \quad (1.34)$$

where  $\mathcal{D}_l(\alpha, \beta)$  is an yet unknown homogeneous polynomial of degree  $l$  of its arguments, we find from (1.12)

$$\left[ L_1, V^l(z) \right] = \lim_{z_{1,2} \rightarrow z} \mathcal{D}_l(\partial_1, \partial_2) \left\{ z_1(z_1\partial_1 + 1) + z_2(z_2\partial_2 + 1) \right\} V(z_1, z_2)$$

which agrees with (1.16) for  $n=1$  iff  $\mathcal{D}_l$  satisfies the differential equation

$$\left( \alpha \frac{\partial^2}{\partial \alpha^2} + \beta \frac{\partial^2}{\partial \beta^2} + \frac{\partial}{\partial \alpha} + \frac{\partial}{\partial \beta} \right) \mathcal{D}_l(\alpha, \beta) = 0 \quad (1.35)$$

which together with (homogeneity and) the normalization condition

$$\mathcal{D}_l(1, -1) = 1 \quad (1.36)$$

yields the operator applied to  $V(z_1, z_2)$  in the right hand side of (1.26). Another way to derive (1.26) is to use the equivalence of (1.26) and (1.27), to apply the operator  $(\partial_1^n (-\partial_2)^n z_{12}^n)$  to both sides of (1.23) and to prove that only the term with  $l = n$  does not vanish in the limit  $z_1 = z_2$ .

We shall use a roundabout road to compute the coefficient to the first term in the right hand side of (1.29) introducing on the way the *Taylor expansion* of  $V(z_1, z_2)$  around the second point

$$V(z_1, z_2) = \sum_{l=0}^{\infty} \frac{z_{12}^l}{l!} J^l(z_2), \quad J^l(z) = \partial_1^l V(z_1, z_2) \Big|_{z_1=z_2=z}. \quad (1.37)$$

**Lemma 1.2** *The fields  $J^l$  satisfy the local CR*

$$\begin{aligned} [J^k(z_1), J^l(z_2)] &= \sum_{\nu \geq 1} \left\{ (-1)^\nu \binom{l}{\nu} J^{k+l-\nu}(z_2) - \binom{k}{\nu} J^{k+l-\nu}(z_1) \right\} \delta^{(\nu)}(z_{12}) \\ &+ c(-1)^{l+1} \frac{k!l!}{(k+l+1)!} \delta^{(k+l+1)}(z_{12}) \end{aligned} \quad (1.38)$$

or, in Fourier–Laurent modes, for

$$J^l(z) = \sum_{n \in \mathbb{Z}} J_n^l z^{-l-n-1},$$

$$\begin{aligned} [J_m^k, J_n^l] &= (lm - kn) J_{m+n}^{l+k-1} \\ &+ \left\{ l(l-1)m(m+2k-1) - k(k-1)n(n+2l-1) \right\} J_{m+n}^{l+k-2} \\ &+ \dots + (-1)^k c k!l! \binom{k+m}{m-l-1} \delta_{m,-n}. \end{aligned} \quad (1.39)$$

The **proof of the lemma** consists in a straightforward computation starting from (1.1) and (1.37) and using identities of the type

$$\left[ \partial_2^l \delta(z_{12}) \right] \frac{z_{12}^{l-n}}{(l-n)!} = (-1)^n \binom{l}{n} \delta^{(n)}(z_{12}), \quad \delta^{(m)}(z_{12}) z_{12}^n = 0 \quad \text{for } n > m. \quad (1.40)$$



**Lemma 1.3** *The quasiprimary fields  $V^l$  are expressed in terms of  $J^l$  and their derivatives by*

$$V^l(z) = \binom{2l}{l}^{-1} \sum_{k=0}^l (-1)^k \binom{l}{k} \binom{2l-k}{l} \partial^k J^{l-k}(z). \quad (1.41)$$

To **prove** (1.41) we express  $-\partial_2$  in (1.27) as  $\partial_1 - (\partial_1 + \partial_2)$ .

The term  $(lm - kn)V_{m+n}^{l+k-1}$  in the right hand side of (1.29) is obtained from the corresponding term in the commutator of  $J$  modes (1.39) by noting that the coefficient to  $J^l$  in the expansion (1.41) is 1. This completes the proof of Theorem 1.1.

There are two obvious advantages in computing correlation functions in terms of quasiprimary fields: the orthogonality relation (1.24) (which has no match for  $J^l$  fields) and the possibility to use Möbius invariance which determines the 3-point functions. The spectrum of the Cartan subalgebra is, however, easier to write down in terms of representatives of the differential operators  $D^l$  (see [KR] and Sec. 2 below).

## 2 The subalgebra $W^{(c)}$ of $\mathcal{A}_c$ and its QFT representations

Let  $W^{(c)}$  be the associative subalgebra of  $\mathcal{A}_c$  spanned by (the vacuum representation of) finite linear combinations of the quasiprimary fields  $V^l$ . It can be viewed as a covering of the (vacuum representation of)  $W_{1+\infty}$ . The CR (1.29) have an important corollary which characterizes  $W^{(c)}$ .

**Proposition 2.1** *The  $(l+1)$ -fold commutator of current components  $J_m = V_m^0$  with  $V^l(z)$  is a  $c$ -number:*

$$\left[ J_{m_0}, \left[ J_{m_1}, \dots \left[ J_{m_l}, V^l(z) \right] \dots \right] \right] = cl! \prod_{\nu=0}^l (m_\nu z^{m_\nu-1}). \quad (2.1)$$

It follows that *the derivative (commutator) action of the Heisenberg algebra generated by  $J_m$  is nilpotent on finite linear combinations of  $V^l$* . This motivates the following **definition**. *An element  $A \in \mathcal{A}_c$  is said to belong to  $W^{(c)}$  iff there exists a positive integer  $N = N(A)$  such that for any choice of the indices  $m_1, \dots, m_N \in \mathbb{Z}$  the  $N$ -fold commutator  $[J_{m_1}, \dots [J_{m_N}, A] \dots]$  vanishes.*

It follows that the associative local chiral algebra corresponding to  $W_{1+\infty}$  is generated by the quasiprimary fields  $V^l$ . We proceed to summarizing the

results of [KR, FKRW] on the unitary *lowest weight modules*  $\mathcal{L}_c$  of  $\widehat{\mathcal{D}}$ . Denote by  $W(z^m f(D))$  the image of the differential operator  $z^m f(D)$  in  $\text{End } \mathcal{L}_c$ . We shall also use this notation for quasipolynomials  $f$  (i.e., for polynomials of  $D$  and of  $e^{\lambda D}$ ). The basic CR can be written in a compact form [KR]

$$\begin{aligned} & \left[ W(z^m e^{xD}), W(z^n e^{yD}) \right] = \\ & (e^{nx} - e^{my}) W(z^{m+n} e^{(x+y)D}) - c \frac{e^{nx} - e^{my}}{e^{x+y} - 1} \delta_{m,-n}. \end{aligned} \quad (2.2)$$

The positive energy unitary irreducible representations of  $\widehat{\mathcal{D}} = W_{1+\infty}$  correspond to positive integers  $c$  and are labeled by  $c$  real charges  $\mathbf{r} = (r_1, \dots, r_c)$ . Let  $|\mathbf{r}\rangle$  be the corresponding minimal energy state satisfying

$$\{W(-D^l) - \lambda_l\}|\mathbf{r}\rangle = 0 = W(-z^m D^l)|\mathbf{r}\rangle \text{ for } m \geq 0. \quad (2.3)$$

The eigenvalues  $\lambda_l$  of Cartan operators are then recovered from the generating function [KR]

$$\Delta_{\mathbf{r}}(x) := \sum_{l=0}^{\infty} \lambda_l \frac{x^l}{l!} = \sum_{i=1}^c \frac{e^{r_i x} - 1}{e^x - 1}. \quad (2.4)$$

We have, in particular,

$$\lambda_0 = \sum_{i=1}^c r_i, \quad \lambda_1 = \frac{1}{2} \sum_{i=1}^c (r_i^2 - r_i), \quad \lambda_2 = \sum_{i=1}^c \left( \frac{r_i^3}{3} - \frac{r_i^2}{2} + \frac{r_i}{3} \right).$$

We should like, to begin with, to reexpress these results in terms of the quasiprimary fields  $V^l$  and their Laurent modes. This is done by using the modes  $J_n^l$  (appearing in (1.39)) as intermediary. We have, according to [KR],

$$J_n^l = -W(z^n [D]_l), \quad [D]_0 = 1, [D]_l = D(D-1)\dots(D-l+1). \quad (2.5)$$

Using the relation (1.41) between  $V^l$  and  $\{J^{l-k}\}_{k=0,\dots,l}$  we find

$$V_n^l = W(z^n f_{ln}(D)) \quad (2.6)$$

where  $f_{ln}$  are polynomials of degree  $l$ :

$$f_{ln}(D) = - \binom{2l}{l}^{-1} \sum_{\nu=0}^l (-1)^{l-\nu} \binom{l}{\nu}^2 [D]_{\nu} [-D-n-1]_{l-\nu} (= -D^l + \dots) \quad (2.7)$$

(we have noted that  $\sum \binom{l}{\nu}^2 = \binom{2l}{l}$ ).

The spectrum of the zero modes  $V_0^l$  can be computed independently starting from the expectation value of the dipole field  $V$  in the ground state  $|\mathbf{r}\rangle$ :

$$\langle \mathbf{r} | V(z, w) | \mathbf{r} \rangle = \frac{1}{z-w} \sum_{i=1}^c \left\{ \left( \frac{z}{w} \right)^{r_i} - 1 \right\}. \quad (2.8)$$

Inserting in it the expansion (1.23) we find

$$\sum_{i=1}^c \left\{ \left( \frac{z}{w} \right)^{r_i} - 1 \right\} = \sum_{l=0}^{\infty} \frac{1}{l!} V_l(\mathbf{r}) \left\{ \left( 1 - \frac{w}{z} \right)^{l+1} F\left( l+1, l+1; 2l+2; 1 - \frac{w}{z} \right) \right\} \quad (2.9)$$

( $F$  being the hypergeometric function (1.32)). The eigenvalues of the zero modes

$$V_l(\mathbf{r}) = \langle \mathbf{r} | V_0^l | \mathbf{r} \rangle \quad (\mathbf{r} = (r_1, \dots, r_c)) \quad (2.10)$$

are thus computed from the recurrence relation

$$\sum_{i=1}^c \frac{r_i(r_i+1)\dots(r_i+n)}{(n+1)!} = \sum_{l=0}^n \binom{n}{l} \frac{n!(2l+1)!}{(l!)^2(l+n+1)!} V_l(\mathbf{r}). \quad (2.11)$$

Another way to compute  $V_l(\mathbf{r})$  is to apply the operator  $(\partial_1^l (-\partial_2)^l z_{12}^l)$  to both sides of (2.8) and to use (1.27). The result is

$$V_0(\mathbf{r}) = \sum r_i \quad (= \lambda_0),$$

$$V_l(\mathbf{r}) = B(l, l+1) \sum_{k=0}^{l-1} \binom{l}{k} \binom{l}{k+1} \sum_{i=1}^c r_i \frac{\Gamma(r_i + l - k)}{\Gamma(r_i - k)}, \quad l \geq 1. \quad (2.12)$$

We find, in particular,

$$V_1(\mathbf{r}) = \frac{1}{2} \sum r_i^2 \quad (= \lambda_1 + \frac{1}{2} \lambda_0), \quad V_2(\mathbf{r}) = \frac{1}{3} \sum r_i^3 \quad (= \lambda_2 - \lambda_1 - \frac{1}{6} \lambda_0),$$

$$V_3(\mathbf{r}) = \frac{1}{4} \sum \left( r_i^4 + \frac{1}{5} r_i^2 \right), \quad V_4(\mathbf{r}) = \sum \left( \frac{1}{5} r_i^5 + \frac{1}{7} r_i^3 \right), \dots$$

$$V_l(-\mathbf{r}) = (-1)^{l+1} V_l(\mathbf{r}). \quad (2.13)$$

The first two eigenvalues giving the total charge and the energy of the ground state justify the use of the term ‘‘charges’’ for the representation labels  $r_i$ . For  $c \geq 1$  a representation of lowest weight  $\mathbf{r}$  is *degenerate* provided some of the differences  $r_i - r_j$  (for  $i \neq j$ ) are integers. It is *maximally degenerate* if all such differences are integers. We shall say that a (reducible) positive energy  $W^{(c)}$  module  $\mathcal{H}_c$  gives rise to a *QFT representation* of  $W_{1+\infty}$  if all its

irreducible positive energy submodules  $\mathcal{L}_c(\mathbf{r})$  are generated from the vacuum sector by  $W^{(c)}$ -primary chiral vertex operators  $\phi_{\mathbf{r}}(z)$  such that

$$\phi_{\mathbf{r}}(z)|0\rangle = e^{zL_{-1}}|\mathbf{r}\rangle \quad (2.14)$$

and if the resulting “field algebra” is closed under fusion. (For non-degenerate representations this means that the lowest weights form an abelian group: if  $|\mathbf{r}\rangle, |\mathbf{s}\rangle \in \mathcal{H}_c$  then also  $|\mathbf{r} + \mathbf{s}\rangle \in \mathcal{H}_c$ . The fusion rules for degenerate representations are also written down in [FKRW].)

### 3 RCFT extensions of $W^{(c)}$ . The case $c = 1$ .

We set the problem of classifying the local extensions of the chiral algebra  $W^{(c)}$  (with the same central charge  $c$ ) which admit a finite number of QFT representations  $\pi_\nu$  such that their specialized characters

$$\text{ch}_\nu(\tau) = \text{tr}_{\pi_\nu} q^{L_0 - \frac{c}{24}}, \quad q = e^{2\pi i\tau}, \quad \text{Im}\tau > 0 \quad (3.1)$$

span a modular (i.e.  $\text{SL}(2, \mathbb{Z})$ ) invariant space.

We shall present the solution to this problem for  $c = 1$  and shall end up with some remarks concerning the general case. We note that the algebra  $W^{(c)}$  with  $c > 1$  can be written as the tensor product of a  $W^{(1)}$  factor and a chiral algebra of central charge  $c - 1$  that involves no  $U(1)$  current in the degenerate case (see [CTZ2], Sec. 3). Thus the  $c = 1$  theory is a necessary ingredient for the solution of the general problem.

The Bose–Fermi ( $\mathbb{Z}_2$ -graded local) chiral algebras of  $c = 1$  containing  $W^{(1)}$  coincide with the (local or Fermi local) extensions of the  $\mathfrak{u}(1)$  current algebra classified in [BMT] and [PT]. Each such extension is generated by a pair of oppositely charged  $\widehat{\mathfrak{u}}(1)$  primary fields  $\psi(z, \pm g)$  with  $g^2 \in \mathbb{N}$  satisfying the  $\mathbb{Z}_2$ -graded (Bose–Fermi) locality

$$z_{12}^{g^2} \left\{ \psi(z_1, g)\psi(z_2, \pm g) - (-1)^{g^2} \psi(z_2, \pm g)\psi(z_1, g) \right\} = 0. \quad (3.2)$$

The condition that  $\psi$  is  $\widehat{\mathfrak{u}}(1)$  primary means that it satisfies the operator Ward identities

$$\left[ J(z_1), \psi(z_2, g) \right] = g\delta(z_{12})\psi(z_2, g) \quad (3.3)$$

$$\left[ T(z_1), \psi(z_2, g) \right] = -\Delta_g \delta'(z_{12})\psi(z_2, g) + \delta(z_{12})\psi'(z_2, g). \quad (3.4)$$

The compatibility of these relations, together with the Sugawara formula

$$T(z) = \frac{1}{2} :J(z)^2:, \quad (3.5)$$

which is a consequence of the  $W_{1+\infty}$  CR (1.29) for  $c = 1$ , yield the relation between charge and conformal dimension

$$\Delta_g = \frac{1}{2}g^2 \quad (\in \frac{1}{2}\mathbb{N}) \quad (3.6)$$

as well as the  $\widehat{u}(1)$  counterpart of the Knizhnik–Zamolodchikov equation [T]:

$$\psi'(z, g) = g:J(z)\psi(z, g): = g\left\{J_{(+)}(z)\psi(z, g) + \psi(z, g)J_{(-)}(z)\right\} \quad (3.7)$$

where

$$J_{(+)}(z) = \sum_{n=1}^{\infty} J_{-n}z^{n-1} = J(z) - J_{(-)}(z). \quad (3.8)$$

The QFT representations of the resulting extended algebra  $\mathcal{A}(g^2) \supset W^{(1)}$  which respect the  $\mathbb{Z}_2$ -graded locality of  $\psi(z, \pm g)$  are generated by a set of  $2g^2$  (multivalued, fractional spin) charge fields  $\phi(z, e_l)$  (including the unit operator) with charges

$$e_l = \frac{l}{2g}, \quad 1 - g^2 \leq l \leq g^2. \quad (3.9)$$

The relation

$$\psi(e^{2\pi i}z, g) = \psi(z, g)e^{2\pi i g J_0} \quad (3.10)$$

implies that  $\psi(z, \pm g)$  are single or double valued in the sector  $\mathcal{H}_l$  with lowest weight vector  $|e_l\rangle$  depending on the parity of  $l$  (since  $e^{2\pi i g e_l} = (-1)^l$ ). If  $\psi(z, \pm g)$  are Bose fields (i.e. if  $g^2$  is even) then  $\mathcal{H}_l$  with odd  $l$  define  $\mathbb{Z}_2$ -twisted sectors. The subset of even  $l$ 's gives rise to a modular invariant local theory in that case. The specialized characters (3.1) in the sectors  $\mathcal{H}_l$  are given by [PT]

$$K_l(\tau, g^2) = \text{tr}_{\mathcal{H}_l} q^{L_0 - \frac{1}{24}} = \frac{1}{\eta(\tau)} \sum_n q^{\frac{1}{2}(ng + \frac{l}{2g})^2} \quad (3.11)$$

where  $\eta(\tau)$  is the Dedekind function

$$\eta(\tau) = q^{\frac{1}{24}} \prod_{\nu=1}^{\infty} (1 - q^\nu) \quad (3.12)$$

and the sum in (3.11) runs over all integers ( $n \in \mathbb{Z}$ ). They span an  $\text{SL}(2, \mathbb{Z})$  invariant space [PT] (for a review – see [FST], Sec. 7). We find as special cases the extensions by a pair of free Fermi fields (satisfying (1.21)) for  $g^2 = 1$ ,

and the level 1  $\mathfrak{su}(2)$  current algebra for  $g^2 = 2$ . In both cases the resulting RCFT has two (untwisted) sectors.

If  $g^2$  is not square free,  $g^2 = k^2 g_0^2$  ( $k, g_0^2 \in \mathbb{N}$ ) then the chiral algebra can be further extended,  $\mathcal{A}(g^2) \subset \mathcal{A}(g_0^2)$ . If we define a  $\mathbb{Z}_k$ -automorphism group of  $\mathcal{A}(g_0^2)$  generated by

$$\zeta^{\frac{J_0}{g_0}} \psi(z, \pm g_0) \zeta^{-\frac{J_0}{g_0}} = \zeta^{\pm 1} \psi(z, \pm g_0), \quad \zeta^k = 1 \quad (3.13)$$

then  $\mathcal{A}(g^2)$  appears as the subalgebra of  $\mathbb{Z}_k$ -invariant elements of  $\mathcal{A}(g_0^2)$  and the corresponding RCFT is named a  $\mathbb{Z}_k$ -orbifold theory of  $\mathcal{A}(g_0^2)$ .

There are known local extensions of  $W^{(c)}$  for arbitrary positive integer  $c$ . These are:

(1) The  $\mathbb{Z}_2$ -graded algebra  $\mathcal{A}_c(1, -1)$  generated by  $c$  pairs of free Fermi fields (see (1.20), (1.21)) or the associated bosonic subalgebra generated by a pair of charge  $\pm 2$  fields and by the level 1  $\mathfrak{so}(2c)$  currents.

(2) Tensor products of  $\widehat{\mathfrak{su}}(c)_1$  theories with any of the above  $\mathcal{A}(g^2)$  algebras.

The (most degenerate) minimal  $W_{1+\infty}$  models can be embedded in a number of rational orbifold theories corresponding to finite subgroups of  $(\mathrm{SU}(c) \subset \mathrm{SO}(2c))$  (cf. [DV<sup>3</sup>]) that are currently under study [KT].

This paper was written during the stay of one of the authors, I.T., at the Department of Mathematics, M.I.T., with the support of a Fulbright grant 19684. I.T. thanks Victor Kac for hospitality and discussions. The work has been supported in part by the Bulgarian Foundation for Scientific Research under contract F-404.

## References

- [AFMO1] H. Awata, M. Fukuma, Y. Matsuo, S. Odake, Phys. Lett. **B332** (1994) 336–344; Commun. Math. Phys. **170** (1995) 151–179; see also H. Awata et al., Lett. Math. Phys. **31** (1994) 289–298.
- [AFMO2] H. Awata, M. Fukuma, Y. Matsuo, S. Odake, *Representation theory of the  $W_{1+\infty}$  algebra*, Kyoto preprint, hep-th/9408158.
- [BHY] B. Bakalov, E. Horozov, M. Yakimov, *Tau-functions as highest weight vectors for  $W_{1+\infty}$  algebra*, Sofia preprint, hep-th/9510211.
- [BMT] D. Buchholz, G. Mack, I. T. Todorov, Nucl. Phys. B (Proc. Suppl.) **5B** (1988) 20–56.

- [BPRSS] E. Bergshoeff, C. N. Pope, L. J. Romans, E. Sezgin, X. Shen, Phys. Lett. **B245** (1990) 447–452.
- [CTZ1] A. Cappelli, C. A. Trugenberger, G. R. Zemba, Nucl. Phys. **B396** (1993) 465–490; Phys. Rev. Lett. **72** (1994) 1902–1905; Nucl. Phys. **B448** [FS] (1995) 470–504, hep-th/9502021.
- [CTZ2] A. Cappelli, C. A. Trugenberger, G. R. Zemba,  $W_{1+\infty}$  *dynamics of edge excitations*, Torino preprint, cond-mat/9407095.
- [DV<sup>3</sup>] R. Dijkgraaf, C. Vafa, E. Verlinde, H. Verlinde, Commun. Math. Phys. **123** (1989) 485–526.
- [FKRW] E. Frenkel, V. Kac, A. Radul, W. Wang, Commun. Math. Phys. **170** (1995) 337–357, hep-th/9405121.
- [FST] P. Furlan, G. M. Sotkov, I. T. Todorov, Riv. Nuovo Cim. **12:6** (1989) 1–202.
- [K] V. G. Kac, *Vertex algebras*, Lecture at this Workshop.
- [KP] V. G. Kac, D. H. Peterson, Proc. Natl. Acad. Sci. USA **78** (1981) 3308–3312.
- [KR] V. Kac, A. Radul, Commun. Math. Phys. **157** (1993) 429–457, hep-th/9308153.
- [KT] V. G. Kac, I. T. Todorov, *Affine orbifolds and rational conformal field theory extensions of  $W_{1+\infty}$* , (to appear).
- [PT] R. R. Paunov, I. T. Todorov, Phys. Lett. **B196** (1987) 519–526.
- [T] I. T. Todorov, Phys. Lett. **B153** (1985) 77–81.