Effective quark Lagrangian in the instanton–gas model

Yu.A.Simonov

Institute of Theoretical and Experimental Physics 117259,Moscow , B.Cheremushkinskaya 25, Russia

March 6, 2018

Abstract

A straightforward derivation of the effective quark Lagrangian is presented for the topologically neutral chiral broken phase of the dilute instanton gas. The resulting quark Lagrangian is a nonlocal NJL – type and contains 4q, 6q,... vertices for any number of flavours. Correspondence of this result with previously known in literature is discussed in detail.

1 Introduction

The instanton gas model has a long history [1,2,3]. Recently lattice measurements of quark and gluon correlators in coordinate space [4] have revealed important contributions of instantons, which gives an additional impetus to the model [5].

On the theoretical side the instantons have been proved to solve the $U(1)$ problem [6], and to generate a specific effective quark Lagrangian (EQL) on a net topological charge [7].

In the topologically neutral system, such as the QCD vacuum, instantons have been proposed as a driving mechanism for the chiral symmetry breaking (CSB) [1,2,8]. The quantitative theory of light quarks (TLQ) in the instanton gas has been suggested in [9] and later elaborated in [10,11,12].

There are a considerable number of papers which use the method for quantitative predictions and comparison with experiment (as an example see [13]).

The essential ingredient of TLQ is the assumption of the dominant role of the quark zero modes on each instanton, and the resulting EQL is proposed in the form of the $2N_f \times 2N_f$ determinant of quark fields with coefficients (vertices) $M(p, p')$ expressed through quark zero modes.

As a result TLQ predicts CSB and the creation of chiral quark mass at all instanton densities; after bosonization the effective bosonic Lagrangian was derived [9, 10] containing expected Goldstone bosons.

In view of numerous successful applications we believe it is important to test the accuracy of the method.

In its modern form [12] the TLQ contains some approximations, first of all the zero-mode approximations (ZMA) introduced already in [9], and the basic role in TLQ is played by the ansatz for EQL [10-12] of the 'tHooft form [7]. The ZMA was questioned in [14], and it was argued there that nonzero modes play an important role in addition to zero modes.

In the present paper a new method is proposed, where all modes are accounted for automatically and we are avoiding standard approximations of TLQ.

In the next section we demonstrate a straightforward and exact way to derive the EQL in the topologically neutral dilute instanton gas. Our results differ from the existing in several points and lay ground for a new formalism of light quarks in the instanton gas.

First of all, our EQL contains a set of quark terms starting from 4q for any number of flavours. The 4q term differs from that of TLQ and is similar instead to the extended NJL type Lagrangian [15]. The structure of our EQL makes it possible to create chiral mass and CSB, and bosonization yields the Nambu-Goldstone boson Lagrangian.

All coefficients in our EQL are explicitly known and this enables one to do quantitative predictions. In particular the appearance of the 4q vertex for any N_f is important from the point of view of theory (such as effective top-condensate model of electroweak interactions [16]) and applications (OZI violation vertex, quark–pair creation, strangeness mixture).

In our formalism there are several ways to find the chiral quark mass entering the averaged quark propagator. One way exploits Dyson–Schwinger equations for our EQL in connection with the mean–field (Hartree) approximation. This is equivalent to the direct summing of leading trms in the $1/N_c$ expansion. In the one–flavour case one obtains an integral equation for the chiral mass, which coincides with the equation obtained earlier in [17] in a different, diagrammatic way. However, instead of the zero-mode approximation made in [17], one can reduce this equation to the one–dimensional integral one, subject to a simple numerical analysis.

This is done in section 3 of the paper. In case of two and more flavours one should use instead the bosonization method and find the chiral mass and chiral condensate from the steepest descent method, which will be a subject of the next paper.

Finally in the last section a detailed discussion is given of the correspondence of our method with TLQ.

2 Effective quark Lagrangians

We consider in this letter the neutral dilute gas of N_+ instantons I and $N_$ antiinstantons \overline{I} , $N_+ = N_- = N/2$, with the superposition principle [1-3]

$$
A_{\mu}(x) = \sum_{i=1}^{N} A_{\mu}^{(i)}(x, \gamma)
$$
 (1)

where γ denote the set of collective coordinates for each instanton $\gamma_i =$ $\{R^{(i)}, \Omega^{(i)}, \rho^{(i)}\}\.$ In what follows we for simplicity fix the instanton size $\rho^{(i)} =$ ρ and neglect perturbative gluon effects together with $II, I\overline{I}$ interaction, assuming that the latter stabilizes gas at small density.

Then the effective quark Lagrangian L_{EQL} can be defined as

$$
Z = \int d\gamma D\psi D\bar{\psi}e^{\sum_{f}^{N_{f}} \int \bar{\psi}_{f}S^{-1}\psi_{f}dx} =
$$

$$
= \int D\psi D\bar{\psi}e^{-\sum_{f}^{N_{f}} \int (\bar{\psi}_{f}S_{0}^{-1}\psi_{f})dx + \mathcal{L}_{EQL}(\psi,\bar{\psi})}
$$
(2)

where

$$
S^{-1} \equiv -i\hat{\partial} - im_f - g \sum_{i=1}^{N} \hat{A}^{(i)}(x, \gamma)
$$
 (3)

$$
S_0^{-1} = -i\hat{\partial} - im_f \tag{4}
$$

and

$$
d\gamma \equiv \prod_{i=1}^{N} (d\Omega_i \frac{d^4 R^{(i)}}{V_4})
$$
\n(5)

The resulting L_{EQL} is a polynomial of $\psi, \bar{\psi}$ of infinitely large power.

To obtain L_{EQL} one has to average in (2) over $d\gamma$ (5), γ entering into $A^{(i)}_{\mu}$ as

$$
A_{\mu}^{(i)}(x,\gamma) = \bar{\eta}_{a\mu\nu} \frac{(x - R^{(i)})_{\nu} \rho^2 \Omega_i^+ \tau_a \Omega_i}{(x - R^{(i)})^2 [(x - R^{(i)})^2 + \rho^2]}
$$
(6)

The straightforward use of the cluster expansion [18] for the averaging over $d\gamma$ yields

$$
Z = \int D\psi D\bar{\psi} e^{-\sum_{f}^{N_f} (\bar{\psi}_f S_0^{-1} \psi_f) dx + \sum_{i=1}^{N} \sum_{n=1}^{\infty} \frac{1}{n!} \ll \theta_i^n \gg} \tag{7}
$$

where we have denoted (f_i) -flavour indices).

$$
\theta_i = \sum_{f_k=1}^{N_f} \theta_i^{f_k}, \theta_i^{f_1} \equiv \int dx \psi_{f_1}^+(x) \hat{A}^{(i)} \psi_{f_1}(x); \tag{8}
$$

and angular brackets imply integration over $d\gamma$, while double brackets stand for cumulants [18] e.g.

$$
\ll \theta^2 \gg_{\Omega_i R_i} = \ll \theta^2 >_{\Omega_i R_i} - \ll \theta >_{\Omega_i R_i} \ll \theta >_{\Omega_i R_i}
$$
\n(9)

Let us look at $\ll \theta_i^n \gg$ more closely. The first term with $n=1$ vanishes due to the integration over $d\Omega_i$. The next term with $n = 2$ will be of most interest to us in what follows. This can be written in the form (see Appendix 1 for details of derivation) $\langle \theta_i^2 \rangle = \sum_{f_1, f_2} \langle \theta_i^{f_1} \theta_i^{f_2} \rangle$, and finally cast in the form,

$$
\sum_{i} < \theta_{i}^{2} > = \int \frac{dpdp_{1}dq}{(2\pi)^{12}} \Gamma(q) \{ (1 - \frac{1}{N_{c}^{2}}) \sum_{i,F} c_{i} J_{i}^{F}(p) J_{i}^{F}(p_{1}) + \n+ \frac{1}{2N_{f}} (1 - \frac{1}{N_{c}^{2}}) \sum_{i} c_{i} J_{i}^{0}(p) J_{i}^{0}(p_{1}) - \frac{1}{q^{2}} \sum_{i,F} c_{i}^{'} \tilde{J}_{i}^{F}(p) \tilde{J}_{i}^{F}(p_{1}) - \n- \frac{1}{2q^{2}N_{f}} \sum_{i} c_{i}^{'} \tilde{J}_{i}^{0}(p) \tilde{J}_{i}^{0}(p_{1}) \}
$$
\n(10)

where c_i, c'_i are given in Appendix 1, and

$$
J_i^n(p) = \psi^+(p)t^n 0_i \psi(p_1 - q), \ J_i^n(p_1) = \psi^+(p_1)t^n 0_i \psi(p + q),
$$

$$
\tilde{J}_i^n(p) = \psi^+(p)\hat{q}t^n 0_i \psi(p_1 - q), \ \tilde{J}_i^n(p_1) = \psi^+(p_1)\hat{q}t^n 0_i \psi(p + q)
$$
 (11)

and $n = F$ or 0, with $F = 1, ...N_f^2 - 1$, and $t^0 = \hat{1}, t^F$ is the flavour group generator.

We have also defined in (10)

$$
\Gamma(q) = \frac{4q^2\varphi^2(q)(4\pi^2\rho^2)^2N}{(N_c^2 - 1)V_4},\tag{12}
$$

and $\varphi(q)$ is the Fourier transform of the instanton field (6) given in Appendix 1, eq $(A.6)$.

The 4q interaction term (10) is of the extended NJL type, containing all possible nonlocal bilinears.

The advantage of (10) over the NJL Lagrangian is that it provides the in-built formfactor $\Gamma(q)$ which makes loop corrections finite since it behaves as

$$
\Gamma(q) \approx \frac{N}{V_4} \frac{(4\pi^2 \rho^2)^2}{(N_c^2 - 1)} \begin{cases} \frac{1}{q^2}, & q \to 0\\ \frac{16}{\rho^4 q^6}, & q \to \infty \end{cases}
$$
 (13)

One can see in (13) that the small q behaviour of $\Gamma(q)$ (which is due to the $1/x^3$ asymptotics of the instanton vector potential(6)) implies massless exchanges between currents $J(q)$ at large distances – a certainly unphysical feature, which is due to the ansatz (1) of noninteracting instanton gas. Taking into account the II and $I\bar{I}$ interactions changes the instanton profile at large distances (possible forms are suggested in [3]) and makes $\Gamma(q \to 0)$ finite in the realistic instanton gas or liquid model.

There are several important properties of the new EQL (7).

First of all, the sum over n is not limited, so therefore it contains vertices containing $4q,6q,...$ etc to infinity for any N_f . This differs from the standard ansatz of [9-12], where only the term with $2N_f$ quark operators appears.

One can observe in (8) the contribution of all quark modes, since one can rewrite θ^i as follows

$$
\theta_i = \int dx \psi^+(x) \hat{A}^i(x) \psi(x) = -\sum_{n,k} \langle \psi^+ | S_0^{-1} | u_n^i \rangle \epsilon_{nk}^i \langle u_k^i | S_0^{-1} | \psi \rangle \tag{14}
$$

with e.g. $\langle \psi^+|S_0^{-1}|u_n^i\rangle \equiv \int \psi^+(x)(-i\hat{\partial} - im)u_n^i(x)d^4x$. and ε_{nk}^i defined as $\varepsilon^i_{nk} = - < u^i_n|S_0\hat A^{(i)}S_0|u^i_k$ (15)

The standard TLQ prescription (9-12) obtains when one keeps in (14) only the zero mode coefficient, ε_{00}^{i} ,

Consider now all terms in (7) with $n \geq 2$ and take the limit $N_c \to \infty$ to perform the averaging over $dΩ$ explicitly.

The EQL (7) is additive for instantons, therefore we consider it for the *i*-th instanton and omit the subscript i in what follows. One has

$$
\mathcal{L}_i = \sum_{n=2}^{\infty} \frac{1}{n!} \ll \theta_i^n \gg, \quad \mathcal{L} = \sum_i \mathcal{L}_i \tag{16}
$$

The double brackets in (16) imply subracting from $\langle \theta^n \rangle$ disconnected pieces which appear while averaging in $d\Omega dR$.

Since each integration dR is accompanied by $1/V_4$ the second term in the difference,

$$
\langle \theta^n \rangle_R - \langle \theta^k \rangle_R \langle \theta^{n-k} \rangle_R, \tag{17}
$$

vanishes in the thermodynamical limit, $N \to \infty$, $V_4 \to \infty$, $\frac{N}{V_4}$ $\frac{N}{V_4} = const.$ Hence the only terms left in $\ll \theta^n \gg$ are of the form

$$
\langle \theta^n \rangle_{R,\Omega} = \langle \langle \theta^n \rangle_{\Omega} - \langle \theta^2 \rangle_{\Omega} \langle \theta^{n-2} \rangle_{\Omega} - \dots \rangle_{R} \tag{18}
$$

Now in averaging over $d\Omega$ one obtains in the leading terms of the $1/N_c$ expansion the set of disconnected (in color indices) products, which are subtracted in (18) and $(n-1)!$ equivalent connected terms, which differ only by permutation of indices. Thus one comes to the EQL defined in (16)

$$
\mathcal{L} = \sum_{n=2}^{\infty} \frac{N}{2nV_4 N_c^n} \prod_{k=1}^n \int \psi_{\alpha_k}^+(p_k) \gamma_{\mu_k} \psi_{\beta_k}(p_k - q_k) \frac{dp_k dq_k}{(2\pi)^8} \times
$$

$$
\times (2\pi)^4 \delta^4 (\sum_{i=1}^n q_i) tr\left(\prod_{k=1}^n A_{\mu_k}^I(q_k)\right) \prod_{i=1}^n \delta_{\alpha_i, \beta_{i-1}} + (I \to \bar{I})
$$
(19)

The EQL (7) and (19) is the central point of this letter and in the following section we use it to derive equations for the chiral mass.

3 Chiral mass, chiral symmetry breaking and the gap equation

The multiquark terms $\ll \theta^n \gg$ in the effective Lagrangian (7) contain all information about possible quark mass creation which automatically signifies CSB. The form of $\langle \theta^2 \rangle$ in (10) tells us that the averaged 4q interaction is of the extended NJL type [15] and is nonlocal.

Nevertheless one can use for the 4q term the same gap equation, derived from the Dyson–Schwinger method, as it was done in the NJL formalism [15]. However the essential difference of our expression (7) from the NJL Lagrangian is that the former contains all powers n , i.e. all n–quark effective vertices, $n = 4, 6, ... \infty$.

Therefore the gap equation, defining the creation of the chiral mass in our case, should contain in general this infinite number of vertices, unless one can find a small parameter suppressing terms with large n . To answer this question we shall consider first the case of one flavour and compute the averaged quark propagator in the instanton gas.

We shall derive it in the large N_c limit and shall obtain the integral equation for the chiral mass, containing contributions of all n–quark vertices.

At this point one can find from (19) the effective quark mass in the mean–field (Hartree) approximation. To this end one should identify in L (19) the structure $-i \int \frac{dp}{(2\pi)^4} \psi^+(p) M(p) \psi(p)$. In the leading order at large N_c one replaces in (19) $(n-1)$ colorless bifermion combinations by their vacuum expectation values, and in Hartree approximation by the effective quark propagator $S(p)$

$$
\bar{S}(p) = \frac{\hat{1}_c}{\hat{p} - iM(p)}\tag{20}
$$

Thus each of the colorless bifermions yields a factor N_c and we obtain

$$
M(p) = -i \frac{N}{2V_4 N_c} \sum_{n=2}^{\infty} \int \prod_{k=1}^{n} \left(\frac{dq_k}{(2\pi)^4}\right) (2\pi)^4 \delta^4(\sum_{i=1}^{n} q_i) tr(\hat{A}(q_1) \bar{S}(p - q_1) ... \hat{A}(q_n)) +
$$

+ $(I \to \bar{I})$ (21)

where the last bracket in (21) refers to the contribution of antiinstantons.

One can rewrite this expression in a simpler form

$$
M(p) = +i\frac{1}{N_c V_4} \sum_{i} \langle p | (\bar{S} - (\hat{A}^i)^{-1})^{-1} | p \rangle_R \tag{22}
$$

Eq.(22) is equivalent to Eq.(7) of [17] when one does Fourier transformation and averaging over dR_i .

Since $S(p)$ contains $M(p)$ as in (20), two coupled equations (20) and (21) define $M(p)$ through $\hat{A}(p)$ and the density parameter $\frac{N}{N_cV_4}$.

One can introduce the "Green's functions" $G(\bar{G})$ for the quark propagation in the instanton (antiinstanton) field $A(A)$ with effective mass $M(p)$, e.g.

$$
(\hat{p} - iM - \hat{A})G(p, p') = (2\pi)^{4}\delta(p - p')
$$
\n(23)

Separating δ -function term from G, one obtains

$$
G(p, p') = \frac{(2\pi)^4 \delta(p - p')}{\hat{p} - iM(p)} + \frac{\xi(p, p')}{(\hat{p} - iM(p))(\hat{p}' - iM(p'))}
$$
(24)

and similarly for \bar{G} and $\bar{\xi}$. Insertion of (24) into (22) yields

$$
M(p) = -\frac{iN}{2N_cV_4}tr \int \frac{d^4q}{(2\pi)^4} \{\hat{A}(p-q)\frac{1}{\hat{q}-iM(q)}\xi(q,p) + (A \to \bar{A}, \xi \to \bar{\xi})\}
$$
(25)

and for ξ one has an equation following from (24)

$$
\xi(p, p') = \hat{A}(p - p') + \int \hat{A}(p - q) \frac{1}{\hat{q} - iM(q)} \xi(q, p') \frac{d^4q}{(2\pi)^4},
$$
 (26)

and similar equation for $\bar{\xi}(p, p')$ with A_{μ} replaced by the antiinstanton field \bar{A}_{μ} .

Since one expects $M(0)$ to be around 200-300 MeV, [9-12, 17], the spectral expansion of G and the ZMA done in [17] which is valid for vanishing $M(0)$ is not convincing. Therefore one has to solve Eqs. (25-26) numerically.

The calculations done above in this section, demonstrate that (19) is the EQL corresponding to the Pobylitsa equation [17] obtained for $N_f = 1$ and $N_c \to \infty$. In case of $N_f > 1$ the EQL is given in (7) and its explicit form for $n = 2$ in (10). For many flavours $(N_f \geq 2)$ it is natural and (easier) to use the bosonized form of EQL, which will be published elsewhere.

Finally we discuss in the section symmetries of our EQL, $\sum_n \frac{\langle \mathcal{L} \mathcal{P}^n \rangle}{n!}$ $\frac{\theta^n \gg}{n!}$. We consider only the case $N_+ = N_-\,$, so that the net topological charge of the system is zero, and there are no global zero modes. Correspondingly the effective 'tHooft Lagrangian [7] with the global zero modes does not occur in this case. Now the 4q term in (7) can be rewritten as in $(A.9)$, the form which respects $U(N_f) \times U(N_f)$ symmetry. The same can be shown for all terms in (7), and therefore violation of $SU(N_f) \times SU(N_f)$ can occur only spontaneously while $U_A(1)$ has anomaly. The explicit example of CSB is given by a nonzero solution of (25-26), derived above in the limit $N_c \to \infty$.

4 Summary and discussion

The main result of the letter is the EQL (7), with explicit form for $n = 2$ and any N_f in (10), and for all n and $N_f = 1$, $N_c \rightarrow \infty$ in (19).

The 'tHooft–type ansatz of EQL of ref. [10-12] is obtained from our EQL (7) when one selects there $n = N_f$ and keeps in the mode expansion (14) only the zero mode contribution. One can check, that in this case automatically appears the determinantal structure of the 'tHooft ansatz. We have not found any arguments why nonzero modes in (14) and all terms with $n \neq N_f$ should be dropped.

It is interesting, that the structure of the 4q term in the EQL Eq. (10), is so much resembling the extended NJL model with all Lorenz group channels (see $(A.12)$), in contrast to the standard TLQ Lagrangian [10-12], where in the leading $1/N_c$ order only scalar and pseudoscalar channels appear.

It is not impossible, however, that the resulting effect of all terms of EQL in the sum over n in (7), will be numerically close to the standard TQL results [10-12] for $M(p)$ and $\langle \bar{q}q \rangle$. As one example of this sort one may quote the equation (22) for the effective quark mass which is obtained above from our EQL in the Hartree approximation and which coincides with that of ref. [17], where diagrammatic analysis for the quark propagator was exploited.

The author of [17] applied the ZMA for (22), valid in the limit of small $M(p)$, and obtained a solution for $M(p)$ closely resembling that of the standard TQL analysis [9-12]. However the resulting value of $M(0) \approx 350 MeV$ is not small and this fact casts some doubts on the whole numerical procedure of [17]. The crucial check of this coincidence would be the exact solution of Eqs. $(25-26)$ for $M(p)$ and its comparison with the standard TQL solution of $M(p)$ [9-12]. This will be a subject of a future publication.

As it is, our EQL (7) is an exact result for any N_c and any N_f , which enables one to calculate different physical effects, from $q\bar{q}$ creation due to the 4q vertex, to chiral symmetry breaking effects $(M(p), < \bar{q}q > \text{etc}).$ The corresponding detailed calculations and comparison with the existing TQL results is planned for the future and will help to clarify the mechanism of chiral dynamics in the QCD vacuum.

The author is grateful to G.'tHooft for a valuable discussion, to D.I.Diakonov for a useful discussion and correspondence, and to B.O.Kerbikov and D.S.Kuzmenko for numerous discussions in the course of the present work.

This work was supported in part by grants RFFI 96-02-19184, RFFI-DFG 96-02-000 88G and INTAS 94-2851.

APPENDIX 1

Derivation of the 4q term of the effective quark Lagrangian

$$
\langle \theta^2 \rangle \equiv \langle \int \psi^{+f_1}(x) \hat{A}(x) \psi^{f_1}(x) \int \psi^{+f_2}(y) \hat{A}(y) \psi^{f_2}(y) dx dy \rangle_{\Omega, R} =
$$

$$
\int (\psi_{\alpha}^{+f_1}(x) \gamma_{\mu_1} \psi_{\beta}^{f_1}(x)) dx \int (\psi_{\beta}^{+f_2}(y) \gamma_{\mu_2} \psi_{\alpha}^{f_2}(y) dy \frac{\Lambda_{\mu_1 \mu_2}(x, y)}{N_c^2 - 1}
$$
 (A.1)

where α, β are color indices, and

$$
\Lambda_{\mu_1\mu_2}(x,y) = \frac{2}{V_4} \int d^4R (\delta_{\mu_1\mu_2}\delta_{\nu\rho} - \delta_{\mu_1\rho}\delta_{\mu_2\nu} \pm \varepsilon_{\mu_1\nu\mu_2\rho})(x - R)_{\nu}(y - R)_{\rho} \times
$$

$$
\times a(x - R)a(y - R) \tag{A.2}
$$

where

$$
a(x) = \frac{\rho^2}{x^2(x^2 + \rho^2)}
$$
 (A.3)

and the sign $+(-)$ refer to the instanton (antiinstanton) case respectively.

In (A.1) one can notice that color indices are intercharged in both quark bilinears, which calls for the application of Fierz transformation, which gives

$$
\gamma_{\mu} \times \gamma_{\mu} = 1 \times 1 - \frac{1}{2} (\gamma_{\mu} \times \gamma_{\mu}) - \frac{1}{2} (\gamma_5 \gamma_{\mu} \times \gamma_5 \gamma_{\mu}) - \gamma_5 \times \gamma_5
$$
 (A.4)

Note that we are using Euclidean γ matrices. i.e.

$$
\gamma_4 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \gamma_i = -i\beta\alpha_i, \gamma_5 = \begin{pmatrix} -1 & 0 \\ -1 & 0 \end{pmatrix}, \gamma_\mu\gamma_\nu + \gamma_\nu\gamma_\mu = 2\delta_{\mu\nu}
$$

For the following it is useful to go over to the momentum space and to this end we need the Fourier transform of the instanton vector potential

$$
A_{\mu a}(x - R) = 2\bar{\eta}_{\mu a\nu}\rho^2 i 4\pi^2 \int \frac{d^4q}{(2\pi)^4} \cdot q_{\nu}\varphi(q)e^{iq(x - R)} \tag{A.5}
$$

where

$$
\varphi(q) = \frac{1}{q^2} \{ K_2(q\rho) - \frac{2}{(q\rho)^2} \}
$$
 (A.6)

Insertion of (A.5) into (A.1) yields

$$
\langle \theta^2 \rangle = \int \frac{dp}{(2\pi)^4} \frac{dp_1}{(2\pi)^4} \frac{dq}{(2\pi)^4} \frac{2(4\pi^2 \rho^2)^2}{(N_c^2 - 1)V_4} (\psi_\alpha^{+f_1}(p)\gamma_{\mu_1}\psi_\beta^{f_1}(p+q))(\psi_\beta^{+f_2}(p_1)\gamma_{\mu_2}\psi_\alpha^{f_2}(p_1-q))
$$

$$
(\delta_{\mu_1\mu_2}q^2 - q_{\mu_1}q_{\mu_2})\varphi^2(q) \tag{A.7}
$$

Using (A.4) and a similar Fierz relation for scalars

$$
1 \times 1 = \frac{1}{4} \{ 1 \times 1 + \gamma_{\mu} \times \gamma_{\mu} + \frac{1}{2} \sigma_{\mu\nu} \times \sigma_{\mu\nu} - \gamma_5 \gamma_{\mu} \times \gamma_5 \gamma_{\mu} + \gamma_5 \times \gamma_5 \} \quad (A.8)
$$

one obtains for (A.7)

$$
\langle \theta_i^2 \rangle = \int \frac{dpdp_1dq}{(2\pi)^{12}} \frac{2(4\pi^2\rho^2)^2\varphi^2(q)}{(N_c^2 - 1)V_4} \{q^2 \sum_{i=1,2,3,4} c_i(\psi^{+f_1}(p)O_i\psi^{f_2}(p_1 - q)) \times
$$

$$
(\psi^{+f_2}(p_1)O_i\psi^{f_1}(p+q)) - \sum_{i=1}^5 c'_i(\psi^{+f_1}(p)\hat{q}O_i\psi^{f_2}(p_1 - q))(\psi^{+f_2}(p_1)\hat{q}O_i\psi^{f_1}(p+q))\}
$$

(A.9)

where O_i and c_i, c'_i are

$$
O_i = 1, \gamma_\mu, \gamma_5 \gamma_\mu, \gamma_5, \sigma_{\mu\nu} = \frac{\gamma_\mu \gamma_\nu - \gamma_\nu \gamma_\mu}{2i} \text{ for } i = 1, 2, 3, 4, 5
$$

$$
c_i = 1, -\frac{1}{2}, -\frac{1}{2}, -1, 0
$$

$$
c'_i = \frac{1}{4}, \frac{1}{4}, -\frac{1}{4}, \frac{1}{4}, \frac{1}{8}
$$
(A.10)

Now we take care of the flavour indices

$$
\sum_{F=1}^{N_f^2-1} t_{fg}^F t_{ed}^F = -\frac{1}{2N_f} \delta_{fg} \delta_{ed} + \frac{1}{2} \delta_{fd} \delta_{ge}, \tag{A.11}
$$

to rewrite the curly brackets in (A.9) as

$$
\{\ldots\} = 2q^2 \sum_{i,F} c_i(\psi^+(p)O_i t^F \psi(p_1 - q))(\psi^+(p_1)O_i t^F \psi(p + q)) +
$$

+
$$
\frac{q^2}{N_f} \sum_i c_i(\psi^+(p)O_i 1^F \psi(p_1 - q))(\psi^+(p_1)O_i 1^F \psi(p + q)) +
$$

$$
-2\sum_{F,i=1}^{5} c'_{i}(\psi^{+}(p)\hat{q}O_{i}t^{F}\psi(p_{1}-q))(\psi^{+}(p_{1})\hat{q}O_{i}t^{F}\psi(p+q)) -
$$

$$
-\frac{1}{N_{f}}\sum_{i=1}^{5} c'_{i}(\psi^{+}(p)\hat{q}O_{i}1^{F}\psi(p_{1}-q))(\psi^{+}(p_{1})\hat{q}O_{i}1^{F}\psi(p+q))
$$
(A.12)

We now can introduce currents

$$
J_i^{F,O}(p) = \psi^+(p)O_i(t^F, 1^F)\psi(p_1 - q)
$$

and similarly for $\tilde{J}_i^{F,O}$, where instead of O_i stands $\hat{q}O_i$, to rewrite all terms in (A.12) as

$$
\{\ldots\} = 2q^2 \sum_{i,F} c_i J_i^F(p) J_i^F(p_1) + \frac{q^2}{N_f} \sum_i c_i J_i^0(p) J_i^0(p_1)
$$

$$
- 2 \sum_{i,F} c'_i \tilde{J}_i^F(p) \tilde{J}_i^F(p_1) - \frac{1}{N_f} \sum_i c'_i \tilde{J}_i^0(p) \tilde{J}_i^0(p_1)
$$
(A.13)

As a result (A.9) assumes the form

$$
\langle \theta_i^2 \rangle = \int \frac{dpdp_1dq}{(2\pi)^{12}} \frac{2(4\pi^2\rho^2)^2 \varphi^2(q)}{(N_c^2 - 1)V_4} \{2q^2(1 - \frac{1}{N_c^2}) \sum_{i,F} c_i J_i^F(p) J_i^F(p_1) +
$$

$$
\frac{q^2}{N_f} (1 - \frac{1}{N_c^2}) \sum_i c_i J_i^0(p) J_i^0(p_1) - 2 \sum_{i,F} c'_i \tilde{J}_i^F(p) \tilde{J}_i^F(p_1) -
$$

$$
- \frac{1}{N_f} \sum_i c'_i \tilde{J}_i^0(p) \tilde{J}_i^0(p_1) \}.
$$
(A.14)

References

- [1] C.G.Callan, R.Dashen and D.J.Gross, Phys. Rev. D17 (1978) 2717, ibid D19(1987) 1826.
- [2] E.V.Shuryak, Nucl. Phys. B203 (1982) 93, 116, 140.
- [3] D.I.Diakonov and V.Yu.Petrov, Nucl. Phys. **B245**(1984) 259.
- [4] M.C.Chu, J.M.Grandy, S.Huang, J.W.Negele, Phys. Rev. Lett. 70 (1993) 225; Phys. ReV D49 (1994) 6039.
- [5] E.V.Shuryak, Lectures at the International E.Fermi School Varenna, 1995. (in press)
- [6] G.'tHooft, Phys. Rep. 142 (1986) 357.
- [7] G.'tHooft, Phys. Rev. D 14 (1976) 3432; Phys. Rev. Lett. 37 (1976)8.
- [8] D.G.Caldi, Phys. Rev. Lett. 39 (1977)121 R.D.Carlitz and D.B.Creamer, Ann. Phys. 188 (1979) 429 C.Carniero and N.A. McDougall, Nucl. Phys. B245 (1984) 293
- [9] D.I.Diakonov and V.Yu.Petrov, Nucl. Phys. B 272(1986) 457.
- [10] D.I.Diakonov and V.Yu.Petrov, Leningrad preprint 1153 (1986)
- [11] E.V.Shuryak, Nucl. Phys. B 302 (1988) 599; M.A.Nowak, J.J.M.Verbaarschot and I.Zahed, Nucl. Phys. B324(1989)1 M.A.Nowak et al. Phys. Lett. B228 (1989) 251
- [12] D.I.Diakonov and V.Yu.Petrov, in Quark Cluster Dynamics, Lecture Notes in Physics, Springer–Verlag (1992) p. 228
- [13] D.I.Diakonov , E.V.Shuryak and J.J.M.Verbaarschot, Nucl. Phys. B410 (1993) 37 A.Gonzalez-Arroyo and Yu.A.Simonov, Nucl. Phys. B460 (1996) 429 T.Wettig, A.Schaefer and H.A.Weidenmuller, Phys. Lett. B367 (1996) 28
- [14] B.O.Kerbikov, D.S.Kuzmenko, Yu.A.Simonov, [hep-ph/9609440](http://arxiv.org/abs/hep-ph/9609440) JETP Letters 65 (1997) 128
- [15] Y.Nambu and G.Jona–Lasinio, Phys. Rev. 122 S.P.Klevansky, Rev. Mod. Phys. 64 (1992) 649
- [16] W.A.Bardeen, C.T.Hill and M.Lindner, Phys. Rev. D41 (1990) 1647 M.Lindner, Int. J. Mod. Phys. A8 (1993) 2167
- [17] P.V.Pobylitsa, Phys. Lett. B226 (1989) 387
- [18] N.G.Van Kampen, Phys. Rep. 24 (1976) 171; Physika 74 (1974) 215, 239