BPS States in F-Theory

Yu. Malyuta and T. Obikhod

Institute for Nuclear Research, National Academy of Sciences of Ukraine 252022 Kiev, Ukraine E-mail: aks@d310.icyb.kiev.ua

Abstract

The spectra of BPS states in F-theory on elliptic fibred fourfolds are investigated.

1 Introduction

Klemm, Mayr and Vafa [1] have shown that F-theory compactification on the Calabi-Yau threefold $X_{24}(1, 1, 2, 8, 12)$ leads to the spectrum of BPS states recorded in Table 1.

m	$n_{0,0,m}$	$n_{1,0,m}$	$n_{2,0,m}$	$n_{3,0,m}$	$n_{4,0,m}$	$n_{5,0,m}$
1	1	252	5130	54760	419895	2587788
2			-9252	-673760	-20534040	-389320128
3				848628	115243155	6499779552

Table 1. Instanton numbers (degeneracies of BPS states) for $X_{24}(1, 1, 2, 8, 12)$.

Other F-theory compactifications on the special fourfolds have been studied by Donagi, Grassi, Witten [2] and Mayr [3] in the context of non-perturbative superpotentials.

The purpose of this work is to investigate the spectra of BPS states in F-theory on elliptic fibred fourfolds. We consider some models which have non-perturbative superpotentials.

2 Donagi-Grassi-Witten model

The fourfold in this case is the elliptic fibration over the base $\mathbf{P}^1 \times S$, where S denotes the del Pezzo surface [2]. The model exhibits the geometrical phase which corresponds to the regular triangulation of the dual polihedron [4]. This phase has the following Mori generators

$$\begin{split} l^{(1)} &= (-3,0;1,1,1,0,0,0,0,0,0,0), \\ l^{(2)} &= (-1,-1;0,0,0,1,1,0,0,0,0,0), \\ l^{(3)} &= (0,-2;0,0,0,0,0,0,1,1,0,0,0), \\ l^{(4)} &= (0,-3;0,0,0,0,0,0,0,0,1,1,1). \end{split}$$

Using prescriptions of the paper [5] we derive the principal parts of the Picard-Fuchs operators

$$L_{1} = 3\theta_{1}^{2} - \theta_{1}\theta_{2} + \theta_{2}^{2},$$

$$L_{2} = \theta_{2}^{2},$$

$$L_{3} = \theta_{3}^{2},$$

$$L_{4} = \theta_{2}^{2} + 4\theta_{2}\theta_{3} + 4\theta_{3}^{2} - 3\theta_{2}\theta_{4} - 6\theta_{3}\theta_{4} + 9\theta_{4}^{2},$$

where $\theta_i = z_i \frac{d}{dz_i}$, z_i are the algebraic coordinates on the complex structure moduli space.

Application of the program INSTANTON [6] gives the spectrum of BPS states recorded in Table 2.

$n_{2,1,3,0,0}$	$n_{2,1,3,0,1}$	$n_{2,1,3,0,2}$		$n_{2,1,3,0,4}$	$n_{2,1,3,0,5}$
	252	$\begin{array}{ccc} 5130 & 54760 \\ (27 \text{ sequences}) \end{array}$		419895	2587788
	$\frac{n_{2,1,3,0,1}}{1 \cdot 252^*}$	$\frac{n_{2,2,3,0,1}}{2 \cdot 5130^*}$	$\frac{n_{2,3,3,0,1}}{3 \cdot 54760^*}$ (27 sequences)	$\frac{n_{2,4,3,0,1}}{4 \cdot 419895^*}$	$\frac{n_{2,5,3,0,1}}{5\cdot 2587788^*}$
$\begin{array}{c c} n_{4,0,3,0,1} \\ 1 \end{array}$	$\frac{n_{4,1,3,0,1}}{252}$	$\frac{n_{4,2,3,0,1}}{5130}$	$\frac{n_{4,3,3,0,1}}{54760}$ (513 sequences)	$\frac{n_{4,4,3,0,1}}{419895}$	$\frac{n_{4,5,3,0,1}}{2587788}$
$n_{5,0,3,0,1}$ 1	$\frac{n_{5,1,3,0,1}}{252}$	$\frac{n_{5,2,3,0,1}}{5130}$	$\frac{n_{5,3,3,0,1}}{54760}$ (702 sequences)	$\frac{n_{5,4,3,0,1}}{419895}$	$\frac{n_{5,5,3,0,1}}{2587788}$
$n_{6,0,3,0,1}$ 1	$\frac{n_{6,1,3,0,1}}{252}$	$\frac{n_{6,2,3,0,1}}{5130}$	$\frac{n_{6,3,3,0,1}}{54760}$ (189 sequences)	$\frac{n_{6,4,3,0,1}}{419895}$	$\frac{n_{6,5,3,0,1}}{2587788}$

Table 2. Instanton numbers (degeneracies of BPS states) for the Donagi-Grassi-Witten model.

We found the tower of infinite sequences of BPS degeneracies. The sequences 1, 252, 5130, ... are well-known. The E_8 partition function is the counting functional of these sequences [7]. But the sequences $1 \cdot 252$, $2 \cdot 5130$, $3 \cdot 54760$, ... (marked by the * in Table 2) are unfamiliar, and physically interesting, as they signal the appearance of additional states.

3 Elliptic fibred fourfold over the \mathbf{P}^1 - bundle on $(\mathbf{P}^1)^2$

The fourfold in this case is the elliptic fibration over the \mathbf{P}^1 -bundle $\mathbf{P}(\mathcal{O}_B \otimes \mathcal{O}_B (f_1 + f_2))$ on $B = (\mathbf{P}^1)^2$, where f_1 and f_2 are fibres of the two projections from B to \mathbf{P}^1 [3][7]. This model exhibits the geometrical phase which has the Mori generators

$$\begin{split} l^{(1)} &= (0; 1, 0, 0, 0, 0, -1, 0, -1, 1), \\ l^{(2)} &= (0; 0, 0, 1, 0, 0, 1, 0, -2, 0), \\ l^{(3)} &= (-6; 0, 0, 0, 2, 3, 0, 0, 1, 0), \\ l^{(4)} &= (0; 0, 1, 0, 0, 0, -1, 1, -1, 0). \end{split}$$

The principal parts of the Picard-Fuchs operators are

$$L_{1} = \theta_{1}^{2},$$

$$L_{2} = \theta_{2}(\theta_{1} - \theta_{2} + \theta_{4}),$$

$$L_{3} = \theta_{3}(\theta_{1} + 2\theta_{2} - \theta_{3} + \theta_{4}),$$

$$L_{4} = \theta_{4}^{2}.$$

With the program INSTANTON we obtain the spectrum of BPS states recorded in Table 3.

$\frac{n_{3,0,0,0,1}}{1}$	$\frac{n_{3,0,0,1,1}}{252}$	$\frac{n_{3,0,0,2,1}}{5130}$	$\frac{n_{3,0,0,3,1}}{54760}$ (1 sequence)	$\frac{n_{3,0,0,4,1}}{419895}$	$\frac{n_{3,0,0,5,1}}{2587788}$
$n_{3,1,0,0,0}$ 1	$\frac{n_{3,1,0,1,0}}{252}$	$\frac{n_{3,1,0,2,0}}{5130}$	$n_{3,1,0,3,0}$ 54760 (1 sequence)	$\frac{n_{3,1,0,4,0}}{419895}$	$\frac{n_{3,1,0,5,0}}{2587788}$
		$n_{3,0,0,2,2}$ -9252	$n_{3,0,0,3,2}$ -673760 (2 sequences)	$\frac{n_{3,0,0,4,2}}{-20534040}$	$\frac{n_{3,0,0,5,2}}{-389320128}$
		$n_{3,2,0,2,0}$ -9252	$n_{3,2,0,3,0} - 673760$ (2 sequences)	$\frac{n_{3,2,0,4,0}}{-20534040}$	$\frac{n_{3,2,0,5,0}}{-389320128}$
			$\frac{n_{3,0,0,3,3}}{848628}$ (3 sequences)	$\frac{n_{3,0,0,4,3}}{115243155}$	$\frac{n_{3,0,0,5,3}}{6499779552}$
			$n_{3,3,0,3,0}$ 848628 (3 sequences)	$\frac{n_{3,3,0,4,0}}{115243155}$	$\frac{n_{3,3,0,5,0}}{6499779552}$
	$\frac{n_{5,1,0,1,0}}{-1 \cdot 252^*}$	$\frac{n_{5,1,0,2,0}}{-2\cdot 5130^*}$	$\frac{n_{5,1,0,3,0}}{-3 \cdot 54760^*}$ (2 sequences)	$\frac{n_{5,1,0,4,0}}{-4 \cdot 419895^*}$	$\frac{n_{5,1,0,5,0}}{-5\cdot2587788^*}$
		$\frac{n_{5,2,0,2,0}}{2 \cdot 9252^*}$	$\frac{n_{5,2,0,3,0}}{3 \cdot 673760^*}$ (2 sequences)	$\frac{n_{5,2,0,4,0}}{4 \cdot 20534040^*}$	$\frac{n_{5,2,0,5,0}}{5\cdot 389320128^*}$
			$ \frac{n_{5,3,0,3,0}}{-3 \cdot 848628^*} $ (2 sequences)	$\frac{n_{5,3,0,4,0}}{-4 \cdot 115243155^*}$	$\frac{n_{5,3,0,5,0}}{-5\cdot 6499779552^*}$

Table 3. Instanton numbers (degeneracies of BPS states) for the elliptic fibred fourfold over the \mathbf{P}^1 - bundle on $(\mathbf{P}^1)^2$.

The sequences

 $\begin{array}{c} 1,\ 252,\ 5130,\ \ldots,\\ -9252,\ -673760,\ -20534040,\ \ldots,\\ 848628,\ 115243155,\ 6499779552,\ \ldots\end{array}$

are well-known. But the sequences

 $\begin{array}{c} -1 \cdot 252, \ -2 \cdot 5130, \ -3 \cdot 54760, \ \ldots, \\ 2 \cdot 9252, \ 3 \cdot 673760, \ 4 \cdot 20534040, \ \ldots, \\ -3 \cdot 848628, \ -4 \cdot 115243155, \ -5 \cdot 6499779552, \ \ldots \end{array}$

(marked by the * in Table 3) are unfamiliar, and physically interesting, as they signal the appearence of additional states.

4 Elliptic fibred fourfold over $\mathbf{P}^1 \times F_1$

The fourfold in this case is the elliptic fibration over $\mathbf{P}^1 \times F_1$, where F_1 is the Hirzebruch surface [3][7]. This model exhibits the geometrical phase which has the Mori generators

$$\begin{split} l^{(1)} &= (-6; 0, 0, 0, 0, 0, 0, 2, 3, 1), \\ l^{(2)} &= (0; 0, 0, 0, 0, 1, 1, 0, 0, -2), \\ l^{(3)} &= (0; 0, 1, 0, 1, 0, -1, 0, 0, -1), \\ l^{(4)} &= (0; 1, -2, 1, 0, 0, 0, 0, 0, 0). \end{split}$$

The principal parts of the Picard-Fuchs operators are

$$\begin{split} L_1 &= \theta_1 (\theta_1 - 2\theta_2 - \theta_3), \\ L_2 &= \theta_2 (\theta_2 - \theta_3), \\ L_3 &= \theta_3 (\theta_3 - 2\theta_4), \\ L_4 &= \theta_4^2. \end{split}$$

With the program INSTANTON we obtain the spectrum of BPS states recorded in Table 4.

	$n_{3,1,0,1,0}$	$n_{3,2,0,1,0}$	$n_{3,3,0,1,0}$	$n_{3,4,0,1,0}$	$n_{3,5,0,1,0}$
	$1 \cdot 252^{*}$	$2 \cdot 5130^{*}$	$\frac{3 \cdot 54760^*}{(1 \text{ sequence})}$	$4 \cdot 419895^{*}$	$5 \cdot 2587788^{*}$
		$\frac{n_{3,2,0,2,0}}{-2 \cdot 9252^*}$	$ \frac{n_{3,3,0,2,0}}{-3 \cdot 673760^{*}} \\ (1 \text{ sequence}) $	$\frac{n_{3,4,0,2,0}}{-4 \cdot 20534040^*}$	$\frac{n_{3,5,0,2,0}}{-5\cdot 389320128^*}$
			$ \frac{n_{3,3,0,3,0}}{3 \cdot 848628^*} \\ (1 \text{ sequence}) $	$\frac{n_{3,4,0,3,0}}{4 \cdot 115243155^*}$	$\frac{n_{3,5,0,3,0}}{5\cdot 6499779552^*}$
$ \begin{bmatrix} n_{6,0,0,1,0} \\ 1 \end{bmatrix} $	$\frac{n_{6,1,0,1,0}}{252}$	$\frac{n_{6,2,0,1,0}}{5130}$		$\frac{n_{6,4,0,1,0}}{419895}$	$\frac{n_{6,5,0,1,0}}{2587788}$
$\begin{array}{c} n_{6,0,0,1,1} \\ 1 \end{array}$	$\frac{n_{6,1,0,1,1}}{252}$	$\frac{n_{6,2,0,1,1}}{5130}$	$\frac{n_{6,3,0,1,1}}{54760}$ (5 sequences)	$\frac{n_{6,4,0,1,1}}{419895}$	$\frac{n_{6,5,0,1,1}}{2587788}$
		$\frac{n_{6,2,0,2,0}}{-9252}$	$\frac{n_{6,3,0,2,0}}{-673760}$ (10 sequences)	$\frac{n_{6,4,0,2,0}}{-20534040}$	$\frac{n_{6,5,0,2,0}}{-389320128}$
		$\frac{n_{6,2,0,2,2}}{-9252}$	$\frac{n_{6,3,0,2,2}}{-673760}$ (10 sequences)	$\frac{n_{6,4,0,2,2}}{-20534040}$	$\frac{n_{6,5,0,2,2}}{-389320128}$
				$\frac{n_{6,4,0,3,0}}{115243155}$	$\frac{n_{6,5,0,3,0}}{6499779552}$
			$\frac{n_{6,3,0,3,3}}{848628}$ (15 sequences)	$\frac{n_{6,4,0,3,3}}{115243155}$	$\frac{n_{6,5,0,3,3}}{6499779552}$

Table 4. Instanton numbers (degeneracies of BPS states) for the elliptic fibred fourfold over $\mathbf{P}^1 \times F_1$.

The sequences

 $\begin{array}{c} 1,\ 252,\ 5130,\ \ldots,\\ -9252,\ -673760,\ -20534040,\ \ldots,\\ 848628,\ 115243155,\ 6499779552,\ \ldots\end{array}$

are well-known. But the sequences

 $1 \cdot 252, 2 \cdot 5130, 3 \cdot 54760, \ldots,$

 $-2 \cdot 9252, -3 \cdot 673760, -4 \cdot 20534040, \ldots,$

 $3 \cdot 848628, 4 \cdot 115243155, 5 \cdot 6499779552, \ldots$

(marked by the * in Table 4) are unfamiliar, and physically interesting, as they signal the appearence of additional states.

References

- A. Klemm, P. Mayr and C. Vafa, BPS States of Exceptional Non-Critical Strings, hep-th/9607139.
- [2] R. Donagi, A. Grassi and E. Witten, A Non-Perturbative Superpotential with E₈-Symmetry, hep-th/9607091.
- [3] P. Mayr, Mirror Symmetry, N = 1 Superpotentials and Tensionless Strings on Calabi-Yau Fourfolds, hep-th/9610162.
- [4] J. De Loera, PUNTOS, ftp.geom.umn.edu/pub/software.
- [5] S. Hosono, A. Klemm, S. Theisen and S.-T. Yau, Mirror Symmetry, Mirror Map and Applications to Complete Intersection Calabi-Yau Spaces, Nucl. Phys. B433 (1995) 501.
- [6] A. Klemm, An updated version of the Mathematica program INSTANTON for complete intersection or hypersurface fourfolds in toric ambient spaces, 1997.
- [7] A. Klemm, B. Lian, S.-S. Roan and S.-T. Yau, Calabi-Yau Fourfolds for M- and F-Theory Compactifications, hep-th/9701023.