

# Modulo 2 periodicity of complex Clifford algebras and electromagnetic field

Vadim V. Varlamov

*Applied Mathematics, Siberian State Academy of Mining & Metallurgy,  
Novokuznetsk, Russia\**

## **Abstract**

Electromagnetic field is considered in the framework of a Clifford algebra  $\mathbf{C}_2$  over a field of complex numbers. It is shown here that a modulo 2 periodicity of complex Clifford algebras may be connected with electromagnetic field.

---

\*E-mail: root@varlamov.kemerovo.su

# 1 INTRODUCTION

It is well-known that algebra introduced by Clifford [1] be simultaneously a synthesis and generalization of Hamilton theory of quaternions and Grassmann algebras. In the other words, Clifford algebra is generalization of quaternion algebra onto a case of manydimensional spaces. Thanks to this fact Lipschitz [2] was determined the tight connection between Clifford algebras and rotation groups of manydimensional spaces. The Clifford algebras becomes the object of fixed attention of physicists after introduction in theory of electron the noncommutative algebra of Dirac  $\gamma$ -matrices.

We shall denote further Clifford algebra as  ${}^l\mathbf{K}_n$ , where  $\mathbf{K}$  is algebraic field of characteristic 0 ( $\mathbf{K} = \mathbf{R}, \mathbf{K} = \mathbf{\Omega}, \mathbf{K} = \mathbf{C}$ );  $\mathbf{\Omega}$  is a field of double numbers (so-called singular field [3]).

In essence, the problem of periodicity of algebras  ${}^l\mathbf{K}_n$  to begin on the first Clifford paper (see ref.1), where the each algebra  ${}^l\mathbf{K}_n$  is considered as a product of the some number of quaternion algebras. After construction of spinor representation of Clifford algebras [4] the structure of periodicity was finally elucidated by Atiyah, Bott, and Shapiro [5]. Namely, the Clifford algebras over a field of real numbers  $\mathbf{K} = \mathbf{R}$  are possess a modulo 8 periodicity. By force of identity  $\mathbf{\Omega} = \mathbf{R} \oplus \mathbf{R}$  the algebras over a field of double numbers are possess the same periodicity. In the case of a field  $\mathbf{K} = \mathbf{C}$  we have a modulo 2 periodicity.

The application of these periodicities in physics to begin on the paper of Coquereaux [6], where the modulo 8 periodicity of real algebras was used to particle physics. In present paper the modulo 2 periodicity of complex algebras is connected with electromagnetic field. This connection is possible by force of realization of the basic notions of electromagnetic field (such that vectors  $\mathbf{E}$  and  $\mathbf{H}$ , Maxwell equations) in the terms of an algebra  $\mathbf{C}_2$ , which by Clifford terminology is called the algebra of hyperbolic biquaternions.

# 2 CLIFFORD ALGEBRAS OVER A FIELD $\mathbf{K}$

A Clifford algebra  $\mathbf{R}_{n-1}$  over a field of real numbers  $\mathbf{R}$  be an algebra with  $2^{n-1}$  basic elements:  $\mathbf{e}_0$  (unit of algebra),  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_{n-1}$  and products  $\mathbf{e}_{i_1 i_2 \dots i_k} = \mathbf{e}_{i_1} \mathbf{e}_{i_2} \dots \mathbf{e}_{i_k}$ . Multiplication in  $\mathbf{R}_{n-1}$  defined by a following rule:

$$\mathbf{e}_i^2 = \mathbf{e}_0 \quad , \quad \mathbf{e}_i \mathbf{e}_j = -\mathbf{e}_j \mathbf{e}_i \quad (1)$$

Furtner on,  ${}^l\mathbf{R}_{n-1}$  be an algebra which we obtain from  $\mathbf{R}_{n-1}$  by substitution (1) onto

$$\mathbf{e}_i^2 = \sigma(i-l)\mathbf{e}_0 \quad , \quad \mathbf{e}_i \mathbf{e}_j = -\mathbf{e}_j \mathbf{e}_i, \quad (2)$$

where

$$\sigma(n) = \begin{cases} -1, & \text{if } n \leq 0 \\ 1, & \text{if } n > 0 \end{cases}$$

The algebra  $\mathbf{R}_{n-1}$  be a particular case of  ${}^l\mathbf{R}_{n-1}$  when  $l = 0$ .

A general element  $\mathcal{A}$  of  ${}^l\mathbf{R}_{n-1}$  represented by a following formal polynomial:

$$\begin{aligned}\mathcal{A} &= a_0\mathbf{e}_0 + \sum_{i=1}^{n-1} a_i\mathbf{e}_i + \sum_{i=1}^{n-1} \sum_{j=1}^{n-1} a_{ij}\mathbf{e}_{ij} + \dots + \sum_{i_1=1}^{n-1} \dots \sum_{i_k=1}^{n-1} a_{i_1\dots i_k}\mathbf{e}_{i_1\dots i_k} + \\ &+ \dots + a_{12\dots n-1}\mathbf{e}_{12\dots n-1} = \sum_{k=0}^{n-1} a_{i_1 i_2 \dots i_k} \mathbf{e}_{i_1 i_2 \dots i_k}.\end{aligned}$$

It is obvious that Clifford algebra  ${}^l\mathbf{R}_{n-1}$  is associative. A center of  ${}^l\mathbf{R}_{n-1}$  consist of unit  $\mathbf{e}_0$  and volume element  $\mathbf{e}_{12\dots n-1}$ . The volume element is belong to the center only if  $n-1$  is odd. Since

$$\begin{aligned}\mathbf{e}_{12\dots n-1}\mathbf{e}_i &= \sigma(i-l)(-1)^{n-i-1}\mathbf{e}_{12\dots i-1i+1\dots n-1}, \\ \mathbf{e}_i\mathbf{e}_{12\dots n-1} &= \sigma(i-l)(-1)^{i-1}\mathbf{e}_{12\dots i-1i+1\dots n-1},\end{aligned}$$

then  $\mathbf{e}_{12\dots n-1}$  is belong to the center if  $n-i-1 \equiv i-1 \pmod{2}$ . When  $n-1$  is even, the element  $\mathbf{e}_{12\dots n-1}$  commutes with elements  $\mathbf{e}_{i_1 i_2 \dots i_{2k+1}}$ , and anticommutes with elements  $\mathbf{e}_{i_1 i_2 \dots i_{2k}}$ .

The transition from  ${}^l\mathbf{R}_{n-1}$  to  ${}^l\mathbf{R}_n$  or  ${}^{l+1}\mathbf{R}_n$  may be represented as the transition from the real numbers in  ${}^l\mathbf{R}_{n-1}$  to complex coordinates  $a + b\omega$ , where  $\omega$  is additional basis element  $\mathbf{e}_{12\dots n}$ . By force of (1)-(2) in the case of transition from  ${}^l\mathbf{R}_{n-1}$  to  ${}^l\mathbf{R}_n$  we have:

$$\mathbf{e}_{12\dots n}^2 = (-1)^{l + \frac{n(n-1)}{2}}$$

and in case  ${}^l\mathbf{R}_{n-1} \rightarrow {}^{l+1}\mathbf{R}_n$ :

$$\mathbf{e}_{12\dots n}^2 = (-1)^{l+1 + \frac{n(n-1)}{2}}.$$

Therefore, in the first case

$$\omega^2 = \begin{cases} -1, & \text{if } n = 4m + 2 \text{ or } n = 4m' - 1, l \text{ is even and} \\ & \text{if } n = 4m' \text{ or } n = 4m' + 1, l \text{ is odd;} \\ +1, & \text{if } n = 4m' \text{ or } n = 4m' + 1, l \text{ is even and} \\ & \text{if } n = 4m + 2 \text{ or } n = 4m' - 1, l \text{ is odd;} \end{cases}$$

and in the second case

$$\omega^2 = \begin{cases} -1, & \text{if } n = 4m + 2 \text{ or } n = 4m' - 1, l \text{ is odd and} \\ & \text{if } n = 4m' \text{ or } n = 4m' + 1, l \text{ is even;} \\ +1, & \text{if } n = 4m' \text{ or } n = 4m' + 1, l \text{ is odd and} \\ & \text{if } n = 4m + 2 \text{ or } n = 4m' - 1, l \text{ is even;} \end{cases}$$

where  $m = 0, 1, \dots; m' = 1, 2, \dots$

Hence it follows that for  $l$  is even we have:

$${}^l\mathbf{R}_{4m'-1} = {}^l\mathbf{C}_{4m'-2}, \quad {}^{l+1}\mathbf{R}_{4m'+1} = {}^l\mathbf{C}_{4m'}, \quad (3)$$

$${}^l\mathbf{R}_{4m'+1} = {}^l\mathbf{\Omega}_{4m'} \quad , \quad {}^{l+1}\mathbf{R}_{4m'-1} = {}^l\mathbf{\Omega}_{4m'-2}, \quad (4)$$

and for  $l$  is odd

$${}^l\mathbf{R}_{4m'+1} = {}^l\mathbf{C}_{4m'} \quad , \quad {}^{l+1}\mathbf{R}_{4m'-1} = {}^l\mathbf{C}_{4m'-2}, \quad (5)$$

$${}^l\mathbf{R}_{4m'-1} = {}^l\mathbf{\Omega}_{4m'-2} \quad , \quad {}^{l+1}\mathbf{R}_{4m'+1} = {}^l\mathbf{\Omega}_{4m'}, \quad (6)$$

where  ${}^l\mathbf{C}_n$  and  ${}^l\mathbf{\Omega}_n$  are Clifford algebras over the field of complex numbers  $\mathbf{C}$  and field of double numbers  $\mathbf{\Omega}$ , respectively. When  $n$  is even, the identities of type (3)-(6) excepted, since in this case volume element  $\omega = \mathbf{e}_{12\dots n}$  is not belong to a center of extended algebra.

In particular case, when  $l = 0$  we have a transition  $\mathbf{R}_{n-1} \rightarrow \mathbf{R}_n$  and

$$\mathbf{e}_{12\dots n}^2 = (-1)^{\frac{n(n-1)}{2}}$$

Therefore,

$$\omega^2 = \begin{cases} -1, & \text{if } n = 4m + 2 \text{ or } n = 4m' - 1, \\ +1, & \text{if } n = 4m' \quad \text{or } n = 4m' + 1. \end{cases}$$

In this case, for  $n$  is odd we have the following identities:

$$\mathbf{R}_{4m'-1} = \mathbf{C}_{4m'-2} \quad , \quad \mathbf{R}_{4m'+1} = \mathbf{\Omega}_{4m'}. \quad (7)$$

### 3 THE ALGEBRAS $\mathbf{R}_3$ , $\mathbf{C}_2$ AND ELECTROMAGNETIC FIELD

Consider now an algebra  $\mathbf{R}_3$ . The squares of units of this algebra are equal to  $\mathbf{e}_i^2 = 1 (i = 0, 1, 2, 3)$ . The general element of  $\mathbf{R}_3$  has a following form:

$$\mathcal{A} = a_0\mathbf{e}_0 + \sum_{i=1}^3 a_i\mathbf{e}_i + \sum_{i=1}^3 \sum_{j=1}^3 a_{ij}\mathbf{e}_{ij} + a_{123}\mathbf{e}_{123}.$$

Further on, for a general element of  $\mathbf{R}_3$  there exists a decomposition  $\mathcal{A}^3 = \mathcal{A}^2 + \omega\mathcal{A}^2$ , where  $\omega = \mathbf{e}_{123} \in \mathbf{R}_3$  and  $\mathcal{A}^2$  is a general element of  $\mathbf{R}_2$ :  $a_0\mathbf{e}_0 + a_1\mathbf{e}_1 + a_2\mathbf{e}_2 + a_{12}\mathbf{e}_{12}$ . In fact

$$\begin{aligned} \mathcal{A}^3 &= \mathcal{A}^2 + \omega\mathcal{A}^2 = a_0\mathbf{e}_0 + a_1\mathbf{e}_1 + a_2\mathbf{e}_2 + a_{12}\mathbf{e}_{12} + \omega(b_0\mathbf{e}_0 + b_1\mathbf{e}_1 + b_2\mathbf{e}_2 + b_{12}\mathbf{e}_{12}) = \\ &= (a_0 + \omega b_0)\mathbf{e}_0 + (a_1 + \omega b_1)\mathbf{e}_1 + (a_2 + \omega b_2)\mathbf{e}_2 + (a_{12} + \omega b_{12})\mathbf{e}_{12} = \\ &= a_0\mathbf{e}_0 + a_1\mathbf{e}_1 + a_2\mathbf{e}_2 - b_{12}\mathbf{e}_3 + a_{12}\mathbf{e}_{12} - b_2\mathbf{e}_{13} + b_1\mathbf{e}_{23} + b_0\mathbf{e}_{123}. \end{aligned} \quad (8)$$

Since in this case  $\omega = \mathbf{e}_{123}$  is belong to the center of  $\mathbf{R}_3$ , and  $\omega^2 = -1$ , then in (8) the expressions  $a_k + \omega b_k$  may be replaced by  $a_k + ib_k$ , where  $i$  is imaginary unit. This way, we have the identity  $\mathbf{R}_3 = \mathbf{C}_2$ , which be a particular case of (7). The algebra  $\mathbf{C}_2$  is called the algebra of complex quaternions (or hyperbolic biquaternions).

Analogously, for the algebra  ${}^3\mathbf{R}_3$  with  $\omega^2 = +1$  we have in accordance with (6) the identity  ${}^3\mathbf{R}_3 = {}^2\mathbf{\Omega}_2$ , where  ${}^2\mathbf{\Omega}_2$  is the algebra of elliptic biquaternions.

Further on, there exists a realization of electromagnetic field in the terms of  $\mathbf{R}_3$ . Let

$$\begin{aligned}\mathcal{A}_0 &= \partial_0\mathbf{e}_0 + \partial_1\mathbf{e}_1 + \partial_2\mathbf{e}_2 + \partial_3\mathbf{e}_3, \\ \mathcal{A}_1 &= A_0\mathbf{e}_0 + A_1\mathbf{e}_1 + A_2\mathbf{e}_2 + A_3\mathbf{e}_3,\end{aligned}\tag{9}$$

where  $\mathcal{A}_0$  and  $\mathcal{A}_1$  are elements of  $\mathbf{R}_3$ . The coefficients of these elements be partial derivatives and components of vector-potential, respectively.

Make up now the exterior product of elements (9):

$$\begin{aligned}\mathcal{A}_0\mathcal{A}_1 &= (\partial_0\mathbf{e}_0 + \partial_1\mathbf{e}_1 + \partial_2\mathbf{e}_2 + \partial_3\mathbf{e}_3)(A_0\mathbf{e}_0 + A_1\mathbf{e}_1 + A_2\mathbf{e}_2 + A_3\mathbf{e}_3) = \\ &= \underbrace{(\partial_0A_0 + \partial_1A_1 + \partial_2A_2 + \partial_3A_3)}_{E_0}\mathbf{e}_0 + \underbrace{(\partial_0A_1 + \partial_1A_0)}_{E_1}\mathbf{e}_0\mathbf{e}_1 + \\ &\quad \underbrace{(\partial_0A_2 + \partial_2A_0)}_{E_2}\mathbf{e}_0\mathbf{e}_2 + \underbrace{(\partial_0A_3 + \partial_3A_0)}_{E_3}\mathbf{e}_0\mathbf{e}_3 + \underbrace{(\partial_2A_3 - \partial_3A_2)}_{H_1}\mathbf{e}_2\mathbf{e}_3 + \\ &\quad + \underbrace{(\partial_3A_1 - \partial_1A_3)}_{H_2}\mathbf{e}_3\mathbf{e}_1 + \underbrace{(\partial_1A_2 - \partial_2A_1)}_{H_3}\mathbf{e}_1\mathbf{e}_2.\end{aligned}\tag{10}$$

The scalar part  $E_0 \equiv 0$ , since the first bracket in (10) be a Lorentz condition  $\partial_0A_0 + \text{div}\mathbf{A} = 0$ . It is easily seen that the other bracket be components of electric and magnetic fields:  $-E_i = -(\partial_iA_0 + \partial_0A_i)$ ,  $H_i = (\text{curl}\mathbf{A})_i$ .

Since  $\omega = \mathbf{e}_{123}$  is belong to the center of  $\mathbf{R}_3$ , then

$$\omega\mathbf{e}_1 = \mathbf{e}_1\omega = \mathbf{e}_2\mathbf{e}_3, \quad \omega\mathbf{e}_2 = \mathbf{e}_2\omega = \mathbf{e}_3\mathbf{e}_1, \quad \omega\mathbf{e}_3 = \mathbf{e}_3\omega = \mathbf{e}_1\mathbf{e}_2.$$

In accordance with these correlations may be written (10) as

$$\mathcal{A}_0\mathcal{A}_1 = (E_1 + \omega H_1)\mathbf{e}_1 + (E_2 + \omega H_2)\mathbf{e}_2 + (E_3 + \omega H_3)\mathbf{e}_3\tag{11}$$

It is obvious that the expression (11) is coincide with the vector part of complex quaternion (hyperbolic biquaternion) when  $\mathbf{e}'_1 = i\mathbf{e}_1$ ,  $\mathbf{e}'_2 = i\mathbf{e}_2$ ,  $\mathbf{e}'_3 = i\mathbf{e}_1i\mathbf{e}_2$ .

Further on, make up the exterior product  $\nabla\mathbf{F}$ , where  $\nabla$  is the first element from (9) and  $\mathbf{F}$  is an expression of type (10):

$$\begin{aligned}\nabla\mathbf{F} &= \text{div}\mathbf{E}\mathbf{e}_0 - ((\text{curl}\mathbf{H})_1 - \partial_0E_1)\mathbf{e}_1 - ((\text{curl}\mathbf{H})_2 - \partial_0E_2)\mathbf{e}_2 - \\ &\quad - ((\text{curl}\mathbf{H})_3 - \partial_0E_3)\mathbf{e}_3 + ((\text{curl}\mathbf{E})_1 + \partial_0H_1)\mathbf{e}_2\mathbf{e}_3 + ((\text{curl}\mathbf{E})_2 + \partial_0H_2)\mathbf{e}_3\mathbf{e}_1 + \\ &\quad + ((\text{curl}\mathbf{E})_3 + \partial_0H_3)\mathbf{e}_1\mathbf{e}_2 + \text{div}\mathbf{H}\mathbf{e}_1\mathbf{e}_2\mathbf{e}_3.\end{aligned}\tag{12}$$

It is easily seen that the first coefficient of the product  $\nabla\mathbf{F}$  be a left part of equation  $\text{div}\mathbf{E} = \varrho$ . The following three coefficients are make up a left part of equation  $\text{curl}\mathbf{H} - \partial_0\mathbf{E} = \mathbf{j}$ , the other coefficients are make up the equations  $\text{curl}\mathbf{E} + \partial_0\mathbf{H} = 0$  and  $\text{div}\mathbf{H} = 0$ , respectively (for more details see [7]).

## 4 MODULO 2 PERIODICITY OF COMPLEX CLIFFORD ALGEBRAS

In dependence from the sign of the square of volume element  $\omega$  all totality of Clifford algebras is divided into two classes. Namely, if  $\omega^2 = 1$  Clifford algebra  ${}^l\mathbf{K}_n$  over a field  $\mathbf{K}$  ( $\mathbf{K} = \mathbf{R}, \mathbf{K} = \mathbf{\Omega}, \mathbf{K} = \mathbf{C}$ ) is called positive, and negative in contrary case ( $\omega^2 = -1$ ). Further on, the each algebra  ${}^l\mathbf{K}_n$  is associated to a vector space  $V$  over a field  $\mathbf{K}$ . This space is endowed with a nondegenerate quadratic form  $Q$ . It is obvious that the dimensionality  $\dim V$  of vector space is equal to a number of units of the algebra  ${}^l\mathbf{K}_n$ . In accordance with this most general definition of Clifford algebra we shall denote this algebra as  ${}^l\mathbf{K}_n(V, Q)$ .

**Theorem 1 (Karoubi [8])** 1) If  ${}^l\mathbf{K}_{n_1}(V_1, Q_1)$  is positive, and if  $\dim V_1$  is even, then

$${}^{l_1+l_2}\mathbf{K}_{n_1+n_2}(V_1 \oplus V_2, Q_1 \oplus Q_2) \cong {}^{l_1}\mathbf{K}_{n_1}(V_1, Q_1) \otimes {}^{l_2}\mathbf{K}_{n_2}(V_2, Q_2).$$

2) If  ${}^l\mathbf{K}_{n_1}$  is negative, and if  $\dim V_1$  is even, then

$${}^{l_1+l_2}\mathbf{K}_{n_1+n_2}(V_1 \oplus V_2, Q_1 \oplus Q_2) \cong {}^{l_1}\mathbf{K}_{n_1}(V_1, Q_1) \otimes {}^{l_2}\mathbf{K}_{n_2}(V_2, -Q_2).$$

**Theorem 2** The algebra  ${}^{l+m}\mathbf{K}_{n+2m}$  is isomorphic to  $M_{2^m}({}^l\mathbf{K}_n)$ .

Indeed, it is well-known that the algebra  ${}^l\mathbf{K}_n$  with even dimensionality ( $n = 2\nu$ ) is isomorphic to a matrix algebra  $M_{2^\nu}(\mathbf{K})$ . Further on, since  ${}^1\mathbf{K}_2$  ( ${}^1\mathbf{R}_2, {}^1\mathbf{\Omega}_2, {}^1\mathbf{C}_2$ ) is positive, then in accordance with theorem 1 we obtain

$${}^{l+1}\mathbf{K}_{n+2} \cong {}^l\mathbf{K}_n \otimes {}^1\mathbf{K}_2 \cong \left\{ \begin{array}{l} {}^l\mathbf{R}_n \otimes {}^1\mathbf{R}_2 \\ {}^l\mathbf{\Omega}_n \otimes {}^1\mathbf{\Omega}_2 \\ {}^l\mathbf{C}_n \otimes {}^1\mathbf{C}_2 \end{array} \right\} \cong \left\{ \begin{array}{l} {}^l\mathbf{R}_n \otimes M_2(\mathbf{R}) \\ {}^l\mathbf{\Omega}_n \otimes M_2(\mathbf{\Omega}) \\ {}^l\mathbf{C}_n \otimes M_2(\mathbf{C}) \end{array} \right\} \cong \left\{ \begin{array}{l} M_2({}^l\mathbf{R}_n) \\ M_2({}^l\mathbf{\Omega}_n) \\ M_2({}^l\mathbf{C}_n) \end{array} \right\}$$

Therefore,

$${}^{l+m}\mathbf{K}_{n+2m} \cong {}^l\mathbf{K}_n \otimes M_2(\mathbf{K}) \otimes \dots \otimes M_2(\mathbf{K}) \cong {}^l\mathbf{K}_n \otimes M_{2^m}(\mathbf{K}) \cong M_{2^m}({}^l\mathbf{K}_n). \quad (13)$$

**Theorem 3** The algebra  $\mathbf{C}_{n+2}$  is isomorphic to  $M_2(\mathbf{C}_n)$ .

It is obvious that in the case of a field  $\mathbf{K} = \mathbf{C}$  we have an isomorphism  ${}^l\mathbf{C}_n \cong \mathbf{C}_n$ . Further on, in accordance with theorem 1 we obtain  $\mathbf{C}_{n+2} \cong \mathbf{C}_n \otimes \mathbf{C}_2$  (the algebra  $\mathbf{C}_2$  is positive if  $\omega = i\mathbf{e}_{12}, \mathbf{e}_1^2 = \mathbf{e}_2^2 = 1$  or if  $\omega = \mathbf{e}_{12}, \mathbf{e}_1^2 = 1, \mathbf{e}_2^2 = -1$ ). Since  $\mathbf{C}_2 \cong M_2(\mathbf{C})$ , hence it follows that

$$\mathbf{C}_{n+2} \cong \mathbf{C}_n \otimes \mathbf{C}_2 \cong \mathbf{C}_n \otimes M_2(\mathbf{C}) \cong M_2(\mathbf{C}_n). \quad (14)$$

From adduced above theorems immediately follows that for any Clifford algebra with even dimensionality ( $n = 2\nu$ ) over a field  $\mathbf{C}$  there exists a decomposition :

$$\mathbf{C}_{2\nu} \cong \underbrace{\mathbf{C}_2 \otimes \mathbf{C}_2 \otimes \dots \otimes \mathbf{C}_2}_{\nu \text{ times}}. \quad (15)$$

In case of odd dimensionality ( $n = 2\nu + 1$ ) the algebra  $\mathbf{C}_{2\nu+1}$  is decomposed to a direct sum of two subalgebras which are isomorphic to  $\mathbf{C}_{2\nu}$ . This decomposition is realized by means of mutually orthogonal idempotents

$$\varepsilon = \frac{1}{2}(1 + \omega), \quad \varepsilon' = \frac{1}{2}(1 - \omega),$$

where  $\omega = \mathbf{e}_{12\dots 2\nu+1}$  and  $\omega^2 = 1$ ; or

$$\varepsilon = \frac{1}{2}(1 + i\omega), \quad \varepsilon' = \frac{1}{2}(1 - i\omega),$$

where  $\omega = \mathbf{e}_{12\dots 2\nu+1}$  and  $\omega^2 = -1$ . It is easily seen that  $\varepsilon + \varepsilon' = 1$ ,  $\varepsilon\varepsilon' = 0$ ,  $\varepsilon^2 = \varepsilon$ ,  $\varepsilon'^2 = \varepsilon'$ . Since in this case volume element  $\omega = \mathbf{e}_{12\dots 2\nu+1}$  is belong to a center of  $\mathbf{C}_{2\nu+1}$ , then the idempotents  $\varepsilon$ ,  $\varepsilon'$  are commutes with all elements of the algebra  $\mathbf{C}_{2\nu+1}$ . Hence it follows that [9]

$$\mathbf{C}_{2\nu+1} \cong \mathbf{C}_{2\nu} \oplus \mathbf{C}_{2\nu},$$

where the each algebra  $\mathbf{C}_{2\nu}$  consist of the elements of type  $\varepsilon\mathcal{A}$  or  $\varepsilon'\mathcal{A}$ , here  $\mathcal{A} \in \mathbf{C}_{2\nu+1}$ . Therefore, in accordance with (15) we have

$$\mathbf{C}_{2\nu+1} \cong \mathbf{C}_2 \otimes \mathbf{C}_2 \otimes \dots \otimes \mathbf{C}_2 \oplus \mathbf{C}_2 \otimes \mathbf{C}_2 \otimes \dots \otimes \mathbf{C}_2. \quad (16)$$

Further on, by force of  $\mathbf{C}_2 \cong \mathbf{M}_2(\mathbf{C})$  we have for a base of spinor representation of  $\mathbf{C}_2$  the following matrices which are correspond to the units  $\mathbf{e}_1$  and  $\mathbf{e}_2$ :

$$E_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad E_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}. \quad (17)$$

Let  $\mathbf{e}_3 = \mathbf{e}_1\mathbf{e}_2$ , then for a vector part of biquaternion (13) we obtain by means of (17) the following matrix:

$$\begin{bmatrix} iF_3 & F_1 - iF_2 \\ F_1 + iF_2 & -iF_3 \end{bmatrix}, \quad (18)$$

here  $F_i = E_i + iH_i$ , since in this case  $\omega \equiv i$ .

This way, *the algebra  $\mathbf{C}_2$  with general element  $\mathcal{A} = F_0\mathbf{e}_0 + F_1\mathbf{e}_1 + F_2\mathbf{e}_2 + F_3\mathbf{e}_3$  is isomorphic to a matrix algebra of type (18), where  $F_0 = \partial_0 A_0 + \text{div}\mathbf{A} \equiv 0$  and  $F_i$  ( $i = 1, 2, 3$ ) are components of a complex electromagnetic field.* We shall denote this matrix algebra as  $\mathbf{M}_2^F(\mathbf{C})$ .

Hence by force of modulo 2 periodicity of complex Clifford algebras from (15) and (16) immediately follows that

$$\mathbf{C}_{2\nu} \cong \mathbf{M}_{2\nu}^F(\mathbf{C}) \cong \mathbf{M}_2^F(\mathbf{C}) \otimes \mathbf{M}_2^F(\mathbf{C}) \otimes \dots \otimes \mathbf{M}_2^F(\mathbf{C}), \quad (19)$$

$$\begin{aligned} \mathbf{C}_{2\nu+1} &\cong \mathbf{C}_{2\nu} \oplus \mathbf{C}_{2\nu} \cong \mathbf{M}_{2\nu}^F(\mathbf{C}) \oplus \mathbf{M}_{2\nu}^F(\mathbf{C}) \cong \\ &\cong \mathbf{M}_2^F(\mathbf{C}) \otimes \mathbf{M}_2^F(\mathbf{C}) \otimes \dots \otimes \mathbf{M}_2^F(\mathbf{C}) \oplus \mathbf{M}_2^F(\mathbf{C}) \otimes \mathbf{M}_2^F(\mathbf{C}) \otimes \dots \otimes \mathbf{M}_2^F(\mathbf{C}). \end{aligned} \quad (20)$$

For example, consider now the algebra  $\mathbf{C}_4$ . In the spinor representation this algebra is isomorphic to a matrix algebra  $\mathbf{M}_4(\mathbf{C})$ , the base of which consist of well-known  $\gamma$ -matrices. In the base of Weyl for these matrices we have

$$\gamma^m = \begin{bmatrix} 0 & \sigma^m \\ \bar{\sigma}^m & 0 \end{bmatrix}, \quad (21)$$

where  $m = 0, 1, 2, 3$  and

$$\begin{aligned} \sigma^0 &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \sigma^1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma^2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma^3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \\ \bar{\sigma}^0 &= \sigma^0, \quad \bar{\sigma}^{1,2,3} = -\sigma^{1,2,3}. \end{aligned}$$

It is easily seen that  $\sigma$ -matrices are make up the base of spinor representation of  $\mathbf{C}_2 \cong \mathbf{M}_2(\mathbf{C})$  and are coincide with matrices  $E_i$  if  $\sigma^3 = \frac{1}{i}E_1E_2$ .

For the algebra  $\mathbf{C}_4$  in accordance with (19) we have

$$\mathbf{C}_4 \cong \mathbf{C}_2 \otimes \mathbf{C}_2 \cong \mathbf{M}_2^F(\mathbf{C}) \otimes \mathbf{M}_2^F(\mathbf{C}).$$

Further on, by general definition, the algebra  $\mathbf{C}_4$  is associated to a complex vector space  $C_4$ . For the vector  $\mathbf{a} = F_i \mathbf{e}_i \in C_4$  in the base (21) we have a matrix  $\mathbf{A} = F_i \gamma^i$ :

$$\begin{bmatrix} 0 & 0 & F_3 & F_1 - iF_2 \\ 0 & 0 & F_1 + iF_2 & -F_3 \\ -F_3 & -F_1 + iF_2 & 0 & 0 \\ -F_1 - iF_2 & F_3 & 0 & 0 \end{bmatrix}.$$

The all matrix algebra  $\mathbf{M}_4^F(\mathbf{C})$  in the base (17) we obtain by means of theorem 3. Indeed, in the case when dimensionality of  ${}^l\mathbf{K}_n$  is even the volume element  $\omega = \mathbf{e}_{12\dots n}$  is not belong to a center of algebra  ${}^l\mathbf{K}_n$ .

However, when  $i \leq 2m$  we have

$$\begin{aligned} \mathbf{e}_{12\dots 2m2m+k} \mathbf{e}_i &= (-1)^{2m+1-i} \sigma(i-l) \mathbf{e}_{12\dots i-1i+1\dots 2m2m+k}, \\ \mathbf{e}_i \mathbf{e}_{12\dots 2m2m+k} &= (-1)^{i-1} \sigma(i-l) \mathbf{e}_{12\dots i-1i+1\dots 2m2m+k} \end{aligned}$$

and, therefore, the condition of commutativity of elements  $\mathbf{e}_{12\dots 2m2m+k}$  and  $\mathbf{e}_i$  is  $2m+1-i \equiv i-1 \pmod{2}$ . Thus, the elements  $\mathbf{e}_{12\dots 2m2m+1}$  and  $\mathbf{e}_{12\dots 2m2m+2}$  are



commute with all elements  $\mathbf{e}_i$  whose indexes are not exceed  $2m$ . Therefore, the transition from algebra  $\mathbf{K}_{2m}$  to algebras  $\mathbf{K}_{2m+2}$ ,  ${}^1\mathbf{K}_{2m+2}$  or  ${}^2\mathbf{K}_{2m+2}$  and from algebra  ${}^l\mathbf{K}_{2m}$  to  ${}^l\mathbf{K}_{2m+2}$ ,  ${}^{l+1}\mathbf{K}_{2m+2}$  or  ${}^{l+2}\mathbf{K}_{2m+2}$  may be represented as transition from the real (complex or double) coordinates of elements of algebras  $\mathbf{K}_{2m}$  and  ${}^l\mathbf{K}_{2m}$  to the coordinates of type  $a + b\phi + c\psi + d\phi\psi$ , where  $\phi$  and  $\psi$  are additional basis elements  $\mathbf{e}_{12\dots 2m2m+1}$  and  $\mathbf{e}_{12\dots 2m2m+2}$ . The elements  $\mathbf{e}_{i_1 i_2 \dots i_k} \phi$  are contain index  $2m+1$  and not contain index  $2m+2$ , and the elements  $\mathbf{e}_{i_1 i_2 \dots i_k} \psi$  are contain index  $2m+2$  and not contain index  $2m+1$ . Respectively, the elements  $\mathbf{e}_{i_1 i_2 \dots i_k} \phi\psi$  are contain both indexes  $2m+1$  and  $2m+2$ . Hence it immediately follows that a general element of  ${}^l\mathbf{K}_{2m+2}$  or  ${}^{l+1}\mathbf{K}_{2m+2}$ ,  ${}^{l+2}\mathbf{K}_{2m+2}$ ,  $\mathbf{K}_{2m+2}$  may be represented as

$${}^l\mathbf{K}_{2m}^0 \mathbf{e}_0 + {}^l\mathbf{K}_{2m}^1 \mathbf{e}_{12\dots 2m2m+1} + {}^l\mathbf{K}_{2m}^2 \mathbf{e}_{12\dots 2m2m+2} + {}^l\mathbf{K}_{2m}^3 \mathbf{e}_{2m+12m+2}, \quad (22)$$

where  ${}^l\mathbf{K}_{2m}^i$  ( $i = 0, 1, 2, 3$ ) are represent the algebras with general element  $\mathcal{A} = \sum_{k=0}^{2m} a^{i_1 i_2 \dots i_k} \mathbf{e}_{i_1 i_2 \dots i_k}$ . When the elements  $\phi = \mathbf{e}_{12\dots 2m2m+1}$  and  $\psi = \mathbf{e}_{12\dots 2m2m+2}$  are satisfy to condition  $\phi^2 = \psi^2 = -1$  we see that the basis  $\{\mathbf{e}_0, \phi, \psi, \phi\psi\}$  is isomorphic to a basis of quaternion algebra. Therefore, in this case from (22) we have

$${}^l\mathbf{K}_{2m+2} \cong M_2({}^l\mathbf{K}_{2m}) = \begin{bmatrix} {}^l\mathbf{K}_{2m}^0 & -i{}^l\mathbf{K}_{2m}^3 & -{}^l\mathbf{K}_{2m}^1 + i{}^l\mathbf{K}_{2m}^2 \\ {}^l\mathbf{K}_{2m}^1 + i{}^l\mathbf{K}_{2m}^2 & {}^l\mathbf{K}_{2m}^0 + i{}^l\mathbf{K}_{2m}^3 \end{bmatrix}.$$

Analogously, when the elements  $\phi$  and  $\psi$  are satisfy to conditions  $\phi^2 = \psi^2 = 1$  or  $\phi^2 = 1, \psi^2 = -1$  we have the transitions from the real (complex or double) coordinates in  ${}^l\mathbf{K}_{2m+2}$  to the anti-quaternionic and pseudo-quaternionic coordinates, respectively (about anti-quaternions and pseudo-quaternions see [10, p. 434]).

In our case by force of isomorphism  ${}^l\mathbf{C}_n \cong \mathbf{C}_n$  we see that any algebra  ${}^l\mathbf{C}_4 \cong \mathbf{C}_4$  ( $l = 0, 1, 2, 3, 4$ ) may be represented by the following quaternion:

$$\mathbf{C}_2^0 \mathbf{e}_0 + \mathbf{C}_2^1 \phi + \mathbf{C}_2^2 \psi + \mathbf{C}_2^3 \phi\psi,$$

where  $\phi = \mathbf{e}_{123}$ ,  $\psi = \mathbf{e}_{124}$  and  $\mathbf{C}_2^i$  the algebras of hyperbolic biquaternions with general element (in our case)  $\mathcal{A} = F^0 \mathbf{e}_0 + F^1 \mathbf{e}_1 + F^2 \mathbf{e}_2 + F^3 \mathbf{e}_{12}$ . In spinor representation for the quaternion basis  $\{\mathbf{e}_0, \phi, \psi, \phi\psi\}$  we have  $\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}, \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix} \right\}$ . Hence it immediately follows that

$$\mathbf{C}_4 \cong \mathbf{C}_2 \otimes \mathbf{C}_2 \cong \mathbf{C}_2 \otimes M_2(\mathbf{C}) \cong M_2(\mathbf{C}_2) = \begin{bmatrix} \mathbf{C}_2^0 - i\mathbf{C}_2^3 & -\mathbf{C}_2^1 + i\mathbf{C}_2^2 \\ \mathbf{C}_2^1 + i\mathbf{C}_2^2 & \mathbf{C}_2^0 + i\mathbf{C}_2^3 \end{bmatrix}.$$

Finally, using (18) we obtain that  $M_4^F(\mathbf{C})$  has a form:

$$\begin{bmatrix} (1+i)F_3 & (1-i)F_1 - (1+i)F_2 & -(1+i)F_3 & -(1-i)F_1 + (1+i)F_2 \\ (1-i)F_1 + (1+i)F_2 & -(1+i)F_3 & -(1-i)F_1 - (1+i)F_2 & (1+i)F_3 \\ -(1-i)F_3 & (1+i)F_1 + (1-i)F_2 & -(1-i)F_3 & (1+i)F_1 + (1-i)F_2 \\ (1+i)F_1 - (1-i)F_2 & (1-i)F_3 & (1+i)F_1 - (1-i)F_2 & (1+i)F_3 \end{bmatrix} \quad (23)$$

Analogously, for the algebra  $\mathbf{C}_6 \cong M_8^F(\mathbf{C})$  we have the anti-quaternion  $\mathbf{C}_4^0 \mathbf{e}_0 + \mathbf{C}_4^1 \phi + \mathbf{C}_4^2 \psi + \mathbf{C}_4^3 \phi\psi$ , where  $\phi = \mathbf{e}_{12345}$ ,  $\psi = \mathbf{e}_{12346}$  and  $\phi^2 = \psi^2 = 1$ . The anti-quaternion basis  $\{\mathbf{e}_0, \phi, \psi, \phi\psi\}$  in spinor representation defined by the following matrices  $\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix} \right\}$ . Using (23) we have

$$2 \begin{bmatrix} iF_3 & F_1 - iF_2 & -iF_3 & -F_1 + iF_2 & F_3 & -F_2 - iF_1 & -F_3 & F_2 + iF_1 \\ F_1 + iF_2 & -iF_3 & -F_1 - iF_2 & iF_3 & F_2 - iF_1 & -F_3 & F_2 + iF_1 & F_3 \\ -F_3 & F_2 + iF_1 & -F_3 & F_2 + iF_1 & iF_3 & F_1 - iF_2 & iF_3 & F_1 - iF_2 \\ -F_1 + iF_1 & F_3 & -F_2 + iF_1 & F_3 & F_1 + iF_2 & -iF_3 & F_1 + iF_2 & -iF_3 \\ iF_3 & F_1 - iF_2 & -iF_3 & -F_1 + iF_2 & F_3 & -F_2 - iF_1 & -F_3 & F_2 + iF_1 \\ F_1 + iF_2 & -iF_3 & -F_1 - iF_2 & iF_3 & F_2 - iF_1 & -F_3 & F_2 + iF_1 & F_3 \\ -F_3 & F_2 + iF_1 & -F_3 & F_2 + iF_1 & iF_3 & F_1 - iF_2 & iF_3 & F_1 - iF_2 \\ -F_1 + iF_1 & F_3 & -F_2 + iF_1 & F_3 & F_1 + iF_2 & -iF_3 & F_1 + iF_2 & -iF_3 \end{bmatrix}.$$

For the algebras with odd dimensionality from (20) we obtain  $\mathbf{C}_5 \cong M_4^F(\mathbf{C}) \oplus M_4^F(\mathbf{C})$ ,  $\mathbf{C}_7 \cong M_8^F(\mathbf{C}) \oplus M_8^F(\mathbf{C})$  and so on.

## References

- [1] W. K. Clifford, *Applications of Grassmann's extensive algebra*, Am. J. Math. **1**, 350 (1878).
- [2] R. Lipschitz, **Untersuchungen über die Summen von Quadraten** (Bonn, 1886).
- [3] N. Salingaros, *Algebras with three anticommuting elements. II. Two algebras over a singular field*, J. Math. Phys. **22**, 2096 (1981).
- [4] R. Brauer and H. Weyl, *Spinors in  $n$  dimensions*, Am. J. Math. **57**, 425 (1935).
- [5] M. F. Atiyah, R. Bott, and A. Shapiro, *Clifford modules*, Topology **3**, (suppl.1), 3 (1964).
- [6] R. Coquereaux, *Modulo 8 periodicity of real Clifford algebras and particle physics*, Phys. Lett. B **115**, 389 (1982).
- [7] V. V. Varlamov, *Physical fields and Clifford algebras*, hep-th/9709051.
- [8] M. Karoubi, **K-Theory** (Springer-Verlag, Berlin, 1979).
- [9] P. K. Rashevskii, *The theory of spinors*, Am. Math. Soc. Transl. (Ser.2) **6**, 1 (1957).
- [10] B. A. Rozenfeld, **Neevklidovi geometrii** (Moscow, 1955).