Comment on "Understanding the Area Proposal for Extremal Black Hole Entropy"

O. B. Zaslavskii

Department of Physics, Kharkov State University, Svobody Sq.4, Kharkov 310077, Ukraine

E-mail: olegzasl@apt.kharkov.ua

A. Ghosh and P. Mitra made the proposal how to explain the area law for the entropy of extreme black holes in some model calculations. I argue that their approach implicitly operates with strongly singular geometries and says nothing about the contribution of regular metrics of extreme black holes into the partition function.

 $04.70. \mathrm{Dy}$

In the recent paper [1] the attempt has been made to explain the area law for extreme black holes. The authors write down the partition function with account for topologies of either extreme or nonextreme black holes: $Z = \sum_{\text{topologies}} \int d\mu(m) \int d\mu(q) e^{-I(m,q)}$ where m is a mass, q is an electric charge. They point out that the action I has different form for nonextreme (I_n) and extreme (I_e) cases:

$$I_n = \beta(m - q\Phi) - \pi(m + \sqrt{m^2 - q^2})^2, I_e = \beta(m - q\Phi)$$
 (1)

where β is the inverse temperature on a boundary (which is supposed in [1] to be situated at infinity), Φ is the difference of potentials between a boundary and horizon. The next step in derivation the area law in [1] is crucial. It consists in the statement that "the nonextremal action is lower than the extremal one for each set of values of q, m" (italic is mine - O.Z.) In other words, for fixed β and Φ

$$I_n(m,q) < I_e(m,q) \tag{2}$$

From eq.(2) authors conclude that it is the nonextreme topology which mainly contributes to Z whence they "explain" the area law for the entropy.

However, eq.(2) implies that m and q in both sides of it are the same. As for nonextreme black holes $m \neq q$ it must hold also for solutions called in [1] "extreme" and whose action is calculated above with the zero entropy term. As I show below this leads to singular behavior of the metric. The metric of a spherically-symmetrical charged black hole can be written as

$$ds^2 = b^2 d\tau^2 + \alpha^2 dr^2 + r^2 d\Omega^2 \tag{3}$$

It follows from the Gauss law and Hamiltonian constraint [2] that

$$\alpha^2 = \left(1 - \frac{2m}{r} + \frac{q^2}{r^2}\right)^{-1} \tag{4}$$

If $m \neq q$ this entails the asymptotics $\alpha^2 \propto (r - r_+)^{-1}$ near an event horizon at $r = r_+$. On the other hand, to distinguish extreme and nonextreme topologies it was used in [1] that $\xi \equiv \lim_{r \to r_+} b'/\alpha = 0$ in the first case in contrast to $\xi = 1$ in the second one. It means that instead of $b^2 \propto (r - r_+)$ typical of nonextreme holes this coefficient behaves in the extreme case as $b^2 \propto (r - r_+)^{1+n}$ with n > 0.

However, both asymptotics $\alpha^2 \propto (r-r_+)^{-1}$ and $b^2 \propto (r-r_+)^{1+n}$ are inconsistent with each other in that the Riemann curvature diverges strongly at the horizon either in Euclidean or Lorentzian version: $R \propto l^{-2}$ where l is a proper distance from the horizon. Although integrals from R entering into the action converge, such behavior of geometry looks unphysical. Moreover, all other curvature invariant diverge as well. For example, $R_{\mu\nu\sigma\rho}R^{\mu\nu\sigma\rho} \propto l^{-4}$. In other words, the "extreme" solutions discussed in [1] correspond to spacetimes for which the whole surface $r=r_+$ consists of singular points. It follows from the above discussion that solutions at hand do not compose some particular, very special class of configurations which may or may not be used in Z at one's will. On the contrary, the properties described above hold true for any solution with $m \neq q$ and $\xi = 0$. One may ask, whether all that means that the result of [1] consists in the proof that the true physical equilibrium cannot be achieved at singular geometries discussed above. This would restrict the significance of the result since it gives no information about the relative contribution of regular extreme topologies but, at least, it would mean that it can be given reasonable

interpretation. Unfortunately, even in this restricted sense the "proof" presented in [1] is incorrect as is argued below.

Let me recall the main points which prescription for the entropy S=0 and temperature $(\beta \text{ is arbitrary})$ of extreme black holes are based on [3]: (i) the condition of the regularity of Euclidean geometry which is satisfied for any β (for nonextreme holes (i) is the same but is satisfied by the only choice of β); (ii) topological meanings, in which the crucial role is played by the property $l=\infty$ between a horizon and any other point outside it. Both (i) and (ii) are not satisfied by "extreme" black holes with $m \neq q$. As far as (i) is concerned, the situation is completely reverse to that in [3]: whereas the Euclidean geometry in [3] is regular for any β , the Euclidean geometry (as well Lorentzian one) in [1] is singular for any β . As regards (ii), l is finite for any $m \neq q$. In view of motivation for S=0 fails for solutions under discussion, the eq.(2) loses its meaning at all (even if singular solutions are admitted) that destroys the "proof" completely. I stress that this takes place irrespectively of how small the difference m-q is.

The crucial point in which the analysis of [1] flaws consists in dealing with metrics for which simultaneously $\xi = 0$ and $m \neq q$. As a result, "explanation" of the area proposal for extreme black holes in [1] is unsound.

The precedent text is published as Comment [4]. Below I give a brief response to Ghosh and Mitra's Reply [5].

1) Ghosh and Mitra state that Eq. (1) "was neither written nor used in [1]". This inequality was not, indeed, written explicitly in [1] as a separate formula (probably, because of its simplicitly). However, it was not only used by Ghosh and Mitra - it is the key point in their "proof": "It is clear from (7) that the nonextremal action is lower than the extremal one for each set of values of q, m. Consequently, the partition function is to be approximated by $e^{-I_{q,m}}(...)$ " (The paragraph after Eq. (8) in [1]). Here $I_{q,m} = I_n(m,q)$ in my notations, Eq. (7) of [1] corresponds to 1 above. As a matter of fact, Ghosh and Mitra repeat (2) in Reply in a slightly different form:

$$I_n(m, q - \varepsilon) < I_e(m, q) \tag{5}$$

with small but nonzero ε . In fact, however, the quantity I_n is the continuous function of its arguments, so one may take the limit $\varepsilon \to 0$ safely and there is no essential difference between (5) and (1). Therefore, all arguments pushed forward as regards 2 retain their validity with respect to 5. That charges in both sides of (5) may slightly different from each other, is not crucial. Much more important is that in the r.h.s. of (5) m and q may be different and this leads to difficulties indicated in Comment.

- 2) It is demonstrated explicitly in [4] that every configuration among those discussed in [1] leads to a singular geometry. Authors of [5] keep silence with respect to the corresponding arguments, thus continuing to state that my objection concerns only some particular classes of metrics (calling the crucial point "the technical observation") without any refutation of the claim made in [4] about the generality of the property under discussion. Instead of it, Ghosh and Mitra emphasize: "these configurations were not explicitly used by us, nor do they need to be used implicitly" as if one might discard at any moment some metrics (in fact, all of them with $m \neq q$ in our case) from the path integral at his own will.
- 3) Inasmuch as the action integral converges, the singular behavior of the metrics under discussion does not bother authors of [1], [5] Such a liberal attitude to singularities conflicts with what is implied to be done with them. For instance, in the conical singularity approach for nonextreme black holes the action integral certainly converges since the singularity even is weaker than that in [1] (delta-like instead of power-like). In spite of it, this singularity is not accepted physically. For higher-curvature Lagrangians the situation becomes worse since in that case even the action integral would diverge under conditions considered in [1]. It is worth noting that the demand of regularity near the horizon imposes general restrictions on a black hole metric [6]. Metrics considered in [1] do not obey the corresponding conditions.

The last claim of [5] that the action integral is calculated directly without any additional assumptions seems to be correct. However, again, such a strange object as an "extreme" black hole with a finite proper distance between any point and the "horizon" and with the

entropy S = 0 is gained due to dropping such a fundamental condition as the regularity of a manifold and, therefore, hardly has any physical sense.

In general, three different topological sectors are involved into competition for given boundary data (β, Φ) : a flat geometry, regular nonextreme metrics with $m \neq q$ and regular extreme ones with m = q. The paper [1] says nothing, however, about their relative contribution and deals with singular "extreme" configurations for which $m \neq q$.

Let me conclude with some general remarks beyond the issues discussed in [4] and [5]. It is quite possible that any estimate of different fractional contributions to the partition function made in the zero- loop approximation (even for regular geometries) will be unphysical. The point is that the temperature of extreme black holes, according to [3], is a finite nonzero quantity, i.e. it differs from the Hawking one. This assumption leads to the divergencies of the stress-energy tensor on an event horizon of a generic extreme black hole [7]. Although this result is obtained in the semiclassical approximation whereas the geometry in the situation considered in [1] fluctuates itself, it is very likely that path integral over geometries each of which is singular itself will diverge. In other words, quantum effects can change the semiclassical thermodynamics of extreme black holes drastically.

- [1] A. Ghosh and P. Mitra, Phys. Rev. Lett. 78, 1858 (1997).
- [2] H.W. Braden, J.D. Brown, B.F. Whiting, and J.W. York, Jr., Phys. Rev. D 42, 3376 (1990).
- [3] S. Hawking, G. Horowitz, and S. Ross, Phys.Rev. D 51, 4302 (1995).
- [4] O. B. Zaslavskii, Phys. Rev. Lett. **80**, 3412 (1998).
- [5] A. Ghosh and P. Mitra, Phys. Rev. Lett. **80**, 3413 (1998).
- [6] P. R. Anderson and C. Mull, Constraints on Black Holes in Classical and Semiclassical Theories of Gravity, preprint gr-qc/9707026.

[7] P.A. Anderson, W.A. Hiscock, and D.J. Loranz, Phys. Rev. Lett. ${\bf 74},\,4365$ (1995).