A Course on: "Modular Localization and Nonperturbative Local Quantum Physics" CBPF, Rio de Janeiro, March 1998

Bert Schroer Institut für Theoretische Physik der FU-Berlin, Arnimallee 14, 14195 Berlin Germany. e-mail: schroer@physik.fu-berlin.de

March 7, 2018

Contents

 The Basics of Quantum Theory Multiparticle Wave Functions, Particle Statistics The Superselection Sectors of CG Illustration of Important Quantum Concepts Measurement and Superselection Rules The Construction of Fock Space. The Bosonic Fock Space The Fermion Fock Space The CCR and CAR Algebras. 	· · · · · · · · ·	 17 17 25 32 37 43 43
 1.1 Multiparticle Wave Functions, Particle Statistics	· · · · · · · · ·	17 25 32 37 43 43
 1.2 The Superselection Sectors of CG	· · · · · ·	25 32 37 43 43
 1.3 Illustration of Important Quantum Concepts	· · · · ·	32 37 43 43
1.4 Measurement and Superselection Rules 2 The Construction of Fock Space. 2.1 The Bosonic Fock Space . 2.2 The Fermion Fock Space . 2.3 The CCR and CAR Algebras. 2.4 Quasifree States .	· · · · ·	37 43 43
2 The Construction of Fock Space. 2.1 The Bosonic Fock Space	 	43 43
 2.1 The Bosonic Fock Space	 	43
 2.2 The Fermion Fock Space	 	
2.3 The CCR and CAR Algebras		52
2.4 Quasifree States		54
		57
2.5 Temperature States and KMS condition		59
2.6 The CCR- and CAR-Functors		62
3 Poincaré Symmetry and Quantum Theory		67
3.1 Symmetry in General Quantum Theory	• •	67
3.2 One Particle Representations	• •	71
3.3 Wigner Theory and Free Fields		86
3.4 The Equivalence Class of a Free Field		95
3.5 A First Look at Modular Localization		101
3.6 Special Features of Zero Mass		106
3.7 Exotic Spin and Localization		112
3.8 Localization and Hawking Temperature		114
3.9 Examples of Dis(order)-Fields		121
3.10 Special Features of m=0, d=1+1 Fields $\ldots \ldots \ldots \ldots$		128
3.11 Counting of Localized Degrees of Freedom		136
3.12 Split Property, Taming of Vacuum Fluctuations		139
3.13 Problems with Entropy		141
4 Perturbative Interactions		147
4.1 Kinematical Decompositions		147
4.2 Perturbative Realization of Interaction		150
4.3 Perturbation and Adiabatic Parametrization		100

	$4.4 \\ 4.5 \\ 4.6$	Invariant Parametrizations, Regularization	163 168 177
5	The 5.1 5.2 5.3 5.4	e General Framework of QFT Model-independent Properties of pointlike Fields	 183 183 188 194 202
6	Nor 6.1 6.2 6.3 6.4 6.5	Introductory Comments	 209 216 217 219 222 225 238 240 247
7	Intr 7.1 7.2 7.3 7.4 7.5 7.6 7.7	Production to Algebraic QFT Some Useful Theorems Abstracting Superselection Principles Starting the Reverse: the DHR Endomorphisms Remarks on Broken Symmetries Chiral Conformal Algebraic QFT Classification of Admissable Particle-Statistics Constructive Aspects of Plektons	 263 263 269 272 280 284 289 296
8	Ten	tative Resume and Outlook	301
A	Mat A.1 A.2 A.3 A.4 A.5 A.6 A.7 A.8 A.9 A 10	thematical Appendix Introduction States on C*-algebras and representations Von Neumann Algebras and their Classification Modular Theory C*-Algebras related with Bimodules Conditional Expectations, Canonical Endomorphisms Deep Inclusions and (spacetime) Geometry Shallow Inclusions and (internal) Symmetry Split Property, Localizing Map and Nuclearity Inverse Problem of Modular Theory	 313 313 313 318 321 325 326 329 329 332 334

4

0.1 Introduction

A new text on an apparently old and established subject as QFT should justify and measure itself relative to the many existing review articles and textbooks. The main pragmatic motivation underlying these notes consists in the desire to unify two presently largely disconnected branches of QFT:

(1) the standard (canonical, functional) approach which is mainly perturbative in the sense of an infinitesimal "deformation" on free fields and

(2) nonperturbative constructions of low-dimensional models as in the formfactorbootstrap approach (which for the time being is limited to factorizable models in d=1+1 space-time dimensions) as well as the non-Lagrangian construction of conformal chiral QFT's.

The synthesis requires a significant step beyond the concepts which were used in order to formulate the two mentioned separate branches of QFT. On the physical side, the S-matrix regains some of its early prominence; however unlike in the old proposals of Heisenberg as well as in the later S-matrix bootstrap approach of Chew et al., it remains subservient to the locality and causality principles encapsulated in the theory of local observables which is the heartpiece of algebraic QFT.

In the new context of the TCP- and the Tomita-reflection operators (explained in detail in these notes), the S-matrix takes the role of a powerful constructive tool in QFT. In theories with a mass gap, the validity of scattering theory immediately provides a natural arena for the nonperturbative description: the Hilbert space of incoming particles which for Fermions/Bosons is a Fock space, i.e. an orthogonal sum of multiparticle spaces which have the anti/symmetric tensor product structure in terms of the Wigner one particle spaces. Whereas the connected Poincaré representations of the incoming and interacting fields are identical¹, those interacting reflection operators which contain the time reflections (T, TP, TCP) differ from their free incoming expressions. According to an old observation of Bisognano and Wichmann this implies that the modular data for the algebra of the wedge regions are known: the modular group is the one parametric group of wedge affiliated Lorentz boosts whereas the modular conjugation differs from its incoming value by the S-matrix. The fundamental significance of the wedge region to the basics of QFT will perhaps not be surprising to physicist who are familiar with Unruh's work on the thermal properties of the horizon of the Rindler wedge as the simplest illustration of Hawking's and Bekensteins thermal and entropical properties of black hole horizons. The point in these notes is that the new concepts of modular localization causes the "outing" of this quasiclassical behavior associated to Killing vectors in CST QFT as a generic property of nonperturbative QFT.

At this place it seems to be natural to make some explanatory remarks about the Tomita-Takesaki modular theory. Whereas the detailed technical aspects will be reserved for the mathematical appendix, some intuitive understanding

 $^{^{1}}$ Note that in general the free incoming hamiltonian is different from the unperturbed hamiltonian of standard perturbation theory. Only for special on shell normalization conditions they agree.

can be obtained by looking at its fascinating history. Mathematically it is a vast generalization of the modular factor which accounts for the difference between right and left Haar measure in the case of non unimodular groups as e.g. SL(2,R). Tomita and later Takesaki succeeded to convert this idea into a powerful tool for the investigation of von Neumann algebras. In fact Alain Connes could not have carried out his path-breaking work on the classification of von Neumann factor algebras without this Tomita-Takesaki theory. Modular theory is also behind the subfactor theory of Vaughn Jones.

The physics side is equally impressive. At the time when Tomita presented his theory, Haag, Hugenholz and Winnink [3] published their fundamental work on (heat bath) thermal aspects of QFT. The "KMS" condition (a name which was coined in that paper) was used before by various physicist (in particular Kubo, Martin and Schwinger) as a clever mathematical trick in order to avoid to compute cumbersome traces in evaluating Gibbs thermal ensembles in relativistic theories. In the hands of Haag Hugenholz and Winnink this formula became the key for their formulation of equilibrium quantum statistical mechanics directly in the thermodynamic limit (in which the Gibbs representation becomes meaningless as a consequence of volume divergencies in its vacuum fluctuations). The generator for the modular operator turns out to be a "thermal Hamiltonian" with two-sided spectrum and finite fluctuations. It acts on the original algebra as well as on its commutant which represents a kind of an non-geometric "shadow world" (corresponding in the above analogy to group theory to the opposite action). The modular conjugation J turns out to be the flip operation which maps the left into the right action. Later Bisognano and Wichmann found another very nontrivial field theoretic illustration of the Tomita-Takesaki theory in their study of the algebra which is generated by fields restricted to the wedge region (which, as already mentioned, corresponds to the Rindler region that played an important role in Unruh's discussion of the Hawking effect), briefly called the wedge algebra. In their case the modular operator turned out to be the wedge-affiliated Lorentz boost and the modular conjugation J was (up to a 180 degree rotation around the boost direction) the field theoretical TCP operator. This time the von Neumann commutant was part of the real world, namely the algebra of the causally disjoint region behind the wedge horizon. This work (as well as some special prior observations on free fields [4]) strongly suggested that there was a deep relation between modular theory and relativistic causality and localization. In more recent times Borchers and Wiesbrock as well as Araki and Zsido showed that with two subalgebras in appropriate modular position, one can build up space-time symmetries (d=1+1)conformal and, with somewhat stronger assumptions, also Poincaré covariance).

This set the stage for asking some fundamental questions about nonperturbative QFT. The framework of algebraic QFT attributes less importance to individual fields and prefers nets of algebras in order to describe the physical content of local quantum physics. On the other hand the conventional introduction of free fields via Wigner's particle theory of positive energy representation of the Poincaré group is known to lead to many free fields which belong to the same (m,s) Wigner representation and live in the same Fock space. They are members of one local equivalence (Borchers-) class of fields and are identical in their physical content; though only very few posses a free Lagrangian L_0 , i.e. can be interpreted as resulting from canonical quantization of a classical field theory. The unicity on the field theoretic level is achieved by a modular construction (explained in chapter 3) which produces the net of interaction free theories directly, i.e. without the intervention of point-like fields and in this way leads to a one to one relation of the (m,s) Wigner representation with one (m,s) algebraic net. Therefore these more fundamental concepts leads not only to a de-emphasis of Lagrangians and quantization ideas, but even delegates the very point-like covariant fields, the apparent heart-piece of QFT, to the status of "field coordinates" for the description of local "nets" of algebras. In this way it re-captures the unicity of the Wigner representation theory which was lost by the existence of a multitude of intertwiners relating the unique Wigner representation to the infinitely many possible covariant representations. This new point of view, which is consistent with fields (and hence incorporates QFT), but places them into a much bigger and richer conceptual arena, will be named Local Quantum Physics [3] or LQP. The name QFT is often linked with quantization, actions, functional integrals etc. which do not appear in our nonperturbative approach but only in the chapter 4 on perturbation theory.

This leads to the interesting question of whether such a construction of fieldcoordinate independent approach is feasible in the presence of interactions. It was natural to start again with the wedge situation and the previously mentioned observation that if one replaces the reference to the Wigner one-particle space by the incoming Fock space of scattering theory, then the modular group will still be the L-boost of the free incoming net but the J operator will deviate from the free one by the true scattering matrix.

In setting up a scenario for a nonperturbative construction of nets in interacting QFT's, an important building block is the "modular Møller operator", a kind of square root out of the S-matrix. In the context of factorizing d=1+1 theories this operator is closely related to the physical representation of the Zamolodchikov-Faddeev algebra. There are good reasons to believe (see chapter6) that such operators exists in general QFT and that they are always associated with semi-local fields without vacuum polarization (just like the Z-F operators in x-space). They are somewhat reminiscent of the elusive light cone fields, but different from those they are not dependent on prerequisites of good short distance behavior for the validity of the canonical formalism and their conceptual auxiliary position relative to the local fields and compactly (double cone) localized algebras is very clear. The absence of pair states after applying them to the vacuum requires that they cannot be localized in smaller compact subregions inside the noncompact wedge. The associated physical picture is that the wedge region with its L-boost invariance group still permits to resolve massive one particle states (a mass gap is assumed)², whereas the regions of double cones are too small in order to still carry particle properties. The construction of

 $^{^{2}}$ However the higher particle scattering states cannot be resolved. The S-matrix cannot be extracted from the interacting wedge algebra alone, but one needs the relation to the relatively extremely nonlocally positioned incoming wedge algebra.

the modular wedge localization spaces is achieved in terms of Unruh-like KMS states on these semilocal generators.

Zero mass situations can give rise to infrared problems which lead to the loss of the reference in-space (the large time limits vanish for example in QED, and scattering cross sections must be computed directly in terms of expectation values and probabilities instead of amplitudes); the associated "infraparticles" do not have a Fock space structure. In that case a Fock reference space can only be associated with compact space-time regions of the interacting theory, e.g. double cones which have no infrared problems. The double cone algebras have a very big "folium" of states which includes the double cone restriction of e.g. free massless states, but on the other hand their modular structure is nongeometric (in the massive case) and less susceptible to physical interpretation.

If the zero mass theories are conformally invariant, the modular objects are again known and turn out to be geometric; however the deviation of the modular involution from its free value is generally not relatable to scattering. This opens the possibility to use standard Fock space methods instead of introducing ad hoc new algebras (example: W-algebras), which are not natural in the setting of particle representations³, with bring in the risk of loosing the Hilbert space setting due to indefinite metric. As a result of *local unitary equivalence* of any conformal quantum algebra to e.g. zero mass Fermions, we always have a (positive) Hilbert reference space for any conformal QFT. In these notes we use the prefix "quantum" only for Hilbert space representations, leaving aside certain indefinite metric representations (which may have some use in statistical mechanics). Despite their conceptual complexity, zero mass situations tend to be analytically simpler, a fact which contributed e.g. to the popularity of chiral conformal QFT.

We emphasized the gain in constructive insight into nonperturbative QFT in order to convince the reader that the times have passed when the contribution of algebraic QFT to problems of high energy physics and the quasiparticle structure of condensed matter physics was "smaller than any preassigned epsilon" (if this ever was true), as some people who worked on long forgotten fashionable subjects of the 60^{ies} used to say at that time.

The gain in understanding of the foundations of QFT is equally impressive. For example certain well-known "technical assumptions" concerning domains of unbounded (smeared) field operators, which appeared first in Wightman's axiomatic formulation, become now related with the basic physical properties of the thermal modular localization subspaces (the domain spaces of the Tomita operator S). The Wightman domain (which appears in Wightman's axiomatics [18]) turns out to be the intersection of all these subspaces and the natural domain for fields restricted to a region is simply the intersection of all modular subspaces corresponding to the double cones which fit into that region. This modular characterization suggests that the basic role of point-like (and hence operator-valued singular functions) fields, as opposed to nets of bounded oper-

³QFT which starts from Wigner's positive energy representation leads to Fock spaces as the arena for interacting QFT. Theorems on positivity are therefore not required (apart from gauge theories) because the embedding Fock space is automatically positive.

ators, is precisely to bring these modular localization properties into the open, i.e. to make them manifest. The space obtained by smearing a field vector $A(x)\Omega$ with arbitrary test functions, i.e. the "one field space", is easily seen in chiral conformal theories to be in one to one correspondence with an irreducible representation space of the *infinite dimensional group generated by all modular* transformations of double cones (or rather intervals in the chiral case)⁴. If this correspondence turns out to be a general property of algebraic field theory, one would have achieved a group theoretical construction of fields in a theory of local observables and in this way equipped them with a new fundamental role which goes far beyond the before mentioned role of being just local coordinatizations of algebras. Compared with the standard introduction of local fields via quantization this would be a major conceptual achievement since the modular properties are the most "quantum" or "noncommutative" in the sense that they do not permit (unlike noncommutative geometry) a commutative version at all. As we will see the modular algebraic properties insure that we maintain the biggest distance to quasiclassical or perturbative concepts, which is just what we want.

In this modular role the "field coordinates" are more than just generators of local algebras which carry information about the range of a local observable algebra applied to the vacuum in their domain property. It is through them that the somewhat hidden modular properties manifest themselves through covariance laws and localization properties. The reason why these modular properties which are so basic to QFT have been discovered so late, lies in the fact that Lagrangians, euclidean methods and functional integrals which dominated the scene of QFT, are too close to the (quasi)classical quantization parallelism. Whereas quantization has played an important intuitive role in the understanding of QFT from the side of quasiclassical concepts and the formalism of renormalized perturbation theory, the totally noncommutative modular structure promises to make systematic inroads into the largely unknown nonperturbative regime.

It would have been nice to use the gain in nonperturbative insight in order to classify free d=2+1 anyons and plektons by constructing their correlators and related nonrelativistic quasiparticles. Unfortunately at the time of writing the study of free plektonic charge carriers had barely begun (chapter 7.7), but from the few indications it appears that the nonrelativistic limit will remain a QFT i.e. there are no fields which applied to the ground state yield a one (quasi)particle state as there are for Fermions/Bosons in the Schrödinger theory. In order to maintain the plektonic relation between spin and statistics in the nonrelativistic limit one needs the presence of virtual particles. This is related to the paradoxical sounding statement that there are no relativistic free fields (on shell, without vacuumpolarization clouds) for the description of free anyons or plektons.

The credo that all the physics of field-coordinate-free QFT is carried by

 $^{^{4}}$ This conjecture about the relation of individual fields with a kind of universal modular group is due to Fredenhagen (private communication) who observed it in chiral conformal QFT [101]

the net of observables has with stood the passing of time and gained growing acceptance, but it still transcends standard text book formalism. I hope in these notes I will convince some of the remaining sceptics about the viability of the idea that all of local quantum physics is already contained in the net of modular localization subspaces by demonstrating how it works in d=1+1 factorizing theories.

At this point it should be abundantly clear that the central theme of these lectures is *localization of states and locality* (Einstein causality in the case of observables) of algebras. These concepts are not only of great physical relevance in condensed matter physics (localized and itinerant states), but also pivotal for the *interpretation* of QFT. Attempts to bypass this and to use instead global concepts from the beginning (as e.g. path integrals, topological field theories) will occasionally receive some critical attention, but do not enter the main line of this notes. Due to the proximity of localization with the notion of temperature, the standard "heat bath" Hamiltonian temperature will be introduced already in chapter 2. The first chapter is reserved for some introductory remarks on general quantum theory. The Wigner theory and free fields in chapter 3 is already presented with the modular hindsight. Subsequently we will learn about the basic distinction between "heat bath"-and "localization"-temperature. In that same chapter we also collect other constructions which stays close to free fields.

The subsequent chapter 5 on the general Wightman and LSZ framework and its euclidean counterpart should be considered as a preparation for the modular nonperturbative approach in chapter 6 which contains a detailed application to the formfactor program of factorizing theories. It also serves as a preparation for the pure algebraic approach in chapter 7. In order to keep the notes reasonably self-contained, a brief collection of mathematical results can be found in an appendix at the end.

It is the proximity of the nonperturbative concepts to TCP symmetry and other fundamental notions in QFT which generate confidence in our approach. The change of paradigm, which accompanies this new approach, is that those structures of QFT which were always thought of as fundamental but somewhat rigid and almost "kinematical" properties as TCP, now move into the center of the "dynamical" stage⁵. In fact in chapter 6 the reader will be led to take notice of the fact that all the interaction resides in the TCP-reflection and that the Smatrix, in addition to its well-known role as the large-time asymptotic operator describing scattering, also plays an important role in modular localization. It is this second property which e.g. explains its fundamental role in the formfactor construction of factorizable QFT's. In this way the latter important concepts used in that construction as crossing symmetry construction becomes intimately connected with the thermal KMS and pair-creation properties of the Rindler-Hawking Unruh issue and with physics near more general horizons.

A new approach is also expected to cast some new light on past successes.

 $^{^{5}}$ Of course such statements should be taken with the full awareness that the cut between "kinematics" and what was hitherto considered as "dynamics" was never as rigid and a priori as the textbooks make us believe.

For this reason we will attempt a rather systematic presentation which includes a substantial part of the standard material of QFT (especially chapter 4), but occasionally somewhat different from the standard textbook treatments by emphasis and interpretation. Algebraic QFT acknowledges the fact that it has a common cradle with other versions of relativistic particle theories which is the renormalized Feynman-Schwinger Tomonaga perturbation theory. In order to maintain and emphasize this common heritage, we present in chapter 4 a simple derivation of the most important effects in QED with a minimum on formalism but emphasis on physical concepts. In the same chapter we also expose the physics of renormalization in greater detail. Our subsequent presentation of perturbative gauge theory uses the linear version of the BRS formalism of cohomological representation formalism of selfinteracting spin 1 theories. This casts a new light on perturbative massive spin 1 theories which in the standard gauge approach were interpreted as spontaneously broken. Our investigations show that the conceptual problems in gauge theories are still ill-understood and many popular ideas of a preliminary nature.

In fact QFT, despite its age, is still in its conceptual infancy, especially from the new nonperturbative viewpoint. This goes contrary to the widespread (but unfounded) belief that the basic equations of the world below the Planck scale are already known and that they predict a great "desert" and that if we would be more clever in our computations (bigger computers etc.), we would have a description of all phenomena below that scale.

The algebraic approach has many aspects in common with the philosophy underlying Wilson's renormalization group. It also rejects the idea of a "theory of everything". Its picture of physical reality is more that of many shells which are hierarchical ordered with respect to decreasing distances. In every shell of this infinite "onion" there is the possibility of having a consistent mathematical theory and the principles of the next shell contain the previous ones as a limiting case. In fact the mathematical description is very much intervoven with the definition of one shell and its separation to the next one. Lagrangians, actions etc. are interpreted as characterizing equivalence classes of theories which are indistinguishable with respect to certain physical properties in the present regime, similar to universality classes in condensed matter physics. Its underlying philosophy is consistent with the use of effective Lagrangians a la Weinberg. However it is less optimistic concerning what one can learn from such phenomenological concepts and its notion of equivalence classes is somewhat different and in some sense sharper. In addition to the well known short distance universality classes with their unique scale invariant representatives there are "long distance" classes (defined via cluster limits of the S-matrix) which have the same charge superselection rules and particle structure but simpler (or no) interactions. In d=1+1 they have factorizing representatives.

The strategy for classifying and constructing models is in a certain sense opposite to that proposed on the basis of renormalization group transformations. Whereas the Wilson formalism approaches the construction of fixed point theories by starting outside the fixed point theories and reaches the latter by iterated renormalization group transformations, the algebraic approach aims directly at an intrinsic construction of those representatives and proposes to explore the more complicated terrain around these simpler theories by perturbative ideas in the vein of Zamolodchikov's picture of massive perturbations on chiral conformal theories.

A nice illustration of the difference in the underlying philosophy is given by chiral conformal theories. In the algebraic approach the (imaginary time) theory of critical indices becomes related to the computational easier classifiable braid group statistics of the associated (real time) chiral QFT and in this way the spectrum of possible critical indices gets related to the superselection rules and their fusion and braid group characteristics of the noncommutative local charges (critical indices modulo 1 = statistical phases). Hence if one wants to compare the philosophy in these notes to the earlier contributions on the connection between Statistical Mechanics and QFT, it would be more appropriate to point towards Kadanoff's use of the Coulomb gas as an analogue system for fusion of superselected charges.

There is however a quite serious discrepancy in points of view related to certain aspects of the post Wilson renormalization group philosophy; notably to the cavalier use of the notion of "cutoffs" in real time QFT⁶. Whereas the algebraic approach does not negate the usefulness of representing field theoretic models as the scaling limit of lattice models (however this only has been achieved for a few cases of d=1+1 factorizing models), it is totally incompatible with the direct use of (relativistic) cutoffs as a method to dump some unknown physics of the next shell in a physically acceptable way. Whereas the proponents of such ideas usually think in terms of cutting off some integrals in their euclidean approach, the algebraic framework is fully aware of the fact that nonlocal "theories" do not permit a self-sustaining physical interpretation (example: the causality origin of the crossing symmetric cluster properties as one needs them for the derivation of scattering theory gets lost together with causality and with it crumbles the whole interpretation). Here it is helpful to remember that almost as long as QFT exists there were attempts to find a physical way to regularize or cut-off light cone singularities or to introduce covariant formfactors in Lagrangians or to add pairs of (conjugate)complex poles into Feynman rules. Each attempt led to a total loss of physical interpretation. LQP takes the 60 years of failure to give physical meaning to such notions as cutoffs elementary length and noncausal interactions very serious and protects the Einstein causality and localization principles of the first half of this century against their present sell out⁷. To put it very bluntly: whereas one can approximate operators as Hamiltonians, it is not possible to approximate principles inasmuch as it is not meaningful in everyday life to use the phrase "partially pregnant". The transcendence of a physical principle is not a "bit of a non-principle", but rather a new princi-

⁶A statistical mechanics with a momentum space cut-off cannot be the analytic continuation of a physically interpretable real time QFT.

⁷The renormalization group approach only maintains its aura of physical modesty as long as it is not directed towards a sellout of the Einstein causality principles by claiming that there is a physically interpretable QFT in the presence of cutoffs and that it can define physically viable nonlocal relativistic QFT.

0.1. INTRODUCTION

ple containing the old one as a limiting case. None of the constructive aspects based on modular localization makes sense without Einstein causality. Their is no chance to manipulate a relativistic cutoff into any of the factorizing models without wrecking irrevocably the entire physical interpretation. From the algebraic point of view one may however entertain the speculative idea that there exists a new principle beyond Einstein causality and that it underlies the still very elusive quantum gravity.

The content of this report rather tries to attract those physicists who, on the one hand still have not lost their conceptual curiosity, and on the other hand feel uneasy about the present predominance of formalism over conceptional insight. Such a reader may enjoy the unique charm and surprise of finding deep physical concepts in rather mundane physical problems, far from big deserts and theories of "everything" and well below the Planck length. This should however not be misunderstood as a return to a mathematical stone age.

The understanding of the material presented in these notes is mathematically as demanding as the investment into geometry and topology, which the typical reader probably already made in good faith and in the hope to cover most of the mathematical structure of quantum physics. However the logical and conceptual structure of local quantum physics is in many respects different from geometry. Some of the new mathematical tools in local quantum physics as the Tomita-Takesaki modular theory and the Vaughn Jones subfactor theory are sketched in the appendix. Many of the important contributions were also made by physicist, but apart from perhaps one exception [3], they have not entered textbooks on QFT. Such a refinement of textbook knowledge would be particularly necessary if it comes to low dimensional QFT. After all, the main physical value of low dimensional models is their use for a better conceptual and structural understanding of general QFT (perhaps also for the purpose of basing universality explanations in condensed matter physics on firm grounds) and only in second place the increase of our mathematical culture. For this reason the presentation of low dimensional models (in particular chiral conformal QFT) may appear different from the way the reader may have met this material in other reviews.

Some ideas which apparently are natural from a differential geometric view and presently enjoy great popularity as e.g. supersymmetry appear somewhat artificial and unphysical from the LQP viewpoint. Apart from the fact that no compelling theoretical principle which leads to such structures is known, supersymmetry reveals itself within the conceptual frame of algebraic QFT as an "accidental symmetry", unlike any other space-time or internal symmetry. This peculiarity one expects to manifest itself e.g. in its thermal behavior.. Indeed, very recent investigations of the thermal behavior of supersymmetry showed [42] that this symmetry is completely unstable in thermal KMS states⁸ and thus confirm old suspicions that one is dealing here with an accidental symmetry. In those low-dimensional supersymmetric models which are soluble (e.g. the con-

⁸The usual (broken) symmetry picture, which holds for locally generates internal and external symmetries in a heat bath does not apply here. Rather supersymmetry suffers a total thermal collapse [42].

formal Ising model and various massive factorizing models), the supersymmetry plays no role in the nonperturbative analysis and its placement within a family of soluble models. The short distance properties of its charge-carrying fields are not better than that of its non supersymmetric neighbors. Whereas usual inner symmetries are deeply related to the charge structure of the representation theory of its observables, the systematic superselection analysis never asks for a supersymmetric encoding. And whereas usual space-time symmetries are profoundly related to the group generated by modular groups of local algebras, supersymmetry points toward nothing but itself. For this reason there is no place for SUSY in these notes.

Needless to say that our approach is rather conservative in its use of existing concepts. The revolutionary aspect to the degree that there is one at all, lies more in precise observations and discoveries than inventions. In particular we will keep to physical notions as equivalence classes of real time fields, TCP- and Tomita J-reflections rather than geometrical concepts, imaginary time formalism, commutative cohomology and S- and T-mirror-reflections etc.

On the other hand the reader will find a detailed presentation of "Haag Duality" which is a pivotal property of nonperturbative local QFT and is known to lead to such fundamental issues as (braid group) statistics and an intrinsic understanding of spontaneous symmetry breaking. In d=1+1 we also study its relation to the QFT formulation of Kadanoff order-disorder Duality which is the local version of the global Kramers-Wannier Duality in lattice systems.

I elaborated this material in the conviction that the best strategy in a time of stagnation and crisis⁹ is to return to the roots of QFT and re-analyze the underlying principles in the light of new concepts. For this reason there is heavy emphasis on Wigner's approach to particles which, long before algebraic QFT came into existence, was the first successful attempt to do relativistic quantum physics in an intrinsic fashion i.e. without recourse to quantization.

The relation of the present nonperturbative approach of LQP to standard textbook QFT and its more recent differential-geometric refinements is somewhat analogous to that of a wave optical treatment of QM relative to proper QM in its own conceptual setting. In both cases one has to redo and relearn the conceptual basis of the theory. The main distinction between the present algebraic method and standard quantization approach (canonical quantization, functional integrals etc.) is that in the first case the properties which are assumed at the start of a construction are really reflected in the result. On the other hand the quantization approach is more artistic: canonical commutation relations outside free fields are almost never valid for the renormalized results (exception: certain super-renormalizable two-dimensional theories) and the same holds for functional integral (Feynman-Kac) representations. Such structures are working hypothesis which serve to start a chain of calculational ideas but which themselves get lost on the way and often lead to other correct properties. It seems to be a general feature of a quantization method that it leads invariably to a connection of the existence of a QFT with a sufficiently good short distance

⁹Instead of a detailed verbal description, I refer to the picture at the end.

0.1. INTRODUCTION

behavior of the underlying basic fields. This is not borne out in our nonperturbative approach which is based on modular localization and uses ideas which are very different from quantization.

Anybody who knows me, understands that my attachment to Brazil (where most of these notes were written) is not only due to its natural beauty, but there are also deep scientific roots. With feelings of nostalgia I remember those wonderful years (1968-1980) of collaboration with J. A. Swieca. These were times of free-roaming scientific endeavour, long before the globalization of physics, which nowadays forces young physicists to build their career around some trendy formal ideas, started to do incalculable harm to QFT. In these notes I remain faithful to the spirit of that collaboration in that I shun away from inventions. All the new developments and ideas the reader finds in this article are discoveries or observations which follow from the principles of local quantum physics. The style is a bit more informal than that of a text book, for example there are many instances of repetitions. With new material originating from a different conceptual framework than that of standard QFT it may actually be an advantage for the reader to have repetitions of the same material from slightly different perspectives.

This presentation should not be misread as a moral judgement against one or the other approach to QFT. It rather is an attempt to revive some of the critical conceptually based Bohr-Heisenberg-Wigner spirit in the present times of: "everything goes" (at least as long as it lives up to that entertaining high caliber scientific journalism which characterizes many contributions to high energy physics, see the hep-th server¹⁰).

CBPF, Rio de Janeiro, March 1998 Bert Schroer

 $^{^{10}\}mathrm{I}$ strongly advice any young person still in search of a permanent position, not to say such things in public.

CONTENTS

16

Chapter 1

The Basics of Quantum Theory

1.1 Multiparticle Wave Functions, Particle Statistics

It is well-known that quantum mechanical multiparticle systems can be obtained by the tensor product construction from one-particle spaces. Let:

$$\psi(x,s_3) \in \mathcal{H}_1 = \mathcal{L}^2\left\{\mathcal{R}^3, \{-s,\ldots,+s\}\right\}$$
(1.1)

be a wave function of a nonrelativistic particle of spin s where, as usual, s_3 is the component of spin with respect to an arbitrarily chosen quantization axis. The L^2 -space has the standard inner product

$$(\psi_2, \psi_1) := \sum_{s_3 = -s}^{s} \int d^3x \psi_2^*(x, s_3) \psi_1(x, s_3)$$
(1.2)

and norm $\|\psi\| = (\psi, \psi)^{\frac{1}{2}}$

The two-particle space:

 $\mathcal{H}_2 = \mathcal{H}_1^a \overline{\otimes_{\mathbf{C}}} \mathcal{H}_1^b \qquad \text{a b..type of particle}$ (1.3)

is simply the closure of the algebraic tensor product over the complex numbers generated by vectors: $\sum_{ik} \psi_i^a \otimes_{\mathbf{C}} \psi_k^b$ equipped with the scalar product (in the following we omit the subscript **C**) induced by the formula:

$$(\psi^a_{i'} \otimes \psi^b_{k'}, \psi^a_i \otimes \psi^b_k) := (\psi^a_{i'}, \psi^a_i) \cdot (\psi^b_{k'}, \psi^b_k)$$
(1.4)

continued by linearity in the right and antilinearity in the left factor. If, as it is usually the case, the one-particle algebra forms a complete (irreducible) set of dynamical variables in the one-particle space, the totality of all observables which are compatible with measurements of subsytem a (form the commutant of the *a*-algebra) inside the composed system a + b are just those which live in the tensor factor b; a and b are stochastically independent in a very strong sense. The generalization to N factors:

$$\mathcal{H}_N = \mathcal{H}_1 \overline{\otimes} \mathcal{H}_1 \dots \overline{\otimes} \mathcal{H}_1 \tag{1.5}$$

is straightforward. The tensorproduct structure is the mathematical formulation of kinematical "statistical independence" in the sense of quantum theory. The dynamical variables of the subsystems (say particles with spin) at a given time factorize:

$$\vec{X}_i = 1 \otimes \dots \otimes \vec{x}_i \otimes 1 \dots \otimes 1$$
and similarly for \vec{P}_i and \vec{S}_i

$$(1.6)$$

They act on N-particle wave functions $\psi(x_1s_1, ...x_Ns_N)$ with s_i (omitting the subscript 3 for brevity) the third component of the i^{th} spin. Global symmetry transformations, as translations and rotations, act multiplicatively:

$$(U(\vec{a})\psi)(x_1s_1,...x_Ns_N) = (U_1(\vec{a})\otimes...\otimes U_1(\vec{a})\psi)(x_1s_1,...x_Ns_N)$$
(1.7)

$$(U(\vec{n},\theta)\psi)(x_1s_1,...x_Ns_N) = (U_1(\vec{n},\theta)\otimes...\otimes U_1(\vec{n},\theta))\psi(x_1s_1,...x_Ns_N)$$
(1.8)

$$U(a) = e^{i\vec{P}\vec{x}}, \qquad \vec{P} = \sum_{i} 1 \otimes \dots \otimes \vec{P}_{i} \otimes \dots \otimes 1$$
(1.9)

 $\begin{array}{ll} U(\vec{n},\theta)=e^{i\vec{J}\vec{n}\theta}, & \vec{J}=\sum_i 1\otimes \ldots \otimes \vec{J_i}\otimes \ldots \otimes 1 \ , \ \vec{J_i}=\vec{x_i}\times \vec{p_i}+\frac{1}{2}\vec{\sigma_i} \\ \text{Here } \vec{\sigma} \text{ are the Pauli-matrices. Operators which implement an interaction} \end{array}$

between particles, as the hamiltonian H, violate this one-particle factorization:

$$U(t) = e^{-iHt}, \quad H = H_0 + H_{int}, \quad H_{int} = \sum_{i < k} 1 \otimes \dots V_{ik} (x_i - x_k) \dots \otimes 1 \quad (1.10)$$

In the last expression the identity operator at the i^{th} and k^{th} place has been replaced by a conventional local pair interaction which, in the exponentiated form looses the pairing property of the infinitesimal generator. Therefore at first sight it appears, that the localized pair-interaction leaves no mark at all on U(t). Fortunately this is not quite correct, its marks are the important "cluster properties". In our context they are :

$$H_N \to H_{N_1} \oplus H_{N_2}, \ \Omega_N \to \Omega_{N_1} \otimes \Omega_{N_2}, \ S_N \to S_{N_1} \otimes S_{N_2}$$

on clustering wave functions:
$$\lim_{a \to \infty} \psi(x_{1,...}x_{n_1}, x_{n_1+1} + a, ...x_{N_1+N_2} + a)$$
(1.11)

Here $\Omega^{\pm} = \lim_{t \to \infty} e^{iH_0 t} e^{-iHt}$ are the Møller operators and S is the S-matrix $S = \Omega^{+*}\Omega^{-}$. Equivalently one may introduce a partial translation $U_{C_{N_2}}(a)$ which translates the particles in the N_2 -cluster C_{N_2} infinitely far away from the rest:

$$\lim_{a \to \infty} U_{C_{N_2}}(a) H_N U^*_{C_{N_2}}(a) = H_{N_1} \oplus H_{N_2},$$
(1.12)

and similarly for e^{iHt} , Ω^{\pm} and S with \otimes instead of \oplus . The cluster property is therefore a kind of asymptotic factorization, expressing statistical independence for long distances. As we will see, it follows from more fundamental locality structures in QFT and the existence of a spatially homogeneous reference state. Although it is trivially satisfied for short-range quantum mechanical interactions, it cannot always be taken for granted if the interactions become long-range. In that case the cluster decomposition requirement is expected to affect the boundary conditions of scattering theory. For example for the relative Aharonov-Bohm interaction between "dyons" (electrically and magnetically charged particles in spacetime dimension d=2+1), the cluster decomposition property is expected to become a *nontrivial imposition on the a priori unknown* scattering boundary conditions \neq plane wave condition, which is the short range interaction boundary condition for stationary scattering on multiparticle scattering states. These quantum theoretical subtleties are of course irrelevant for the calculation of the Aharonov-Bohm phase shift, which can be understood in entirely quasiclassical terms. In the tentative applications to braid group statistics of dyons however, the boundary condition is apriori not known. In the presence of an A-B long range interactions that scattering boundary condition is the correct one, which yields the cluster decomposition property of the scattering amplitude, thus assuring that the N-particle S-matrix passes to the previously determined (N-1)-particle S-matrix upon removal of one particle to infinity (plane wave scattering conditions for long range interactions generally destroy this property). In this way the N-particle problem becomes related to the (N-1)-particle problem via the cluster property, even though there is no actual creation of particles. Hence what appeared at the beginning as a standard quantum mechanical problem, turns out to be a problem in which particle states for all N enter, i.e. a problem with a field theoretic aspect. Presently not even a description of nonrelativistic anyons exists since no solution of the N-particle Aharonov-Bohm scattering with cluster properties has been elaborated.

In local QFT, cluster properties are a consequence of the general framework and do not have to be imposed on interactions as in nonrelativistic physics. In fact it is appropriate to think of the cluster decomposition property as the relic of LQP (Haag) in the nonrelativistic limit. The statistics of particles must have this inclusive structure lelating N - 1 to N and the fact that the identity of particles is expressed by the braid group B_{∞} which is the inductive limit of ... $\subset B_i \subset B_{i+1} \subset ...$ (the same holds for the permutation group S_{∞} which is a special quotient of the braid group) reflects precisely the inclusive "russian matrushka" structure of the clustering property on purely algebraic level with the spacetime localization removed. In fact, as will become clearer in a later chapter, this removal in LQP corresponds to a theory with full localization "flesh" too a combinatorical theory of intertwiners together with a natural tracial state. That theory of "bones" in the geometric approach is called "topological field theory".

It is well-known that the dynamical variables obtained as tensor products form an irreducible system (\simeq validity of Schur's lemma) if the single particle variables have this property in the one-particle Hilbert space (true in our example of $\vec{p}, \vec{x}, \vec{s}$). In that case the requirement of identity of particles permits only one-dimensional (abelian) representations of the permutation group i.e. fermionic or bosonic representations. In order to understand this, we have to incorporate the notion of indistinguishability into our tensor products. Writing symbolically $\psi_n(1, 2, ...n)$ as a short hand for our *x*- and spin-dependent wave functions, a permuted wave function (as usual in QM, the permutations act on the indices as P^{-1} in order to comply with the group composition, a behaviour well known from the action of the rotation group on spatial coordinates):

$$\begin{aligned} (U(P)\psi)_n (1,2,...n) &:= \psi_n^P (1,2,...,n) := \\ &= \psi_n (P^{-1}(1), P^{-1}(2), ...; P^{-1}(n)) \end{aligned}$$
 (1.13)

must describe the same physical state¹, even though the permutation leads to a different vector in Hilbert space. This distinction between vectors and physical states (in the sense of defining expectation values on the algebra of observables) is crucial for the understanding of the (later presented) superselection rules in the later sections. In terms of observables, this simply means that the U(P)'s commute with the observables. If, as in standard Schrödinger theory, the observables form an irreducible set of operators, the representation U(P) of the permutation group S_n must be abelian and hence:

$$\psi_n^P(1,2,...n) = \omega(P)\psi_n(1,2,...n)$$

$$\omega(P) = \begin{cases} 1 & Bose \\ sign(P) & Fermi \end{cases}$$
(1.14)

Mathematically this statement is a tautology and physically there is no reason to assume irreducibility. In fact the occurrence of nonabelian representations of S_n is equivalent to the appearance of reducible quantum mechanical observable algebras.

In order to construct such a nonabelian example, a rudimentary knowledge of the representation theory of the permutation group is helpful. The equivalence classes of irreducible representations of S_N are characterized by partitions with height n:

$$N_1 \ge N_2 \ge \dots \ge N_n \,, \quad \sum_i N_i = N \tag{1.15}$$

They are pictured by so called Young tableaus, an array of N boxes in rows of nonincreasing size (the admissibility condition for tableaus). Young tableaux are also in a one to one relation to irreducible representations of SU(N).

The representation of S_N corresponding to each tableau of depth d is most conveniently described by decomposing the natural representation on the *N*fold tensor product of a d-dimensional complex vector space $V^{\otimes N}$ (of dimension $d \cdot N$) into irreducibles. Up to equivalencies, one obtains a subset of irreducible

¹i.e. the same expectation values in permuted wave functions for quantum mechanical observables. Later we will introduce a more general concept of "state" which does not require a Hilbert space but rather leads to one.

components by applying S_N to the reference vector:

$$e_1^{\otimes (N_1 - N_2)} \otimes (e_1 \wedge e_2)^{\otimes (N_2 - N_3)} \otimes \dots \otimes (e_1 \wedge e_2 \dots \wedge e_d)^{\otimes N_d}$$

$$e_1 \dots e_d \text{ basis in } V$$

$$(1.16)$$

with the tensor product action:

$$U(P)\xi_1 \otimes \xi_2 \otimes \dots \otimes \xi_N = \xi_{P(1)} \otimes \xi_{P(2)} \dots \otimes \xi_{P(N)}, \qquad (1.17)$$
$$\xi_i \in \{e_1 \dots e_d\}$$

The cyclically generated subspace is the irreducible representation space corresponding to (1.16). Here the \wedge designates the wedge product leading to completely antisymmetric tensors. Clearly the vectors (1.16 are in a one-to-one correspondence with partitions (1.15) with height n = d, which are also cyclic vectors with respect to irreducible SU(d) representations. An "admissible" numbering of a given tableaux is any numbering which is increasing in each row and column. The different admissible numberings correspond to the multiplicity with which the irreducible representation occurs in the regular representation (which has a larger dimension than the above natural representation) of the group algebra $\mathbf{C}S_N$ (see next section).

Another method to construct the representation theory is the inductive construction of tableaus according to Schur's rules of adding a small box to a previously constructed tableaux corresponding to S_{n-1} :

$$\pi_T \times box = \oplus_{adm.} \pi_{T+box} \tag{1.18}$$

The admissible ways to add a box are such that the resulting tableaus are admissible in the aforementioned sense. But now there is no maximal height d and instead of the natural representation on tensor products on d-dim. vectorspaces we are inducing the so called regular representation. The iteration starting from n = 1 gives a reducible representation of S_N with $m(T_N)$ =multiplicity of occurrance of the tableaux T_N :

$$\oplus_{T_N} m(T_N) \pi_{T_N} \tag{1.19}$$

The multiplicity is the same as that of T_N in the group algebra $\mathbf{C}S_N$ (next section). It is obviously equal to the number of possibilities to furnish admissible numbering (the box numbering inherited from the inductive Schur construction). It agrees with the dimensionality of the representation. Furthermore the regular representation contains all equivalence classes of irreducible representations whose number, as will be shown in the next section for general finite groups, equals the number of conjugacy classes.

Now we are able to sketch a (rather trivial) counterexample against the irreducibility of the observable algebra. Imagine that we have Bose particle which carry besides spin a "hidden" quantum number ("flavor" or "color") i.e. an internal degree of freedom which can take d values. Assume that the measurability is restricted to flavor neutral operators:

$$\underline{\vec{x}} = \frac{1}{d} \sum_{a=1}^{d} \vec{x}^{(a)}, \quad \underline{\vec{p}} = \dots, \underline{\vec{s}} = \dots$$
(1.20)

Clearly the flavor averaged multiparticle observables act cyclically on a smaller "neutral" reduced Hilbert space which may be described in the following way:

$$\mathcal{H}_N^{red.} = \mathcal{H}_1 \otimes \dots \mathcal{H}_1 \otimes \mathbf{C}S_N \tag{1.21}$$

Here the one-particle wave function spaces H_1 have no flavor degree of freedoms and $\mathbb{C}S_N$ is the N! dim. representation space of the regular S_N representation. We connect the reduced inner product with the one in the flavor description:

$$\begin{pmatrix} \Phi \otimes P^{-1}, \Psi \otimes Q^{-1} \end{pmatrix}_{red} = \sum_{i_1 \dots i_N} \frac{1}{d^N} \left(\varphi_1 \otimes e_{i_{P(1)}} \otimes \dots \varphi_N \otimes e_{i_{P(N)}}, \psi_1 \otimes e_{i_{Q(1)}} \otimes \dots \otimes \psi_N \otimes e_{i_{Q(N)}} \right)_{sym}$$
(1.22)

Here the $\varphi's$ and $\psi's$ are the spatial wave functions without flavor and the e's are from the basis in d-dimensional flavor space V. The inner product is the natural scalar product in the symmetrized tensor space $(H \otimes V)_{symm.}^{\otimes N}$. On the left hand side Φ and Ψ are the tensor products of the φ_i and ψ_i . The reduction is implemented through averaging over the flavor degrees of freedom. The orthonormality relations of the e's allow to simplify the result to:

$$\sum_{i_1\dots i_N} d^{-N} \sum_S \prod_j \left(\varphi_j, \psi_{S(j)}\right) \left(e_{i_{P(j)}}, e_{i_{Q(j)}}\right) = \sum_S \prod_j \left(\varphi_j, \psi_{S(j)}\right) \varphi^{(N)}(PSQ^{-1})$$
(1.23)

 $\varphi^{(N)}$ denotes a tracial linear functional on CS_N which on the basis elements P is given by:

$$\varphi^{(N)}(P) = \frac{1}{d^N} Tr U_N(P) \equiv tr(p) \tag{1.24}$$

where $U_N(P)$ is the previously introduced natural representation of S_N on $V^{\otimes N}$ and Tr stands for the natural matrix trace.

The memory on the averaged flavor is completely absorbed in the multiplicity factor. Note that flavored fermions would have given a similar result with the only difference of signP factors.

The conceptually aware reader will note, that this "cooked up" parastatistics illustration is precisely what an experimentalizer confronts, if in a nonrelativistic atomic problem the electron spin would have no dynamical manifestation (neglegible spin-orbit coupling) or if for a neutron-proton system he would not be able to measure electric charge. Internal symmetry is a very clever theoretical invention which trades the unpleasant nonabelian Young-tableaux against the more physical (*more local!*) standard compact internal symmetry-group description. However it is not universally applicable, see the later discussion of attempts to encode nonabelian braid group statistics into a "quantum symmetry" concept.

22

1.1. MULTIPARTICLE WAVE FUNCTIONS, PARTICLE STATISTICS 23

It is now easy to see that the normalized trace of the natural representation has an intrinsic characterization in terms of a tracial state φ (a positive linear function) on the group algebra $\mathbb{C}S_{\infty}$. Here S_{∞} is the inductive limit of the S_N groups:

$$S_2 \subset \dots \subset S_N \subset S_{N+1} \subset \dots \subset S_{\infty} \tag{1.25}$$

The normalized tracial functional $\varphi^{(N)}(P)$ on CS_N (the extension from S_N to the group algebra CS_N is by linearity) has a natural extension φ to the inductive limit CS_{∞} . It is characterized by the following three properties:

$$\varphi(x)$$
 is tracial state on CS_N (1.26)

i.e.
$$\varphi(x^*x) \geq 0, \ \varphi(1) = 1, \ \varphi(xy) = \varphi(yx)$$

 φ fullfills the Markov-property: $\varphi(P_1P_2) = \varphi(P_1)\varphi(P_2)$ (1.27)

$$\varphi(P = \text{transposition}) = \pm \frac{1}{d}$$
, for flavoured fermions (1.28)

where $P_1 = P_1(\tau_1, \tau_2, ..., \tau_{N_1-1})$ is a permutation involving the first $N_1 - 1$ generators and

 $P_2 = P_2(\tau_{N_1+1},...,\tau_{N-1})$ is a permutation involving the remaining generators except τ_{N_1} .

The generators τ_i (transpositions) are most conveniently pictured as crossings of the i^{th} strand with its neighbor i + 1, whereas all the other strands are running parallel (say upward).

These generators are subject to three relations:

$$\tau_i \tau_j = \tau_j \tau_i, \quad if \quad |i-j| \ge 2 \tag{1.29}$$

$$\tau_i \tau_{i+1} \tau_i = \tau_{i+1} \tau_i \tau_{i+1}, \quad \tau_i^2 = 1 \tag{1.30}$$

The first relation by its own is most appropriately pictured by allowing overand under-crossings in the τ_i i.e.by introducing $\tau_i^{\pm} = \{\tau_i, \tau_i^{-1}\}$ see Fig 2. If one adds to this Artin relation the second one $\tau^2 = 1$, the braid group (which even for finite number of strands is an infinite group) B_{∞} passes to the permutation group S_{∞} . Both groups owe their physical relevance to the fact that they are crucial for the understanding of particle statistics and normal commutation relations between charge-carrying fields. Their natural inclusive structure should be seen as an algebraic counterpart of the physical cluster property. As we already pointed out, the S_{∞} representations described by Young tableaus occur in the centralizer algebras of tensor products of group representations. Take e.g. the algebra generated by the d-dimensional defining matrix representation π of SU(n):

$$\pi(g) \otimes \pi(g) \dots \otimes \pi(g) \quad \text{in } V^{\otimes_N} \tag{1.31}$$

In this case the commutant (or centralizer of the group representation) of these operators does not only contain the algebra $\mathbf{C}S_N$ (this is evidently the case for all tensor representations of groups) but this algebra is even identical to the centralizer. The following two questions are relevant for the statistics classification: (1) Can one argue that the indistinguishability requirement of particles (or other localizable objects) together with other physical principles leads to representations which are multiples of the above tracial representations of S_{∞} ?

(2) Is it natural to interpret the appearance of a tracial state with the Markov property in terms of an inner symmetry, and does one gain anything by introducing the symmetry multiplicities (Heisenberg's isospin and its "flavor" generalization) manifestly into the formalism?

Both questions have an affirmative answer, i.e.statistics and internal symmetry are inexorably linked. The relevant theorem is the following:

Theorem 1 (following Doplicher, Roberts [43]) The most general statistics allowed in d=3+1 dimensions is abelian (Bose-Fermi) together with a compact internal symmetry group. The Boson-Fermion alternative is related to that of integer versus halfinteger spin (the spin-statistics relation). The parastatistics associated with nonabelian Young-tableaux (and described mathematically by a Markov trace on the infinite permutation group algebra CS_{∞}) can be traded for a compact internal symmetry group G and Bosons/Fermions. The latter description leads to a more local and less noncommutative description of QFT which is also more susceptible to a "quantization interpretation" with natural semiclassical limits.

Note that in the DR formulation the statistics problem is not only linked with the spin, but also with the internal group symmetry.

Although the tracial nature and the Markov property can be derived from a properly adapted (to the indistinguishability principle) cluster decomposition property, the natural place for its understanding and proof is relativistic QFT (as is the case with other structural properties as spin and statistics). The QFT locality and positive energy requirements naturally imply the cluster property and the inclusive picture relating N particles with N+1. We will give a derivation in a later chapters 5 and 7 on QFT in connection with the theory of superselection sectors. It is this picture, and not the N-particle quantum mechanics for fixed N (the standard quantum mechanical "proof" for the Fermi-Bose alternative in most books on QM is a tautology) which is responsible for the results on statistics. This continues to be true in the case of the new quantizations of statistical dimensions for braid group statistics which one finds in $d \leq 2 + 1$ dimensional QFT.

Superselection rules appeared first in the 1952 work of Wick, Wightman and Wigner[3]. These authors pointed out that the unrestricted superposition principle of quantum mechanics or equivalently, the unrestricted identification of self adjoint operators with observables (as formulated by von Neumann) suffers a limitation through the appearance of superselection rules. Their main example was standard quantum theory which describes integer as well as halfinteger spin. Its Hilbert space is a sum of \mathcal{H}^{\pm} where (-)+ corresponds to (half)integer spin. A linear combination of vectors from both spaces changes its relative sign under 2π -rotation:

$$\psi = \alpha \psi_{-} + \beta \psi_{+} \rightarrow \psi^{2\pi} := U(2\pi)\psi = -\alpha \psi_{-} + \beta \psi_{+}$$
(1.32)

Whereas the projective nature allows state vectors to suffer phase changes (the quantum mechanical origin of halfinteger spin!), observables and states (in the sense of expectation values) are unchanged under such a 2π rotation and in this sense behave classically. The following calculation shows that this is only possible iff the observables have vanishing matrix elements between \mathcal{H}^- and \mathcal{H}^+

$$(\psi, A\psi) = (\psi^{2\pi}, A\psi^{2\pi}) \iff (\psi_{-}, A\psi_{+}) = 0$$
 (1.33)
for all observables $A \in \mathcal{A}$

The proof just follows by inserting the above linear combinations. This selection rule is called the "univalence rule". In contradistinction to e.g. the $\Delta l = \pm 1$ angular momentum selection rules of atomic physics which suffer changes in higher order radiative corrections, *superselection* rules are universally valid. The vector state ψ above cannot be distinguished from a density matrix ρ :

$$(\psi, A\psi) = tr\rho A \quad with \ \rho = |\alpha|^2 |\psi_-\rangle \langle \psi_-| + |\beta|^2 |\psi_+\rangle \langle \psi_+| \tag{1.34}$$

The formal generalization for the Hilbert spaces and observables is obviously:

$$\mathcal{H} = \oplus_i \mathcal{H}_i \,, \quad A = \oplus_i A_i \,, \quad A_i = A \mid_{\mathcal{H}_i} \tag{1.35}$$

Such observable algebras in block form have a nontrivial center given by the block projections. In the following we will illustrate this decomposition theory by a simple but rich mathematical example, the superselection rules of the group algebra.

1.2 The Superselection Sectors of CG

As a mathematical illustration of superselection rules we are going to explain the representation theory of the group algebras.

Let G be a (not necessarily commutative) finite group. We affiliate a natural \mathbb{C}^* -algebra, the group-algebra $\mathbb{C}G$ with G in the following way:

• (i) The group elements $g \in G$ including the unit e form the basis of a linear vectorspace over **C**:

$$x \in \mathbf{C}G, \quad x = \sum_{g} x(g)g, \quad with \ x(g) \in \mathbf{C}$$
 (1.36)

• (ii) This finite dimensional vector space $\mathbf{C}G$ inherits a natural convolution product structure from G:

$$\left(\sum_{g \in G} x(g)g\right) \cdot \left(\sum_{h \in G} y(h)h\right) = \sum_{g,h \in G} x(g)y(h)g \cdot h = \sum_{k \in G} z(k)k \quad (1.37)$$

with $z(k) = \sum_{h \in G} x(kh^{-1})y(h) = \sum_{g \in G} x(g)y(k^{-1}g)$

• (iii) A *-structure, i.e. an antilinear involution:

$$x \to x^* = \sum_{g \in G} x(g)^* g^{-1} , \quad i.e.x^*(g) = x(g^{-1})^*$$
 (1.38)

Since :

$$(x^*x)(e) = \sum_{g \in G} |x(g)|^2 \ge 0, \quad (=iff \quad x = 0)$$
(1.39)

this *- structure is nondegenerate and defines a positive definite inner product:

$$(y,x) \equiv (y^*x)(e)$$

• (iv) The last formula converts $\mathbf{C}G$ into a Hilbert space and hence, as a result of its natural action on itself, it also gives a C^* norm (as any operator algebra):

$$||x|| = \sup_{\|y\|=1} \|xy\|, \quad C^* - condition: \ ||x^*x|| = ||x^*|| \, ||x|| \tag{1.40}$$

A C^* -norm on a *-algebra is necessarily unique (if it exists at all). It can be introduced through the notion of spectrum (mathematical appendix).

It is worthwhile to note that (iii) also serves to introduce a tracial state on $\mathbf{C}G$ i.e.a positive linear functional φ with the trace property:

$$\varphi(x) := x(e), \quad \varphi(x^*x) \ge 0, \quad \varphi(xy) = \varphi(yx) \tag{1.41}$$

This state (again as a result of (iii)) is even faithful, i.e. the scalar product defined by:

$$(\hat{x}, \hat{y}) := \varphi(x^*y) \tag{1.42}$$

is nondegenerate. On the left hand side the elements of $\mathbf{C}G$ are considered as members of a vector space. The nondegeneracy and the completeness of the algebra with respect to this inner product (a result of the finite dimensionality of $\mathbf{C}G$) give a natural representation (the regular representation of $\mathbf{C}G$) on this Hilbert space:

$$x\hat{y} := \widehat{xy} \tag{1.43}$$

The norm of these operators is identical to the previous one.

This construction of this "regular" representation from the tracial state on the C^* -group-algebra is a special case of the general Gelfand-Neumark-Segal (GNS-)construction presented in a later section.

Returning to the group theoretical structure, we define the conjugacy classes K_g and study their composition properties.

$$K_q := \{hgh^{-1}, h \in G\}$$
(1.44)

In particular we have $K_e = \{e\}$. These sets form disjoint classes and hence:

$$G = \bigcup_{i} K_{i}, \ |G| = \sum_{i=0}^{r-1} |K_{i}|, \ K_{e} = K_{0}, \ K_{1,\dots}K_{r-1}, \ r = \#classes$$
(1.45)

26

1.2. THE SUPERSELECTION SECTORS OF CG

We now define central "charges":

$$Q_i := \sum_{g \in K_i} g \in \mathcal{Z}(\mathbf{C}G) := \{ z, [z, x] = 0 \quad \forall x \in CG \}$$
(1.46)

It is easy to see that the center $\mathcal{Z}(\mathbf{C}G)$ consists precisely of those elements whose coefficient functions z(g) are constant on conjugacy classes i.e. $z(g) = z(hgh^{-1})$ for all h. The coefficient functions of Q_i :

$$Q_i(g) = \begin{cases} 1 & if \ g \in K_i \\ 0 & otherwise \end{cases}$$
(1.47)

evidently form a complete set of central functions. The composition of two such charges is therefore a linear combination of the r independent $Q'_i s$ with positive integer valued coefficients (as a result of the previous formula (1.46)):

$$Q_i Q_j = \sum_l N_{ij}^l Q_l \tag{1.48}$$

The fusion coefficients N can be arranged in terms of r commuting matrices

$$\mathbf{N}_{j}, \quad with \quad \left(\mathbf{N}_{j}\right)_{i}^{l} = N_{ij}^{l} \tag{1.49}$$

The associativity of the 3-fold product QQQ is the reason for this commutativity, whereas the abelianess of the algebra (only valid for abelian groups!) results in the i - j symmetry of the fusion matrices:

$$Q_i \ central \ \curvearrowright \ Q_i Q_j = Q_j Q_i \ \curvearrowright N_{ij} = N_{ji}$$

Functions on conjugacy classes also arise naturally from characters χ of representations π :

$$\chi^{\pi}(g) = Tr\pi(g), \quad \chi^{\pi}(g) = \chi^{\pi}(hgh^{-1})$$
(1.50)

This applies in particular to the previously defined left regular representation λ with $(\lambda_g x)(h) = x(g^{-1}h)$. Its decomposition in terms of irreducible characters goes hand in hand with the central decomposition of **C**G:

$$\mathbf{C}G = \sum_{l} P_{l}\mathbf{C}G, \qquad Q_{i} = \sum_{l} Q_{i}^{\pi_{l}}P_{l}$$
(1.51)

The central projectors P_l are obtained from the algebraic spectral decomposition theory of the $Q'_i s$ by inverting the above formula. The "physical" interpretation of the coefficients is: $Q_i^{\pi_l} = \pi_l(Q_i)$ i.e. the value of the i^{th} charge in the l^{th} irreducible representation. The central projectors P_l are simply the projectors on the irreducible components contained in the left regular representation. Since any representation of G is also a representation of the group algebra, every irreducible representation must occur in $\lambda(\mathbf{C}G)$. One therefore is supplied with a complete set of irreducible representations, or in more intrinsic terms, with a complete set of r equivalence classes of irreducible representations. As we met the intrinsic (independent of any basis choices) fusion rules of the charges, we now encounter the intrinsic fusion laws for equivalence classes of irreducible representations.

$$\pi_k \otimes \pi_l \simeq \sum_m \tilde{N}_{kl}^m \pi_m \tag{1.52}$$

Whereas the matrix indices of the N's label conjugacy classes, those of \tilde{N} refer to irreducible representation equivalence classes. The difference of these two fusions is typical for nonabelian groups and corresponds to the unsymmetry of the character table: although the number of irreducible representations equals the number of central charges (= # conjugacy classes), the two indices in $\pi_k(Q_a)$ have a different meaning. With an appropriate renormalization this mixed matrix which measures the value of the i^{th} charge in the l^{th} representation we obtain the unitary character matrix $S_{ka} \equiv \sqrt{\frac{|K_a|}{|G|}} Tr\pi_k(g_a)$ (Tr is the normalized trace) which diagonalizes the commuting system of N's as well as $\tilde{N}'s$:

$$\frac{S_{ka}}{S_{0a}} \cdot \frac{S_{la}}{S_{0a}} = \sum_{m} \tilde{N}_{kl}^{m} \frac{S_{ma}}{S_{0a}}, \qquad (1.53)$$

$$\frac{\sqrt{|K_{a}|}S_{ka}}{S_{k0}} \cdot \frac{\sqrt{|K_{b}|}S_{kb}}{S_{k0}} = \sum_{c} N_{ab}^{c} \frac{\sqrt{|K_{c}|}S_{kc}}{S_{k0}}$$

The surprise is that S shows up in two guises, once as the unitary which diagonalizes this $N(\tilde{N})$ -system, and then also as the system of eigenvalues which can be arranged in matrix form. We will not elaborate on this point.

In passing we mention that closely related to the group algebra $\mathbf{C}G$ is the so-called "double" of the group (Drinfeld):

$$D(G) = C(G) \bowtie_{ad} G \tag{1.54}$$

In this crossed product designated by \bowtie , the group acts on the functions on the group $\mathbf{C}(G)$ via the adjoint action:

$$\alpha_h(f)(g) = f(h^{-1}gh)$$
 (1.55)

The dimension of this algebra is $|G|^2$ as compared to dim $\mathbb{C}G = |G|$. Its irreducible representations are labeled by pairs $([\pi_{irr}], K)$ of irreducible representation and conjugacy class and therefore their matrices N and S are *selfdual*. In this sense group doubles are "more symmetric" than groups. In chapter 7 we will meet selfdual matrices S which cannot be interpreted as a double of a group and which resemble the S of abalian groups.

Finally we may notice that the equivalence classes of irreducible representations appear with the natural multiplicity:

$$mult(\pi_l \ in \ \lambda_{reg}) = dim\pi_l \tag{1.56}$$

The results may easily be generalized to compact groups where they are known under the name of Peter-Weyl theory.

Since group algebras are very special, some remarks on general finite dimensional algebras are in order.

Any finite dimensional C^* -algebra \mathcal{R} may be decomposed into irreducible components.and any finite dimensional irreducible C^* -algebra is isomorphic to a matrix algebra $Mat_n(\mathbf{C})$. If the irreducible component $Mat_{n_i}(\mathbf{C})$ occurs with the multiplicity m_i , the algebra \mathcal{R} has the form is isomorphic to the following matrix algebra:

$$\mathcal{R} = \bigoplus_{i} Mat_{n_{i}}(\mathbf{C}) \otimes 1_{m_{i}} \quad in \ \mathcal{H} = \oplus \mathcal{H}_{n_{i}} \otimes \mathcal{H}_{m_{i}}$$
(1.57)

and the multiplicities are unrelated to the dimensionalities of the components. The commutant of \mathcal{R} in H is:

$$\mathcal{R}' = \bigoplus_i \mathbb{1}_{n_i} \otimes Mat_{m_i}(\mathbf{C}), \quad Z := R \cap R' = \bigoplus_i \mathbf{C} \cdot \mathbb{1}_{n_i} \otimes \mathbb{1}_{m_i}$$
(1.58)

The last formula defines the center.

Let us conclude with some remarks on states over (finite dim.) C^* -algebras. Since finite dimensional C^* -algebras decompose into irreducible components (this decomposition agrees with the central decomposition), it suffices to look at states on irreducibles i.e.on the matrix algebra $Mat_n(\mathbf{C})$. The linear functionals can be identified with these matrix spaces since one can use the unique normalized tracial state $\varphi(A) = \frac{1}{n}TrA$ to define a nondegenerate inner product which does this identification. By restricting these linear functionals to be positive and normalized one obtains the well known representation of states in terms of density matrices:

$$\varphi_{\rho}(A) = tr\rho A, \quad \rho \ge 0, \quad tr\rho = 1 \tag{1.59}$$

States are simply positive normalized linear functionals φ on (in the present case finite dimensional) *-algebras \mathcal{A} . From the definition one immediatly abstracts two properties:

$$\varphi(A^*) = \overline{\varphi(A)}$$
$$|\varphi(A^*B)|^2 \leq \varphi(A^*A)\varphi(B^*B), \,\forall A, B \in \mathcal{A}$$
(1.60)

The reality property follows directly from the positivity $\varphi((A+cB)^*(A+cB)) \geq 0$ for arbitrary complex numbers c and the second (Cauchy-Schwartz) inequality follows from the positivity of the sesquilinear form $\varphi((cA+dB)^*(cA+dB))$ in c,d. The validity of the above density matrix representation of φ follows from the linearity which on $Mat_n(\mathbf{C})$ which is spanned in terms of matrix units $e_{ik}, \varphi(A) = \sum a_{ik}\varphi(e_{ik})$ and the positivity which in terms of $\varphi(e_{ik})$ leads to the positivity of the matrix $(\rho)_{ik} \equiv \varphi(e_{ik})$.

In the simplest case of $Mat_2(C)$, which corresponds to the spin algebra generated by the Pauli matrices together with the identity matrix, the convex space

of states (the space of density matrices ρ) is a 3-dim.ball:

$$\rho = \frac{1}{2} (1 + \vec{r}\vec{\sigma}), \quad \vec{r}^2 \le 1$$

$$\sigma_1 = \begin{pmatrix} 0 & 1\\ 1 & 0 \end{pmatrix}, \sigma_2 = \begin{pmatrix} 0 & -i\\ i & 0 \end{pmatrix}, \sigma_3 = \begin{pmatrix} 1 & 0\\ 0 & -1 \end{pmatrix},$$
(1.61)

Besides the normalization we used the positivity of ρ (which requires positive eigenvalues and hence a positive determinant) to derive the inequality. The pure states correspond to one-dimensional projectors and therefore cover the surface of the ball. Note that pure states can be defined intrinsically (i.e.without referring to one-dimensional projectors or state vectors) by the property of indecomposability of φ :

$$\not\exists \varphi_1, \varphi_2 \ s.t. \quad \varphi = \alpha \varphi_1 + \beta \varphi_2, \ \alpha, \beta \ge 0, \quad \alpha + \beta = 1, \ \varphi_1 \neq \varphi_2 \tag{1.62}$$

Starting from $Mat_3(\mathbf{C})$, one meets the new phenomenon of a stratification of the surface into convex subregions called "faces". Density matrices with 3 different eigenvalues correspond to faithfull states inside the convex state space. If one of the eigenvalues vanishes, one looses faithfulness and although these states are "purer", they are not pure in the sense of indecomposability. They form a "face" which looks precisely like the previous ball. In higher dimensions one finds lower dimensional faces inside higher dimensional ones. Once one is on a face, any further purification takes place inside this face. The facial structure is important for recognizing that a normalized convex space is the state space of a C^* -algebra. Returning to the situation of a general finite dimensional \mathbf{C}^* (one for each central component) with normalized total trace.

In order to appreciate the structural differences to classical observables and states, one should remember that the classical observable algebra consists simply of (continuous) functions on phase space and pure states are represented by Dirac δ -functions whereas the Liouville measure and the Radon-Nikodym derivatives with respect to it represent the mixtures and may be viewed as continuously smeared out δ -functions. Since the notion of coherent superposition is absent, a classical analogon of the states for the $Mat_n(\mathbf{C})$ toy models would be a higher dimensional simplex. The pure states are the vertices and every mixed state is obtained by a unique convex combination of the vectors corresponding to the vertices. This is quite different from the ball-shaped region of quantum physics where the representation of a point in the interior in terms of pure states on the surface is highly nonunique. In fact this structure of states is, more than anything else, the most characteristic property of quantum physics. The presence of superselection rules tells us that at least partially there exists a classical structure within quantum physics: the central decomposition of a C^* algebra is unique (similar to the classical case), and the unrestricted superposition principle holds only within each component. In fact this classical aspect of the central decomposition plays an important role in the measurement process and the notion of "commuting histories".

30

The concepts needed in the infinite dimensional case are more subtle and will be presented in a mathematical appendix.

The reader may ask the question of why superselection rules despite their importance are rarely mentioned in quantum mechanics. The reason is the validity of the Stone-von Neumann uniqueness-theorem on irreducible (regular) representations of the Schrödinger theory. In QM of finite degrees of freedom it is only only through topological nontriviality of configuration- or phase-space that inequivalent representations may enter the quantization procedure. Let us look at a typical example, the quantum mechanics of a particle in a circle. The geometric argument in favour of finding many inequivalent representations is as follows. Diagonalizing first the algebra of the position operator we represent the state vectors by (periodic) wave functions $\psi(\varphi)$ on S^1 . In order to fulfill the Heisenberg-Weyl commutation relations, the most general form for the momentum is $p = i \frac{\partial}{\partial \varphi} + f(\varphi)$. Wheras on **R** the f can be (gauge-)transformed away, the best we can do on a circle is to transform to a constant. In n dimension the n-component f_i has to fulfill the flatness condition $\partial_i f_i - \partial_j f_i = 0$. the can be transformed The p only commutes with itself iff the real function f is a constant (in higher dimensions the p would be like a gauge covariant derivative, and the constancy of f like the flatness of a connection or the vanishing of the field strength associated with a vector potential). In our one dimensional example on S^1 , the p which generates the circular translation obeys the famous θ -angle formula:

$$p = i\frac{\partial}{\partial\varphi} + \theta, \ \theta \ mod \ 2\pi \tag{1.63}$$

Hence there are many Schrödinger theories corresponding to different selfadjoint extensions of the nonselfadjoint hermitian differential operator parametrized by a theta-angle which have different physical content (e.g. the spectrum of $H_0^{(S_1)}$). An equivalent form is obtained by keeping the standard Schrödinger-form of the operator p but accepting quasiperiodic wave functions. The " θ -obstruction" is intrinsic, it can be shifted from the algebra to the states, but only in a simply connected space it can be removed alltogether by a nonsingular operator transformation; this is the case in ordinary Schrödinger theory.

This geometric viewpoint has one disadvantage: there is no natural way to consider the different θ -theories as just different manifestations of one system, rather they are different geometrical objects (generally inequivalent vectorial fibre bundles). Here the *algebraic view* of superselection rules is *physically superior*: the different θ -theories are simply different representations of one abstract C^* algebra (in our case the "rotational algebra").

The mathematical example of a particle on a circle is closely related to the Aharonov-Bohm effect. As long as the solenoid has not passed to the infinite thin limit, the A-B system falls into the ordinary Schrödinger description. It is only through the *limiting overidealization* that the θ -dependent circular mechanics with its nonsimple C*-algebra enters. There is a general message in this example: all topologically nontrivial quantum mechanics result from an overidealization of Schrödinger theory. Only for infinite degrees of freedom in QFT it becomes possible to encounter superselection rules which have a natural

fundamental origin (e.g. a phase transition). We refer to section 5 of the last chapter where in particular the globalization through the universal observable algebra (which mimics the geometrical compactification) comes close to topologically nontrivial models of QM if one only retains the global (nonlocalizable) degrees of freedom.

One lesson to be drawn for QFT from these illustrations is that one is not limited by "quantization" methods. Rather one may use the superselection idea and try to classify and construct QFT's by studying representations of observable algebras instead of quantizing classical physics. It was realized by Haag, Araki, Borchers and others [3] already at the beginning of the 60's that the principle of locality makes such a formulation very consistent and structurally rich. But the path from those early studies to the more recent advances in e.g. properties of low dimensional QFT with new surprising nonperturbative insights was very thorny indeed. Our approach to QFT and even these remarks on QM are strongly influenced by this "algebraic QFT". In particular the problem of particle statistics will be presented as part of the understanding of superselection charges.

1.3 Illustration of Important Quantum Concepts

In this section some additional quantum-physical concepts will be introduced in a finite dimensional setting (for simplicity). For the much richer infinite dimensional concepts used to analyse C^* - and von Neumann algebras, we refer to the mathematical appendix.

We start with the GNS-construction. It associates in a canonical way with a given \mathbb{C}^* -algebra A and a state ω on it a so called GNS-tripel $(\mathcal{H}_{\omega}, \pi_{\omega}(\mathcal{A}), \Omega_{\omega})$ which consists of a representation space \mathcal{H}_{ω} with a distinguished vector Ω_{ω} on which A through its representation $\pi_{\omega}(\mathcal{A})$ acts cyclically. The construction is analogous to that of the regular representation of $\mathbb{C}G$. Again one uses the state ω in order to construct a positive semidefinite sesquilinear form on the linear space of A:

$$(\psi_A, \psi_B) := \omega(A^*B) = \overline{\omega(B^*A)} \tag{1.64}$$

Here we use a notation which distinguishes the vectors from the elements of the algebra. The reason is that whereas the tracial state φ on the CG was faithfull i.e. the sequilinear form was strictly positive, a general state ω on A leads to a nontrivial nullspace N_{ω}

$$\mathcal{N}_{\omega} = \{A \in A \mid \omega(A^*A) = 0\}$$

$$(1.65)$$

Fortunately this nullspace N_{ω} is also a left ideal (the Gelfand ideal) of A i.e. with $A \in \mathcal{N}_{\omega}$ also $BA \in \mathcal{N}_{\omega}$ for any $B \in \mathcal{A}$. This follows from the Cauchy-Schwartz inequality for states (1.60):

$$\left|\omega(A^*B)\right|^2 \le \omega(A^*A)\omega(B^*B) \tag{1.66}$$

if we write the latter in the adapted form:

$$|\omega((BA)^*BA)|^2 \le \omega(C^*C)\omega(A^*A), \quad C := ((BA)^*B)^*$$
(1.67)

In this form it is obvious that the left hand side must vanish if $A \in \mathcal{N}_{\omega}$. The proof of the Cauchy-Schwartz inequality for states is identical to that for scalar products of vectors in Hilbert space.

If one now defines the state vectors ψ as elements of A mod N_{ω} and the action of A on these vectors as:

$$\pi_{\omega}(B)\psi_A := \psi_{BA}, \quad B \in \mathcal{A} \tag{1.68}$$

then one obtains the desired relation between the state and the representation:

$$(\Omega_{\omega}, \pi_{\omega}(A)\Omega_{\omega}) = \omega(A) \tag{1.69}$$

Here Ω_{ω} is the distinguished vector in \mathcal{H}_{ω} which corresponds to **1** (vanishing Gelfand ideal on Ω_{ω}). The only additional step for infinite dimensional algebras is to form the closure of the linear space and to continue the definition of $\pi_{\omega}(A)$ to this Hilbert space closure of $\mathcal{A} \mod \mathcal{N}_{\omega}$. Since a dense set of vectors is obtained by applying A to Ω_{ω} , the proof is finished. It remains to be added that every other cyclic representation $\pi_{\omega}(A)$ with the same $\omega(A)$ turns out to be unitary equivalent to the canonical GNS representation.

As a preparation for the next topic let us use a tracial state as a reference state. On a factor $Mat_n(C) \otimes \underline{1}_m$ in the central decomposition of a semisimple algebra a tracial state is unique and has the standard form:

$$\tau(A) = \frac{1}{n} TrA \equiv trA, \quad TrA = \sum_{i} A_{ii}$$
(1.70)

On a semisimple algebra there is a family of tracial states parametrized by $\lambda_i \ge 0$, $\sum_i \lambda_i = 1$:

$$\tau_{(\lambda_1 \lambda_2 \dots \lambda_k)}(A) = \sum_{i=1}^k \lambda_i \tau(A_i)$$
(1.71)

$$\tau_{\left(\frac{n_i}{N},\frac{n_2}{N}\dots\frac{n_k}{N}\right)}\left(A\right) =: \tau(A) = \frac{1}{N}TrA, \quad N = \Sigma_i n_i \tag{1.72}$$

where \mathcal{A}_i is the *i*th central component of \mathcal{A} and τ is the standard faithful tracial state. We have seen before that the most general state φ may be described in terms of a density matrix ρ :

$$\varphi(A) = \tau(\rho A) \tag{1.73}$$

This formula is just a concrete realization of the GNS construction via the trace formalism (in which case the nullspace vanishes and the correspondence between matrices and vectors in the representation space is one to one):

$$\tau(A^*B) = (\Omega_\tau, \pi_\tau(A^*)\pi_\tau(B)\Omega_\tau)$$

$$\varphi(A) = (\Omega_\varphi, \pi_\varphi(A)\Omega_\varphi) = \tau(\rho^{\frac{1}{2}}A\rho^{\frac{1}{2}})$$

$$\Omega_\omega = \rho^{\frac{1}{2}}\Omega_\tau, \quad \Omega_\tau = \underline{1} \quad \text{in space of matrices}\mathcal{H}_\tau$$

(1.74)

Note the analogy of this construction with the regular representation of $\mathbf{C}G$: in both cases the algebra in its role as a space, together with the trace, served as the the arena for GNS faithful representations. Such a Hilbert space \mathcal{H}_{τ} may be written in a more suggestive L^2 -notation:

$$\mathcal{H}_{\tau} = L^2(Mat_N, \tau), \quad \Omega_{\tau} = \underline{1} \tag{1.75}$$

In addition to the "left" GNS representation $\pi_l = \pi_\tau = \pi$, we introduce an antilinear right representation:

$$\pi_r(B)\psi_A = \psi_{AB^*} = \pi_\tau(A)\underline{1}\pi_\tau(B^*)$$
(1.76)

In the group case, the right action unravels the multiplicity structure of the irreducible representations (the irreducible representations occur according to the Peter-Weyl theory with a multiplicity identical to the dimension of the representation as a result of the complete symmetry between left and right action on $\mathbb{C}G$). In our present more general setting, the existence of the two commuting left-right actions furnish the germ of a deep and general theory: the Tomita-Takesaki modular theory. One first defines the antiunitary "flip" operator J:

$$J\psi_A = \psi_{A^*} \frown j(\pi(A)) := J\pi(A)J = \pi_r(A)$$
(1.77)

In this case the J not only implements the flip, but it also transforms the vector $A\Omega_{\tau}$ into $A^*\Omega_{\tau}$. In the more general φ -representation one has two different involutive operators S and J:

$$S\pi(A)\Omega_{\varphi} := \pi(A^*)\Omega_{\varphi} \quad \Omega_{\varphi} = \rho^{\frac{1}{2}} \underline{1}$$

$$J\pi(A)J = \pi_r(A)$$
(1.78)

Fom this one reads off the new operator S in terms of J and ρ (1.73):

$$S = J\Delta^{\frac{1}{2}}, \quad \Delta = \pi(\rho)\pi_r(\rho^{-1}) \tag{1.79}$$

since the identity: $S\pi(A)\rho^{\frac{1}{2}}\underline{1} = J\rho^{\frac{1}{2}}\pi(A)\rho^{\frac{1}{2}}\underline{1}\rho^{-\frac{1}{2}} = \pi(A^*)\rho^{\frac{1}{2}}\underline{1}$ follows from the definitions. The formula for S agrees with the polar decomposition formula into an "angular" part J and positive "radial" part $\Delta^{\frac{1}{2}}$. The above formulas require ρ to be invertible i.e. the φ -representation to be faithful (or equivalently Ω_{φ} to be a separating vector in \mathcal{H}_{τ}). Remembering the notation of the von Neumann commutant \mathcal{A}' and the algebraic role of these operators, the following important properties are an easy consequence:

$$j(\mathcal{A}) = \mathcal{A}', \quad \sigma_t(A) := \Delta^{it} A \Delta^{-it} \in \mathcal{A}$$
 (1.80)

 $j(\cdot)$ is the modular conjugation and $\sigma_t(\cdot)$ is called the modular group. Δ^{it} commutes with J and hence the action of the latter on the commutant \mathcal{A}' is obtained by replacing Δ^{it} by its inverse. Now every ρ may be written as a Gibbs formula in terms of a (ad hoc) hamiltonian H:

$$\rho = \frac{1}{Z} e^{-\beta H}, \quad Z = T r e^{-\beta H} \tag{1.81}$$

In this case $\Delta^{it} = \pi_l(e^{-i\beta tH})\pi_r(e^{i\beta tH})$ and the modular automorphism on \mathcal{A} is apart from a stretching factor $-\beta$ equal to the hamiltonian automorphism. Note that the infinitesimal generator \mathbf{H} of the time translations is not simply H but rather: $\mathbf{H} = \pi_l(H) \otimes \mathbf{1} \oplus \mathbf{1} \otimes \pi_r(-H)$. This fact becomes important in the realistic (infinite dimensional) case, since the heat bath fluctuations of the hamiltonian H become infinitly large in the thermodynamical limit. The generator of the modular automorphism acts on the algebra as well as (in opposite fashion) on its commutant. In the context of thermal physics: the generator of time translation is the sum of the hamiltonian and its opposite. Only in this way the formalism controlls the fluctuations of \mathbf{H} , wheras those of the hamiltonian H have the well-known volume divergencies.

It is therefore not surprising that the modular theory was discovered independently (and at the same time) by physicists in the study of temperature states on bosonic or fermionic algebras.

In our toy case, instead of writing down the Gibbs formula, one may also characterize the faithful state φ by the so-called KMS condition:

$$\varphi(A\sigma_t(B)) = \varphi(\sigma_{t+i\beta}(B)A) \tag{1.82}$$

In the finite dimensional case the analytic dependence of σ on t is automatic whereas for the general case the analyticity of a function $F_z(A, B) := \varphi(A, \sigma_z(B))$ (for A, B from a dense subalgebra of \mathcal{A}) and the formulas:

$$F_{z}(A; B) \text{ analytic in strip: } 0 \leq \operatorname{Im} z \leq \beta$$

with $F_{z} = \begin{cases} \varphi(A\sigma_{t}(B)) \text{ for Im } z = 0 \text{ i.e. } z = t + i0 \\ \varphi(\sigma_{t+i\beta}(B)A) \text{ for Im } z = t + i\beta \end{cases}$ (1.83)

constitute part of the *definition* of the KMS property of φ . This characterization of thermal equilibrium states is more general than the Gibbs formula, since the latter looses its meaning as a result of the volume divergencies in the thermodynamic limit $V \rightarrow \infty$ with the particle densities kept fix. In addition, even in case of finite volume, the calculation of φ by KMS boundary condition is much easier than by calculating traces. This practical advantage was the reason why Kubo, Martin and Schwinger introduced this condition, whereas the mathematical physics connection was made much later by Haag, Hugenholtz and Winnink [3].

Besides the notion of states and representation, the concept of inclusions of von Neumann algebras will play an important role in later sections. Here we will only present a "baby" version. Suppose that $Mat_2(\mathbf{C})$ acts not on its natural irreducible space \mathbf{C}^2 but by left action on the 4-dim Hilbert space $\mathcal{H}(Mat_2(\mathbf{C}), \frac{1}{2}Tr)$. In that space the commutant is of equal size and consists of $Mat_2(\mathbf{C})$ acting in the opposite order from the right which will be shortly denoted as $Mat_2(\mathbf{C})^{opp}$. Explicitly the realization of \mathcal{H} as \mathbf{C}^4 may be defined as

$$\begin{pmatrix} \xi_{11} & \xi_{12} \\ \xi_{21} & \xi_{22} \end{pmatrix} \rightarrow \begin{pmatrix} \xi_{11} \\ \xi_{21} \\ \xi_{12} \\ \xi_{22} \end{pmatrix}$$
(1.84)

and the action of $\mathcal{A} = Mat_2(\mathbf{C})$ takes the following form:

$$a = \begin{pmatrix} a_{11} & a_{12} & 0 & 0\\ a_{21} & a_{22} & 0 & 0\\ 0 & 0 & a_{11} & a_{12}\\ 0 & 0 & a_{21} & a_{22} \end{pmatrix} \simeq \begin{pmatrix} a_{11} & a_{12}\\ a_{21} & a_{22} \end{pmatrix} \otimes \underline{1}$$
(1.85)

The most general matrix in the commutant $a' \in \mathcal{A}'$ has evidently the form:

$$a' = \begin{pmatrix} a'_{11} & 0 & a'_{12} & 0\\ 0 & a'_{11} & 0 & a'_{12}\\ a'_{21} & 0 & a'_{22} & 0\\ 0 & a'_{21} & 0 & a'_{22} \end{pmatrix} \simeq \underline{1} \otimes \begin{pmatrix} a'_{11} & a'_{12}\\ a'_{21} & a'_{22} \end{pmatrix}$$

The norm $\|\xi\| = \left(\frac{1}{2}Tr\xi^*\xi\right)^{\frac{1}{2}}$ is invariant under the involution $\xi \to \xi^*$ which in the \mathbb{C}^4 representation is given by the isometry:

$$J = \begin{pmatrix} K & 0 & 0 & 0 \\ 0 & 0 & K & 0 \\ 0 & K & 0 & 0 \\ 0 & 0 & 0 & K \end{pmatrix}, \quad K : \text{natural conjugation in } \mathbf{C}$$
(1.86)

We have:

$$j(\mathcal{A}) := J\mathcal{A}J = \mathcal{A}', \quad \text{antilin. map } \mathcal{A} \to \mathcal{A}'$$
 (1.87)

which may be rewritten in terms of a linear anti-isomorphism:

$$a \to Ja^*J, \quad \mathcal{A} \to \mathcal{A}'$$
 (1.88)

Consider now the trivial algebra $\mathcal{B} = \mathbf{C} \cdot \mathbf{1}_2$ as a subalgebra of $\mathcal{A} = Mat_2(\mathbf{C})$. In the \mathbf{C}^4 representation the B-algebra corresponds to the subspace:

$$\mathcal{H}_{B} = \left\{ \begin{pmatrix} \xi \\ 0 \\ 0 \\ \xi \end{pmatrix}, \xi \in \mathbf{C} \right\}, \quad \mathcal{H}_{B} = e_{B}\mathcal{H}, \quad e_{B} = \begin{pmatrix} \frac{1}{2} & 0 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \frac{1}{2} & 0 & 0 & \frac{1}{2} \end{pmatrix} \quad (1.89)$$

The projector e_B commutes clearly with \mathcal{B} i.e. $e_B \in \mathcal{B}'$. We now define a measure for the relative size of $\mathcal{B} \subset \mathcal{A}$ the Jones index:

$$[A:B] = \tau_{B'}(e_B)^{-1}, \quad \tau: \text{ normalized trace in } \mathcal{B}'$$

In our example $\tau(e_B) = \frac{1}{4}(\frac{1}{2} + 0 + 0 + \frac{1}{2}) = \frac{1}{4}$ i.e. the satisfying result that the Jones index is 4. The same method applied to the inclusion:

$$Mat_4(\mathbf{C}) \supset Mat_2(\mathbf{C}) \otimes \mathbf{1}_2 = \left\{ \begin{pmatrix} X & 0 \\ 0 & X \end{pmatrix}, X \in Mat_2(\mathbf{C}) \right\}$$
(1.90)

also gives the expected result:

$$[A:B] = \frac{\dim Mat_4(\mathbf{C})}{\dim Mat_2(\mathbf{C})} = 4$$
(1.91)
If, as in the previous cases B is a finite dimensional subfactor (i.e.a full matrix algebra) of A, the Jones index is the square of a natural number. For inclusions of finite dimensional semisimple algebras the index takes on more general values. For example:

$$Mat_{2}(\mathbf{C}) \oplus \mathbf{C} = \begin{pmatrix} X \\ X \\ x \end{pmatrix} \subset Mat_{2}(\mathbf{C}) \oplus Mat_{3}(\mathbf{C}) \quad (1.92)$$
$$X \in Mat_{2}(\mathbf{C}), \quad x \in \mathbf{C}1$$

Here the index is 3. It is easy to see that instead of the projector formula one may also use the incidence matrix formula:

$$\left[\mathcal{A}:\mathcal{B}\right] = \left\|\Lambda_n^m\right\|^2 \tag{1.93}$$

The incidence matrix Λ is describable in terms if a bipartite graph. The number of say white vertices correspond to the number of full matrix component algebras for the smaller algebra and the black vertices labelled by the size of the components to the analogously labelled irreducible components of the bigger algebra. A connecting line between the two sets of vertices indicates that one irreducible component of the smaller is included into one of the bigger algebra. In our case:

$$\Lambda = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \quad ||\Lambda||^2 = 3 \tag{1.94}$$

From a sequence of ascending graphs one obtains important infinite graphs (Bratteli diagrams) which are very useful in the "subfactor theory" [26] which will appear in the mathematical appendix. In the infinite dimensional case the inclusion of full matrix algebras corresponds to the inclusion of von Neumann factors i.e. the "subfactor problem". In that case the spectrum of inclusions shows a fascinating and unexpected quantization phenomenon, the Vaughn Jones quantization formula for index ≤ 4 . AFD (almost finite dimensional) C*-algebras obtained by sequences of ascending Bratteli diagrams equipped with tracial states enter LQP via the intertwiner algebra of charge transporters. A special case are the combinatorial theories which result from Markov-traces on selfintertwining transporters which contain the braid group and mapping class group generators for arbitrary genus. They are better known under the name "topological field theories" (their differential geometric name). Their physical role in 3-dimensional plektonic LQP will be briefly mentioned in chapter 7.

1.4 Measurement and Superselection Rules

The interpretation of quantum theory requires an observer which also may be a registration apparatus outside the observed system. Therefore notions like the "state or wave function of the universe" have to be handled with great care and are in some cases meaningless, at least within the standard physical interpretation of QFT. Somewhere between the (generally microscopic) observed system

and the observer a "cut" is needed. As already pointed out by Heisenberg, this cut may be somewhat shifted, but it must be there somewhere.

According to von Neumann the observed system is described by a selfadjoint operator and the measured values are the eigenvalues α of the "observable" A (with suitable mathematical adaptation in case of continuous spectrum). The state immediately after the measurement is obtained by a "quantum jump" i.e. cannot be computed via the Schrödinger time evolution of the observed system. Taking for A a projector P whose eigenvalues are just 0 and 1, the new state created by the measurement according to von Neumann can be described as follows:

$$\omega_{after}(A) = \omega_{before}(PAP) + \omega_{before}((1-P)A(1-P)),$$

 $A \in \mathcal{A}$, algebra of observables (1.95)

If ω_{before} was a pure state described by a state vector ψ , ω_{after} corresponds to the density matrix:

$$\rho_{after} = |P\psi\rangle \langle P\psi| + |(1-P)\psi\rangle \langle (1-P)\psi| =$$

$$= p_1 |\psi_1\rangle \langle \psi_1| + p_2 |\psi_2\rangle \langle \psi_2|$$
(1.96)

$$\psi_{1} = \frac{P\psi}{||P\psi||}, \ p_{1} = ||P\psi||^{2},$$

$$\psi_{2} = \frac{(1-P)\psi}{||(1-P)\psi||}, \ p_{2} = ||(1-P)\psi||^{2}$$
(1.97)

The last formula represents the mixed state associated with ρ_{after} as a sum of two orthogonal minimal projectors. However, as stressed before, an impure state permits myriads of decompositions into minimal projectors. If we could find a physical argument in favor of an orthogonal decomposition (as for the one above in terms of ψ_1, ψ_2) then uniqueness follows. But no such principle is known. In addition, to have a change of states as the above reduction of wave packet (or "quantum jump") which is outside the unitary time development of the Schrödinger equation is somewhat mysterious and in order to sharpen this seemingly paradoxical situation Schrödinger replaced the pointer of a measuring apparatus by a cat with the more dramatic alive/dead state replacing the two pointer positions whose coherent superposition defies common sense. Any hamiltonian dynamics leading to unitary propagation in time, necessarily preserves the purity of states.

The important role of superselection rules for a resolution of these mysterious aspects has been first pointed out and illustrated by a mathematical model in the work of Hepp. The essential idea is that the macroscopic measuring apparatus has superselection sectors² which, as we have learned, is a generic phenomenon for systems with infinitely many degrees of freedom. Hepp's idea

 $^{^{2}}$ The physically realizable observable algebra which is restricted by locality may be genuinly smaller than the (more global) mathematical algebra. Therefore the superselection rules in the measurement process should be understood in the "effective" sense.

is that although the complete system including the apparatus is governed by a unitary time development :

$$\alpha_t(A) = e^{iHt} A e^{-iHt}, \ A \in \mathcal{A}$$
(1.98)

the limit for $t \to \infty$ may well be only *a positive map* instead of an automorphism of the observable algebra. In terms of Hilbert space concepts the limit of e^{iHt} may only be an isometric mapping of the total Hilbert space on a subspace. Therefore the initial state which may be a pure state implemented by a vector in one superselection sector leaves the coherent subspace in the limit and acquires components in other superselection sectors. Although it remains formally a vector in the total space, it describes physically a mixture since it has projections to several coherent subspaces.

$$\omega_{before} \to \omega_{after} = \sum_{i} \lambda_i \omega^i_{after} \tag{1.99}$$

Note that this central decomposition is completely intrinsic. It is a special case of a partial orthogonal decomposition. Whereas the latter is only unique within one superselection sector, the former is unique in general and in this sense behaves like a classical decomposition.

Although the more modern treatment of the measuring process for good reasons has discarded Hepp's idea about the importance of the superselection rules in the Hilbert space of the apparatus and the $t \to \infty$ limit [44], the crucial role of superselection rules and the uniquely defined central decompositions has remained. In the modern treatment these properties are generated by the "environment" of the apparatus i.e. the apparatus is considered as an open system which remains in contact with its infinite degree of freedoms surroundings (which also may consist of the unobserved internal degrees of freedom). Therefore the total Hilbert space consists of three tensor factors: the system to be observed and the apparatus, both localized, and the infinite degree of freedom environment which is best described by methods of QFT. As an infinite degree of freedom system, the environment has natural superselection sectors (in distinction to topological nontrivial QM's, which owes its superselection structure to unnatural physical overidealizations viz. the Aharonov-Bohm system) and the mathematical physics method is similar to that of determining the various possible phases in statistical mechanics models of phase transitions. The measurement interaction correlates first the system S with the apparatus A and then with the environment E, in case of a two-valued spin system

$$(\alpha |\uparrow\rangle + \beta |\downarrow\rangle) |A_I\rangle \otimes |E_I\rangle \rightarrow ((\alpha |\uparrow\rangle |A_\uparrow\rangle + \beta |\downarrow\rangle) |A_\downarrow\rangle) \otimes |E_I\rangle$$

$$(1.100)$$

$$\rightarrow \alpha |\uparrow\rangle |A_\uparrow\rangle \otimes |E_\uparrow\rangle + \beta |\downarrow\rangle |A_\downarrow\rangle \otimes |E_\downarrow\rangle \equiv |\Psi\rangle$$

Interactions which accomplish such a Schrödinger transition can be constructed by suitable idealizations of the apparatus A and the environment E. The last step which destroys the purity of the system S + A is the averaging over the degrees of freedom of the environment E (to observe S is to not observe E):

$$\rho_{S,A} = Tr_E \left| \Psi \right\rangle \left\langle \Psi \right| \tag{1.101}$$

The trace stands symbolically for an averaging which is better described (the trace is only well defined for type II von Neumann algebras) in terms of a conditional expectation. It is important to note that the collapse of the wave packet is not instantaneous [45]. In order to find the nonunitary time development of S we assume that the total initial state is a product state $\omega = \tilde{\omega} \otimes \sigma$

Here the von Neumann factualization of the measurement has been accomplished by the nonobservation of E. Since the crucial role is played by the superselection sectors of the apparatus + environment (the observed system Sthrough its coupling with A + E only activates the superselection sectors of the latter), the question about the origin of the superselection sectors arises. The standard sectors of infinite degrees of freedom models of statistical mechanics are identical to the various phases in the theory of phase transition. As mentioned before, in the environmental case a detailed consideration reveals that the relevant observable algebra of E (resp.E + A) is not the algebra of a full tensor factor H_E , but rather a certain C^* -subalgebra of $B(H_E)$, since the operators which are nonlocal with respect to the observer (those which could monitor phase relations with a state of which one component is spatially far separated from the rest) are missing [44]. So the restriction of the E-observables due to locality turns out to be crucial for the understanding of the von Neumann collapse of the wave packet.

In the environmental approach the "reduction of the wave packet" is achieved in the limit $t \to \infty$, i.e. it is a time-dependent process. Assume that the initial state is a product state $\omega = \tilde{\omega} \otimes \sigma$ on the total observable algebra $\mathcal{A} = \mathcal{A}_{SA} \otimes \mathcal{A}_E$. Restricting ω to the subalgebra $\mathcal{A}_{SA} \equiv \mathcal{A}_{SA} \otimes 1_E$ we can compute the time development of $\tilde{\omega}$ from the original unitary automorphism α_t as:

$$\tilde{\omega}_t(A) = \omega(\alpha_t \left[A \otimes 1_E \right]), \quad A \in \mathcal{A}_{SA} \tag{1.102}$$

One can define a reduced time development on the restricted state as:

$$\tilde{\omega}_t(A) = \tilde{\alpha}_t^*(\tilde{\omega}) \tag{1.103}$$

This can be recast by "dualization" into a nonautomorphic time development in terms of a positive map $\tilde{\alpha}_t$ on \mathcal{A}_{SA}

$$\tilde{\alpha}_t [A] = E_{\sigma}(\alpha_t [A \otimes 1_E])$$

$$E_{\sigma}(A \otimes B) = \sigma(B)A, \quad E_{\sigma}^2 = E_{\sigma}, \quad \omega \circ E_{\sigma} = \omega$$
(1.104)

Here E_{σ} is a ω -preserving conditional expectation which is defined on all $\mathcal{A} \to A_{SA} \subset \mathcal{A}$ by linear extension of the above definition on tensor products. Typical properties, as the above projection property, the formula $E_{\sigma}(ABC) = AE_{\sigma}(B)C$ if $A, C \in \mathcal{A}_{SA}$ as well as the positivity: $E_{\sigma}(A^*A) \geq E_{\sigma}(A^*)E_{\sigma}(A) \geq 0$ follow from this definition. In order to calculate decoherence times τ_D which measure the duration of the collapse, we need a model. An interesting phenomenological class of models which has been used for the first very recent experiments [46] on Schrödinger cat states (rather on Schrödinger "kittens", because the apparatus contained only a few number of photons) is that proposed by Zurek and

others[45]. The effective equation of motion projected onto the one particle density matrix is the master equation:

$$\frac{d\rho}{dt} = -\frac{i}{\hbar} \left[H,\rho\right] - \gamma(x-x') \left(\frac{\partial\rho}{\partial x} - \frac{\partial\rho}{\partial x'}\right) - \frac{2m\gamma k_B T}{\hbar^2} (x-x')^2 \rho \qquad (1.105)$$

The second term is a dissipation term which results from an interaction of a particle with a field which represents the infinite degrees of freedom of the environment and γ is the relaxation rate resulting from such an interaction. T is the temperature of this field. One finds a relaxation time $\tau_D \simeq \tau_R \frac{\hbar^2}{2mk_B T(\Delta x)^2}$ with $\tau_R = \gamma^{-1}$ where Δx is the nondiagonal separation for the initial pure state which has been described by coherent superposition of two gaussian wave functions $\psi(x) = \psi^+(x) + \psi^-(x)$ whose peaks have a distance Δx . Therefore the initial density matrix is $\rho(x, x') = \psi(x)\psi^*(x')$ and the decoherence time is the time in which the loss of the coherent off diagonal terms i.e. the collapse has taken place.

Closely related to the measurement process and the role of superselection sectors is the problem of why molecules have a "nuclear frame", i.e. why the nuclei do not have fuzzy positions corresponding to coherent superpositions of position eigenstates. This "Gestaltproblem" is mathematically connected to the validity and interpretation of the Born-Oppenheimer approximation. For its conceptual understanding the environment and the superselection rules play a similar role as in the measurement process [47].

We have seen that the important concepts in the measurement process are open systems (the environment) and locality (responsible for the superselected \mathcal{A}_E subalgebra of $B(H_E)$). In QM these concepts have to be enforced from the outside (by forming tensor products etc.) whereas in QFT there is an intrinsic concept of openness. For example the split property discussed in the last chapter and the affiliated statistical independence as well as the relativistic localization concept and superselaction sectors are intrinsic natural properties of QFT which allow to view the outside environment as a split tensor factor of one QFT. So the proper arena for the analysis of the measurement process seems to be QFT. But apart from a beautiful structural discussion of the significance of the Bell inequalities [48], there has been no series attempt to base the maesurement investigation on the characteristic modular properties of QFT which recently have given consderable insight into thermal and entropical aspects of modular localization (see chapter 3).

Literature for chapter 1

R. Haag, "Local Quantum Physics" Springer Verlag, Berlin Heidelberg 1992 O. Bratteli and D. W. Robinson, "Operator Algebras and

F. Goodman, P. de la Harpe and V. F. R. Jones, "Coxeter graphs and towers of algebras, MSRI Publications (Springer) **14** (1989)

V. F. R. Jones, "Subfactors and Knots", CMBS Number 80, American Mathematical Society.

V.F. Jones and V.S. Sunder, "Introduction to Subfactors", London Mathematical Society, Lecture Note Series 234.

42

Chapter 2

The Construction of Fock Space.

2.1 The Bosonic Fock Space

There are several reasons for combining *N*-particle spaces together with a onedimensional "no-particle" space (vacuum) into a big "Master" space, the so called Fock-space:

$$\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \dots$$
(2.1)

One obvious reason is that relativistic local interactions do not conserve the particle number but only total charges (i.e. particle-antiparticle creation is allowed as long as it obeys the energy-momentum conservation). This is also valid in the infinite volume limit (the so called thermodynamic limit) in nonrelativistic systems for which the ground state (which has generally a finite density of particles) becomes the reference state for generally number-nonconserved (but charge-conserved) "quasiparticle" excitations. Another less obvious reason is that the Fock space together with locality is also the natural setting for the formulation of cluster-properties which are the most important imprints of local quantum physics on QM. The textbook treatment of e.g. scattering theory in QM strictly speaking applies to short range interactions because in that case the cluster requirements on the S-matrix are automatically taken care of by the plane wave boundary conditions, but not so for long range interactions. For this reason the use of e.g. the Aharonov-Bohm interaction between "dyons" in order to produce exotic (anyonic) statistics is not treatable as a standard quantum mechanical problem for fixed particle number n. Rather the main unsolved problem is the elaboration of stationary scattering boundary conditions which makes the n-particle problem for the S-matrix physically consistent with the clustering of the n+1 particle scattering for all n. It is doubtful that such nonrelativistic problems without real (mass shell) particle creation, but with the effective particle number change caused by shifting particles to infinity, can be solved without the use of a field theoretic framework.

The bosonic Fock-space $\mathcal{H}^{\mathcal{B}}$ is obtained by projecting the full *N*-particle spaces onto their symmetrized subspaces \mathcal{H}_N^B . In the following we will introduce the creation and annihilation operator formalism in x-space, having in mind wave functions in Schrödinger theory. If we interpret the formulas in momentum space however, there will be no difference between the relativistic and nonrelativistic formalism apart from possible changes in the normalization. A general vector in Fock-space is given by a finite norm sequence of symmetric wave functions:

$$\mathcal{H}_{F}^{B} \ni \Psi = (\psi_{0}, \psi_{1}(\vec{x}), \psi_{2}(\vec{x}_{1}, \vec{x}_{2}), \dots) (\Psi, \Psi) := |\psi_{0}|^{2} + \sum_{n=1}^{\infty} (\psi_{n}, \psi_{n}) < \infty$$
(2.2)

The creation operator depends linearly on the wave function f and the n-particle component after its application on Ψ is defined as:

$$(a^*(f)\Psi)_n(\vec{x}_1,....,\vec{x}_n) = \frac{1}{\sqrt{n}} \sum_{i=1}^n f(\vec{x}_i)\psi_{n-1}(\vec{x}_1...\vec{x}_i...\vec{x}_n), \quad n \ge 1,$$

$$(a^*(f)\Psi)_0 = 0$$
(2.3)

Here the "roof" $\hat{}$ indicates deletion of the i-th coordinate. The formula for the hermitian adjoint annihilation operator a(f) follows from the defining property:

$$\left(\Psi, a^*(f)\Psi'\right) = \left(a(f)\Psi, \Psi'\right), \text{ namely} (a(f)\Psi)_n (\vec{x}_1....\vec{x}_n) = \sqrt{n+1} \int d^3x \bar{f}(\vec{x})\psi_{n+1}(\vec{x}, \vec{x}_1....\vec{x}_n)$$
(2.4)

The annihilation operator depends antilinear on f. In particular for the vacuum $\Omega = (1, 0, 0...)$

$$(a(f)\Omega)_n = 0 \quad \forall n \tag{2.5}$$

i.e. a(f) annihilates Ω .

The multiple application of these operators leads to lengthy formulas, however the commutators are very simple:

$$\begin{aligned} [a(g), a^*(f)] &= (g, f) \, 1, \quad \text{with} \quad (g, f) = \int \bar{g}(\vec{x}) f(\vec{x}) d^3x \\ [a(f), a(g)] &= 0 = [a^*(g), a^*(f)] \end{aligned}$$
 (2.6)

This simplicity was the reason for the choice of normalization in the definition of a^* .

The number operator **N** is defined to be that positive semidefinite operator which multiplies each N-particle vector with N. Its commutation rules with $a^{\#}$ (this notation is used if we do not want to distinguish between a and a^*) is:

$$[\mathbf{N}, a(f)] = -a(f), \quad [\mathbf{N}, a^*(f)] = a^*(f)$$
(2.7)

In terms of an orthonormal basis it looks as $\mathbf{N} = \sum_{i} a^{*}(f_{i})a(f_{i})$.

It is convenient to liberate the formalism from the wave functions by introducing operator-valued distributions $a^{\#}(\vec{x})$:

$$a^{*}(f) = \int a^{*}(\vec{x})f(\vec{x})d^{3}x, \quad a(f) = \int a(\vec{x})\bar{f}(\vec{x})d^{3}x$$

with $[a(\vec{x}), a^{*}(\vec{y})] = \delta(\vec{x} - \vec{y})$ etc. (2.8)

One can then introduce the improper basis (vector-valued distributions) in Fock-space:

$$|\vec{x}_1, ..., \vec{x}_N\rangle = \frac{1}{\sqrt{N!}} a^*(\vec{x}_1) ..., a^*(\vec{x}_N) |0\rangle, \quad |0\rangle := \Omega$$
 (2.9)

$$|\Psi\rangle = \psi_0 |0\rangle + \sum_N \int \psi_N(\vec{x}_1 \vec{x}_N) |\vec{x}_1 \vec{x}_N\rangle d^3 x_1 d^3 x_N$$
(2.10)

The action of the $a^{\#}$ on the basis vectors is (as always) contragredient to that on the wave functions. It is more common to use the former.

Of frequent use (especially in the application of Fock-space in statistical mechanics) is the so called occupation number representation. One chooses an orthonormal set of wave functions f_i , $i = 1, 2, ...\infty$ and defines a basis in Fock-space by:

$$|n_{i_1}, n_{i_2}, \dots, n_{i_r}\rangle = \frac{1}{\sqrt{n_{i_1}!}} \dots \frac{1}{\sqrt{n_{i_r}!}} a^* (f_{i_1})^{n_{i_1}} \dots a^* (f_{i_r}) |0\rangle, \quad \sum_{k=1}^r n_{i_k} = N$$
(2.11)

Often (in particular in Stat. Mech.) one encloses the system in a box and uses the discrete set of plane waves as the orthonormal system for the occupation number representation.

The creation and annihilation operators in x-space are useful for rewriting the Schrödinger theory into the Fock space formalism. One easily verifies the validity of the following formulas:

$$\mathbf{H} = \int \mathbf{H}(x) d^3 x \quad , \quad \mathbf{H}(x) = \mathbf{H}_{\mathbf{0}}(x) \quad + \quad \mathbf{V}(x)$$
$$\mathbf{H}_0(x) = \frac{1}{2m} \vec{\partial} a^*(x) \cdot \vec{\partial} a(x)$$
$$\mathbf{V}(x) = \frac{1}{2} a^*(x) \int d^3 y V(x - y) a^*(y) a(y) a(x)$$
(2.12)

applied to the previously introduced N-particle state $|\Psi\rangle$ give the N-particle Schrödinger-operator:

$$\mathbf{H} |\Psi(t)\rangle = i\frac{\partial}{\partial t} |\Psi(t)\rangle, \quad \Leftrightarrow \\ \left(\sum_{i=1}^{N} \frac{1}{2m} \Delta_i + \sum_{i < j} V(x_i - x_j)\right) \psi_N(\vec{x}_1 \dots \vec{x}_N; t) = i\frac{\partial}{\partial t} \Psi_N(\vec{x}_1 \dots \vec{x}_N; t)$$
(2.13)

The verification only uses the commutation relations of the $a^{\#}(x)$ and the annihilation property of a(x) applied to the vacuum. The various terms in the Nbody Schrödinger operator result from the following commutators which arise in the process of moving **H** through the $N a^{*}(x)'s$ onto the no-particle state:

$$[\mathbf{H}, a^*(\vec{x})] = -\frac{1}{2m} \Delta_x a^*(x), \quad [\mathbf{V}, a^*(\vec{x})a^*(\vec{y})] = V(\vec{x} - \vec{y})a^*(\vec{x})a^*(\vec{y}) \quad (2.14)$$

The hamiltonian **H** in Fock space is used to define time-dependent operators:

$$a(\vec{x},t) = e^{i\mathbf{H}t}a(\vec{x})e^{-i\mathbf{H}t}$$
(2.15)

Only in the case $\mathbf{H} = \mathbf{H}_0$ and for "external interactions"

$$\mathbf{H} = \mathbf{H}_0 + \int V(x)a^*(x)a(x)d^3x \qquad (2.16)$$

is the time dependent a(x,t) linear in a(x).

Before we consider an application some remarks on the mathematical status of the $a^{\#}$'s and related operators are helpful. Since the N-particle states for arbitrary N form a dense set of states, the $a^{\#}$ are densely defined. Using the number operator it is easy to compute:

$$\left\| a(f) \mathbf{N}^{-\frac{1}{2}} \right\|_{H_F^{\perp}} = (f, f), \quad H_F^{\perp} = \text{subspace of } H_F \perp \Omega$$
(2.17)

We remind the reader that the norm of an operator A is related to the vector norm in the Hilbert space:

$$||A|| = \sup_{\psi} \frac{||A\psi||}{||\psi||}$$
(2.18)

The technique for computing such norms is always the same: first one uses the defining formula for a(f) in order to prove the inequality and then one exhibits a particular vector for which the equality sign holds. In our case $\int f(\vec{x})a^*(\vec{x})\Omega$ is such a vector. The norm of the adjoint is the same $||A|| = ||A^*||$.

The relative boundedness with respect to N e.g. $||a^*(f)\psi|| \leq ||\mathbf{N}\psi||^{\frac{1}{2}}$ may be used to show that these unbounded operators are closable and hence admit e.g. a polar decomposition.

Most physicist's calculations do not touch these fine points. They only check equations for densely defined bilinear forms . In case of the above formulas for \mathbf{H} this means that one checks this formula for $\langle \Psi, \mathbf{H}\Phi \rangle$ with the vectors running through the dense set of smooth states of finite particle number. The extension to a relation between densely defined closable or selfadjoint operators is in most physically relevant cases possible and follows a standard scheme (Reed-Simon). In those cases we will be satisfied with the check for matrix elements which is easily done with the commutation relations for $a^{\#}$ and the above "pulling through onto the vacuum" rule. All perturbative calculation in Fock space are done with these rules and this applies also to the derivation of Feynman rules in the real-time operator setting of relativistic QFT¹.

In order to illustrate the application of bosonic (symmetrized) Fock space techniques to coherent states, we first convince ourself that the $a^{\#}$ - formalism for an oscillator is a special case of the present formalism (specialization to one degree of freedom). For a one-dimensional space $\mathcal{H}_1 = \mathbf{C}$, all the tensor product

46

¹An important exception to the applicability of this cavalier attitude of physicists (concerning dense set of vectors in a Hilbert space) is the modular localization theory of section 5, chapter 3 and chapter 6. In that case the relevant dense sets of vectors can be parametrized in terms of real closed subspaces which describe the physical localization and carry the main physical content of the theory in their net properties of real Hilbert spaces.

2.1. THE BOSONIC FOCK SPACE

spaces are also one dimensional and the "single degree of freedom" operator $a^{\#}$ does not require any additional label. An arbitrary vector may be written as:

$$|\Psi\rangle = \psi_0 |0\rangle + \psi_1 |1\rangle + \psi_2 |2\rangle + \dots, \quad |n\rangle = \frac{a^{*n}}{\sqrt{n!}} |0\rangle, \quad a |0\rangle = 0 \quad (2.19)$$

Writing instead of the $a^{\#'}$ s standard dynamical variables of QM p and q (with natural units $\hbar = 1$):

$$a^* = \sqrt{\frac{\omega}{2}}x + i\sqrt{\frac{1}{2\omega}}p \tag{2.20}$$

the $a^{\#}$ - commutation relations go over into the Heisenberg c.r. and the standard oscillator hamiltonian takes the form:

$$H_{osc} = \frac{1}{2}p^2 + \frac{\omega^2}{2}x^2 = \omega(a^*a - \frac{1}{2})$$
(2.21)

The x-space wave functions $\langle x | n \rangle$ of the eigenstates $|n\rangle$ turn out to be the well-known Hermite functions. Coherent states are obtained by asking for the eigenstates of the perturbed hamiltonian:

$$H = H_{osc} + \lambda(a + a^*) = H_{osc} + \lambda\sqrt{2\omega}x \qquad (2.22)$$

Apart from an uninteresting c-number term, the linear perturbation can be obtained by applying a spatial translation by $a = \frac{\lambda}{\sqrt{2\omega}}$ to H_{osc} . In terms of $a^{\#}$ this translation U(a) is:

$$U(\frac{\lambda}{\sqrt{2\omega}}) = e^{-\frac{\lambda}{\omega}(a^* - a)}$$
(2.23)

Different from the previous use of Fock space formalism, the number operator for the oscillator quanta $\mathbf{N} = a^*a$ does not commute with the perturbation. Hence the eigenstates of H do not have a well defined number of such quanta. In order to obtain explicit formulas for $U | n \rangle$ we use the Campbell-Baker-Hausdorff formulas:

$$e^{A}e^{B} = e^{A+B+\frac{1}{2}[A,B]+\dots}$$
(2.24)

where the.... stands for higher commutator terms. This is easily established for matrices and (modulo domain problems) by perturbative arguments in the general case. Due to the vanishing of higher commutators we get:

$$e^{-\frac{\lambda}{\omega}a^*}e^{\frac{\lambda}{\omega}a} = e^{-\frac{\lambda}{\omega}(a^*-a) + \frac{1}{2}\frac{\lambda^2}{\omega^2}}$$
(2.25)

Therefore the ground state of H is an "eigenstate" of the annihilation operator:

$$|\Psi_0\rangle = U |0\rangle = e^{-\frac{1}{2}\frac{\lambda^2}{\omega^2}} e^{-\frac{\lambda}{\omega}a^*} |0\rangle, \quad a |\Psi_0\rangle = -\frac{\lambda}{\omega} |\Psi_0\rangle$$
(2.26)

Here in the first step we used the BCH formula to separate the annihilation part of U to the right (where an $e^{\alpha a}$ factor on $|0\rangle$ becomes the identity). For the eigenvalue equation use the translation property. On the higher eigenstates $U\left|n\right\rangle$, the application of a leads to an additive modification of the eigenvalue relation by $U\left|n-1\right\rangle$. The probability distribution of the oscillator quanta follows the Poisson distribution:

$$\left|\left\langle n \mid \Psi_0 \right\rangle\right|^2 = \frac{e^{-\frac{\lambda^2}{\omega^2}}}{n!} \left(\frac{\lambda}{\omega}\right)^{2n} \tag{2.27}$$

Physically the perturbed oscillator may be thought of as resulting from a constant electric field:

$$H = H_{osc} - eEx \tag{2.28}$$

This field causes the expectation values of the $a^{\#}$'s to be nonvanishing:

$$\left\langle \Psi_n \mid a^{\#} \mid \Psi_n \right\rangle \sim E \tag{2.29}$$

The free time development on the state vectors leads to the classical oscillatory behaviour of expectation values:

$$\langle \Psi_n(t) | x | \Psi_n(t) \rangle \sim E \cos(\omega t - \varphi) \langle \Psi_n(t) | p | \Psi_n(t) \rangle \sim E \sin(\omega t - \varphi)$$
 (2.30)

This oscillatory behaviour would be the result of a sudden switching on of the field:

$$H(t) = \begin{cases} H & for \ t < 0\\ H_{osc} & for \ t \ge 0 \end{cases}$$
(2.31)

by which the coherent states are created. The following classical behaviour of expectation values of functions in $a^{\#}$ is characteristic for coherent states:

$$\left\langle \Psi_{0}\left|f(a^{\#})\right|\Psi_{0}\right\rangle = f\left(\left\langle \Psi_{0}\left|a^{\#}\right|\Psi_{0}\right\rangle\right)$$

$$(2.32)$$

In case of a time dependent source:

$$H(t) = H_{osc} + H_{int}(t), \quad H_{int}(t) = -eE(t)x$$
 (2.33)

we are dealing with time dependent unitary transformations which implement the time dependent canonical transformations:

$$U(t)a^{\#}U^{*}(t) = a^{\#} - \sqrt{2\omega}E$$
(2.34)

which lead from H_{osc} to H(t). In this simple case the U(t) has the same form as in the stationary case except that the constant in front of the $a - a^*$ term in the exponential is now time dependent. It is useful to have a more systematic method which also works for cases for which the U(t) is less simple. Such a method go back to Dirac (and flourished in QFT thanks to Dyson). It treats the time-dependent problems in the "interaction picture" which is between the Heisenberg picture and the Schrödinger picture. All these pictures agree on the level of physical states i.e. in their expectation values, but they differ in how the total time development is distributed between operators and state vectors. Whereas in the Heisenberg - and Schrödinger-picture the full time development

2.1. THE BOSONIC FOCK SPACE

is on the operators respectively on the state vectors, the interaction picture is characterized by the property that the operators only suffer the free time development and the rest (the interaction picture time development V(t)) is dumped on the vectors. According to this definition the interaction operator $H_{int}(t)$ becomes:

$$H_W(t) = e^{iH_0 t} H_{int}(t) e^{-iH_0 t}$$
(2.35)

The time development operator $V(t_2, t_1)$ which propagates the vector state (or wave function) from one time to a later time is:

$$V(t_2, t_1) = T e^{-i \int_{t_1}^{t_2} H_W(t) dt}$$
(2.36)

It is a solution of the Schrödinger equation in the interaction picture:

$$i\frac{d}{dt}V(t,t') = H_W(t)V(t,t')$$
 (2.37)

The time- (or path-) ordering is defined as:

$$TA_{1}(t_{1})A_{2}(t_{2})...A(t_{n}) = A_{i_{1}}(t_{i_{1}})A_{i_{2}}(t_{i_{2}})...A_{i_{n}}(t_{i_{n}})$$

for $t_{i_{1}} \ge t_{i_{2}} \ge ... \ge t_{i_{n}}$ (2.38)

and the above time ordered exponential is defined by the power series with time ordered integrands or as the limit of subsequent products with decreasing length of the time intervalls as:

$$Te^{-i\int_{t_1}^{t_2}H_W(t)dt} =$$

$$\lim_{\Delta t \to 0} Te^{-i\int_{t_{n-1}}^{t_n}H_W(t)dt} \dots Te^{-i\int_{t_2}^{t_3}H_W(t)dt} Te^{-i\int_{t_1}^{t_2}H_W(t)dt}$$
(2.39)

The proof consists in rewriting the Heisenberg time development operator U(t,s). Just as in the case of time independent interactions, this operator can be factorized into $e^{-iH_0(t-s)}$ and a remaining interaction V(t,s):

$$U(t,s) = e^{-iH_0(t-s)}V(t,s) \quad i.e. V(t,s) = e^{iH_0(t-s)}U(t,s)$$
(2.40)

The Schrödinger equation for U(t, s) is then equivalent to the following differential equation for V(t, s):

$$i\frac{d}{dt}V(t,0) = e^{iH_0t} (H - H_0) e^{iHt} = e^{iH_0t} (H_{int}) e^{-iH_0t} e^{iH_0t} e^{-iHt}$$

= $H_W(t)V(t,0)$ (2.41)

The rest of the proof consists in deriving the time-ordered representation from the formal integration of this differential equation. One first converts this into an integral equation (using V(0,0) = 1 as an initial condition):

$$V(t,s) = 1 - i \int_{s}^{t} dt' H_{W}(t') V(t',s)$$
(2.42)

Clearly the perturbative solution is the geometric series:

$$V(t,s) = 1 + (-i)\int_{s}^{t} H_{W}(t')dt' + (-i)^{2}\int_{s}^{t} dt_{2}\int_{s}^{t_{2}} dt_{1}H_{W}(t_{2})H_{W}(t_{2}) + \dots$$
(2.43)

where the n^{th} term is integrated over the simplex $s \leq t_1 \leq t_2 \leq \ldots \leq t_n \leq t$. The use of the (nonlocal, by its very definition) time-ordering prescription allows to convert the integration over a simplex into one over the n-dim. hypercube $s \leq t_i \leq t, i = 1...n$:

$$V(t,s) = 1 + (-i) \int_{s}^{t} H_{W}(t')dt' + \frac{1}{2!} \int_{s}^{t} \int_{s}^{t} TH_{W}(t_{2})H_{W}(t_{1})dt_{1}dt_{2} + \dots + \frac{1}{n!} \int \dots \int \dots + \dots$$
(2.44)

which has the desired exponential time-ordered form. These somewhat formal manipulations may be mathematically justified in two different ways. Either one finds a bound for the n^{th} term, or one shows the equivalence of the time-ordered expression with an exact unitary transformation which, like the one at the beginning of this section is a "dressing transformation" i.e. applied to the free hamiltonian it generates the interaction. Let us briefly explain this for the infinite degrees of freedom analog of the perturbed oscillator: a bosonic field system under the influence of an external source described by the hamiltonian:

$$H(t) = H_0 + a(j_t) + a^*(j_t), \quad H_0 = \int (\frac{\vec{p}^2}{2m} + \mu) a^*(\vec{p}) a(\vec{p}) d^3p, \quad (2.45)$$

where $j_t(\vec{x}) = j(\vec{x}, t)$ and we have added a chemical potential term (in order to avoid infrared divergencies of the p-integrals in subsequent calculations). The dressing transformation is:

$$U(t) = e^{a(g_t) - a^*(g_t)}, \quad g(\vec{x}, t) = \frac{1}{(2\pi)^3} \int \tilde{j}(\vec{p}, t) (\frac{\vec{p}^2}{2m} + \mu)^{-1} e^{i\vec{p}x} d^3p \qquad (2.46)$$

The connection with the time development U(t, s) is evidently (since it dresses the free operator):

$$U(t,s) = U(t)e^{-iH_0(t-s)}U^*(t) \quad or \quad V(t,s) = e^{iH_0(t-s)}U(t)e^{-iH_0(t-s)}U^*(t)$$
(2.47)

The direct calculation of the time-ordered representation for V uses the previously mentioned infinite product representation:

$$V(t,s) = \lim_{\Delta t \to 0} \prod_{ord} e^{-i \int_{t_i}^{t_{i+1}} H_w(t')dt'}$$
(2.48)

where the product is path-ordered (with ascending times going to the left), but the time ordering within each factor is omitted. This formula is similar to the famous Trotter product formula; in integrals over shrinking intervalls the difference between the time ordered and the unordered expression disappears. To this product form we may apply the BCH-formula in order to collect all

50

operators within one unordered exponential. The BCH series ceases after the quadratic term in H_w :

$$V(t,s) = \lim_{\epsilon \to 0} \exp(-i\sum_{j=0}^{n-1} \int_{s+j\epsilon}^{s+(j+1)\epsilon} H_w(t')dt' - \frac{1}{2}\sum_{j\neq k} \int_{s+j\epsilon}^{s+(j+1)\epsilon} dt' \int_{s+k\epsilon}^{s+(k+1)\epsilon} dt'' [H_w(t'), H_w(t'')])$$
(2.49)

Since the commutators are c-numbers, the result is of the form:

$$V(t,s) = \exp(-ia(j_{t,s}) - ia^*(j_{t,s}) - i\frac{\alpha}{2})$$
(2.50)

where α is a numerical phase (resulting from the commutator) and $j_{t,s}(\vec{x})$ is the result of time propagation of the original source function in H_{int} . Hence we obtain agreement between the two methods. Furthermore we learn that the time ordered exponential leads to a phase factor which is not present in the dressing approach. Specializing now to the limit $t \to \infty, s \to -\infty$ (assuming that the interaction only extends over a finite time or that the integrals over time in V converge) we define the S-operator as the full interaction picture transition operator V which relates the free system before and after the interaction:

$$S = \lim_{t,s \to \pm \infty} V(t,s) \tag{2.51}$$

Clearly the application of

$$S = exp(-ia(g) - ia^{*}(g) - i\frac{\alpha}{2}), \ g = lim_{t,s \to \pm \infty}(j_{t,s})$$
(2.52)

onto the vacuum Ω gives a coherent state vector:

$$S(g)\Omega = e^{-i\frac{\alpha}{2}}\Omega(ig), \qquad (2.53)$$

The successive action of sources leads to the composition law:

$$S(f)\Omega(ig) = c(f,g)\Omega(ig+if), \quad |c| = 1$$
(2.54)

Therefore even if we eliminate (by using the projective nature of QT) the phase factor in the definition of $S\Omega$, it will reappear in form of a so called 2-cocycle in the composition law. As in the case of the oscillator, the source generates a coherent distribution of say photons with a Poisson probability distribution. On a coherent state vector $\Omega(ig)$ the action of S(f) will change the mean particle number to:

$$\Delta N = (\Omega(if + ig), \mathbf{N}\Omega(if + ig)) - (\Omega(ig), \mathbf{N}\Omega(ig)) = ||f + g||^2 - ||g||^2 = ||f||^2 + 2Re(f, g)$$
(2.55)

The interference term 2Re(f,g) describes induced absorption or emission depending on the sign. Many important results of laser physics may be developed in this formalism.

2.2 The Fermion Fock Space

The antisymmetric N-particle space was obtained by acting with the antisymmetric projector P_a on the full N-fold tensor product H_N of one-particle spaces:

$$H_N^{(-)} = \pi(P_a)H_N, \qquad P_a = \frac{1}{N!} \sum_{P \in S_N} sign(P)P$$
 (2.56)

Here $\pi(P) \ P \in S_N$ stands for the natural representation of S_N on the full tensor space H_N . The fermionic Fock space is simply the direct sum of all antisymmetrized N-particle spaces augmented by the one dimensional no-particle state.

$$H^{(a)} = H_0 + H_1 + \sum_{N=2}^{\infty} H_N^{(a)}$$
(2.57)

The only difference to the bosonic case (besides the antisymmetry of the wave functions) is the sign appearing in the formula for the creation operator:

$$(a^*(f)\psi)_n(\vec{x}_1,...,\vec{x}_n) = \frac{1}{\sqrt{n}} \sum_i (-1)^{i+1} f(\vec{x}_i)\psi_{n-1}(\vec{x}_1,..,\vec{x}_i,..,\vec{x}_n)$$
(2.58)

Here the roof on the \vec{x}_i indicates omission of this variable. In a completely analogous fashion we obtain the anticommutation relations:

$$\{a(f), a^*(g)\} = (f, g), \quad \{a(f), a(g)\} = 0, \quad \{a^*(f), a^*(g)\} = 0$$
(2.59)

and its pointlike form by removing the wave packets:

$$\{a(\vec{x}), a^*(\vec{y})\} = \delta(\vec{x} - \vec{y}), \quad \{., .\} = 0 \text{ in all other cases}$$
(2.60)

There is no change in the formulas which carry the Schrödinger theory on antisymmetric N-particle wave functions to the Fock space (at least if one writes **H** exactly in the same order in the $a^{\#}$'s). A significant difference to the bosonic case begins to show up, if one realizes that as a consequence of:

$$\{a(f), a(f)\} = 2a(f)^2 = 0 \tag{2.61}$$

and the hermitian adjoint relation, we obtain the Pauli exclusion principle: in an occupation number representation any quantum level can maximally be singly occupied:

$$|n_1, n_2, ...\rangle = \frac{a_1^{*n_1}}{\sqrt{n_1!}} \frac{a_2^{*n_2}}{\sqrt{n_2!}} |0\rangle, \quad n_i = 0, 1$$
 (2.62)

This principle holds only if all quantum numbers of a particle (including spin and possible internal charges) have been taken into account. Closely related is the ability of fermion-systems to form a new reference state by simply *occupying* a given set of levels (orthonormal one-particle vectors $f_i \ i = 1...N$):

$$|\Psi_0\rangle = a_1^*...a_N^* |0\rangle, \quad a_i^* \equiv a^*(f_i)$$
 (2.63)

2.2. THE FERMION FOCK SPACE

This vector is annihilated by the new annihilation operators:

$$b(f) = \begin{cases} a^*(f) & if \ f \in H(f_1, \dots f_N) \\ a(f) & if \ f \in H^{\perp}(f_1, \dots f_N) \end{cases}$$
(2.64)

Here $H(f_1...f_N)$ is the subspace of the one-particle space spanned by the system of vectors f_i . Note that the $b^{\#}s$ obey the same commutation relations as the $a^{\#}s$. The annihilation property $b |\Psi_0\rangle = 0$ is an easy consequence. Note that the hermitian adjoint $b^*(f_i)$ creates holes in $|\Psi_0\rangle$. This ability to design states which are annihilated by transformed Fermion variables $b^{\#}$ through occupying levels, is typical for CAR. On the other hand for coherent states (relevant in e.g.laser physics) and Poisson-distributions one needs Bosons. Mathematically the CAR- structure (canonical anticommutation relations) is well behaved since the $a^{\#}(f)$ are bounded operators:

$$\begin{aligned} (\Phi, \{a(f), a^*(f)\} \Phi) &= (f, f) (\Phi, \Phi) \\ i.e. \quad ||a(f)\Phi||^2 &\le (f, f) (\Phi, \Phi) \end{aligned}$$
(2.65)

By taking $\Phi = a^*(f)\Omega$ we establish saturation (=) and therefore ||a(f)|| = (f, f) for the operator norm. The counterpart of the one-dimensional oscillator is:

$$\sigma_x = a + a^* \quad \sigma_y = i (-a + a^*) \quad \sigma_z = aa^* - a^*a$$
 (2.66)

with σ_i being the Pauli-matrices i.e. the smallest irreducible representation of the Clifford algebra structure defined by $a^{\#}$ is in terms of Pauli matrices. This observation can be generalized:

Theorem 2
$$Alg(a_i^{\#}, i = 1..., N) \equiv Cliff(C^N) = \otimes^N Mat_2(C) = Mat_{2^N}(c)$$

The proof consists in starting with a generating system of matrix units for the N-fold tensor product of $Mat_2(\mathbf{C})$:

$$e_{ij}^{(k)} = \underline{1} \otimes ... \underline{1} \otimes e_{ij} \otimes \underline{1} ... \underline{1}$$

$$(2.67)$$

Clearly the four 2×2 matrix units e_{ij} are linear combinations of the four Paulimatrices and the system commutes for different k. The step towards anticommutation requires the introduction of the famous Jordan-Pauli transformormation. With the help of:

$$\mu_k = \prod_{i=1}^k (e_{11}^{(k)} - e_{22}^{(k)}) = \prod_{i=1}^k (1 - 2a^*a)$$
(2.68)

we define $a_i = \mu_{i-1} e_{12}^{(i)}$ and its hermitean adjoint. Commuting $e_{12}^{(i)}$ factor in a_i through the $(e_{11}^{(i)} - e_{22}^{(i)})$ -factor in $a_j^{\#}$ for j > i leads to the -sign. The relation between the matrix units and the $a^{\#}s$ can be inverted and the generated algebras are identical.

In the "Paulion" formalism, the filling operation is described by the unitary:

$$U = \sigma_1 \otimes \sigma_1 \otimes \dots \otimes \sigma_1, \quad \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$
(2.69)

From this one reads off the filling operator in the $a^{\#}$ representation.

Clearly the filling mechanism is as typical for Fermions as the coherent state properties are for Bosons. Without the former there would be no periodic table (nor we) and without the latter no laser.

The commutation relation of the charge-carrying fields in local quantum physics are inexorably linked with the physical and mathematical properties of the algebras which they generate, a fact which will be presented in detail in the last section on algebraic QFT. There is only one exception: in massive d=1+1 theories the statistics (either in the sense of particles or field commutation relations) loses its intrinsic physical meaning, e.g. a periodic table of elements in a d=1+1 world may also be described in terms of bosons with long range interactions. In that case the inexorable link between the charge properties (superselection sectors) and the particle statistics is lost and particles become statistical "schizons" with well defined (solitonic) charge properties [32] but ill-defined commutation relations (bosons, fermions or "plektons") of their interpolating fields. We will take up the investigation of such 2-dimensional specialities in the next chapter.

2.3 The CCR and CAR Algebras.

Whereas in the case of the Fermion Fock space an abstraction to a C^* -algebra is straightforward (just take the C*-algebra generated by the $a^{\#}s$ subject to the anticommutation relations, uniqueness will be shown later), a construction of a C^* -algebra from unbounded operators generally meets serious obstacles. Following Weyl, one formally converts the $a^{\#}$ into unitary operators:

$$W(f) = e^{i\Phi(f)}, \quad \Phi(f) = \frac{1}{\sqrt{2}}(a(f) + a^*(f))$$
 (2.70)

The application of the BCH-formula leads to:

$$W(f)W(g) = e^{-i\frac{1}{2}\sigma(f,g)}W(f+g) = e^{-i\sigma(f,g)}W(g)W(f)$$
(2.71)

where $\sigma(f,g) = Im(f,g)$ is a non degenerate symplectic form. Formally the unbounded operators $a^{\#}(f)$ may be reobtained by multiplying f with a parameter t and differentiating the modified W with respect to t at t=0. We now take this Weyl relation or rather $Alg(W(f), f \in H)$ as our basic definition of the Boson algebra. This algebra is clearly an infinite degree of freedom generalization of the well-known Heisenberg-Weyl algebra which underlies standard QM:

$$U(\vec{\alpha}) := e^{i\vec{\alpha}\vec{p}}, \quad V(\vec{\beta}) := e^{i\vec{\beta}\vec{q}}$$

$$W(\vec{\gamma}) := e^{i\frac{1}{2}\vec{\alpha}\vec{\beta}}U(\vec{\alpha})V(\vec{\beta}), \quad \vec{\gamma} = \vec{\alpha} + i\vec{\beta}$$
(2.72)

54

One easily checks with BCH, that the W fulfills the above Weyl relation with $f = \gamma \in C^N$ and the symplectic form being the form of standard type known from 2N dimensional phase space of classical mechanics. The following theorem collects the important structural properties of the CAR and CCR(Weyl) algebras.

Theorem 3 The CAR and CCR algebras are simple (no ideals) C^* -algebras generated by the CAR resp. CCR commutation relations.

We only indicate the proof and refer to Bratteli-Robinson Vol 2 for details.

In the CAR case we know from the previous consideration, that for finite degrees of freedom Fermions $a_i^{\#}$ i = 1...N can be replaced by "Paulions". For infinite degrees of freedom we take a basis f_i , $i = 1...\infty$ in the one particle space H. The uniqueness of the limiting algebra follows from the continuity resulting from ||a(f)|| = ||f||. The full algebra is in fact an inductive limit of finite degree of freedom algebras and its separability is inherited from the Mat_N -algebras.

The proof in the CCR case is somewhat different. In this case one reduces the problem to a projective unitary representation of an (infinite) abelian group H (associated with the linear space H) with the multiplier $exp - i\sigma(f, g)$ being a character. In this way the problem is reduced to that of uniqueness of C^* -group algebras. The triviality of the ideal is established by showing that the kernel of every representation is trivial. However separability and the inductive uniform limit property do not hold.

As in the CAR case one may ask about the uniqueness of irreducible representations (up to unitary equivalence). This indeed holds in the important class of "regular" representations i.e. representations π for which the unitaries $\pi(W(tf))$ are strongly continuous in t.

Theorem 4 (Stone-von Neumann uniqueness theorem). Every regular irreducible representation of the Heisenberg-Weyl algebra for a finite number of degrees of freedom is unitarily equivalent to the Schrödinger representation, or in other words: the algebraic structure of standard Q.M. has no nontrivial superselection rules.

The proof uses the infinitesimal generators Φ , which thanks to the regularity property turn out to have a densely defined domain which allows to construct the $a(f)^{\#}$ and the number operator $\mathbf{N} = \sum a_i^* a_i$. The positivity of the latter requires the existence of a "lowest" vector which is the required reference state for the annihilation operators. If on the other hand we are dealing with infinite degrees of freedom (i.e.dimH= ∞), the sum in \mathbf{N} need not to converge. In such representations the number operator does not exist. Examples are easily given.

Bosonic illustration If the shift function c(x) in $a(x) \rightarrow b(x) = a(x) + c(x)$ is not square integrable (physically because of short distance [ultraviolet] or long distance [inf rared] divergencies), then **N** does not exist in the corresponding representation and the formal expression for the unitary implementer U(c) (see (2.77) below) cannot be given a meaning.

Fermionic illustration If the "occupied" Hilbertspace is infinite dimensional, no unitary implementer can exist. The reason is that a vector $\Phi = \prod_i a_i^* \Omega$ is orthogonal on each basis vector of the particle number representation:

$$(\Phi, \Psi(n_1, n_2 \dots n_N)) = 0,$$

$$\Psi(n_1, n_2 \dots n_N) = a^{*n_1} a^{*n_2} \dots a^{*n_N} \Omega, \quad n_i = 0 \text{ or } 1$$
(2.73)

This is because for any arbitrary large N the Φ contains infinitely many creation operators which remain uncompensated. The formal expression for Φ cannot be given a meaning in Fock space.

There is another way of looking at this illustration. The infinite sequence of 0 and 1 in (n_1, n_2, \dots) may be considered as a binary fraction. Whereas the Fockbasis consists of binary fractions with $n_i = 0$ for sufficiently large *i* (which may become arbitrarily large), the binary fraction for the above Φ is the constant sequence (1, 1, 1, ...). This sequence is not in the vacuum class associated to the representative (0, 0, ..0, ...), where "class" here means the set of sequences which deviate from each other only in an arbitrary large but finite number of places. Each class belongs to a basis in a seperate Hilbert space and the different basis elements $\Psi(n_1, n_2, ...)$ are obtainable from one reference element in the class by the application of a finite (but arbitrarily large) number of $a^{\#'s}$. The various irreducible representation spaces obtained from the different classes are orthogonal subspaces of an inseparable unwieldy (and unphysical) Hilbert space generated by all binary fractions (which form a continuous set). The same idea of classes of sequences works for bosons. In that case the n_i run through all natural numbers including zero. One obtains myriads of inequivalent irreducible representations, and this construction is not even exhaustive. Most of them are physically uninteresting, and one needs a physical selection principle. Many of the physically interesting ones are in the subset of "quasifree" states explained in the next section.

We close this section by commenting on automorphisms of the CCR and CAR C^* -algebras which are linear maps of the algebra onto itself which preserving the algebraic structure. In physical terms they are symmetry transformations which preserve the Weyl resp. CAR relation. Of particular interest are the Bogoliubov automorphisms. They are induced by (anti-) linear invertible transformations of the underlying linear wave function space H. In the CCR case they are required to leave the symplectic form σ on H invariant and map the Weyl generators as follows:

$$\sigma(Tf, Tg) = \sigma(f, g), \quad \alpha(W(f)) = W(Tf) \tag{2.74}$$

In the CAR case we must use (anti-)unitary operators in order to preserve the anti-commutator:

$$(Uf, Ug) = \begin{cases} \frac{(f,g)}{(f,g)} & \text{unitary} \\ antiunitary \end{cases}$$
(2.75)

A slightly more general automorphism is obtained by combining these two possibilities:

$$\alpha(a(f)) = a(Uf) + a^*(Vf)$$

$$UU^* + VV^* = 1 = U^*U + V^*V, \quad V^*U + U^*V = 0 = UV^* + VU^*$$
(2.76)

where U is linear and V antilinear. The previous case is obtained by specialization V = 0 resp. alternatively U = 0. Clearly the earlier occupation transformation corresponds to the automorphism $\alpha(a(f)) = a^*(Vf)$. The crucial question is now whether the automorphism is really a bona fide symmetry i.e. implementable by a unitary transformation. Take as an example the shift $a^*(\vec{x}) \to a^*(\vec{x}) + c(\vec{x})$ which is formally implemented by the unitary:

$$Ua^*(\vec{x})U^* = a^*(\vec{x}) + c(\vec{x}), \quad U = e^{a(c) - a^*(c)}$$
(2.77)

But without the condition $\int |c(\vec{x})|^2 d^3x < \infty$ the formal expression for U would not define a bona fide unitary operator in Fock space. Physical intuition would tell us to expect that Bogoliubov tranformations in the one particle space Hneed to be sufficiently close to the identity in order to have an implementation in Fock space. This is indeed the case; the deviation from <u>1</u> should be in the Hilbert-Schmidt class in order not to be thrown out of the representation class (folium). Since Bogoliubov transformations leave the property of "quasifreeness" invariant, the natural place for presenting the relevant implementation formulas is the next section.

2.4 Quasifree States

The most convenient way to obtain representations of the CCR and CAR \mathbb{C}^* algebras is through states ω on these algebras. We have seen that the GNSconstruction gives a canonical association of a representation π_{ω} with a state ω . Since there are too many inequivalent states and associated representations on infinite algebras which nobody has been able to classify, we need some limitation. It turns out that the class of quasifree states and their representations can be classified completely. They are defined by their two-point functions together with a combinatorial formula which expresses their *n*-point functions in terms of the given two-point functions. On the generators $a^{\#}$ we specify the state ω by giving first its two-point functions:

$$\omega(a^{*}(f)a^{*}(g)), \quad \omega(a(f)a^{*}(g)),
 or \quad \omega(a^{*}(x)a^{*}(y)), \quad \omega(a(x)a^{*}(y))$$
(2.78)

The remaining two-point function $\omega(a(g)a(f))$ is (according to the reality properties of states following from their positivity) just the complex conjugate of $\omega(a^*(f)a^*(g))$ and $\omega(a^*(g)a(f))$ may be obtained by (anti-)commutation from $\omega(a(f)a^*(g))$. The higher correlation functions of ω are given in terms of the two-point function by the following combinatorical formula:

$$\omega(a^{\#}(f_1)a^{\#}(f_2)....a^{\#}(f_{2n})) = \sum_{pairings P} signP \prod_{i_k < i_{k+1}} \omega(a^{\#}(f_{i_k})a^{\#}(f_{i_{k+1}}))$$

$$\omega(a^{\#}(f_1)a^{\#}(f_2)....a^{\#}(f_{2n+1})) = 0$$
(2.79)

We have to prove that ω is positive on the polynomial algebra generated by the $a^{\#}$'s:

$$\omega(A^*A) \ge 0 \quad A = polyn(a^{\#}) \tag{2.80}$$

For the CAR-algebra the bound from the anticommutation relations:

$$\omega(a(f)a^*(f)) \le ||f||^2 \tag{2.81}$$

gives immediately:

$$\omega(a(f)a^*(g)) = (f, Tg) \quad 0 \le T \le 1$$
(2.82)

The positivity on monomials $A = a^{\#}(f_1)...a^{\#}(f_n)$ is a result of the basic twopoint positivity:

$$||(a(f) + a^*(g)) \Omega||^2 = (2.83)$$

$$\omega (a^*(f)a(f) + a(g)a^*(g) + a^*(f)a^*(g) + a(g)a(f)) \ge 0$$

(The latter holds as the result of the positivity of T and the Cauchy-Schwarz inequality) and the combinatorial definition of the n-point function.

A particular subclass of quasifree states are the gauge invariant quasifree states. By definition only those correlation functions are nonvanishing which contain the same number of a and a^* . Instead of working with unbounded operators one prefers to define the quasifree gauge invariant states directly in the Weyl algebra:

$$\omega(W(f)) = \exp(-\frac{1}{4} ||f||^2 - \frac{1}{2} \left| \left| T^{\frac{1}{2}} f \right| \right|^2)$$
(2.84)

The standard Fock-representation reemerges as the special case T=0.

Quasifree states are regular (by construction) so that we can return to unbounded $a^{\#}$ in a similar manner as we introduce Lie algebra generators and their enveloping algebras in noncompact group representation theory. The corresponding GNS representation is most conveniently written in terms of an auxiliary "doubled Fock space", for CAR:

$$a_{\omega}(f) = a(\sqrt{1-T}f) \otimes 1 + \gamma \otimes a^*(K\sqrt{T}f), \quad T \le 1$$

$$a^*(f) = a^*(\sqrt{1-T}f) \otimes 1 + \gamma \otimes a(K\sqrt{T}f)$$
(2.85)

For CCR we obtain the analogous formula:

$$a_{\omega}(f) = a(\sqrt{1+T}f) \otimes 1 + 1 \otimes a^*(K\sqrt{T}f)$$

$$a_{\omega}^*(f) = a_{\omega}^*(\sqrt{1+T}f) \otimes 1 + 1 \otimes a(K\sqrt{T}f)$$
(2.86)

58

Here K is the standard conjugation satisfying (Kf,Kg)=(g,f) and γ (only defined in the CAR case) is the unitary operator which implements the \mathbb{Z}_2 gauge transformation (distinguishes even from odd numbers of Fermions) in Fock space. The proof consists in a simple calculation of the two-point function in the vector $\Omega_{double} = \Omega \otimes \Omega$.

The irreducibility condition for gauge invariant quasifree representations is that T is a projector T = P. The equivalence criterion for two gauge invariant quasifree representations is:

Theorem 5 Two irreducible representations given in terms of P and Q are equivalent iff $||P - Q||_{H.S.}^2 < \infty$. Here the H-S norm of K is defined as

$$\|K\|_{HS}^2 \equiv TrK^*K < \infty \tag{2.87}$$

2.5 Temperature States and KMS condition

For a finite quantization box (i.e. a discrete energy spectrum), finite temperature states on the CCR or CAR-algebra are described in terms of the Gibbs formula:

$$\rho = \frac{1}{Z} e^{-\beta H}, \quad Z = Tr e^{-\beta H}, \quad H = H_0(\mu) + H_{int}$$
(2.88)

since $e^{-\beta H}$ is then a trace class operator. Here H_0 includes the chemical potential μ :

$$H_0(\mu) = \int \omega(\vec{p}) a^*(\vec{p}) a(\vec{p}) d^3 p, \quad \omega(\vec{p}) = \frac{\vec{p}^2}{2m} - \mu$$
(2.89)

The box-enclosed version is of course a sum over discrete momenta which results by extending the Laplace operator on smooth functions with support in the volume V in a selfadjoint manner to square integrable functions in V. The various ways of doing this correspond to the various boundary conditions. The physical role of the chemical potential is that the ground state energies for different particle numbers can be adjusted in such a way that the averaged particle n(x) and energy h(x) density:

$$\bar{n} = Tr\rho n(\vec{x}) \quad \bar{\epsilon} = Tr\rho h(\vec{x}) \tag{2.90}$$

remain finite in the thermodynamical limit $V \to \infty$ and hence can be expressed in terms of the two parameters β and μ . For the ideal Fermi or Bose gas ($H_{int} = 0$) the approach of the (quasifree) Gibbs state to the limit KMS state is obvious by explicite calculation:

$$\lim_{V \to \infty} \omega_V(a^*(f)a(g)) = \omega(a^*(f)a(g))$$

$$\omega_V(a^*(f)a(g)) = \frac{1}{Z}Tre^{-\beta \mathbf{H_0}_V}a^*(f)a(g) = (g, T_V f)$$

$$T_V = (\exp -\beta H_{0V})(1 + \exp -\beta H_{0V})^{-1}$$
(2.91)

Here the (non-bold) H_0 are the one-particle operators acting on wave functions whereas \mathbf{H}_0 acts in Fock-space. The ω in the thermodynamic limit is also of the quasifree form with H_0 replacing H_{0V} . The simplest way of proving these relations is to use the KMS property²:

$$\omega_V(\sigma_t(a^*(f))a(g)) = \omega_V(a(g)\sigma_{t+i\beta}(a^*(f)))$$

$$\sigma_t(\cdot) = e^{it\mathbf{H}_0} \cdot e^{-it\mathbf{H}_0}$$
(2.92)
(2.93)

which for t=0 together with the (anti)commutation relation leads to:

$$\omega_V(a^*(f)a(g) \pm a^*(e^{-\beta H_{0V}}f)a(g)) = (g, e^{-\beta H_{0V}}f)$$
(2.94)

We used that the hamiltonian automorphism σ_t is of the Bogoliubov type. We rewrite this equation as

$$\omega_V(a^*((1 \pm e^{-\beta H_{0V}})f)a(g)) = (g, e^{-\beta H_{0V}}f)$$
(2.95)

Clearly this relation is solved by:

$$\sigma_t(a^*(f)) = a^*(e^{-\beta H_{0V}}(1 \pm e^{-\beta H_{0V}})^{-1}f)$$
(2.96)

After Fourier-transformation, $e^{-\beta H_{0V}}$ becomes a multiplication operator $\exp -\beta(\frac{\vec{p}^2}{2m} - \mu)$ on the momentum space wave functions and hence:

$$n(\vec{p}) = \exp{-\beta\omega(\vec{p})(1 \pm \exp{-\beta\omega(\vec{p}))^{-1}}} \\ \omega_V(a^*(\vec{x})a(\vec{y})) = \frac{1}{(2\pi)^3} \int e^{i\vec{p}(\vec{x}-\vec{y})}n(\vec{p}), \quad \omega(\vec{p}) = \frac{\vec{p}^2}{2m} - \mu$$
(2.97)

Since the + case belongs to the Fermions, one obtains for $\beta \to \infty$ the expected occupation for the finite density ground state:

$$\lim_{\beta \to \infty} n(\vec{p}) = \begin{cases} 1, & \text{if } \vec{p}^2 < \mu \\ 0, & \text{if } \vec{p}^2 > \mu \end{cases}$$
(2.98)

The main difference between the finite volume expression and the thermodynamic limit is that in the former case the p-values are discrete and that in the latter case the trace class property of $\exp -\beta \mathbf{H}$ is lost and therefore the numerator and denominator in the Gibbs formula (and hence the Gibbs formula itself) become meaningless. It is not difficult to establish the thermodynamic limit for large classes of \mathbf{H}_{int} .

With the help of the KMS condition one may avoid the finite quantization box and study statistical mechanics directly in the infinite system. It is interesting to note that the KMS condition is equivalent to the stability of the state under appropriately formulated local perturbations and to the second law of thermodynamics (see Haag's book). The GNS construction with a KMS state gives a GNS triple with a reference state Ω which, in addition of being cyclic, also has the separating property i.e. an operator from the algebra A which annihilates Ω must itself be zero. In fact the hamiltonian, or more generally

 $^{^{2}}$ The analytic continuation to imaginary times in the operator expressions can be shown to exist on a dense set of operators in their operator algebra.

the KMS automorphism, is the Tomita automorphism of the associated modular theory and vice versa, the Tomita modular automorphism is characterized by its KMS property. Let us illustrate this for quasilocal states on the CARalgebra. Writing:

$$\langle \Omega | a(f)a^*(g) | \Omega \rangle = (f, Tg), \quad 0 \le T \le 1$$
(2.99)

the separability of Ω is warranted if T has no eigenvalues 0 and 1 and the representation of the CAR algebra is even factorial if the multiplicity of the eigenvalue $\frac{1}{2}$ is finite. The previous considerations suggest that the modular operator is related to S by:

$$T = \frac{\Delta_T}{1 + \Delta_T} \tag{2.100}$$

and the GNS representation may be most naturally be described by "doubling" i.e. in a Fock space $\mathcal{H}_{double} = \mathcal{H}_F \otimes \mathcal{H}_F$ associated with the doubled one particle space $h_{double} = h \oplus h$:

The quasifree states in the doubled description are pure on the tensor product algebra (and its representation is irreducible) since P is a projector. But its restriction S to the first factor (which is the image of the original CAR algebra under the doubling) is impure and reducible. For this reason the doubling is also called "purification" ³. In the application to KMS states of statistical mechanics, the second factor in the doubling is a "shadow world" i.e. a copy of the original one (corresponding to the algebra of the previously discussed right action) which has no spatial localization. Later we will also meet examples of the modular theory for which the commutant algebra has a complementary space-time localization. In those cases the modular theory has a deep relation to TCP symmetry (the particle-antiparticle issue) and the Hawking temperature.

In the previous section we have described the von Neumann algebras associated with quasifree states on CCR or CAR algebras in terms of (2.85) of generators in tensor product form. In this form the commutant and the Tomita involution J may be easily read off. The modular operator Δ^{it} is determined in terms of the KMS property and can be easily written in terms of T. E.g.in the CAR case it is the inversion of (2.100): $\Delta^{it} = T^{it}(1-T)^{-it}$. The information about the type of von Neumann algebra is contained in the structure of T or Δ . Generically, i.e. without further conditions we have the hyperfinite type III_1 (see mathematical appendix). In the CAR case all types of hyperfinite factors (factor condition: dim ker $(T - \frac{1}{2}) = 0, \infty, even$) occur. If the restriction $T_{[0,1]}$ of T to the spectrum [0, 1] is of trace-class (Gibbs like behavior) the factor

³The inverse mechanism, namely the incoherent mixture through "nonobservation" of degrees of freedom, is important in the environmental approach to the measurement problem (section 4, chapter 1)

is I_{∞} , whereas for a Hilbert-Schmidt operator $(T^{\frac{1}{2}} - (\frac{1}{2})^{\frac{1}{2}})$ the factor type is II_1 . The type II_{∞} arises from a combination of the H-S property for $(T^{\frac{1}{2}}_{[0,c]} - (\frac{1}{2})^{\frac{1}{2}})$ and the trace property for $T_{[0,c]}$ with $0 < c < \frac{1}{2}$. The remaining cases are type III. Generically one obtains hyperfinite III_1 ; in order to get also hyperfinite III_0 and $III_{\lambda} T$ must fulfil more subtle spectral conditions. In the CCR case the hyperfinite type III_1 factors are also generic, but regular states on Weyl algebras do not yield type II factors. The local relativistic algebras (see next chapter) belonging to spacetime regions with a nontrivial causal disjoint are always hyperfinite type III_1 .

2.6 The CCR- and CAR-Functors

In section 3 we introduced the CCR and CAR C^* -algebras as maps of Hilbert spaces of functions into C^* -algebras. In particular the Fock-representations of these C^* -algebras define functors from the category of Hilbert spaces into von Neumann algebras.

Let us first look at the CCR-functor. Starting from a Hilbert space (always complex unless stated otherwise) with a scalar product $f, g \to (f, g)$, we first describe the associated bosonic Fock space in the following way. Let e^f be the suggestive notation for the vector in the the Fock space $\mathcal{H}_F^{sym} \equiv e^H$ associated to H with the following n-particle components and inner product:

$$e^{f} = 1 \cdot \Omega + \sum_{n} \frac{1}{\sqrt{n!}} \underbrace{f \otimes \dots \otimes f}_{n}, \quad (e^{f}, e^{g}) = e^{(f,g)}$$
(2.102)

In this notation the vacuum is $\Omega = e^0$. These special vectors are linear independent as well as "total" (i.e. they form a dense set) in e^H . The Weyl operator W(f) is defined on this dense set as:

$$W(f)e^{g} = e^{-\frac{(f,f)}{4}}e^{-(f,g)}e^{f+g}$$
(2.103)

The unitarity of W and hence the extension to the whole space H follows from this formula. The isomorphic map $H \to e^H$ carries subspaces of H into subspaces of e^H as well as the direct sum decompositions into tensor products decompositions. Furthermore linear densely defined maps A between one particle spaces $H \xrightarrow{A} K$ go over into $e^H \xrightarrow{e^A} e^K$ with the computational rules:

$$e^{A}e^{h} = e^{Ah}, \quad (e^{A})^{*} = e^{A^{*}}$$

 $e^{A} = e^{U}e^{|A|}, \quad A = U|A|$ (2.104)

the latter describing the fate of the polar decomposition under the map.

In order to use the Weyl-operators W as a functor from the category of linear spaces to von Neumann algebras, we need to understand a particular family of real subspaces of H. Let M be a set of vectors in H. Define the symplectic complement of M:

$$M' = \{ f \in H \mid \text{Im}(f, g) = 0 \,\,\forall g \in M \}$$
(2.105)

Then M' is a closed real subspace (the use of the symplectic form Im(f,g) requires the restriction to real linear combinations). The following list of properties follows directly from the definition:

$$M \subset N \curvearrowright N' \subset M'$$

$$M \text{ dense in } H \curvearrowright M' = \{0\}$$

$$(M + iM)' = M' \cap iM'$$

$$(2.106)$$

As for von Neumann algebras, the two-fold application of the '-operation i.e. $M \to M''$ gives the (in this case symplectic) completion i.e. the smallest closed real space generated by the set M. The following definition strengthens the analogy with von Neumann algebras.

Definition 1 A real closed subspace M is called "standard" if M + iM is dense and $M \cap iM = \{0\}$. Every standard M defines a "canonical involution" \mathfrak{s} via $\mathfrak{s}(f + ig) = f - ig$ where $f, g \in M$.

In other words, standard M's are +1 eigenspaces of an (unbounded) involution antilinear s. We need its polar decomposition:

$$\mathfrak{s} = j\delta^{\frac{1}{2}}, \quad j^2 = 1, \quad j\delta^{\frac{1}{2}} = \delta^{-\frac{1}{2}}j \qquad (2.107)$$

$$\mathfrak{s}h = h^* \quad \text{on dense set } \mathcal{D}_M = M + iM$$

with * referring to the reality concept defined by M. The important relations:

$$j(M) = M', \quad \delta^{it}M = M \tag{2.108}$$

are rather direct consequences of the definitions.

We now define a von Neumann algebra R(M) associated with the real subspace M:

$$R(M) = \{W(f) \mid f \in M\}'' = alg\{W(f) \mid f \in M\}$$
(2.109)

The double commutant of a *-symmetric family of operators in a Hilbert space is identical to the von Neumann algebra generated by this family (see appendix).

Note that although M is real, the von Neumann Algebras R are always complex. The map:

$$M \to R(M) \tag{2.110}$$

turns out to be an "orthocomplementary functor" from the category of Hilbert spaces H and their standard real subspaces into the $B(H_F)$ operator algebra on Fock space and von Neumann subalgebras in standard position. Orthocomplementary means that the complement M' corresponds to the commutant R(M)'i.e. the validity of the following "duality":

$$R(M') = R(M)'$$
(2.111)

The importance of this functor in QFT in quantum physics results from the fact that the R(M)' describes all observables which are compatible (simultaneously measurable) with an observable from R(M), where in important QFT

cases M describes a space of real (classical) functions localized in some region \mathcal{O} in Minkowski space and $M(\mathcal{O})' = M(\mathcal{O}')$ where \mathcal{O}' denotes the causal disjoint region to \mathcal{O} . So the functor relates classical localization regions with the quantum notion of simultaneous measurability. The process of passing from classical functions with a symplectic structure to operator algebras is often referred to as "quantization". Since this word creates the misleading impression that quantum physics is founded on a parallelism to classical physics and in particular that localization needs a classical function space, we prefer to avoid it alltogether (Bohr's "correspondence principle" is the reverse, namely to recover classical physics in some special limiting situations). In some way algebraic QFT is the investigation of those structures which cannot be obtained by "quantization" methods as Lagrangian canonical- and pathintegral-methods.

The most interesting remaining problem is the connection between the properties of \mathfrak{s}, j and δ and their Fock space counterparts $S = e^{\mathfrak{s}}, J = e^{j}$ and $\Delta^{it} = e^{\delta^{it}}$. As a result of:

$$SW(f)\Omega = W(-f)\Omega = W(f)^*\Omega$$
(2.112)

S is Tomita's (unbounded) involution:

$$SA\Omega = A^*\Omega, \quad A \in R(M)$$
 (2.113)

$$S = J\Delta^{\frac{1}{2}} \tag{2.114}$$

The Tomita-Takesaki theory states (see appendix) that any von Neumann \mathcal{A} algebra acting in a Hilbert space \mathcal{H} with a cyclic ($\mathcal{H} = \overline{\mathcal{A}\Omega}$) and separating vector Ω ($A\Omega = 0 \curvearrowright A = 0$) has a closable operator S defined as above which admits a polar decomposition with $\Delta^{\frac{1}{2}}$ being the radial part and the antiunitary J the angular part. The Δ^{it} implements the modular automorphism:

$$\sigma_t(A) \equiv \Delta^{it} A \Delta^{-it} \in \mathcal{A}$$

$$JAJ \in \mathcal{A}'$$

$$(2.115)$$

with \mathcal{A}' the commutant of \mathcal{A} in \mathcal{H} . In other words a pair (\mathcal{A}, Ω) in "general position" leads to modular objects (Δ, J) with the above properties.

In our case $\mathcal{A} = R(M) \subset B(H)$ and the *M*-dependence of *S* is solely encoded in its dense domain (whereas for \mathfrak{s} the star depends on *M*). It is the simple part of the Tomita-Takesaki theory that *S* and the operators *J* and Δ which result from polar decomposition thereof always exist for general von Neumann algebras \mathcal{R} in standard position i.e. pairs { \mathcal{R}, Ω } with $\mathcal{R} \in B(H)$ and $\Omega \in H$ cyclic and separating. Tomita's deep theorem is the formula (2.115)

It is not difficult to see that in our case $R(M) \cap R(M)' = \mathbb{C}1$ (i.e. R(M) is a factor) iff $M \cap M' = \{0\}$. This suggests the definition:

Definition 2 A real subspace M is called factorial if $M \cap M' = \{0\}$.

The family of standard von Neumann algebras which are in the range of this functor are a subset of all standard von Neumann algebras in $B(H_F)$.

There exists another functor which maps H into H_F^{antis} and the standard real subspaces M of H into von Neumann algebras generated by CAR operators:

$$\{a(g), a^*(f)\} = (g, f)\mathbf{1}$$
 (2.116)

$$\{a^{\#}(g)a^{\#}(f)\} = 0$$

$$CAR(M) = alg \{ A(f) = a^*(f) + a(f) \mid f \in M \}$$
(2.117)

where a(f) is the Fock space annihilation operator: $a(f)\Omega = 0$.

The functorial constructions of the CAR appear somewhat simpler (and more natural) if one follows Araki and interprets the complex Hilbert space H as a "doubled" real Hilbert space. This is achieved by taking two copies H_{\pm} and introducing an antiunitary involution Γ :

$$\Gamma \begin{pmatrix} f_+ \\ f_- \end{pmatrix} = \begin{pmatrix} \bar{f}_- \\ \bar{f}_+ \end{pmatrix}, \quad f = \begin{pmatrix} f_+ \\ f_- \end{pmatrix} \in K = H_+ \oplus H_-, \quad (2.118)$$

$$PK = H_+$$

$$f_{\pm} \rightarrow \bar{f}_{\pm} \text{ conjugation in } H_{\pm}, \quad (f,g) = (f_+,g_+) + (f_-,g_-)$$

The selfconjugate subspace

$$\operatorname{Re} K = \{ f \in K \mid \Gamma f = f \}$$

$$(2.119)$$

inherits on the one hand a real inner product and on the other hand this real subspace is isomorphic with K_+ considered as a real space with the isomorphism being:

$$f \to \sqrt{2Pf}, \quad f \in \operatorname{Re} K$$

 $\operatorname{Re} K$ admits the following complex structure (P as above):

$$if := iPf - i(1-P)f, \quad f \in \operatorname{Re} K \tag{2.120}$$

This description of one particle spaces K is the same for both functors. The only difference is in the interpretation: instead of the symplectic complement M' one uses the "i-symplectic" complement: $\tilde{M}' = iM'$. This could also be called the real orthogonal complement. The relation with the vanishing anticommutator is:

$$\{A(f), A(g)\} = 0, \forall g \in M, \curvearrowright f \in \tilde{M}'$$

$$(2.121)$$

An important distinction between the CCR and the CAR functor shows up if one looks at the Tomita-Takesaki theory. In the CAR case one finds:

$$S = J\Delta^{\frac{1}{2}}, \quad J = Te^j, \Delta = e^{\delta} \tag{2.122}$$

T is the so-called Klein twist, a transformation which is only defined in \mathcal{H}_F^{antis} and not in H:

$$T = \frac{1 + ie^{i\pi\mathbf{N}}}{1 + i}, \quad \mathbf{N} : \text{number op. in } \mathcal{H}_F^{antis}$$
(2.123)

The general setting does not tell which of the two functors one must take in concrete situations. In QFT this additional physical information is supplied by localization properties.

Literature to Chapter 1 and 2:

Rudolf Haag : "Local Quantum Physics", Fields, Particles, Algebras. Springer-Verlag 1992

Ola Bratteli and Derek W.Robinson : "Operator Algebras and Quantum Statistical Mechanics" Vol.1 and 2, Springer-Verlag 1979

J.H.Roberts in "The Algebraic Theory of Superselection Sectors, Introduction and Recent Results" Ed. D.Kastler, World Scientific 1990.

The detailed presentation of the Weyl functor is taken from some unpublished notes of P. Leylands, J.Roberts and D.Testard [4].

Chapter 3

Poincaré Symmetry and Quantum Theory

3.1 Symmetry in General Quantum Theory.

In QM and elementary QFT the generator of a symmetry operation is described by an hermitian operator ("charge", in case of inner symmetry) which commutes with the Hamiltonian.. Usually, in particular for continuous symmetries, this operator has a geometric origin in terms of the quantization of a Noether "current".

Since in relativistic QFT there is a characteristic causality and covariance principle, a more intrinsic approach would suggest to avoid objects which depend on the reference frame as the Hamiltonian H and instead to use concepts which are closer to LQP than those symmetry concepts obtained through that parallelism to classical field theory referred to as "quantization".

Let ψ a vector in a coherent subspace of a Hilbert space of a quantum theory (example: an irreducible representation space of a CCR- or CAR-algebra). The corresponding physical state, in the sense of expectation values as defined previously, corresponds to the unit ray:

$$\underline{\psi} = \left\{ e^{i\alpha}\psi \mid \alpha \in [0, 2\pi], (\psi, \psi) = 1 \right\}$$
(3.1)

The probability for a "source" state ψ containing a "measured" state φ is:

$$w(\underline{\varphi}, \underline{\psi}) = |(\varphi, \psi)|^2 \tag{3.2}$$

and does not depend on the representing vectors. A symmetry transformation \underline{S} is defined to be a transformation of unit rays:

$$\underline{\psi} \longrightarrow \underline{\psi}'$$
 with $w(\underline{\varphi}', \underline{\psi}') = w(\underline{\varphi}, \underline{\psi})$ (3.3)

The physical full significance of such \underline{S} only becomes evident through its action on local observables, an issue which we will take up in a later section.

It is comforting to know, that this projective definition may be reduced to the standard situation of (anti-)unitary operators in Hilbert space:

Theorem 6 (Wigner): Any ray representation \underline{S} may be rewritten in terms of a (anti-)unitary vector representation S:

$$\psi = S\psi' \quad \text{with } (\varphi', \psi') = \begin{cases} (\varphi, \psi) & \text{unitary} \\ (\psi, \varphi) & \text{antiunitary} \end{cases}$$
(3.4)

In the antiunitary case, S may be written in terms of any conjugation K (an antilinear operator which flips the bras and ket of a inner product) as S = UK with U unitary. A proof of this fundamental theorem of quantum theory unfortunately only found its way into few QM textbooks. The reader finds a fairly explicit proof in e.g. in [61] Antiunitary operators appear in quantum theory exclusively in symmetry transformations which contain the operation of time reversal. For physical reasons one does not want a symmetric spectrum, since the energy of systems at zero temperature should be bounded below in order to avoid instabilities due to transitions into arbitrarily negative energy eigenstates (the same reason why Dirac filled the negative energy Dirac sea).

The time reversal T flips the direction of time and therefore:

$$Te^{iHt}\psi = e^{-iHt}T\psi \tag{3.5}$$

Taking T unitary and ψ an energy eigenvector (the use of the spectral representation for H would be more rigorous), one would obtain a symmetric energy spectrum which is in conflict with the existence of a ground state (but not with the structure of finite temperature states). Conversely, if T is antiunitary, then by (3.5) it commutes with H and preserves its spectrum.

If the symmetry S is part of a symmetry group whose group manifold is connected (i.e. every element is continuously deformable into the identity), evidently only unitary representor can occur.

The presence of superselection rules limits the previous consideration to coherent subspaces. In the total space

$$\mathcal{H} = \sum_{\oplus i} \mathcal{H}_i \tag{3.6}$$

the phases between the S's in the subspaces are arbitrary and without physical significance. Symmetries not related continuously with the identity, as the various reflections: P, T, PT, as well as discrete symmetries not related to spacetime e.g. charge conjugation C, can in principle transform one subspace into another. The various possibilities require a careful discussion [61]

If we apply the above consideration to symmetry groups, the two operators $U(g_2)U(g_1)$ and $U(g_2g_1)$ need only to be identical up to a phase factor:

$$U(g_2)U(g_1) = e^{i\varphi(g_2,g_1)}U(g_2g_1)$$
(3.7)

This relation ("ray representation") follows from the fact that $U(g_2g_1)$ and $U(g_2)U(g_1)$ act the same way on observables (i.e. like a classical symmetry) and here we assume that the latter generate an irreducible operator algebra in each \mathcal{H}_i , i.e. one for which Schur's Lemma holds. The associativity of the threefold composition yields a consistency condition for the phase which depends on two group elements. It is called a 2-cocycle condition. It is important to know under what circumstances this phase may be absorbed into a redefinition of the U's, i.e. under what circumstances the "cocycle is a coboundary". In passing from classical to quantum theory, physicists are usually familiar with two "obstructions": phase factors coming from the topology of groups, as e.g. the phase factor -1 in half-integer spin representations (which becomes a projective representation if one considers SO(3) and not SU(2) the represented group), and central extensions of Lie algebras which, after exponentiation, lead to unremovable phase factors in the associated groups. A famous physical illustration of the physical relevance of central extensions is the Galilei group in Schrödinger theory. For semisimple Lie-groups the absence of Lie-algebraic central extensions follows from the absence of a 2-cohomology $H^2(Lie-algebra) = 0$. In this case every ray representation is equivalent to a vector representation of the universal covering group. The only known group in physics which has nontrivial central extensions (and necessarily is not semisimple) is the Galilei group. The physical interpretation is the noncommutativity of the momentum operator \vec{p} with the infinitesimal generator of the velocity (Galilei) transformation $G = \vec{pt} - m\vec{x}$. This group emerges from the Poincaré group by a process called "contraction". More surprisingly, it also appears in a 8-parametric subgroup of the 10-parametric Poincaré group if one splits the generators into longitudinal and transversal as suggested by the concept of modular wedge localization which will be introduced in a later section.

Theorem 7 (Wigner, Bargmann) The projective unitary representations of the Poincare group \mathcal{P} are equivalently described by vector representations of its universal covering $\widetilde{\mathcal{P}}$.

Physicists refer to the representations of the covering group sometimes as multi-valued representations. From a topological point of view the two-fold covering of \mathcal{P} happens already inside the rotation subgroup SO(3) whose covering is SU(2) i.e. the phenomenon of halfinteger spin.

As usual in Lie group theory, one describes representations in terms of infinitesimal generators fulfilling Lie algebra relations. The best known case in physics is the unitary representation theory of the $SU(2) = \widetilde{SO(3)}$. If we characterize the rotation by an angle Θ and axis \overrightarrow{n} , we have:

$$U(\overrightarrow{n},\theta) = e^{i\theta \,\overrightarrow{n} \cdot \overrightarrow{J}} \tag{3.8}$$

where \vec{J} is the quantum mechanical rotation operator with the Lie-algebra:

$$[J_i, J_j] = i\epsilon_{ijk}J_k \tag{3.9}$$

The unitary irreducible representations are all finite dimensional and are explicitly given by the following well-known matrices . For $\overrightarrow{J^2} = s(s+1)$:

$$\langle s, m | J_3 | s, m \rangle = m \langle s, m+1 | J_+ | s, m \rangle = \langle s, m | J_- | s, m+1 \rangle = [(s-m)(s+m+1)]^{\frac{1}{2}}$$
 (3.10)

Here $-s \leq m \leq s, J_{\pm} = J_1 + J_2$, and we have listed only the nonvanishing matrix elements in the 2s + 1 dimensional representation space.

The distinction between a group and its covering does not show up in the Lie algebra, but it can be seen in e.g. irreducible representations by looking at the values of the Casimir (invariant) operators; in the present case the distinction between halfinteger and integer spin in the eigenvalues of \vec{J}^2 . The finite dimensional irreducible representations of the Lorentz group are constructed in complete analogy to the rotation group. The following construction shows that the knowledge of the above formalism suffices. Choosing generators :

$$\Lambda = e^{\frac{1}{2}iM_{\mu\nu}\theta^{\mu\nu}} = e^{i\overrightarrow{M}\overrightarrow{n}\theta + i\overrightarrow{K}\overrightarrow{e}\chi}
\overrightarrow{M} \text{ and } \overrightarrow{K} \text{ related to } M^{\mu\nu} \text{ as } \overrightarrow{B} \text{ and } \overrightarrow{E} \text{ to } F^{\mu\nu}$$
(3.11)

the Lie algebra relations are :

$$\left[M_{i}^{\pm}, M_{j}^{\pm}\right] = i\epsilon_{ijk}M_{k}^{\pm}, \qquad \left[M_{i}^{+}, M_{j}^{-}\right] = 0 \qquad \text{with } M_{j}^{\pm} = \frac{1}{2}(M_{j} \pm iK_{j})$$
(3.12)

Our previous SU(2) representation of the rotation extends to the SL(2, C) (two-valued) representation space of the Lorentz group in terms of (undotted or dotted [3]) two-component spinors is:

$$\alpha(\Lambda) = e^{i\frac{1}{2}\overrightarrow{\sigma} \cdot \overrightarrow{n}\theta + \frac{1}{2}\overrightarrow{\sigma} \cdot \overrightarrow{e}\chi} = \begin{cases} \cos\frac{\theta}{2} + i\sin\frac{\theta}{2} \cdot \overrightarrow{n}\overrightarrow{\sigma} , \chi = 0\\ \cosh\frac{\chi}{2} + sinh\frac{\chi}{2} \cdot \overrightarrow{e}\overrightarrow{\sigma} , \theta = 0 \end{cases} (3.13)$$

$$\beta(\Lambda) \equiv \alpha(\Lambda^{-1})^* = i\sigma_2 \overline{\alpha(\Lambda)} (i\sigma_2)^{-1}$$
(3.14)

First we convince ourselves that SL(2, C) matrices (i.e. complex matrices with determinant =1) allow an exponential parametrization in terms of a complex linear combination of the three traceless Pauli-matrices i.e. $\alpha = \exp i(\vec{c_1} - i\vec{c_2})\frac{\vec{\sigma}}{2}$. The interpretation of the $\vec{c_i}s$ in terms of the Lorentz group parameters $(\vec{n}, \theta; \vec{e}, \chi)$ follows from the relation of the L_+^{\uparrow} and SL(2, C) groups via the conversion formula:

$$\begin{array}{rcl}
x^{\mu} & \rightarrow & \underset{\sim}{x} = x^{\mu} \sigma_{\mu} \\
x & \rightarrow & \alpha(\Lambda) & \underset{\sim}{x} & \alpha^{*}(\Lambda) & \frown & x^{\mu} \rightarrow \Lambda^{\mu}_{\nu} x^{\nu}
\end{array}$$
(3.15)

The matrix x transforms therefore like a mixed spinor with two spinor indices, one transforming with α and the other one with $\bar{\alpha}$.

3.2. ONE PARTICLE REPRESENTATIONS

Using the notation \overrightarrow{J} and \overrightarrow{K} for the representors of \overrightarrow{M} and \overrightarrow{K} in the respective representations we find for this spinor representation :

$$\overrightarrow{J} = \frac{\overrightarrow{\sigma}}{2} , \ \overrightarrow{K} = -i\frac{\overrightarrow{\sigma}}{2} \quad \text{or} \quad \overrightarrow{J^+} = \frac{\overrightarrow{\sigma}}{2} , \ \overrightarrow{J^-} = 0 \quad \text{with} \ \overrightarrow{J^\pm} = \frac{1}{2} \left(\overrightarrow{J} \pm i\overrightarrow{K} \right)$$
(3.16)

The standard notation for this fundamental (undotted) $SL(2,{\mathbb C})$ spinor representation is :

$$D^{\left[\frac{1}{2},0\right]}(\Lambda) = e^{i(\theta \overrightarrow{n} - i\chi \overrightarrow{e})\overrightarrow{J^{+}}} = \alpha(\Lambda) \qquad J^{+} = \frac{\overrightarrow{\sigma}}{2}$$
(3.17)

Similarly for the "dotted" spinors :

$$D^{\left[0,\frac{1}{2}\right]}(\Lambda) = e^{i(\theta \overrightarrow{n} + i\chi \overrightarrow{e})\overrightarrow{J^{-}}} = i\sigma_2\overline{\alpha}(\Lambda)(-i\sigma_2) = \beta(\Lambda) \qquad J^{-} = \frac{\overrightarrow{\sigma}}{2} \qquad (3.18)$$

The equivalence transformation with the Pauli-matrix $\varepsilon \equiv i\sigma_2$ assures that both representations are identical if restricted to SU(2); this ε plays a similar role in the spinor calculus as the metric tensor $g_{\mu\nu}$ for the tensor calculus. It allows to pass from lower (un)dotted indices to upper. $D^{[0,\frac{1}{2}]}(\Lambda)$ refers to upper dotted indices and the x matrix is a mixed spinor of the type x $\sim_{\alpha,\dot{\beta}}$.

Note that in these representations (as well as in all finite dimensional representations of the Lorentz group) the generators K are not hermitean i.e. the associated D's are not unitary. The general irreducible finite dimensional representation are characterized in terms of two (half)integers $\frac{n_{\pm}}{2}$ which denote the formal J_{\pm} angular momenta.

$$D^{\left[\frac{n_{+}}{2},\frac{n_{-}}{2}\right]}(\Lambda) = e^{i(\theta\overrightarrow{n} - i\chi\overrightarrow{e})\overrightarrow{J^{+}}} \otimes e^{i(\theta\overrightarrow{n} + i\chi\overrightarrow{e})\overrightarrow{J^{-}}}$$
(3.19)

Here $\overrightarrow{J^{\pm}}$ are the previously defined matrices of size $(2\frac{n_{\pm}}{2} + 1) \times (2\frac{n_{\pm}}{2} + 1)$. In the spirit of the spinor calculus, one should envisage these operators to act on tensor products of (un-)dotted spinors. The J_{\pm} act as Pauli matrices on $\otimes_{n_{\pm}} \operatorname{Mat}_{2}(\mathbf{C})$:

$$\overrightarrow{J}_{i} = \sum_{i=1}^{n} 1 \otimes \dots \otimes \left. \frac{\overrightarrow{\sigma}_{i}}{2} \otimes \dots \otimes 1 \right|_{symm.}$$
(3.20)

The symmetrization in the n_{\pm} spinorial indices assures the irreducibility. The difference in undotted and dotted spinors is in the action of L-boosts, whereas the rotations act the same way. The entries of the matrices (3.19) can be expressed in terms of Jacobi functions.

3.2 One Particle Representations

We are now ready to study unitary representations of the covering of the Poincaré $\widetilde{\mathcal{P}}$ and its subgroup $\widetilde{SO(3, 1)}$. The infinitesimal generators for non-compact groups are necessarily unbounded operators. The domain problems

for unbounded Lie-generators (common dense domain etc.) have been studied, and we will ignore them unless they are of direct physical significance (as e.g. in the relation between the Tomita-Takesaki modular theory and symmetries described in a later section).

The commutation relations of the Poincaré generators follow from the composition property for the two-fold covering $\widetilde{\mathcal{P}}$

$$(a_2, \alpha_2) \cdot (a_1, \alpha_1) = \left(a_2 + \alpha_2 a_1 \alpha_2^{-1}, \alpha_2 \alpha_2\right)$$
(3.21)

The second translational term is simply the Lorentz-transformed vector $\Lambda_2 a_1$. From the special case

$$(0, \alpha^{-1})(a, 0)(0, \alpha) = (\Lambda^{-1}a, 1)$$
(3.22)

one abstracts for infinitesimal translations:

$$U^{-1}(\alpha)P^{\mu}U(\alpha) = \Lambda^{\mu}_{\nu}P^{\nu} \quad \text{with} \quad U(a) = e^{iaP}$$
(3.23)

Analogously the transformation of the operator $U(\alpha)^1 = e^{iM^{\mu\nu}\theta_{\mu\nu}}$ by another Lorentz transformation yields the tensor transformation property of $M^{\mu\nu}$

$$U^{-1}(\alpha)M^{\mu\nu}U(\alpha) = \Lambda^{\mu}_{\kappa}\Lambda^{\nu}_{\lambda}M^{\kappa\lambda}$$
(3.24)

In order to avoid clumsy notation, it is convenient to suppress the unimodulars α inside unitaries and write simply $U(\Lambda)$ with the understanding that Λ denotes an element of $\tilde{\mathcal{P}}$. Only for matrices (i.e. finite dimensional representations) the notational distinction matters. The Lie-algebra relations are obtained from the above transformation laws by expanding $U(\alpha) \equiv U(\Lambda)$ retaining only linear terms in $\theta_{\mu\nu}$:

$$\begin{bmatrix} M^{\alpha\beta}, P^{\mu} \end{bmatrix} = i(g^{\alpha\mu}P^{\beta} - (\alpha \longleftrightarrow \beta)) [M^{\mu\nu}, M^{\rho\sigma}] = i\{g^{\nu\sigma}M^{\mu\rho} - g^{\mu\rho}M^{\nu\sigma} - (\mu \longleftrightarrow \nu)\}$$
(3.25)

The last relation is the tensor form of the previous J^{\pm} commutation relations.

Approaching the Wigner theory via the infinitesimal generators P^{μ} , $M^{\mu\nu}$, one first looks for the Casimir (invariant) operators which take on characteristic values in irreducible representations :

$$P^{\mu}P_{\mu}, \quad W^{\mu}W_{\mu} \tag{3.26}$$

The first invariant is the mass operator and the second one is usually referred to as the Pauli-Lubanski invariant. It is formed with the Pauli-Lubanski vector:

$$W_{\mu} = \frac{1}{2} \varepsilon_{\mu\alpha\beta\gamma} M^{\alpha\beta} P^{\gamma}$$

Its commutation properties follow from those of the Poincare generators $M^{\mu\nu}$ and P^{μ} :

$$[W_{\mu}, P_{\nu}] = 0 \qquad [W_{\mu}, M_{\nu\kappa}] = i(g_{\nu\mu}W_{\kappa} - g_{\kappa\mu}W_{\nu}) \qquad [W_{\mu}, W_{\nu}] = i\epsilon_{\mu\nu\kappa\lambda}W^{\kappa}P^{\lambda}$$
(3.27)

¹Following habits of physicist, we will often write $U(\Lambda)$ instead of $U(\alpha(\Lambda))$.
Since $P_{\mu}W^{\mu} = 0$, there is no nontrivial third invariant. The interpretation of W_{μ} and W^2 in terms of intrinsic angular momentum becomes manifest, if we specialize to so called positive energy representations.

Wigner classified the irreducible representations according to their transitive p-space orbits (submanifolds of momentum space traced out by the action of \mathcal{L} to a given vector):

- (i) $p^{\mu}p_{\mu} = m^2 > 0$ $p_0 > 0$
- (ii) $p^{\mu}p_{\mu} = 0$ $p_0 > 0$
- (iii) $p_{\mu} = 0$

and the corresponding orbits with negative energies $p_0 < 0$, as well as the spacelike orbits $p_{\mu}p^{\mu} < 0$. The first two exhaust the positive energy representations. In order to construct them explicitly, we look at the stability group ("little group") of a point on the orbit. Without loss of generality we may specialize to the stability group of a selected reference momentum, since the stability group for other momenta are equivalent (by Lorentz-boosts). In the case (i) we choose $p_R = (m, 0)$ which yields the SO(3) respectively. its covering SU(2) as the quantum theoretically relevant little group. On the other hand the little group of the lightlike reference vector which is chosen to be $p_R = (1, 0, 0, 1)$ turns out to be the euclidean group E(2) in two dimensions. Only the rotation around the 3-axis is geometrically obvious, the interpretation of the two euclidean "translations" is somewhat hidden and will be presented later. Let us now look in detail at the massive case (i). We start with a 2s+1 dimensional representation of the little group. This irreducible representation induces a unitary irreducible positive energy representation of the Poincaré group \mathcal{P} as follows. We first chose the momentum in rest $p_R = (m, 0)$ as the reference vector on the orbit $p^2 > 0, p_0 > 0$. The action on (improper, like plane waves) reference basis vectors is:

$$P^{\mu} | p_{R}, s_{3}; \gamma \rangle = p_{R}^{\mu} | p_{R}, s_{3}; \gamma \rangle ,$$

$$W_{0} | p_{R}, s_{3}; \gamma \rangle = 0 ,$$

$$W_{k} | p_{R}, s_{3}; \gamma \rangle = \frac{m}{2} \epsilon_{k\mu\nu0} M^{\mu\nu} | p_{R}, s_{3}; \gamma \rangle$$
(3.28)

The last relation connects the spatial components of W with the Wigner spin i.e. with the angular momentum in the rest frame:

$$W_k |p_R, s_3; \gamma\rangle = \frac{m}{2} \epsilon_{kij} M^{ij} |p_R, s_3; \gamma\rangle = m J_k |p_R, s_3; \gamma\rangle$$
(3.29)

Since an invariant operator can be evaluated on any state vector, we have $W^2 = -m^2 \vec{J}^2$ and therefore in an irreducible representation: $W^2 = -m^2 s(s+1)$. In this approach irreducibility just means the absence of an additional degeneracy label, say γ (such labels, which go beyond spacetime characteristics as momentum and spin, are related to internal symmetries and called charges.). One now uses a distinguished family of Lorentz-transformations which link p_R

with a general point p on the $p^2 = m^2$ orbit. One chooses the family of rotational free Lorentz-transformations ("boosts") to relate the p-eigenstates:

$$|p, s_3\rangle \equiv U(L(p)) |p_R, s_3\rangle, \quad \langle p', s_3' | p, s_3\rangle = 2p^0 \delta(\vec{p}' - \vec{p})$$
(3.30)

$$L(p) = \frac{1}{m} \begin{pmatrix} p, & p \\ \vec{p}, m\delta_{ik} + \frac{p^i p^k}{p^0 + m} \end{pmatrix} = \Lambda(\vec{e}, \chi), \ \alpha(L(p)) = \sqrt{\frac{p^\mu \sigma_\mu}{m}} (3.31)$$

with:
$$\vec{e} = \frac{\vec{p}}{|\vec{p}|}$$
 $ch\chi = \frac{p^0}{m}$ (3.32)

The invariant δ -function which appears on the right hand side of the inner product of relativistic (improper) momentum eigenstates corresponds to the relativistic measure $d^3p/2\omega$ with $\omega = p^0 = \sqrt{\vec{p}^2 + m^2}$. For later use, we also wrote the positive unimodular matrix $\alpha(L(p))$ which the spinor calculus affiliates with the boost.

We now are able to describe the $[m_+, s]$ Wigner representation in global terms as follows. The one- particle Hilbert space is:

$$H_{[m,s]}^{(1)} = \left\{ \int \sum_{s_3} \tilde{\psi}(\vec{p}, s_3) \left| p, s_3 \right\rangle \frac{d^3 p}{2\omega} \left| \int \sum \left| \tilde{\psi} \right|^2 \frac{d^3 p}{2\omega} < \infty \right\}$$
(3.33)

and the transformation properties on the basis (3.30) are:

$$U(\Lambda) |p, s_3\rangle = U(L(\Lambda p))U(R(\Lambda, p)) |p_R, s_3\rangle \quad \text{with } R(\Lambda, p) = L^{-1}(\Lambda p)\Lambda L(p)$$
(3.34)

and
$$U(R(\Lambda, p)) |p_R, s_3\rangle = \sum_{s'_3} \left| p_R, s'_3 \right\rangle D_{s'_3 s_3}(R(\Lambda, p))$$
 we obtain: (3.35)

$$U(\Lambda) |p, s_3\rangle = \sum_{s'_3} \left| \Lambda p, s'_3 \right\rangle D_{s'_3 s_3}(R(\Lambda, p)) \text{ and for translations :}$$
$$U(a) |p, s_3\rangle = e^{ipa} |p, s_3\rangle$$
(3.36)

The successive transformations by a boost, followed by a Lorentz transformation Λ and the inverse boost yields a transformation $R(\Lambda, p)$ which leaves p_R invariant and therefore is called the Wigner rotation. The appearance of pdependent unitary matrices is typical for relativistic quantum theory. It prevents a simple minded transition to x-space covariant localizable functions via Fourier transformation. As well known, one can rewrite the transformations from the basis vectors in $H_{[m,s]}^{(1)}$ to the wave functions on which one finds the contragredient action:

$$\left(U(\Lambda)\tilde{\psi}\right)(\vec{p},s_3) = \sum_{s'_3} D_{s_3s'_3}(R(\Lambda,\Lambda^{-1}p)\tilde{\psi}(\Lambda^{-1}\vec{p},s'_3)$$
(3.37)

Besides this Wigner "canonical" representation, there exists the closely linked "helicity" representation for which the spin quantization axis is identified with

the direction of the spatial momentum of the particle. Calling the magnetic quantum number with respect to this direction λ we define:

$$|p,\lambda\rangle \equiv \sum_{s'_{3}} \left| p, s'_{3} \right\rangle D_{s'_{3}\lambda}(R_{p,\tilde{p}}) \text{ with } R_{p\tilde{p}} = Rot(\varphi,\theta):$$
(3.38)

being the "minimal" rotation which changes the z-direction into $\vec{n} = \frac{\vec{p}}{|\vec{p}|}$ i.e. a rotation around the y-axis with latitude θ followed by a φ -rotation around the z-axis. In the helicity basis the Wigner rotation is modified:

$$\check{R}(\Lambda, p) = R_{\Lambda p, \Lambda \vec{p}}^{-1} R(\Lambda, p) R_{p, \vec{p}} \text{ leaves } \check{p} = (\sqrt{p^2 + m^2}, 0, 0, |\vec{p}|) \text{ invariant.}$$
(3.39)

The evaluation of W_0 on the helicity reference state gives:

$$W_{0} |\check{p}, \lambda\rangle = \vec{J} \cdot \vec{p} |\check{p}, \lambda\rangle = |\vec{p}| \lambda |\check{p}, \lambda\rangle \quad \text{or with} \quad h \equiv \frac{\vec{J} \cdot \vec{p}}{|\vec{p}|} , \quad h |\check{p}, \lambda\rangle = \lambda |\check{p}, \lambda\rangle$$
(3.40)

The column vectors of $D^{(s)}(R_{p,\tilde{p}})$ furnish a complete set of eigenstates of the helicity operator h, e.g. for $s = \frac{1}{2}$ we have:

$$w^{\pm} = D^{\left(\frac{1}{2}\right)}(R_{p,\vec{p}})\chi^{\pm} , \ \chi^{\pm} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} i.e.$$
$$D^{\left(\frac{1}{2}\right)}(R_{p,\vec{p}}) = \begin{pmatrix} w_{1}^{+} & w_{2} \\ w_{2}^{+} & w_{2}^{-} \end{pmatrix}, \ \frac{1}{2}\vec{\sigma}\cdot\vec{p}\,w^{\pm} = \pm\frac{1}{2}w^{\pm}$$
(3.41)

The advantage of the helicity basis is that one may take the limit $m \to 0$. As expected, the helicity rotation matrix $D(\check{R})$ approaches a diagonal limit in terms of a "Wigner phase" $\varphi(\Lambda, p)$ e.g. for $s = \frac{1}{2}$:

$$\lim_{m \to 0} D^{\left(\frac{1}{2}\right)}(\check{R}) = \begin{pmatrix} e^{i\frac{1}{2}\varphi(\Lambda,p)} & 0\\ 0 & e^{-i\frac{1}{2}\varphi(\Lambda,p)} \end{pmatrix},$$

$$e^{i\frac{1}{2}\varphi(\Lambda,p)} = \sqrt{\frac{p^0}{(\Lambda p)^0}} \langle w^+(\Lambda p) | \alpha(\Lambda) | w^+(p) \rangle$$
(3.42)

In the massless limit, the (2s + 1)- component representation decomposes into 2s+1 one-component representations.

A direct approach a la Wigner to the m = 0 case would start with the representation theory of the stability group of a light-like vector. In this situation there is no such natural choice as before. Choosing a light-like vector in the z-direction $p_R = (1, 0, 0, 1)$, one obtains the following matrix realization of the 3-parametric euclidean group E(2) in 2 dimensions:

$$G(\alpha,\beta) = \begin{pmatrix} 1+\frac{1}{2}|\rho|^2 & \alpha & \beta & -\frac{1}{2}|\rho|^2 \\ \alpha & 1 & 0 & -\alpha \\ \beta & 0 & 1 & -\beta \\ \frac{1}{2}|\rho|^2 & \alpha & \beta & 1-\frac{1}{2}|\rho|^2 \end{pmatrix}, \quad R(\theta) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\theta & \sin\theta & 0 \\ 0 & -\sin\theta & \cos\theta & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$
(3.43)

The first matrix is a Lorentz-transformation which leaves p_R invariant and transforms the time axis into $Gp = (1 + \frac{1}{2} |\rho|^2, \alpha, \beta, \frac{1}{2} |\rho|^2)$, $\rho = \alpha + i\beta$, where the transformed vector, whose time component increases, has been conveniently parametrized. Any other transformation having this property can only deviate from $G(\rho)$ by a transformation which leaves the two vectors p_R and the time axis invariant i.e. a x-y rotation $R(\theta)$. Obviously they generate a stability group which is easily checked to be isomorphic to E(2), the euclidean translations corresponding to $G(\rho)$. To be more precise, since the euclidean group has to be considered as a subgroup of the covering of the Poincaré group, only the two fold covering $\tilde{E}(2)$ is relevant.

The description in terms of the corresponding subgroup of SL(2,C) and the reference vector is somewhat simpler:

$$\alpha(\rho,\theta) = \begin{pmatrix} e^{i\frac{1}{2}\theta} & \rho\\ 0 & e^{-i\frac{1}{2}\theta} \end{pmatrix}, \ p_R \sim \begin{pmatrix} 2 & 0\\ 0 & 0 \end{pmatrix}$$
(3.44)

The unitary representation theory of this noncompact group is somewhat more complicated than that of SU(2). But it is obvious that the representations fall into two classes; the "neutrino- photon" class with $U(G(\rho)) = 1$ i.e. trivial representation of the euclidean translations, and the remaining "continuous spin" representation with $U(G(\rho)) \neq 0$. The difference also shows up in the spectrum of the operator W^2 . Whereas in the first case $W^2 = 0$ (in fact $W^{\mu} = hP^{\mu}$), the value of W^2 in the second case can be any negative number which is responsible for the name. These representations of $\tilde{E}(2)$ are infinite dimensional. They are usually discarded as a result of the apparent absence of such particles in nature. We will later return to this representations, since a theoretician should use theoretical arguments i.e. point to a property which makes this positive energy representation appear less physical than the others.

As a curious side remark we mention that the two "translations" inside the homogenous Lorentz group behave as transversal Galilei transformations if we split Minkowski space into two longitudinal direction $e_{\pm} = (1, 0, 0, \pm 1)$ and the remaining transversal spacelike unit vectors $e_{x,y}$. This also simplifies the calculation of the Wigner phases $\theta(\Lambda, k)$.

It is comforting to know that the [m, s] representations admit an extension of the Poincaré group which includes the reflections, without enlarging the representation space. One obtains the well-known formulas for the parity \mathcal{P} and the time reversal T:

$$\mathcal{P}|p,s_{3}\rangle = \xi_{P}|p_{0},-\vec{p},s_{3}\rangle \qquad T|p,s_{3}\rangle = \xi_{T}\sum_{s_{3}'} D_{s_{3},s_{3}'}^{(s)}(i\sigma_{2})\left|p_{0},-\vec{p},s_{3}'\right\rangle \quad (3.45)$$

Here the $\xi's$ are undetermined phase factors. This result follows by first writing down the action of \mathcal{P} and T on the reference vectors $|p_R, s_3\rangle$ (the antiunitarity of T brings in the "spin-flip" matrix $D(i\sigma_2)$ which represents the later appearing charge conjugation). The rest follows from the commutation relation of the reflections with the boost:

$$\mathcal{R}_{\lambda}L(p)\mathcal{R}_{\lambda}^{-1} = L(p_0, -\vec{p}), \quad R_{\lambda} = \mathcal{P} , \ T \ or \ \mathcal{P}T$$
(3.46)

3.2. ONE PARTICLE REPRESENTATIONS

The corresponding operator relation may contain phase factors D_{λ} i.e.

$$\mathcal{R}_{\lambda}U(\Lambda)\mathcal{R}_{\lambda}^{-1} = D_{\lambda}(\Lambda)U(R_{\lambda}\Lambda R_{\lambda}^{-1})$$
(3.47)

These phase factors must form a representation of the Lorentz group. But since there are no 1-dimensional representations, we have $D(\Lambda) = 1$. The above phases can be fixed. For unitary reflections we can achieve $\mathcal{R}^2_{\lambda} = 1$, whereas for antiunitaries $\mathcal{R}^2_{\lambda} = \pm 1$. In the above special case we find:

$$\mathcal{T}^2 = (-1)^{2s} \tag{3.48}$$

The formulae for the [0, s] representations are different as a result of the different p_R and its stability group which contain the arbitrary z-direction :

$$\mathcal{P} |p,s\rangle = \xi_P e^{\pm i\pi s} |p,-\vec{p},-s\rangle \qquad T |p,s\rangle = \xi_T e^{\pm i\pi s} |p,-\vec{p},s\rangle \tag{3.49}$$

The \pm sign depends on the sign of p_y (see Weinberg [61]), and this phase factor is only relevant if the states of opposite helicity are not separated by a superselection rule.

The original motivation of Wigner was to classify relativistic wave equations up to physical equivalence. Disregarding the continous spin class, the classification of wave equations associated with finite positive energy representations is as follows. We first present the three special cases $s=0,\frac{1}{2}, 1.m>0$

• s=0

The Fourier transformation leads to covariant x-space wave function :

$$\psi(x) = \int e^{-ipx} \tilde{\psi}(p) \frac{d^3 p}{2\omega} \quad \text{with } (U(\Lambda)\psi)(x) = \psi(\Lambda^{-1}x)$$
(3.50)

The x-space function is a positive frequency solution of the Klein-Gordon equation:

$$\left(\partial^{\mu}\partial_{\mu} + m^2\right)\psi\left(x\right) = 0 \tag{3.51}$$

• $\mathbf{s} = \frac{1}{2}$

Here one has to convert the Wigner representation into a covariant one involving (un)dotted spinors which transform covariantly even under boost transformations. This is achieve by:

$$\tilde{\Phi}_{a}(p) := \sum_{s_{3}} \alpha_{a,s_{3}}(L(p))\tilde{\psi}(p,s_{3}) \quad \text{with} \quad \alpha(L(p)) = \sqrt{p^{\mu}\sigma_{\mu}} / \sqrt{m} \\
\text{a positive matrix, in short:} \quad \tilde{\Phi} = \alpha(L(p))\tilde{\psi}$$
(3.52)

$$\sigma_{\mu} \equiv (\underline{1}, \overrightarrow{\sigma}) \tag{3.53}$$

As the notation suggests, $\tilde{\Phi}$ transforms like a (undotted) spinor, a fact which follows by transforming the $\tilde{\psi}$ with the Wigner rotation and then using its representation in terms of boosts :

$$\alpha(L(p))\alpha\left(R(\Lambda,\Lambda^{-1}p)\right) = \alpha(L(p))\alpha\left(L^{-1}(p)\alpha(\Lambda)\alpha\left(L(\Lambda^{-1}p)\right)\right)$$

= $\alpha(\Lambda)\alpha(L(\Lambda^{-1}p))$
i.e. $\left(U(\Lambda)\tilde{\Phi}\right)(p) = \alpha(\Lambda)\tilde{\Phi}(\Lambda^{-1}p)$ (3.54)

For later purpose it is helpful to rewrite the action of $\alpha(L(p))$ on $\tilde{\psi}$ in terms of the column vectors of the boost matrix:

$$\tilde{\Phi}(p) = \sum_{s_3} u(p, s_3) \tilde{\psi}(p, s_3) , \quad u_a(p, s_3) := \alpha_{a, s_3}(L(p))$$
(3.55)

 $\alpha(L(p))$ and hence its column vectors u has the intertwining property between the Wigner and the covariant representation :

$$\alpha(L(p))D^{\left(\frac{1}{2}\right)}(R(\Lambda,\Lambda^{-1}p)) = D^{\left[\frac{1}{2},0\right]}(\Lambda)\alpha(L(\Lambda^{-1}p))$$
(3.56)

A similar intertwining relation is valid between the complex conjugate of the Wigner representation D^* and the same covariant $D^{\left[\frac{1}{2},0\right]}$. In this case the intertwining matrix is $\alpha(L(p))i\sigma_2$ and its columns are called *v*-spinors.

Fourier transformation gives the x-space wave function:

$$\Phi(x) := \int \tilde{\Phi}(p) e^{-ipx} \frac{d^3 p}{2\omega} , \quad (U(\Lambda)\Phi)(x) = \alpha \left(\Lambda\right) \Phi(\Lambda^{-1}x) , \qquad (3.57)$$
$$\left(\partial^{\mu} \partial_{\mu} + m^2\right) \Phi(x) = 0$$

In order to make contact with the Dirac theory, one defines a dotted spinor

$$\tilde{\chi}^{\cdot}(p) := \frac{1}{m} p^{\mu} \tilde{\sigma}_{\mu} \tilde{\Phi}(p) = \alpha (L^{-1}(p)) \tilde{\psi}(p)$$

$$\tilde{\sigma}_{\mu} \equiv (\underline{1}, -\overrightarrow{\sigma})$$
(3.58)

As indicated in the notation, $\tilde{\chi}$ transforms as an upper dotted spinor i.e. with a matrix $\beta(\Lambda) \equiv \alpha(\Lambda^{-1})^{\dagger}$. This is a result of the relation:

$$(\Lambda p)^{\mu} \tilde{\sigma}_{\mu} \alpha(\Lambda) = \alpha (\Lambda^{-1})^{\dagger} p^{\mu} \tilde{\sigma}_{\mu}$$
(3.59)

This leads to the covariant transformation law for the corresponding x-space wave function $\chi^{.}(x)$:

$$(U(\Lambda)\chi)^{\cdot}(x) = \beta(\Lambda)\chi^{\cdot}(\Lambda^{-1})$$
(3.60)

Defining a 4-component Dirac spinor, we immediately read off its properties:

$$\tilde{\Psi}(p) = \sqrt{m} \begin{pmatrix} \tilde{\Phi}(p) \\ \tilde{\chi}(p) \end{pmatrix}, \text{ with } \left(p^{\mu} \gamma_{\mu} - m\right) \tilde{\Psi} = 0 \text{ and } \gamma_{\mu} = \begin{pmatrix} 0 & \sigma_{\mu} \\ \tilde{\sigma}_{\mu} & 0 \end{pmatrix}$$
(3.61)

3.2. ONE PARTICLE REPRESENTATIONS

The first two components of the Dirac equation are identical to the definition of $\tilde{\chi}$ in terms of $\tilde{\Phi}$ and the remaining equation is the Klein Gordon identity for $\tilde{\Phi}$:

$$p^{\mu}\tilde{\sigma}_{\mu}\tilde{\Phi} = m\tilde{\chi} , \ p^{\mu}\sigma_{\mu}\left(p^{\mu}\tilde{\sigma}_{\mu}\right)\tilde{\Phi} = m^{2}\tilde{\Phi}$$
(3.62)

Rewriting the inner product in terms of Ψ we obtain:

$$(\psi_2,\psi_1) = \frac{1}{2m} \int \tilde{\Psi}_2^* \tilde{\Psi}_1 \frac{d^3 p}{2\omega} = \frac{1}{2m} \int \overline{\tilde{\Psi}} \gamma_0 \tilde{\Psi} \frac{d^3 p}{2\omega}$$
(3.63)

with Ψ : = $\Psi^* \gamma_0$ the Dirac adjoint

where we have introduced the Dirac conjugate on the 4-spinors by a bar on top of the symbol (sorry for the possible confusion with the complex conjugate). Since the gamma matrices transform as a 4-vector, the Dirac formalism permits to form tensors. In x-space we have:

$$\Psi(x) = \int e^{-ipx} \widetilde{\Psi}(p) \frac{d^3p}{2\omega} , \quad \left(i\gamma_{\mu}\partial^{\mu} - m\right)\Psi(x) = 0, \qquad (3.64)$$

$$\overline{\Psi}(x)\Psi(x) = scalar, \ \overline{\Psi}(x)\gamma_{\mu}\Psi(x) = vector \ etc. \tag{3.65}$$

There are altogether 16 independent tensorial densities which one can form in this way from products of $\gamma's$.

Dirac's inner product is conveniently expressed in terms of the conserved current:

$$j_{\mu} = \overline{\Psi}_{2} \gamma_{\mu} \Psi_{1} , \ \partial^{\mu} j_{\mu} = 0, \quad \left(\overline{\Psi}_{2} \Psi_{1} \right) := \int j_{0} d^{3} x = 2m(\psi_{2}, \psi_{1})$$
(3.66)

The 4-component description allows a local matrix realization of the parity symmetry:

$$\left(\mathcal{P}\Psi\right)(x) = \gamma_0 \Psi(x_0, -\vec{x}), \quad \gamma_0 \gamma_i \gamma_0^{-1} = -\gamma_i \tag{3.67}$$

It is helpful to define a fifth γ -matrix as the product of all four: $\gamma_5 := \gamma_0 \gamma_1 \gamma_2 \gamma_3$. This matrix is block- diagonal and behaves like a pseudoscalar under parity. Therefore densities involving γ_5 are pseudo-scalars, -vectors etc. Finally we mention the u- and v-intertwiners:

$$u(p, s_{3}) = S(L(p))u(p_{R}, s_{3}), \quad u(p_{R}, \pm \frac{1}{2}) = \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \end{pmatrix}, \quad (3.68)$$
$$S(L(p)) = \begin{pmatrix} \sqrt{p^{\mu}\sigma_{\mu}} & 0 \\ 0 & \sqrt{p^{\mu}\tilde{\sigma}_{\mu}} \end{pmatrix}$$
$$v(p, s_{3}) = Cu^{*}(p, s_{3}), \quad C = i\gamma_{2} \quad (3.69)$$

It is easy to check that u and v intertwine the $s = \frac{1}{2}$ Wigner representations $D^{\left(\frac{1}{2}\right)}(R)$ resp. $D^{\left(\frac{1}{2}\right)*}(R)$ with $D^{\left[\frac{1}{2},0\right]} \oplus D^{\left[0,\frac{1}{2}\right]}$ which is implemented by the matrices $S(\Lambda)$. The so-defined v fulfills:

$$\left(-p^{\mu}\gamma_{\mu}-m\right)v=0; \quad \overline{u}u=2m, \quad \overline{v}v=-2m, \quad \overline{u}v=0 \tag{3.70}$$

It is an interesting historical side remark that Dirac found his equation in a more formalistic way. In order to overcome what he considered as a serious shortcoming of the scalar Klein Gordon equation, Dirac searched for a first order matrix differential operator which is a kind of square root of the K-G. operator, i.e. $(i\partial^{\mu}\gamma_{\mu} - m)(-i\partial^{\mu}\gamma_{\mu} - m) = \partial^{\mu}\partial_{\mu} + m^2$. The necessary and sufficient condition are the Clifford algebra conditions:

$$\left\{\gamma_{\mu},\gamma_{\nu}\right\} = 2g_{\mu\nu} \tag{3.71}$$

There is only one finite-dimensional irreducible representation, it has dimension equal to 4 (the Clifford algebra for a 2*n*-dim. space has a 2^{*n*}-dimensional irreducible representation). Our group-theoretical approach has provided us with the so called chiral representation in which γ_5 is diagonal and which for m $\rightarrow 0$ decomposes naturally into the two Weyl equations:

$$p^{\mu}\widetilde{\sigma}_{\mu}\Phi = 0, \quad p^{\mu}\sigma_{\mu}\chi^{\cdot} = 0 \tag{3.72}$$

There are many equivalent representations which are useful for other purposes. We will mention two of them. There is the representation used first by Dirac:

$$\gamma_0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \gamma_i = \begin{pmatrix} 0 & \sigma_i \\ -\sigma_i & 0 \end{pmatrix}$$
(3.73)

This representation is useful in calculations involving the nonrelativistic limit as in the hydrogen-problem. On the other hand for the field theoretic application to selfconjugate $s = \frac{1}{2}$ particles and fields the following Majorana representation is useful (with purely real $i\gamma_{\mu}$ i.e. a real Dirac operator).

$$\gamma_0 = \begin{pmatrix} 0 & \sigma_2 \\ \sigma_2 & 0 \end{pmatrix}, \quad \gamma_1 = \begin{pmatrix} i\sigma_3 & 0 \\ 0 & -i\sigma_3 \\ \sigma_2 & 0 \end{pmatrix}, \quad \gamma_3 = \begin{pmatrix} -i\sigma_1 & 0 \\ 0 & i\sigma_1 \end{pmatrix}, \quad (3.74)$$

• s=1

In this case there are several low dimensional covariant intertwining possibilities:

$$D^{(1)}(R) \to \begin{cases} D^{[1,0]} \\ D^{[0,1]} \\ D^{[\frac{1}{2},\frac{1}{2}]} \end{cases}$$
(3.75)

the first two have three components and the last is the 4-component vector description which, if restricted to the rotation group decomposes as follows:

$$D^{\left[\frac{1}{2},\frac{1}{2}\right]}(R) \cong D^{(1)}(R) \oplus D^{(0)}(R)$$
(3.76)

For an explicit description we apply the boost to the three spatial coordinate vectors e_1, e_2, e_3 :

$$(e_i(p))_{\mu} \equiv e_{\mu}(p,i) := L^{\nu}_{\mu}(p)e_{\nu}(0,i) \qquad e(0,i) := e_i \tag{3.77}$$

Remembering the definition of the Wigner rotation, the transformation law is (suppressing the vector indices):

$$\Lambda e_i(p) = L(\Lambda p)R(\Lambda, p)e_i = \sum_{i'} L(\Lambda p)e_{i'}R_{i'i}(\Lambda, p) = \sum_{i'} e_{i'}(\Lambda p)R_{i'i}(\Lambda, p)$$
(3.78)

The covariant vector like wave functions are then:

$$\tilde{V}_{\mu}(p) = \sum_{i} e_{\mu}(p,i)\tilde{\varphi}(p,i), \quad V_{\mu}(x) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int e^{-ipx} \sum_{i} e_{\mu}(p,i)\tilde{\varphi}(p,i) \frac{d^{3}p}{2\omega}$$
(3.79)

As always they fulfill the Klein-Gordon equation but, as a result of the transversality $p^{\mu}e_{\mu}(p,i) = 0$ which expresses the absence of the scalar component $D^{(0)}$, they also are divergenceless:

$$\left(\partial^{\kappa}\partial_{\kappa} + m^2\right)V_{\mu} = 0, \quad \partial^{\mu}V_{\mu} = 0 \tag{3.80}$$

Both equations can be combined into a so called Proca-Wentzel equation:

$$\left(\partial^{\kappa}\partial_{\kappa} + m^2\right)V_{\mu} - \partial_{\mu}\partial^{\kappa}V_{\kappa} = 0 \tag{3.81}$$

The covariance of this equation incorporates the transformation properties of the field (just like for the Dirac equation) and is the Euler-Lagrange equation of the Proca-Wentzel Lagrangian. Although Euler-Lagrange fields exist for any spin (e.g. for $s=\frac{3}{2}$ the Rarita-Schwinger equations), the Wigner approach, in contrast to the canonical or functional integral approach, does not provide a preferential status to Lagrangian fields.

From the definition one reads off the completeness relation:

$$\sum_{i=1}^{3} e_{\mu}(p,i)e_{\nu}(p,i) = -g_{\mu\nu} + \frac{p_{\mu}p_{\nu}}{m^2}$$
(3.82)

A limit $m \to 0$ does not exist i.e. there is no possibility to intertwine the [m = 0, s = 1] Wigner representation with $D^{\left[\frac{1}{2}, \frac{1}{2}\right]}$. The Maxwell description in terms of field strength $F_{\mu\nu}$ corresponds to $D^{[1,0]}$ or $D^{[0,1]}$. This restriction together with the demand that vector potentials are indispensable for describing the long-range electromagnetic interaction in the context of quantum theory (in classical physics vector potentials can be avoided) forces one to look for a compromise slightly outside the Wigner scheme which will be presented in the sequel.

Remark 1 The chosen covariant representations for $s=0,\frac{1}{2},1$ are "Eulerian". This means that they obey (multicomponent) spacetime differential equations

such that the Poincaré transformation properties follow from the covariance of the (matrix-valued) differential operator. For the Dirac operator this argument is well-known and can be found in any QFT textbook. The matrices which appear in the infinitesimal Lorentz-transformation are products of the matrices in the Euler equation. In fact the above equations are even "Euler-Lagrange" because they are the Euler equations of an action principle associated with a (free) quadratic Lagrangian. In fact the Wigner theory always admits (precisely) one such covariant description; for e.g. $(m,s=\frac{3}{2})$ it is the Rarita-Schwinger equation. In this case the Wigner wave functions for (m,s) particles and anti-particles correspond to the totality of all (positive and negative frequency) solutions of this E-L equation. This also prevails to the massless case, e.g. for the case below of (m=0,h=1) we obtain the Maxwell equations. However since in the present approach we are not going to use quantization methods, the existence of such classically preferred covariant fields will be of no special interest to us. All covariant representations of (m,s) are equally well suited to serve as local field coordinates for the same local net of algebras; the intrinsic carriers of the physical properties are not these field coordinates but the unique local (free) (m.s)-net of algebras which each of them generates. This is not only the correct underlying philosophy of our nonperturbative approach in chapter 6 but it also is behind our on shell (Bogoliubov-Shirkov-Epstein-Glaser) causal perturbation theory which completely avoids off shell Lagrangians, actions and functional integrals and only uses on-shell invariant interaction operators W (special elements of the free field Borchers equivalence class).

This point is the guiding thread through these notes and we will come back to it and elaborate it on many occasions. It will become clear that whereas the various quantization approaches start to enter QFT on the commutative (quasi)classical side and develop their main strength in deforming around free solutions, the more powerful nonperturbative method of chapter 6 starts from modular theory which is extremely noncommutative in the sense that it cannot even be formulated (and has no counterpart) in the context of commutative algebraic structures, i.e. it is based on one of the rare properties of LQP which are pure quantum and have no quasiclassical limit.

• s=1 m=0

In order to obtain a formalism similar to the previous case of vector mesons, one extends the two "polarization vectors" $e_i = 1,2$ in x-and y-direction by two orthogonal light-like vectors:

$$e^{\mu}_{\pm} = e^{\mu}_{0} \pm e^{\mu}_{3} = (1, 0, 0, \pm 1) \tag{3.83}$$

We choose e_+ as the reference vector k_R from which to start the boost $L(k, k_R)$. The latter consists of a rotation of the z-axis into the momentum direction $\vec{n} = \frac{\vec{k}}{\omega}$ (fixed uniquely by the standard prescription in terms of two Euler angles) and a subsequent L-boost along this direction:

$$\begin{pmatrix} 1\\ \vec{n} \end{pmatrix} \to k = \omega \begin{pmatrix} 1\\ \vec{n} \end{pmatrix}$$
(3.84)

The decomposition of a general Lorentz transformation Λ in terms of its little group component $H(\Lambda, k)$:

$$\Lambda = L(\Lambda k, k_R) H(\Lambda, k) L^{-1}(k, k_R), \qquad H(\Lambda, k) \in \tilde{E}(2)$$
(3.85)

the twofold covering of the euclidean group in two dimensions which, as explained before, is generated by two translations α , β and one rotation θ , where all euclidean parameters are functions of Λ and k which can be computed from the previous formula. One defines two transversal polarization vectors:

$$\epsilon(k,\lambda) = L(k,k_r) \begin{cases} \frac{1}{\sqrt{2}} (e_1 + ie_2), & \lambda = + \\ \frac{1}{\sqrt{2}} (-e_1 + ie_2), & \lambda = - \end{cases}$$
(3.86)

They are used as intertwiners in the attempt to define a vectorial wave function:

$$\tilde{A}_{\mu}(k) = k_{\mu}\tilde{\Phi}(k) + \sum_{\lambda=\pm} \epsilon_{\mu}(k,\lambda)\tilde{\varphi}(k,\lambda)$$
(3.87)

Here the longitudinal first component is not determined by the Wigner theory. We cannot consistently set it equal to zero, since the intertwiners generate such an additive term under the action of the $\tilde{E}(2)$ "translations":

$$G(\rho)\epsilon(k_R,\lambda) = \epsilon(k_R,\lambda) + \begin{cases} -\frac{1}{2}(\bar{\rho},0,0,\bar{\rho}), & \lambda = + \\ +\frac{1}{2}(\rho,0,0,\rho), & \lambda = - \end{cases} \qquad \rho = \alpha + i\beta \quad (3.88)$$

whereas under x-y rotations the ϵ picks up the standard Wigner phase factor. The polarization vectors do not behave as 4-vectors since they are not invariant under the euclidean translations in $\tilde{E}(2)$, as one would have expected for a (nonexisting!) bona fide intertwiner from the [0, h = 1] Wigner representation to the $D^{\left[\frac{1}{2}, \frac{1}{2}\right]}$ covariant representation. Rather the intertwiner only has Lcovariance up to additive gauge transformations i.e. up to affine longitudinal terms (for more details see section 6):

$$(U(\Lambda)A)_{\mu}(k) = \Lambda^{\nu}_{\mu}A_{\nu}(\Lambda^{-1}k) + k_{\mu}G(k)$$
(3.89)

This peculiar manifestation of the (0, h = 1) little group $\dot{E}(2)$ is the cause for the appearance of the local gauge issue in local quantum physics. Unfortunately this quantum origin is somewhat hidden in the quantization approach, where it remains invisible behind the geometrical interpretation in terms of fibre bundles.

In the covariant quantization approach, contrary to the Wigner theory, the gauge aspect becomes completely decoupled from the L-transformations. This close relation to classical fibre bundles is only obtained at the expense of leaving the realm of quantum physics and entering the world of "ghosts", i.e. mathematical tricks which allow to maintain the benefits of the standard formalism of pointlike fields in a situation which physically is already outside that formalism. In fact it is my contention, that this is one of the potential points of possible fruitful clashes between the requirements of (classical) geometry and local quantum physics. In the presence of electromagnetic interactions this problem will

appear again in our perturbative treatment of QED. Although we will try to sharpen this apparent clash between the Wigner and the quantization framework in the following section, the solution of the associated conceptual problem is not yet known. But at least one has a recipe which works in perturbation theory.

Whereas the covariantization of the canonical Wigner (m = 0, h = 1) representation can be done in terms of covariant field strength, the requirement that the scalar product be expressible in terms of a local tensorial formula necessitates the introduction of the above vector potential A_{μ} . The Lorentz (gauge) invariant inner product for the \tilde{A}_{μ} is now only positive semidefinite on individual A_{μ} (but positive definite on gauge classes):

$$\begin{pmatrix} A, A' \end{pmatrix} = -\int \tilde{A}_{\mu}^{*} \tilde{A}'^{\mu} \frac{d^{3}p}{2 |\vec{p}|} = \sum_{\pm} \int \tilde{\varphi}^{*} \tilde{\varphi}' \frac{d^{3}p}{2 |\vec{p}|}$$

$$p^{\mu} \tilde{A}_{\mu} = 0$$

$$(3.90)$$

The connection of this space H_A with the positive definite Wigner H_W space of the $\tilde{\varphi}$ -wave functions is give by factoring out the null-space H_0 :

$$H_W = \frac{H_A}{H_0} \tag{3.91}$$

This factor space representation of the Wigner space has however one drawback: the subspaces have no natural behavior under multiparticle tensoring; the well known impossibility to use a local condition $\partial^{\mu}A_{\mu}\Phi = 0$ for the characterization of physical equivalence of vectors in the Gupta-Bleuler treatment (and the necessity to use matrix elements) is a consequence. It turns out that a BRSlike cohomological extension based on a nilpotent operator yields a more natural relation of the Wigner space to the multiparticle tensor spaces. However the strongest arguments in favor of such a cohomological approach comes from the perturbative approach to renormalizable interactions of massive spin 1 particles. It turns out that without this cohomological trick there is no renormalizable solution at all. Although by its use the problem becomes soluble, the solution is extremely restrictive and essentially unique. This is in contrast to the classical situation, where the increase of spin (components) leads to an increasing number of interaction terms. The classical restriction principle in the case of massless spin 1 fields is the gauge principle. In our approach in the next chapter it is not imposed on the QFT by quantization, but rather this (quasi)classical restriction is obtained from the more fundamental (perturbative) QFT which obeys the aforementioned severe consistency restrictions characteristic for higher spin. This Bohr correspondence principle point of view is opposite to the quantization point of view of gauge theory. It leads to significant revisions about the Schwinger-Higgs mechanism and other concepts of perturbative gauge theories.

As pointed out by Weinberg[61], this gauge aspect is common to all $\left[0, h = \frac{n}{2}\right]$ representations for $\frac{n}{2} \geq 1$. There are simply no intertwiners from this Wigner representation to $D^{[A,A]}$ symmetric tensors, rather the possibility of intertwining

is restricted to $D^{[A,B]}$ with |A - B| = h (h the Wigner helicity). The covariant vector potentials for s = 1 and the covariant symmetric tensor $g_{\mu\nu}$ for s = 2 of the classical general relativity can at the quantum level only be introduced at the prize of nonlocal (noncompactly localizable) $\psi^{[A,A]}$ fields or by the cohomological trick.

The case of general [m, s] intertwiners u is a routine exercise in Clebsch-Gordan gymnastics. One uses the intertwining relation for u:

$$u(p)D^{(s)}(R(\Lambda, p)) = D^{[A,B]}(\Lambda)u(\Lambda^{-1}p)$$
(3.92)

for the calculation of the *u*'s. Here we found it convenient to interpret the intertwiner *u* as a rectangular matrix with 2s+1 columns and (2A+1)(2B+1) rows. The first step consists in analyzing this equation for $p = p_R$ (Weinberg) with the result that the $u(\vec{0})$ is proportional to the Clebsch-Gordan coefficients:

$$u(\vec{0}) \sim C_{AB}(s, s_3; a, b)$$
 (3.93)

The second step consists in an application of a boost:

$$u(p) = D^{[A,B]}(L(p))u(0)$$
(3.94)

For details we refer to Weinberg [61].

The mathematical method behind Wigner's representation theory, together with the Frobenius theory of induced representations of finite groups, was extended by G. Mackey into the general theory of induced representations which in turn is a special case of a theory of inductions and reductions of (von Neumann) algebras by M. Rieffel and others (see mathematical appendix).

For the case at hand, the starting point is the semi-direct product of two locally compact groups of which one is abelian and denoted by T and the other say K acts as automorphisms on T:

$$G \equiv K \times_{\alpha} T : (k_2, t_2) \cdot (k_1, t_1) = (k_2 k_1, t_2 + \alpha_{k_2}(t_1))$$

The action of K on the dual group \tilde{T} , which is defined as:

$$(k \circ \tilde{t})(t) \equiv \tilde{t}(\alpha_{k^{-1}}(t))$$

foliates the \tilde{T} into transitive K-orbits \mathcal{O} . Each point \tilde{t} on one \mathcal{O} defines a stability group:

$$K_{\tilde{t}} \equiv \left\{ k \in K \mid k \circ \tilde{t} = \tilde{t} \right\}$$

and apart from possibly singular points, these "little groups" for different \tilde{t} are all inner equivalent (i.e. inside G). Often one finds a geometrically preferred reference point \tilde{t}_0 (e.g. the restframe momentum on the mass hyperboloid). Let π be an irreducible representation of $K_{\tilde{t}}$ and $U_{\pi}(k, \tilde{t})$ the associated $(K, O, U(H_{\pi}))$ -cocycle i.e.:

$$U_{\pi}(k_{2}, k_{1} \circ t)U_{\pi}(k_{1}, t) = U_{\pi}(k_{2}k_{1}, t)$$
$$U_{\pi}(k, \tilde{t}_{0}) = \pi(k)$$

Introducing a Borel map $L: \mathcal{O} \to K$ (the family of boosts in Wigner's theory), one may take:

$$U_{\pi}(k,\tilde{t}) = \pi(L(k\circ\tilde{t})^{-1}kL(\tilde{t}))$$

The Mackey induction "machine" associates to every pair (\mathcal{O}, π) an irreducible representation of G on the Hilbert space $H \equiv L^2(\mathcal{O}, H_\pi)$:

$$(V(k,t)\psi)(\tilde{t}) = \tilde{t}(t)U_{\pi}(k,k^{-1}\circ\tilde{t})\psi(k^{-1}\circ\tilde{t})$$

Here we assumed that there exists an K-invariant measure on \mathcal{O} ; for a quasi-invariant measure the right hand side has to be corrected by a Radon Nykodym derivative. For more details I refer to [104]

3.3 Wigner Theory and Free Fields

We now use the Wigner representation theory in order to construct fields in bosonic or fermionic Fock spaces. The creation operators in momentum space should transform in the same way as the one particle states since their application to the vacuum vector creates the latter.

$$U(\Lambda)a^{*}(p,m)U^{*}(\Lambda) = \sum_{m'} a^{*}(\Lambda p,m')D_{m'm}^{(s)}(R(\Lambda,p))$$
(3.95)

For computational convenience we identify the Wigner rotation with its unimodular matrix representation:

$$R(\Lambda, p) \longrightarrow \left(\sqrt{\frac{1}{m} \left(\Lambda p\right)^{\mu} \sigma_{\mu}}\right)^{-1} \alpha(\Lambda) \sqrt{\frac{1}{m} p^{\mu} \sigma_{\mu}}$$
(3.96)

The corresponding relation for the annihilation operator contains the complex conjugate matrix D^* which is equivalent to D:

$$D(i\sigma_2)D^*(R)D(-i\sigma_2) = D(R)$$
(3.97)

If the particles are charged, there are also operators $b^{\#}(p,m)$ which describe annihilation and creation of antiparticles with the same mass and spin and hence the same transformation property as $a^{\#}(p,m)$. In order to obtain covariant operators one uses the intertwiners u and v introduced in the previous section. Interpreting these intertwiners as p-dependent rectangular matrices of size $N \times (2s+1)$ with N= dimension of the representation space on which the matrices $D^{[A,B]}(\Lambda)$ act, we have:

$$D^{[A,B]}(\Lambda)u(\Lambda^{-1}p) = u(p)D^{(s)}(R^{-1}(\Lambda,p))$$

$$D^{[A,B]}(\Lambda)v(\Lambda^{-1}p) = v(p)D^{(s)*}(R^{-1}(\Lambda,p)), \quad v(p) = u(p)D^{(s)*}(i\sigma_2)$$

(3.98)

Therefore we find the following covariant creation and annihilation operators:

$$\begin{aligned}
A^{(*)}(p) &= \sum_{m} v(p,m)a^{*}(p,m) \\
B^{(*)}(p,) &= \sum_{m} v(p,m)b^{*}(p,m) \\
A(p) &= \sum_{m} u(p,m)a(p,m) \\
B(p) &= \sum_{m} u(p,m)b(p,m)
\end{aligned}$$
(3.99)

We have added a bracket to the * in order to indicate that the covariant creation operator is not exactly the hermitian adjoint of the covariant annihilator and we used m (magnetic quantum number) instead of the cumbersome s_3 notation.. The Fourier transform preserves covariance:

$$\psi_A^{(-)}(x) := \frac{1}{(2\pi)^{\frac{3}{2}}} \int e^{-ipx} A(p) \frac{d^3 p}{2\omega} , \qquad \psi_A^{(+)}(x) := \frac{1}{(2\pi)^{\frac{3}{2}}} \int e^{ipx} A^*(p) \frac{d^3 p}{2\omega} \psi_B^{(-)}(x) := \frac{1}{(2\pi)^{\frac{3}{2}}} \int e^{-ipx} B(p) \frac{d^3 p}{2\omega} , \qquad \psi_B^{(+)}(x) := \frac{1}{(2\pi)^{\frac{3}{2}}} \int e^{ipx} B^*(p) \frac{d^3 p}{2\omega}$$
(3.100)

obey the covariant transformation law:

$$U(\Lambda)\psi(x)U^*(\Lambda) = D^{[A,B]}(\Lambda)^{-1}\psi(\Lambda x)$$
(3.101)

We want to construct local covariant fields i.e. covariant fields which (anti)commute for spacelike distances. The physical motivation is Einstein causality for local observables. Prominent local observables associated with charged fields are e.g. currents. Since they are typically second or higher even degree polynomials in the fields, the (anti)commutation of the fields is sufficient for the Einstein causality (spacelike commutativity) of the local observables. Fields which are themselves observables as e.g. the Maxwell field, must obey spacelike commutation relations.

It is well known that support properties in momentum space as the restriction to the forward light cone prevent support properties of (anti)commutators in x-space. The former give rise to analytic properties of the latter. The standard example is the Fourier transform of a function with support in the positive half-axis which is the boundary value of a function analytic in the upper half-plane. According to the Schwarz reflection principle, such function cannot vanish in a dense real subset without vanishing identically. The above Fourier transforms are multidimensional counterparts in which the halfline is replaced by the forward light cone and the upper half plane by a tube $z_{\mu} = x_{\mu} + iy_{\mu}$ with y in the dual cone i.e. the backward light cone. We therefore make the following Ansatz for local fields:

$$\psi_A(x) = \psi_A^{(-)}(x) + \psi_A^{(+)}(x), \quad \psi_B(x) = \psi_B^{(-)}(x) + \psi_B^{(+)}(x) \psi(x) = \psi_A^{(-)}(x) + \psi_B^{(+)}(x)$$
(3.102)

Complex coefficients in this linear combination bring no gain in generality, since they can be absorbed into redefinitions. The following calculations show that all these combinations between different frequency parts are local covariant fields. The first two combinations are only physically useful if A and B would be (accidentally equal mass and spin) selfdual particles. If on the other hand, there is a charge superselection rule between A and B i.e. B is the antiparticle of A, then we are forced to take the ψ combination because otherwise we would not be able to form local (Einstein-causal) neutral observables. In this sense causality and the superselection principle require the existence of anticharged particles of the same (m,s).

Returning to our notation for indices of irreducible finite dimensional representations for the Lorentz-group, we find the following relation between the spin and the spacelike (anti)commutativity:

Theorem 8 (Spin-statistics for free fields) :

$$\left[\psi_{a,b}^{[A,B]}(x),\psi_{a',b'}^{[A,B]}(y)\right]_{\pm} = 0 \quad for \quad (x-y)^2 < 0 \tag{3.103}$$

where the +sign i.e. the anticommutator is to be taken for A+B=halfinteger.

The proof consists in calculating the vacuum expectation value of the product of two fields in the two different orders. Each two-point function is the Fourier transform of a sesquilinear expression in the u resp. v intertwiners, e.g.

$$\left\langle \psi_{a,b}^{[A,B]}(x), \psi_{a',b'}^{[A,B]*}(y) \right\rangle_{0} = \frac{1}{(2\pi)^{3}} \int e^{-ip(x-y)} \sum_{m} u_{ab}^{[A,B]}(p,m) \overline{u_{a'b'}^{[A,B]}}(p,m) \frac{d^{3}p}{2\omega}$$
(3.104)

The computation of the *m*-sum in the integrand is (as the computation of the intertwiners) a purely group theoretic problem and it yields a $(2A + 1) \cdot (2B + 1) \times (2A + 1)(2B + 1)$ matrix with $P_{a,b;a',b'}(p)$ covariant *p*-polynomial entries. Therefore in x-space we may write the correlation function in terms of the scalar two-point function $i\Delta^{(+)}(\xi)$ (studied in detail below):

$$\left\langle \psi_{a,b}^{[A,B]}(x), \psi_{a',b'}^{[A,B]*}(y) \right\rangle_{0} = P_{a,b;\,a',b'}(i\partial_{x})i\Delta^{(+)}(x-y)$$
 (3.105)

The polynomial matrix is even/odd under the transposition of matrix indices together with $x \leftrightarrow y$ for A + B =integer/halfinteger. Since the scalar function $i\Delta^{(+)}(\xi)$ is symmetric in ξ for spacelike ξ , only the difference (commutator)/sum (anticommutator) vanishes for spacelike distances, i.e integer/halfinteger \leftrightarrow commutator/anticommutator. We mention that the commutation relation between two $\psi's$ must follow the same \pm rule as the above $\psi - \psi^*$. Since the two-point function vanishes for two fields with the same charge, the proof requires the use of the 4-point function and will only be mentioned in the case of interacting fields.

It is helpful to illustrate this spin-statistics connection with free fields for $s = 0, \frac{1}{2}, 1$.

3.3. WIGNER THEORY AND FREE FIELDS

With
$$\psi(x) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int \left(e^{-ipx} A(p) + e^{ipx} B^*(p) \right) \frac{d^3p}{2\omega}$$
 we obtain:

$$\left[\psi(x),\psi(y)\right]_{\pm} = \frac{1}{(2\pi)^{\frac{3}{2}}} \int \left(e^{-ip(x-y)} \pm e^{ip(x-y)}\right) \frac{d^3p}{2\omega} = i\Delta^{(+)}(x-y)\pm i\Delta^{(-)}(x-y)$$
(3.106)

Here $i\Delta^{(-)}(\xi) := i\Delta^{(+)}(-\xi)$ and the momentum space integrals may be expressed in terms of Hankel functions. One first uses the fact that $i\Delta^{(+)}(\xi)$ is analytic in the tube $\xi \to \zeta = \xi - i\eta$ with $\eta \in \bar{V}^+$, the closed forward light cone as a result of the spectrum property $p \in V^+$. This means that the euclidean vector $(\xi_4 = i\xi_0, \vec{\xi})$ is in the analyticity region at least if $\xi_0 \ge 0$. This analytic continuation is part of the so called Wick-rotation. In this euclidean domain one now rewrites the integral for $i\Delta^{(+)}$ in terms of an euclidean contour integral:

$$i\Delta^{(+)}(x) = \frac{1}{(2\pi)^4} \int_C e^{ipx} \frac{d^4p}{p^2 + m^2}$$
(3.107)

The contour C in the complex p_0 -plane is the imaginary ξ_0 -axis or the new (Wick-rotated) $\xi_4 = i\xi_0 + \epsilon$ axis. The proof of this claim follows simply by closing the contour by an infinitely large half-circle in the upper half plane on which the integrand vanishes sufficiently and the subsequent application of the residuum theorem to the pole at $p_4 = i\sqrt{\vec{p}^2 + m^2}$. Since the Minkowski metric has disappeared and there is no restriction on the Wick-rotated ξ_4 (the euclidean representation achieved an analytic continuation to all real ξ_4), the remaining task is to perform a euclidean Fourier-integral with a rotational invariant rational integrand. The d-dimensional integration in polar coordinates requires the same amount of work as d=4.

$$\frac{1}{(2\pi)^d} \int e^{ip\xi} \frac{d^d p}{p^2 + m^2} = \frac{1}{(2\pi)^d} \frac{2(\sqrt{\pi})^{d-1}}{\Gamma(\frac{d-1}{2})} \int_0^\infty \frac{p^{d-1}}{p^2 + m^2} dp \int_0^\pi \sin^{d-2\theta} d\theta e^{ipr\cos\theta}$$
(3.108)

$$=\frac{2\left(\sqrt{\pi}\right)^{d}}{\left(2\pi\right)^{d}}\int\frac{2^{\frac{d}{2}-1}}{\left(rp\right)^{\frac{d}{2}-1}}\frac{1}{p^{2}+m^{2}}J_{\frac{d}{2}-1}(pr)p^{d-1}dp$$
(3.109)

or the Bessel functions J as well as a formula linking the Hankel function of the first type to an integral over a Bessel function. The Hankel function $H_{\nu}(z)$ is analytic in the cut z-plane with a cut running from $-\infty$ to zero i.e. K(z) has a cut for $z^2 \leq 0$. Specializing to d =4, we obtain the following representation of the free field two-point function as a boundary value of an analytic function:

$$i\Delta^{(+)}(\xi) = \lim_{\epsilon \to 0} \frac{1}{4\pi^2} \frac{m}{\sqrt{-(\xi_0 - i\epsilon)^2 + \vec{\xi}^2}} K_1(m\sqrt{-(\xi_0 - i\epsilon)^2 + \vec{\xi}^2}) \qquad (3.110)$$

As expected, the space-like points are (together with the euclidean points) in the analytic domain. The distributional boundary value prescription becomes important only in the time-and light-like region where the transcription of the Hankel H or Kelvin K function in terms of J- and N-functions and the subsequent performance of the ϵ -limit gives the physical boundary values in terms of Schwartz distributions:

$$i\Delta^{(+)}(\xi) = \frac{1}{4\pi}\epsilon(\xi_0)\delta(\xi^2) + \frac{m}{\sqrt{-\xi^2}}\frac{i}{8\pi}\left(J_1(m\sqrt{\xi^2})\epsilon(\xi_0) + iN_1(m\sqrt{\xi^2})\right), \ \xi^2 \ge 0$$
(3.111)

The strength of singularity on the light cone (determined by the singularities of K_1 or N_1) is independent of the mass and given by the zero mass two-point correlation function:

$$iD^{(+)}(\xi) = \lim_{\epsilon \to 0} \frac{1}{4\pi^2} \frac{1}{-(\xi_o - i\epsilon)^2 + \vec{\xi}^2}$$
(3.112)

The particular ε -prescription is not only the consequence but even equivalent to the positive energy property of its Fourier transform, a fact which is often tends to be overlooked. The strength of the x-space short distance behavior is independent of the state, e.g. the two-point function in the ground state and that in any other vector or density matrix state (e.g. in a temperature state, as considered later) on the same *-algebra have the same leading light-cone behavior.. This also applies to the generalization to curved space-time (chapter 4). The next-to leading behavior (the log-term) in K_1 does however depend on the mass. The fact that correlation functions have analyticity properties in space-like regions is however very specific for the vacuum state; other states in the same "folium" of states have only spacelike smoothness but no analytic behavior in their correlation functions. The dependence of the singularities on the space-time dimension follows from the properties of the K_{ν} functions. It is conveniently encoded into the notion of "operator dimensions" of the fields e.g. one says e.g. that dimA = 1 (in mass units) for d = 3 + 1 and $dimA = \frac{1}{2}$ for d = 2 + 1 if the two-point function has the leading singularity $(-\xi^2)^{-\dim \hat{A}}$

It turns out that the correlation functions of the higher spin free fields can all be expressed in terms of $i\Delta^{(+)}$ and its zero mass limit $iD^{(+)}$ with matrix valued differential operators in front. We again look at the important special cases $s = \frac{1}{2}$, 1 before we sketch properties of general free fields.

•
$$\mathbf{s} = \frac{1}{2}$$

The ansatz for the positive and negative frequency parts for the local spinor field (in analogy to the previous scalar field) is (using the condensed notation from the beginning of this section for A(p) and B(p)):

$$\psi(x) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int \left(e^{-ipx} A(p) + e^{ipx} B^*(p) \right) \frac{d^3p}{2\omega}$$
(3.113)

For the (anti)commutators of the covariant creation and annihilation operators one needs to know the completeness relations for the u-and v-spinors:

$$\sum_{s_3} u(p, s_3) \bar{u}(p, s_3) = p^{\mu} \gamma_{\mu} + m = 2m\Lambda_+$$
(3.114)

3.3. WIGNER THEORY AND FREE FIELDS

$$\sum_{s_3} v(p, s_3) \bar{v}(p, s_3) = p^{\mu} \gamma_{\mu} - m = -2m\Lambda_-$$
(3.115)

where Λ_{\pm} are projectors $\Lambda_{+} + \Lambda_{-} = 1$ on the \pm frequency subspaces in the 4-dim Dirac spinor space and the bar on the 4-spinors denotes the Dirac's conjugate (3.63). With:

$$\left\langle \psi_{\alpha}(x)\overline{\psi_{\beta}}(y) \right\rangle = \frac{1}{(2\pi)^3} \int e^{-ip(x-y)} \sum_{s_3} u_{\alpha}(p,s_3) \bar{u}_{\beta}(p,s_3) \frac{d^3p}{2\omega} (3.116)$$
$$= \left(-i\partial_x^{\mu}\gamma_{\mu} + m \right)_{\alpha,\beta} i\Delta^{(+)}(x-y)$$

$$\langle \overline{\psi}_{\beta}(y)\psi_{\alpha}(x)\rangle = \frac{1}{(2\pi)^{3}} \int e^{ip(x-y)} \sum_{s_{3}} \overline{v}_{\beta}(p,s_{3})v_{\alpha}(p,s_{3}) \frac{d^{3}p}{2\omega}$$
(3.117)
$$= -\left(i\partial_{y}^{\mu}\gamma_{\mu} + m\right)_{\alpha,\beta} i\Delta^{(+)}(y-x)$$

we obtain with $\psi^{\#} = \psi \operatorname{or} \overline{\psi}$

$$\left\{\psi^{\#}(x),\psi^{\#}(y)\right\} = 0 \ if \ (x-y)^2 < 0$$
 (3.118)

$$\left\{\psi(x),\overline{\psi}(y)\right\} = \left(-i\partial_x^{\mu}\gamma_{\mu} + m\right)i\Delta(x-y)$$
(3.119)

whereas the commutator is nonvanishing for spacelike distances. We get the first glimpse at the spin-statistics connection.

The present construction of local $\psi's$ also sheds some light on the physical interpretation of the *v*-spinors in connection with the charge conjugation symmetry. The latter transformation is defined in Fock-space by :

$$\mathcal{C}A(p,s_3)\mathcal{C}^* = B(p,s_3) \tag{3.120}$$

Its action on the local fields is local and the transformation law involves a matrix C in Dirac space:

$$\psi^C := \mathcal{C}\psi\mathcal{C}^* = C\psi^* \tag{3.121}$$

In the helicity representation used here, the matrix is $C = \gamma_2$, whereas in the Majorana representation one finds C = 1. This matrix transforms the *u*-spinors into the *v*'s and vice versa and therefore is the image of $D^{(s)}(i\sigma_2)$ under the intertwining map into the Dirac spinor space. It is an additional fringe benefit that via the Dirac doubling all global Fock-space symmetries as \mathcal{P} , T and \mathcal{C} have local representations with constant matrices on Dirac spinors. Furthermore the Dirac description goes over into the two decoupled Weyl equation in the zero mass limit. Finally we notice that dim $\psi = \frac{3}{2}$.

• s=1,m≠0

The massive (vector meson) case is straightforward. The local field is :

$$V_{\mu}(x) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int \sum_{i} e_{\mu}(p,i) \left(e^{-ipx} a(p,i) + e^{ipx} b^{*}(p,i) \right) \frac{d^{3}p}{2\omega}$$
(3.122)

Its 2-point function results from the completeness relation of the polarization vectors :

$$\langle V_{\mu}(x)V_{\nu}(y)\rangle = \left(-g_{\mu\nu} - \frac{\partial_{\mu}\partial_{\nu}}{m^2}\right)i\Delta^{(+)}(x-y)$$
(3.123)

It is obvious that only the commutator can vanish for spacelike distances. Different from the previous case, vector meson fields does not permit a zero mass limit. Therefore we should not be surprised to meet some peculiarities in the vectorial description of photons. Finally we have dim $V_{\mu} = 2$ in contrast to the expected classical dim_{class} $V_{\mu} = 1$

• s=1,m=0

Here we may use a physical description in terms of nonlocal (semiinfinite string localized) vector potentials in the physical Fock space which under the unitary Lorentz transformations suffer a affine transformation law (3.89) which will be analyzed in more details in section 6 of this chapter. In the following we briefly describe the traditional Gupta-Bleuler photon formalism For a formally local² description in terms of vector fields, the longitudinal part which the stability group transformation behavior of wave functions demands (see previous section) is not enough; one also needs "scalar photons":

$$\tilde{A}_{\mu}(k) = e_{\mu}^{(+)}(k)c_{+}(k) + e_{\mu}^{(-)}(k)c_{-}(k) + \sum_{\lambda=\pm}\epsilon_{\mu}(k,\lambda)a(k,\lambda)$$
(3.124)

$$A_{\mu}(x) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int \left(e^{-ikx} \tilde{A}_{\mu}(k) + e^{ikx} \tilde{A}_{\mu}^{*}(k) \right) \frac{d^{3}k}{2\left| \vec{k} \right|}$$
(3.125)

Here the $e^{(\pm)}$ are obtained by boosting the light-like vectors $(1, 0, 0, \pm 1)$, i.e. $e^{(+)}$ is the old longitudinal part. We obtain the covariant two point function :

$$\langle A_{\mu}(x)A_{\nu}(y)\rangle = \frac{-g_{\mu\nu}}{(2\pi)^3} \int e^{-ik(x-y)} \frac{d^3k}{2\left|\vec{k}\right|}$$
 (3.126)

from the completeness relation of the four vectors:

$$\sum_{\lambda=\pm} \epsilon_{\mu}(k,\lambda)\epsilon_{\nu}^{*}(k,\lambda) + \frac{1}{2} \left(e_{\mu}^{(+)}(k)e_{\nu}^{(-)}(k) + e_{\mu}^{(-)}(k)e_{\nu}^{(+)}(k) \right) = -g_{\mu\nu} \quad (3.127)$$

This is the case iff the $a^{\#'s}$ behave in the standard way and the c's have a nondiagonal inner product which, as a result of the presence of the scalar photons c_{-} corresponds to a genuine indefinite (not just semidefinite) metric:

$$\left\langle c_{+}(k)c_{-}^{*}(k')\right\rangle = 2\left|\vec{k}\right|\delta(\vec{k}-\vec{k}')$$
 (3.128)

²Local here means pointlike, i.e. fields which can be smeared with unrestricted Schwartz test functions. Without the unphysical components, we would have to restrict vector-valued test functions f_{μ} by demanding transversality.

The *a*'s mix with the *c*'s under L-transformations viz. the comments on gauge transformations in the wave function discussion of the last section. But whereas in the wave function treatment of photons in terms of vectors only an extension by c_+ "photons" was necessary, in our present Fock space description of formally local point like vector potentials we need in addition the negative metric causing c_- scalar contribution (or alternatively a notion of pseudo-adjoint different from the bona fide adjoint³). Only in the weak sense of matrix elements the condition of absence of scalar "photons" can be enforced in terms of a *local condition*:

$$\langle \psi \left| \partial^{\mu} A_{\mu} \right| \varphi \rangle = 0 \tag{3.129}$$

Whereas in the case of the Wigner space it was possible to have the transversality condition as a defining property of the Wigner representation space, it is not possible to have transversal pointlike free vector potentials in Fock space but rather only transversal matrix elements between physical multiparticle states are compatible. This is equivalent to the use of a nonlocal condition on vectors in terms of the annihilation part of A: $\partial^{\mu}A^{(+)}_{\mu}|\varphi\rangle = 0$. In the interacting case this Gupta-Bleuler formalism only works because one can show that $\partial^{\mu}A_{\mu}$ continues to fulfill the free wave equation. In order to preserve the transversality condition under the tensor product formation of multiparticle states, we will use in section 5 of the next chapter a cohomological BRS-like formalism. The main reason why we postpone the introduction of that more stable (under interactions) formalism is that the main argument in its favor is not so much the incorporation of zero mass but rather the requirement of renormalizability of massive and massless higher spin ≥ 1 fields under the deformations by interaction implementing local functions W of free fields. We return to the description of the Gupta-Bleuler formalism.

As we mentioned already in the previous section, all these problems are absent (apart from the fact that the Wigner inner product does not permit a local rewriting in terms of local field strength amplitudes), if we describe the photons in terms of field strength instead of vector potentials. In that case we only deal with physical photons:

$$F_{\mu\nu}(x) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int \sum_{\lambda=\pm} \left(e^{-ikx} u_{\mu\nu}(k,\lambda) a(k,\lambda) + h.a. \right), \quad (3.130)$$
$$u_{\mu\nu}(k,\lambda) = ik_{\mu}\epsilon_{\nu} - \{\mu \leftrightarrow \nu\}$$

But in order to formulate interacting QED with its specific long range interaction⁴ through the local renormalizable coupling of free fields, the vector potential has been indispensable. The fact that we do not know how to employ string localized vector potentials $A_{\mu}(x, n)$ with n being the spacelike vector along the

³If one retains the positive metric, the Lorentz generators will be only pseudo selfadjoint and the Poicaré symmetries do not operate as automorphisms on *-algebras, i.e. both descriptions lead away from *-algebras. Whereas for the free case the metric can be changed at will, the situation for interacting theories become mathematically uncontrollable and physically questionable.

 $^{^4\}mathrm{i.e.}$ the quantum counterpart of the minimal external electromagnetic coupling.

string direction, is the origin for certain conceptual complications in spin ≥ 1 LQP and the explanation for the special role of the formalism of local gauge theories. In the standard indefinite metric method, the descend from the unphysical vectorial description defined by a free field with the two-point function 3.126 to the physical photons in the sense of Wigner is done with the help of the Gupta-Bleuler method. By the above transversality constraint one eliminates the scalar c_+ " photons". This step leads from the indefinite metric "Fock"-space to a *positive semidefinite* subspace \mathcal{H}_{ps} which still contains the zero norm longitudinal "photons". The elimination of the latter can only be accomplished through descend to a factor space (defined by equivalence classes):

$$\mathcal{H}_{phys.} = \frac{\mathcal{H}_{ps}}{\mathcal{H}_{ps}^{(0)}}, \quad \mathcal{H}_{ps}^{(0)} = nullspace \ of \ zero \ norm \ vectors \tag{3.131}$$

The Gupta-Bleuler method (as well as its BRS generalization) has a certain formal geometrical elegance in renormalized perturbation theory, but its conceptual physical aspects leave a lot to be desired. Of course the conceptual and mathematical troubles start only with interactions. I do not know any controllable mathematics for indefinite metric algebras which could be used for structural investigations i.e. spin&statistics, localization etc. The alternative to stay in physical and introduce nonlocal vectorpotentials has not been seriously considered because the standard perturbative framework requires pointlike fields. In the net approach this problem seems to be related to finding a natural algebraic analogon of semi-infinite "axial gauges" (3.171). More remarks on such ideas will appear in a later section (3.5).

The higher spin cases are treated analogously. We only give a brief sketch, the details may be found in Weinberg's book. Using the completeness relations of the general [m, s] intertwiners one finds a two point function of the form (3.105) with P a covariant polynomial in the derivatives. Again one observes the possibility of a matrix realization of \mathcal{P} , T and \mathcal{C} if one uses the "doubling" $D^{[A,B]} \oplus D^{[B,A]}$. The requirement of locality leads to the spin-statistics connection of the previous theorem.

The zero mass case leads to a severe restriction between A, B and the helicity h = s namely |A - B| = h. For h = 2 the analogy with classical general relativity and the long range nature of the graviton interaction again demands to side step this rule by using a gauge theoretic description in terms of a symmetric tensor $g_{\mu\nu}$. in analogy (but more complicated) with the vector potential for h = 1. The massive $s = 0, s = \frac{1}{2}$ and s = 1 fields as well as their massless helicity counterparts are "Eulerian" i.e. the transformation property is a consequence of the matrix form of the differential operator which is the 4×4 Dirac or the 4×4 s = 1 Proca-Wentzel operator. Also for higher spins there are such Eulerian operators e.g. the Rarita-Schwinger operator for $s = \frac{3}{2}$. But most of the covariantizations of the Wigner representations are not "Eulerian" and can not be used for Lagrangians and canonical quantization procedures (in particular all minimal i.e. 2s+1 component descriptions for s > 0). But this does not make them less physical or useful. As already mentioned the helicity restriction of zero

mass fields only puts in evidence a problem which is also looming behind the spin ≥ 1 massive fields. As a result of their high operator dimensions dim $\psi \geq 2$ (there is no covariant $\psi^{[A,B]}$ with operator dimension below this value), a simple minded trilinear coupling analysis for the interaction operator W reveals that $\dim W \geq 5$, and hence W is formally nonrenormalizable. The existence of the "magic cohomological trick" (section 5) reveals that the idea that only local functions of local fields with $\dim W \leq 4$ (i.e. elements of the free field Borchers class) can produce local deformations is not quite correct. As long as one avoids the vector potentials and their higher spin generalizations, the cohomological approach, which restricts the consistent expressions for W severely, produces deformed local expressions whose operator dimensions is only off by logarithmic terms from their free field dimensions. The best way to deal with zero mass problems is to view them as zero mass limits of the massive cohomological approach. As explained in section 5, one obtains the classical gauge picture as the semiclassical manifestation of this more fundamental LQP formalism in agreement with Bohr's correspondence principle (which is the opposite of quantization).

Finally we make the following important observation. Despite the fact that the Wigner theory gives a unique description for each mass and spin, we completely loose this uniqueness on the level of local fields. We obtain a countable covariant local family of fields which all share the same Fock-space operators but differ in their u and v intertwiners. This is true for any spin; even in case of s = 0 we may use vectors or tensors which of course turn out to be just derivatives of the standard scalar field. In the next section we will show that these different fields generate the same local algebras. With respect to those algebras they behave like different coordinates in geometry. The intrinsic physical information is in the "net" of local algebras. As in geometry, it is of course not wrong to use field coordinates in LQP.

3.4 The Equivalence Class of a Free Field

We have seen that the Wigner representation theory together with the locality principle leads to a multitude of (m, s) fields. Actually the set of physically equivalent descriptions is even much larger. Let us understand this first in the case of a neutral scalar field:

$$A(x) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int \left(e^{-ipx} a(p) + e^{ipx} a^*(p) \right) \frac{d^3p}{2\omega} = A^{(-)}(x) + A^{(+)}(x) \quad (3.132)$$

Such operator-valued distributions cannot be pointwise multiplied as classical functions can. In order to find a substitute for classical pointwise multiplication, one studies first the matrix elements of products of A at different points e.g.

$$\langle \Omega | A(x_1) A(x_2) \dots A(x_n) | \Omega \rangle \tag{3.133}$$

Clearly the terms which become singular for coalescent points (more generally if one of the difference vectors $x_i - x_j$ becomes light like) results from "Wickcontractions":

$$A^{(-)}(x_i)A^{(+)}(x_j) = i\Delta^{(+)}(x_i - x_j) + A^{(+)}(x_j)A^{(-)}(x_i)$$
(3.134)

 $i\Delta^{(+)}$ which are generated by commuting the annihilation components $A^{(-)}$ through the $A^{(+)}s$ to the right vacuum. The resulting terms in which the annihilators are on the right of creators i.e. operator products of the form:

$$A^{(+)}(x_{i_1})....A^{(+)}(x_{i_k})A^{(-)}(x_{i_{k+1}})....A^{(-)}(x_{i_n})$$
(3.135)

have vanishing vacuum expectation values and finite matrix elements between finite (but arbitrarily large) particle number vectors in Fock space. In those "Wick-ordered" products the limit $x_i \to x$ of colliding points can be taken without peril. Therefore one defines local functions of the local field A(x) in the sense of Wick-ordering as:

$$: A^{n}(x) := \sum_{k-partitions} A^{(+)}(x) \dots A^{(+)}(x) A^{(-)}(x) \dots A^{(-)}(x)$$
(3.136)

i.e. the terms which result by simply ignoring the contractions. These are the equal point limits of "split-point" Wick-products:

$$: A(x_1)...A(x_n) := \sum_{k-partitions} A^{(+)}(x_{i_1})...A^{(+)}(x_{i_k})A^{(-)}(x_{i_{k+1}})...A^{(-)}(x_{i_n})$$
(3.137)

where the order inside the k-partition is the original order. The usefulness of the Wick-ordering results from the fact that despite their nonlocal origin in terms of frequency separation, the resulting operators are local or multilocal. This is because the above definition is equivalent to the following obviously local inductive definition:

where the lower brackets represent the Wick- "contracted" pairs and the sum goes over all m-pairings and finally over all m. Clearly this formula provides an inductive definition of Wick ordering (the right hand sum only involves ordered products with a lower number of operators). The proof that the previous frequency-ordering definition leads to this inductive formula is elementary and left to the reader. The multi-localized (at $x_1...,x_n$) product obviously approaches the one-fold localized Wick power of the free field. Here the word "local" has a classical as well as a quantum meaning. Classically it means that one only has to know the A's around the spacetime point x in order to compute : $A^n(x)$:, whereas the operational quantum meaning is that this pointlike composite commutes with all the A's whose localization is spacelike with respect to x (locality in the sense of Einstein causality, which in local Quantum Theory means simultaneous measurability). The best way to reconcile the classical with the quantum aspects is the notion of Borchers class defined below. In order to get a feeling for the properties of local composites, let us look at two point functions of n^{th} Wick powers:

$$\langle : A^{n}(x) :: A^{n}(y) : \rangle = n! \left(i\Delta^{(+)}(x-y,m^{2}) \right)^{n}$$

$$= \int_{m^{2}}^{\infty} \rho^{(n)}(\kappa^{2}) i\Delta^{(+)}(x-y,\kappa^{2}) d\kappa^{2}$$
(3.139)

here we indicated the dependence on mass parameters. According to a wellknown and easy to prove statement of Kallén and Lehmann, any two-point function of a Lorentz covariant scalar field in a theory with positive energy conditions has a spectral representation in terms of a positive ρ -function. In the present case the ρ is the n-fold convolution of the forward mass shell distribution $\theta(p_0)\delta(p^2 - m^2)$. This only involves integrations over finite regions of *p*-space and may be carried out explicitly with the result that the $\rho's$ are polynomial bounded (depending on n) functions.

The family of pointlike Wick-ordered composites is bigger than the above illustrations; also derivatives as : $\partial_{\mu}A(x)\partial_{\nu}A(x)$: etc. are included. It is very gratifying that also the inverse is true:

Theorem 9 The set of fields in Fock space which commute for spacelike distances with the free field A(x):

$$\mathcal{B}(A) := \left\{ B \mid [B(x), A(y)] = 0 \quad for \quad (x - y)^2 < 0 \right\}$$
(3.140)

is called the Borchers equivalence class $\mathcal{B}(A)$ and consists precisely of the local composites which are generated by Wick powers.

The equivalence class aspects will be discussed in a later chapter in the context of interacting fields. At the end of this section we will give a proof of this theorem. It is important (e.g. for the derivation of the Feynman rules) to be able to Wick-order products of local composites of free fields. Let us look at examples:

$$: A^{4}(x) ::: A^{4}(y) ::=: A^{4}(x)A^{4}(y) :+ 4^{2}i\Delta^{(+)}(x-y) : A^{3}(x)A^{3}(y) :+ + 4^{2}3^{2} \left(i\Delta^{(+)}(x-y)\right)^{2} : A^{2}(x)A^{2}(y) :+ (4!)^{2} \left(i\Delta^{(+)}(x-y)\right)^{3} : A^{3}(x)A^{3}(y) :+ + (4!)^{2} \left(i\Delta^{(+)}(x-y)\right)^{4}$$

$$(3.141)$$

$$: \bar{\psi}(x)\gamma_{\mu}\psi(x):: \bar{\psi}(y)\gamma_{\nu}\psi(y)::::\bar{\psi}(x)\gamma_{\mu}\psi(x)\bar{\psi}(y)\gamma_{\nu}\psi(y): +:\bar{\psi}(x)\gamma_{\mu}iS^{(+)}(x-y)\gamma_{\nu}\psi(y): +Tr\left\{iS^{(-)}(y-x)\gamma_{\mu}:\psi(x)\bar{\psi}(y):\gamma_{\nu}\right\} + Tr\left\{iS^{(-)}(y-x)\gamma_{\mu}iS^{(+)}(x-y)\gamma_{\nu}\right\}$$
(3.142)

For a good understanding of the Wick-formalism of local functions a knowledge of the following statements is indispensable.

Statement 1: Powers of the two-point functions are well-defined distributions (singular functions), e.g. $F(x) = (i\Delta^{(+)}(x))^n$ is again a distribution with momentum space support properties. This is a multidimensional generalization of the well-known statement that singular functions F in one variable, whose Fourier transform \tilde{F} have support on the half line, can be freely multiplied. The reason is that (as a result of the support property) $i\Delta^{(+)}(x)$ is the boundary value on the real axis of an analytic function holomorphic in the upper half plane and therefore this property is inherited by its n^{th} power F. Equivalently the convolutions of \tilde{F} only extend over a compact region. In the multidimensional version the half lines are to be replaced by conic regions. In standard QFT momentum space correlation functions are well behaved functions, which at most have singularities at small momenta (infrared problems). Their asymptotic increase is responsible for the x-space singularities on the light cone.

Statement 2: The Noether conservation laws of classical field theory also hold for the corresponding Wick-orderd objects in the free field Borchers class.

We provide two typical illustrations involving a Dirac field ψ and a scalar field A:

$$\partial^{\mu} j_{\mu}(x) = 0, \quad j_{\mu}(x) =: \overline{\psi}(x)\gamma_{\mu}\psi(x): \quad \psi = \text{Dirac-field}$$

$$T_{\mu\nu}(x) =: \partial_{\mu}A(x)\partial_{\nu}A(x): -g_{\mu\nu}\frac{1}{2}: (\partial^{\kappa}A(x)\partial_{\kappa}A(x) - m^{2}A^{2}(x)): \quad (3.143)$$

$$\partial^{\mu}T_{\mu\nu}(x) = 0$$

As in the classical case the covariant divergence hits both of the fields and lead to operations on the u-and v-intertwiners which thanks to certain identities (e.g. the vanishing of the momentum space Dirac operator on these intertwiners) give the desired conservation law. In no stage of the argument does one need the canonical formalism or the Euler-Lagrange form of equation of motions, one only needs identities on intertwiners u and v which are an immediate consequence of their definition.

Statement 3: In the relation between local "currents" and global "charges":

$$Q = \int d^3x j_0(x), \quad P_\mu = \int d^3x T_{\mu 0}(x), \quad (3.144)$$

the phenomenon of vacuum polarization enforces a nonclassic subtlety which is explained in the following.

A composite of a free field is more singular than the free field. In particular for $d \geq 2 + 1$ it does not fit into the framework of canonical equal time (anti)commutation relation, but rather has to be smeared with test functions in d dimensions (in our case d = 3 + 1). This can already be seen by using the previously calculated two-point function of the composite current operator j , e.g.

$$\langle j_{\mu}(x)j_{\nu}(y)\rangle = \int (g_{\mu\nu}\partial^2 - \partial_{\mu}\partial_{\nu})i\Delta^{(+)}(x-y,\kappa^2)\rho(\kappa^2)d\kappa^2$$
(3.145)

Since $\int \rho(\kappa^2) d\kappa^2 = \infty$, the smearing with test functions supported on a spacelike hypersurface i.e. of the form $f(x) = \hat{f}(\vec{x})\delta(t)$ does not give a finite answer, one rather needs smoothness in time as well. As in the classical case, one tries to

obtain the global charge $Q = \int (a^*(p)a(p) - b^*(p)b(p))\frac{d^3p}{2\omega}$ as a limit of "partial" charges referring to a finite region:

$$\begin{aligned} Q(g.h) &:= \int j_0(x) g(\vec{x}) h(t) d^4 x, \quad \text{sup p g} \ \subseteq V + \delta V, \ \text{supp h} \subseteq \{ |t| \le \epsilon \} \\ g \equiv 1 \text{ in 3-volum V}, \quad \int h dt = 1 \end{aligned}$$

(3.146) V ---hish haa

In words: g is a characteristic function of the 3-dim. volume region V which has been smoothened outside, whereas h(t) is a smoothened δ -function. It is easy to see that:

$$[Q(g,h),B] = [Q,B], \text{ for locB in completion of V}$$
(3.147)

i.e. for operators B localized in the causal completion of V (smearing functions with support in completion of V) the commutator is already independent of g,h (and identical to the global charge). However on the vacuum vector Ω the partial charge has such strong vacuum fluctuations (resulting from the presence of $a^* - b^*$ terms) that:

$$\lim_{V \to \infty} ||Q(g,h)\Omega||^2 = \infty, \quad \text{but } \lim_{V \to \infty} (\psi, Q(g,h)\Omega) = 0$$
(3.148)

Here ψ is from the dense domain on which the local functions of the free field are defined i.e. the polynomial domain. The vacuum fluctuations were discovered in the early days of QFT by Heisenberg and their physical significance was studied by Weisskopf. One such manifestation is a contribution to the Lamb-shift (see next chapter). This quantum phenomenon has no counterpart in quantum mechanics and it has far going structural consequences, e.g. it makes the local algebras of QFT essentially different from the quantum mechanical Heisenberg-Weyl algebras (the former admit no pure states or minimal projectors).

We now indicate the proof of the previous theorem on the structure of the free Borchers class [18]. We start with the general Wick expansion of a translation covariant field B(x) in terms of Wick products of the free field A. The first step consists in "peeling off" iteratively the lower Wick monomials so that the new relatively local field say $B_{m+n}(x)$ after n+m steps starts with Wick monomials of degree n+m and higher i.e.

$$\langle \Omega \mid: A(y_1)...A(y_{m'}) : B_{m+n}(0) : A(x_1)...A(x_{n'}) : \mid \Omega \rangle$$

$$= \langle \Omega \mid A(y_1)...A(y_{m'})B_{m+n}(0)A(x_1)...A(x_{n'}) \mid \Omega \rangle$$

$$(3.149)$$

for $m' + n' \ge m + n$ and all matrix elements ("formfactors") of B_{n+m} with m' + n' < m + n vanish. We start the induction with $B_0(0) \equiv B(0)$ and first subtract the constant vacuum expectation of B in order to arrive at a $B_1(0)$ in Bo(A). Since the various monomials do not mix if commuted with the free field A(x), the commutator of the first Wick degree contribution $B_1^{(1)}(0)$ in $B_1(0)$ with the free field A(x) yields a c-number which (as a result of causality) must be proportional to $i\Delta(x)$, possibly involving derivatives. But this means that $B_1^{(1)}(0) = (P(\partial)A)(0)$. We then subtract this local one-particle contribution

 $B_1^{(1)}$ and obtain B_2 and $B_2^{(2)}$ etc. The induction consists in showing, that if $B_{i-1}^{(i-1)}$ is a local Wick monomial (possibly including derivatives of) of degree i-1 in A, then $B_i(0) \in \operatorname{Bo}(A)$. We then show that the lowest term $B_i^{(i)}(0)$ has local formfactors (particle matrix elements in the A-particle basis between n, m-particle vectors) identical to those of a i=(n+m) degree Wick polynomial. This follows from the structure of the multiple *i*-fold commutator $K^{(i)}$ which fulfills a Klein-Gordon equation in every free field coordinate. Therefore K obeys a multi Cauchy initial value representation on the $t_j=0$ space-like hypersurface (3.149):

$$K(\underline{x},\underline{y}) := \left\langle \left| \left[\dots \left[\left[\dots \left[B_{n+m}^{(i)}(0), A(x_1) \right], \dots, A(x_n) \right], A(y_1) \right] \dots, A(y_m) \right] \right| \right\rangle$$

which is just a linear combination of (n + m)-degree formfactors. The Cauchy initial value problem representation for the free fields A(.) leads to the following structure:

$$K(\underline{x},\underline{y}) = \prod_{i \le n, j \le m} \int \dots \int \Delta(x_i - x'_i) \Delta(y_j - y'_j) \overleftrightarrow{\partial'_i} \overleftrightarrow{\partial'_j} f(\underline{x}',\underline{y}') \mid_{\substack{x'_0 = x_0 \\ y'_0 = y_0}} d^3 \underline{x} d^3 \underline{y}$$
(3.150)

Since the initial values f on the equal time hypersurface is a product of δ -functions and derivatives (here the causality of K is used), K is identical to the multiple commutator of a local Wick monomial $B_i^{loc} = mon.^{(i)}(A)$ i.e. we have the formal representation:

$$B(x) = \sum_{i} B_i^{loc}(x) \tag{3.151}$$

Since local fields are operator-valued Schwartz (tempered) distributions with power singularities on the light cone for $d \ge 1 + 2$, the series must be finite and B is a Wick polynomial Q.E.D..

It is an interesting question whether this result still holds if one does not know that B(x) is relatively local to the free field A(x), but only has an information about the absence of interaction in the sense of a trivial scattering operator $S_B = 1$. In this case one would take for A the free incoming field provided by scattering theory. Although there is no direct information from locality, the information provided by analytic properties of p-space formfactors of B between incoming ket and outgoing bra state vectors (see chapter 6) leads to the same result (3.151). In the present framework of equivalence classes this means that a "weakly local" equivalence class consisting of all fields with the same S-matrix in the special case S = 1 contains only one local Borchers class, namely. the standard free field class. This suggests that the S-matrix is a very precise indicator of interactions. It also hints at the existence of a general unique inverse scattering structure of Local Quantum Physics.

3.5 A First Look at Modular Localization

Recently it turned out that the Wigner representation theory contains information on localization which allows a direct access to the local algebras, thus avoiding the use uf nonunique field coordinates [2]. The starting point is the abelian subgroup of Lorentz boosts belonging to a wedge, say the standard t-x wedge x > |t|. Note that wedges can be also characterized in terms of two light rays which for the standard wedge are $e_{\pm} \sim (1, \pm 1, 0, 0)$. The Wigner theory also provides an anti-unitary operator which reflects the standard wedge into its opposite wedge. In the simplest case of irreducible representation for scalar neutral particles, this reflection j differs from the TCP operation by a π -rotation around the x-axis:

$$(\Theta\varphi)(p) = \bar{\varphi}(p) \quad (j\varphi)(p) = \bar{\varphi}(p_0, p_1, -p_2, -p_3) \tag{3.152}$$

Define now an unbounded positive closed operator δ by functional calculus from the selfadjoint standard (x - t) boost generator K:

$$\delta = e^{-K}, \quad \delta j = j\delta^{-1}, \quad \text{since } e^{iK\chi}j = je^{iK\chi} \tag{3.153}$$

With the help of the Tomita-like unbounded involutive operator $\mathfrak{s} := j\delta^{\frac{1}{2}}$ we define a closed "real" subspace H_R of the Wigner representation space H:

$$H_R = \{\varphi(p) \in H \mid \mathfrak{s}\varphi = -\varphi\}, \quad \mathfrak{s} = j\delta^{\frac{1}{2}}$$
(3.154)

The \pm eigenspaces (since \mathfrak{s} is antilinear, only real linear combinations are possible) of the closed operator \mathfrak{s} can easily be shown to form a dense set in H and the above definition is also the unique polar decomposition of \mathfrak{s} . To be more specific, \mathfrak{s} acts as:

$$\mathfrak{s}: \quad h+ik \to -h+ik, \quad h,k \in H_R \tag{3.155}$$

Here $H_R + iH_R$ is a dense subspace of the Wigner space (it is only dense, even though H_R is closed). It is the modular localization space for the standard wedge W^{sta} . Using the standard mathematical trick of introducing the graph norm affiliated with $\Delta^{\frac{1}{2}}$, this dense space becomes a Hilbert space $H_{\rm mod}$ in its own right. As we will see in the next section, the Wigner inner product restricted to the modular localization space can be rewritten as a (Hawking) thermal inner product in this new Hilbert space $H_{\rm mod}$ associated with the modular localization.

A more explicit description of H_R is obtained by introducing the wedgeaffiliated "rapidity" θ :

$$p = m(q \cosh\theta, q \sinh\theta, n_2, n_3), \quad q = \sqrt{1 + n_2^2 + n_3^2}$$
 (3.156)

The domain of the operator $\Delta^{\frac{1}{2}}$ (and hence of \mathfrak{s}) in terms of rapidity-dependent wave functions consists of boundary values of analytic functions which are holomorphic in the θ -strip $0 < \theta < i\pi$ and H_R , the real subspace of \mathfrak{s} with -1 eigenvalue, is the closed real space of wave functions which are analytic in the strip and fullfil the boundary condition:

$$\overline{\varphi(\theta + i\pi)} = -\varphi(\theta) \tag{3.157}$$

where we suppressed the dependence on n_i . Let us call this the "s-reality property". It is somewhat surprising that this concept did not seem to have appeared in mathematical physics, e.g. it is absent from the various books (including those by Reed and Simon).

For massive spin s representations the *s*- reality property reads:

$$D^{(s)}(i\sigma_2)lim_{\chi \to i\pi} D^{(s)}(R(p,\Lambda^{sta}(\chi))\overline{\varphi(\theta+i\pi)} = -\varphi(\theta)$$
(3.158)

If particles are not selfconjugate, the 2s+1 component φ must be doubled and the action of J on the direct sum involves a flip-operation on the two Hilbert spaces. For zero mass, the rapidity parametrization for the standard wedge is defined by $k = \Lambda^{sta}(\theta)k_0$ with $k_0 = (1, \vec{n})$ and the Wigner rotation $R(k, \Lambda)$ is to be replaced by the helicity representation in terms of the Wigner phase factor of the euclidean group E(2).

Looking at the geometric interpretation of this construction, one conjectures that the subspace H_R of these momentum space wave functions has something to do with localization in the standard wedge (or in the opposite wedge in case of the -subspace). In fact the intertwiner formalism of the previous sections allows to write H_R (or equivalently iH_R) in terms of vector valued W-supported testfunctions f i.e. the 2s+1 component wave function:

$$u^*(p)f(p)\mid_{p\in m.h.}$$

where u^* is u written as a row vector with complex conjugate entries and the subscript indicates the restriction of p to the forward mass shell. Using the analyticity properties of \tilde{f} which result from the wedge support and the charge conjugation properties of u, the eigenvalue equation for \mathfrak{s} can be checked easily. The wedge localization can be confirmed without the intertwiner formalism (the intertwiners are not available in all positive energy representations, see later) by studying coherence properties of the net of real wedge spaces generated via Poincaré transformations g on the standard wedge :

$$H_R(W) \equiv U(g)H_R, \quad W = gW^{sta}$$

$$H_R = H_R(W^{sta})$$
(3.159)

For localization in the quantum sense, one needs a concept of "outside". In Schrödinger theory as well as in the relativistic work of Newton and Wigner, one uses the orthogonality in wave function space: one calls f localized in a 3-dim. region R if a spatial translation which carries R into its geometric complement transforms the wave function into the orthogonal complement. This is of course the "Born-localization" based on Born's probability interpretation of Schrödinger wave functions at a fixed time. It incorporates the fluctuation (uncertainty relation) aspect of states in QT and leads to finite extension for bound states i.e. to a distinction elementary versus bound.. For relativistic wave functions this idea unfortunately (much to the dismay of Wigner) cannot be extended from equal time localization to spacelike localization (apart from localization in an "effective" sense i.e. modulo Compton tails). Fortunately for relativistic local QFT there exists another more geometric notion of localization⁵. It is that one which underlies the pointlike fields of standard QFT. Its transcription to wave functions is related to the following notion of symplectic "orthogonality" (duality):

$$H'_{R} = \{h' \mid Im(h', h) = 0\}, \quad H'_{R}(W) = U(g)H'_{R}$$
(3.160)

It then follows that:

 $H'_{R}(W) = H_{R}(W)', \text{ and } H'_{R}(W) = H_{R}(W'), W' \equiv W^{opp}$ (3.161)

where the last nontrivial equality is a consequence of:

$$\mathfrak{s}_W^* = \mathfrak{s}_{W'} \quad \longleftrightarrow \quad j_W = j_{W'} \ , \ \ \delta_W^{-1} = \delta_{W'} \tag{3.162}$$

which in turn follows from the commutation relation of the standard (x-t) reflection j (which sends the wedge W into W^{opp}) and the Lorentz-boost δ^{it} . Again one ends up with real Hilbert spaces which are standard and factorial in the sense of chapter 2, section 6.

Thus we arrived at a covariant net of wedge spaces and now we want to show that this net is isotonous i.e. that if a wedge is contained in another one, the same is true for the associated spaces. But in such a situation the second wedge is obtained from the first by two lightlike translations which carry it inside, so we have to show isotony for lightlike translations. For such translations we have:

$$\mathfrak{s}_{\hat{W}} \subset \mathfrak{s}_W, \quad \hat{W} = g(\lambda l) W \subset W, \quad \lambda > 0$$

$$(3.163)$$

where $g(\lambda l)$ is a translation along the lightlike vector l. In order to show that \mathfrak{s}_W extends $\mathfrak{s}_{\hat{W}}$ we rewrite this relation as :

$$U(\lambda l)j_W \delta_W^{\frac{1}{2}} U(\lambda l)^* \subset j_W \delta_W^{\frac{1}{2}}$$
(3.164)

For the bounded antilinear operator j_W this gives the covariance law, whereas for unbounded δ the required relation results from the commutation relation of the lightlike translation with the standard Lorentz-boost $U(\chi)$:

$$U(\lambda l)U(\chi) = U(\chi)U(e^{\chi}\lambda l)$$
(3.165)

One can show that the isotony is quite generally equivalent to the positivity of the energy.

⁵Even this more "algebraic" localization does not always agree with the geometric one. The vector potential for the (m = 0, s = 1) representation and the continuous spin representation are free field examples for such cases, not to mention d = 2 + 1 anyons.

Wedge localization is too weak for a physical interpretation of the theory (e.g. for the derivation of statistics and scattering theory). The localization underlying standard (e.g. Lagrangian) theory is compact localization which in our context means (K stands for double cone):

$$H_R(K) = \bigcap_{W \supset K} H_R(W) \tag{3.166}$$

$$H_R(K) + iH_R(K)$$
 dense in H , $H_R(K) \cap iH_R(R) = \{0\}$ (3.167)

 $H_R(K) \cap H'_R(K) = \{0\}$

This property which previously (in section 2.6) was called standard and factorial can be shown for all Wigner (m, s) representations and even for m = 0 (with the exception of the continuous spin representations, which do not permit such a localization, since in this case the $H_R(K)$ spaces turn out to be trivial). As a consequence the spaces fulfill the following duality property:

$$H'_R(W) = H_R(W'), \quad H'_R(K) = H_R(K')$$
 (3.168)

In case of (m, s)-representations one can prove this even for disconnected and non simply connected regions in Minkowski-space.

The abstract CCR functor introduced in the previous chapter converts the real localization subspaces directly (i.e. without the interference of pointlike fields) into a net of von Neumann algebras in Fock space. This and the analogous statement for CAR is the only precise meaning of the word second "quantization". Therefore the quantization approach to relativistic QFT is limited to these cases and their formal Lagrangian perturbation which also happen to be those to which differential geometric concepts as fibre bundles are applicable. Beyond the word "quantization" has an intuitive artistic connotation. But this territory beyond these functors and their perturbative Lagrangian extension is precisely the region where the algebraic approach takes over. There is however one important message in these localization functors which will be studied in the sequel.

With these remarks we have entered the central issue of these notes: a formulation of QFT which is independent of the choice of "field co-ordinates" and refers directly to the map between localization regions in Minkowski-space and observable algebras. Among the myriads of pointlike fields there is of course no complete democracy. E.g. Noether currents are physically distinguished. On the observational side it appears that all prominent measured quantities can be represented in terms of matrix elements of such currents. It is deeply gratifying that algebraic QFT attributes a special role to such currents (via the "split property" see[3]). The rich physical harvest resulting from this new point of view outweighs by far our present modest motivation in this section which was to reconcile the multitudes of free fields with the uniqueess of the positive energy Wigner representations. We hope to be able to convince the reader about this in the rest of these notes.

It is interesting and in the spirit of this course to note that the global spacetime symmetry can be encoded into the inclusions and intersections of modular localization subspaces. Let us illustrate this by two examples.

- Suppose we shift the standard wedge W_{stan} into itself by applying a light like translation parallel to its longitudinal light ray. This gives a "halfsided modular inclusion" with one of the boundary surfaces of the shifted wedge W_{shif} lying on the corresponding boundary of the standard wedge. We note that the Lorentz transformation associated with W_{stan} compresses W_{shif} into itself if we choose the sign of the Lorentz rapidity χ correctly. Since the modular group is equal to the appropriately parametrized Lorentz boost this justifies the terminology halfsided modular inclusion: $\delta_W^{\pm it} H_R(W_{shif}) \subset H_R(W_{shif})$ is $\pm hs$. In fact a simple computation inside the Poincaré group shows that $\frac{1}{2\pi}(ln\delta_{W_{shif}})$ $ln\delta_{W_{stan}}$) is the positive spectral generator of the unitary light cone translation. The converse is also true. If we start from a say a +hs modular inclusion of real Hilbert spaces $H_1 \subset H_2$, we note that $\delta_2^{\frac{1}{2}} \leq \delta_1^{\frac{1}{2}}$ and assuming that is essentially selfadjoint on $D(\ln\delta_1) \cap D(\delta_2)$ we may apply the Trotter product formula $U(a) = expia(ln\delta_1 - ln\delta_2) = s - lim_{n\to\infty}(\delta_2^{\frac{-ia}{n}}\delta_1^{\frac{+ia}{n}})^n$. The last step namely to prove that in this abstract situation the so defined U(a) together with the δ_2^{it} and the modular reflection j define a 2-parametric group with $\delta^{it} U(a) \delta^{-i\tilde{t}} = U(e^{-2\pi t}a)$ and jU(a)j = U(-a), requires a bit more of the analytic techniques of Borchers [7].
- By intersecting the standard wedge with another L-transformed wedge W which has one light ray l in common with W_{stan} (and not a hypersurface as above), one finds by computations within the Poincaré group that the intersection $W_{stan} \cap W$ is halfsided modular included in W_{stan} as well as in W. This leads again to a positive translation as above however this translation is not a geometric translation inside the Poincare group; a reflection of the fact that the modular group associated to the intersection $W_{stan} \cap W$ is not geometric. But this time the W's are not included and the corresponding "translation" (the boost transformation transformations remain inside the homogeneous Lorentz group) in the above formula $(\ln \delta_W^{\frac{1}{2}} - \ln \delta_{W_{sta}}^{\frac{1}{2}})$ has no positive generator (it is a difference of the two previous nongeometric translations in which the unknown nongeometric modular operator of $W_{stan} \cap W$) cancel out). In fact these transformations are easily identified as one of the translations of the previously encountered little group E(2). The other one results from a modular intersection in which the other light ray of W_{stan} is shared with a second wedge. The two "translations" are precisely the afore mentioned transversal Galilei generators G^{\pm} .
- In fact the intimate relation between modular inclusions and intersections with spacetime symmetries poses the question whether the full Poincaré group may not be encoded ito a finite number of modular data. This is indeed the case[8], but the better context for explaining such results is the later presented general framework of algebraic QFT.

3.6 Special Features of Zero Mass

The above duality situation continues to be valid if one replaces K by non simply connected regions, but not so for the $s \ge 1$ zero mass representations. For example in the case of photons (m = 0, h = 1), one finds a duality violation for the toroidal "corona" region. Let T be the causal completion of a spatial torus which we call the "corona". A convenient mathematical description is to start from a unit double cone in the longitudinal z-t plane centered at $\rho = \sqrt{x^2 + y^2} = a > 1, z = 0, t = 0$ and rotating it in the transversal x-y- plane once around. The causal complement of \mathcal{T} is causally multiply connected. Then one obtains the following proper corona- inclusion:

$$H^{R}(\mathcal{T}) \subset H^{R}(\mathcal{T}')'$$

$$(3.169)$$

$$\sim \mathcal{A}(\mathcal{T}) \subset \mathcal{A}(\mathcal{T}')'$$

where $H^R(\cdot)'$ denotes the previously defined symplectic complement and the A's denote the corresponding von Neumann-algebras as obtained from the $H^R(\cdot)$ by the Weyl construction.

This obstruction (formally related to the appearance of δ' in the electricmagnetic canonical commutation relation) can be physically understood in terms of a suitably regularized magnetic flux through a surface which stretches from a circle inside the torus into the space-like separated region inside. This flux does not change if one passes through another spacelike surface subtended from the same circle. Hence such a flux, although not being localizable within the 4-dim. toroidal region nevertheless belongs to the symplectic complement of the spacelike complement of the corona which is multiply spacelike connected. Via the CCR functor this entails the above violation of Haag duality for the corresponding algebras.

The issue of modular localization of (m=0,h=0) representations can also be discussed in the A_{μ} description (3.90). We can remove the nullmodes explicitely in a fixed Lorentz frame by introducing a directional dependent potential $A_{\mu}(x,n)$ by the following formulas:

$$\tilde{A}_{\mu}(k,n,\pm) = \frac{n^{\nu} \tilde{F}_{\nu\mu}(k,\pm)}{n \cdot k + i\varepsilon}, \quad \tilde{F}_{\nu\mu}(k,\pm) = k_{\{\nu} \varepsilon_{\mu\}}(k,\pm) \psi(k,\pm)$$

$$\sim \int |\psi(k,\pm)|^2 \frac{d^3k}{2\omega} = \int \overline{\tilde{A}}_{\mu}(k,n,\pm) \tilde{A}^{\mu}(k,n,\pm) \frac{d^3k}{2\omega}$$

$$\omega = \left|\vec{k}\right| \tag{3.170}$$

where wavy brackets denote the antisymmetrization in μ, ν and $\varepsilon_{\mu}(k, \pm)$ are the polarization vectors and the last formula results from (3.130). The singularity in k-space corresponds to the semiinfinite line integral along n in x-space.

$$A_{\mu}(x,n) = \int_{0}^{\infty} n^{\nu} \tilde{F}_{\nu\mu}(x-ns) ds \qquad (3.171)$$
$$= \int (e^{-ikx} \sum_{i=\pm} \tilde{A}_{\mu}(k,n,i) + h.c.) \frac{d^{3}k}{2\omega}$$

This vector potential has the following obvious properties under the Wigner representation $U(\Lambda)$:

$$\partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu} = F_{\mu\nu}$$

$$(U(\Lambda)A)_{\mu}(x,n) = \Lambda^{\nu}_{\mu}A_{\nu}(\Lambda^{-1}x,n')$$

$$= \Lambda^{\nu}_{\mu}A_{\nu}(\Lambda^{-1}x,n) + \partial_{\mu}G(x)$$

$$G(x) = \lim_{\varepsilon \searrow 0} \int e^{ikx} \frac{1}{(kn - i\varepsilon)(kn' - i\varepsilon)} n \cdot \tilde{F}(\Lambda^{-1}k) \cdot n' \frac{d^{3}k}{2\omega}$$

i.e. the Lorentz transformation which acts on the Wigner wave function resp. on the $F_{\mu\nu}$ tensor, transforms the potential covariantly except for an additive nonlocal "gauge" term i.e. adds a nullmode contribution. The nonlocality and the lack of covariance of this description in terms of an affine transformation law is manifest. This peculiar "gauge" behavior is a consequence of the nonfaithful helicity representation of the noncompact little group E(2). In particular as we already emphasized in section 2 and 3 of this chapter, the quantum origin of gauge, and gauge invariance in the sense of quantization of classical gauge theories are somewhat different. Unlike in classical theory we either must work with noncovariant vector potentials or (if we work with transverse vector wave functions) or abandon the idea of pointlike transverse quantum vectorpotentials.

In the presence of interactions one presently does not know a formulation which uses the ghostfree semiinfinite stringlike localized Wigner potential $A_{\mu}(x, n)$. All the perturbative formulations of gauge theories use covariant vectorpotentials which involve ghosts. We will return to this issue in chapter 4. Here we only note that the nonlocal and noncovariant behavior causes no problems if we could build up the net of localized subspaces from wedges. The reason is that for spacelike vector n in the standard x-t wedge plane, the vector potentials corresponding to wedge supported field strength F are also wedge supported since the semiinfinite spacelike line stays inside W_{st} :

$$A_{\mu}(x,n) \in H_R(W_{st}) + iH_R(W_{st}), \ if \ support \ F \subset W_{st}$$

$$(3.173)$$

The vector potentials are however not belonging to the simply connected compact modular localization spaces (e.g. double cones) which result from intersections except for closed spacelike contour integrals over A which loose their semiinfinite extension and can be expressed in terms of field strength inside that localization region at least as long as the region stays simply connected. The aforementioned peculiar behavior of the non-simply connected corona region expressed in terms of vector potentials means that the difference in the geometrical versus the quantum localization i.e. the obstruction against Haag duality for nonsimply connected regions can be accounted for in terms of a circular line integral over the vector potential around the corona which looses its noncompact tail. Note that in the covariant indefinite metric description the corona situation behaves like for massive vector mesons i.e. there is no duality obstruction. The duality obstruction only reappears after getting rid of ghosts. Hence the mathematical locality and formal duality is another indication of the unphysical nature of the "ghostly" description which only makes sense as an auxilary intermediate bookkeeping.

In passing we remark that different from the massive case, there are many positive energy nonvacuum representations of free photons. The appearance of such non Fock "infrared representations" is characteristic for zero mass. They show up in a richer representation theory of the associated Weyl algebra. What we previously called Weyl functor should now be called the Fock-Weyl functor. As before, the latter carries the net of real Hilbert spaces into the vacuum observable net, but for zero mass there are other functors from the same spaces to non Fock space algebras. Whereas free massive nets have a unique positive energy representation, free zero mass nets have inequivalent "infravacua" [88]. They also possess electrically or magnetically charged "infraparticle" states.

Note that there is no problem in formulating the Weyl algebra based on the vector potential description since only the imaginary part (symplectic form) and not the full inner product is used in its definition. It just does not possess a Fock representation in which the Lorentz group automorphisms of the Weyl C^{*}-algebra possess a unitary representation. It is not clear if in this description there are physically relevant infrared representations which activate the zero "modes" i.e. the longitudinal components of the vector potential.

Let us briefly return to a formal aspects of the corona obstruction. A simple way to restore the harmony between the geometrical (classical) and the quantum(modular) localization is to extend the corona localization space by one type of object:

$$\oint_{C \subset \mathcal{T}} A_{\mu}^{reg}(x, n) dx^{\mu}$$

$$A_{\mu}^{reg}(x, n) = \int \rho(\vec{x} - \vec{y}) A_{\mu}(\vec{y}, x_0, n) d^3y$$
(3.174)

This explains why the defect dimension of the two real Hilbert spaces in (3.169) is:

$$\dim\left[H^{R}(\mathcal{T}')':H^{R}(\mathcal{T})\right] = 1 \tag{3.175}$$

The integration in the first formula is over a closed path C inside \mathcal{T} and we regularized the vector potential with a smooth function of small support $supp \rho \in B_{\varepsilon}$ so that one maintains the normalizability of the wave function and remains inside \mathcal{T} . The line integral represents the class of expressions of this kind, any two such elements differ only by field strength localized in \mathcal{T} . The line integral is a Linvariant and may be expressed in terms of a magnetic flux through any surface S with the C boundary. It is precisely this floating surface stretching beyond C, which in the quantum setting of commutativity (or symplectic orthogonality) prevents the affiliation with $H^{R}(T)$ and makes it a member of the nongeometric $H^{R}(T')'$.

It is a much more difficult question as to what becomes of this topological obstruction in the presence of interactions. It is tempting to interpret this
obstruction as indicating the necessity of an interaction 6 i.e. of the presence of non-vanishing electric or magnetic (or both) currents.

$$\partial^{\mu}F_{\mu\nu}(x) = j_{\nu}(x), \ \partial^{\mu}\tilde{F}_{\mu\nu}(x) = \tilde{j}_{\nu}(x)$$
 (3.176)

One idea is that interactions are necessary to *restore perfect Haag duality* (i.e. for all multiple connected regions) which is violated in the free theory. Such a point of view would attribute a very distinguished role to electromagnetic duality and link it to Haag duality on a very fundamental level. But lack of nonperturbative insight prevents a clear-cut resolution of this duality connection. This is of course related to the non-understood problematizing the notion of "magnetic field" on the same level of depth as the notion of "charge" in the DHR superselection theory.

In low dimensional QFT the analogous issue of order-disorder duality and the connection with Haag duality is much better understood. There, even in free theories (see last section of this chapter), it is not possible to have *no* charge sectors with both order and disorder. In the massive case one charge comes out and the presence of both charges only occurs in d=1+1 in the zero mass limit. The previous idea of maintaining corona duality would enforce interactions and bring the interacting Maxwell-like theories closer to the 2-dim. situation. A good understanding of modular localization aspects of electromagnetic interactions (in the vein of the remarks about interactions in chapter 6) seems to be essential for future progress.

As mentioned before, the corona inclusion may be constructed solely in terms of the Wigner theory supplemented by the modular theory for wedges which works with subspaces rather than with pointlike covariant amplitudes like $F_{\mu\nu}$. Helicities $h \geq 1$ present similar gauge problems and corona obstructions as a result of the zero mass E(2) little group. As in the electromagnetic case, these duality obstructions are inexorably liked with the gauge aspects of these massless theories. In the quantization approach of the text books, this requires the introduction of ghosts and the use of the BRS formalism including the cohomological control of the physical factor space.

There remains in Wigner's list the positive energy zero mass representations with so called "continuous helicity" which require infinite component momentum space wave function. They are usually dismissed by saying that "they are not realized in nature". On the theoretical side there still remains the question of whether their much weaker modular localization is the theoretical reason why they may not behave as genuine particles. Irreducibility (indecomposibility) and positive energy of the representation of (space-time) symmetry is the only prerequisite for particles. Already Wigner knew that this was insufficient and that one needs an appropriate relativistic localization. But how far beyond wedge localization can one go? It turns out that the localization properties are rather similar to those of d=2+1 anyons i.e. the best possible modular localization.

⁶This speculative remark is taken from [3], page 147.

Furthermore this analogy also suggests that the corresponding "free field theory" does not have the Fock space structure. More comments will be presented in the last chapter.

In such a situation one must first understand the physical consequences before one rushes to the above dismissal. After all there are strange particlelike objects as quarks which certainly cannot be identified with the standard massive and massless Wigner particles.

Let us make some more remarks about the massive case. The principle of locality requires to study intersection of wedges. Intersecting the translated standard wedge W_{a}^{st} with the opposite wedge W_{opp}^{st} leads to an x - t double cone which is cylindrically extended in the y - z direction. Since the modular localization in W_{opp}^{st} corresponds to a s-reality condition in the negative θ -strip, the intersection of both gives rise to a new "edge of the wedge" problem i.e. a Hilbert space $H_R(W_a^{st} \cap W_{opp}^{st})$ of analytic functions which are meromorphic in both strips and fulfil a matching condition on the real θ -axis in which the translation enters. Again $H_R(W_a^{st} \cap W_{opp}^{st})$ is standard in the sense of the definition given in the section on CCR and CAR functors.

The analytic situation for intersections of non coplanar wedges as one needs them for double cones in (3.166) becomes very rich and is essentially unexplored. In d=3+1 theories with halfinteger spin QFT of free fields indirectly yield the information that the corresponding real subspaces are standard and factorial.

If we apply this localization concepts to halfinteger spin, we find a very interesting discrepancy by a factor i between the action of j and that of the π -rotation of the wedge caused by the SU(2) transformation law of the spin. Through this obstruction the Wigner theory already takes notice of the spin-statistics.

We now explain the direct conversion of the net of Wigner subspaces into a net of CCR- and CAR-algebras using the functorial formalism in section 6, chapter 2.

Consider first the case of integral spin. The application of the Weyl-functor to the subspace $H_R(W)$ gives the von Neumann-algebra:

$$\mathcal{F}: H \to \mathcal{A}(H), \quad f \longmapsto W(f)$$
 (3.177)

$$\mathcal{A}(W) = \text{v.Neumann Alg.} \{W(f) | f \in H_R\}$$

$$= \mathcal{F}(H_R(W))$$
(3.178)

which inherits the following properties from the Hilbert spaces:

isotony :
$$\mathcal{A}(W) \subset \mathcal{A}(\tilde{W})$$
, for $W \subset \tilde{W}$ (3.179)
Haag duality : $\mathcal{A}(W') = \mathcal{A}(W)'$, $W' = W^{opp.}$
covariance : $U(g)\mathcal{A}(W)U^*(g) = \mathcal{A}(gW)$, $g \in \mathcal{P}$

In the halfinteger spin case we take the CAR functor $\psi^{\#}(f)$:

$$\mathcal{F}: H \to \mathcal{A}(H), \quad f \to \psi(f) \in B(\mathcal{H}_F) = \mathcal{A}(H)$$
 (3.180)

$$\mathcal{A}(W) = \text{v.Neumann Alg.} \left\{ \psi^{\#}(f) \mid f \in H_R(W) \right\}$$
(3.181)
$$= \mathcal{F}(H_R(W))$$

Different from the bosonic case, the operators J and S of this algebra are not given by the application of the previous functor \mathcal{F} but the J contains the famous Klein twist K which changes geometrically causally disjoint localized (hence anticommuting) operators into commuting ones which one needs in the Tomita-Takesaki theory:

$$J = K\mathcal{F}(j), \quad \Delta^{it} = \mathcal{F}(\delta^{it}), \quad S = J\Delta^{\frac{1}{2}}$$
(3.182)

This is the T-T modular theory for wedge subalgebras of the CAR-algebra.

The same modular formalism can be used in order to construct relativistic KMS states on free field algebras. In complete analogy to chapter 2.5, the thermal two-point functions have the form $(z=e^{\beta\mu})$:

$$\begin{split} \left\langle \psi(x)\psi(y)\right\rangle_{e,\beta,\mu} &= \\ & \frac{1}{(2\pi)^{\frac{3}{2}}}\int e^{-ip(x-y)}\sum_{s_3}u(p,s_3)\bar{u}(p,s_3)\frac{1}{1\mp ze^{-\beta p\cdot e}}\frac{d^3p}{2\omega}3) \\ & +\frac{1}{(2\pi)^{\frac{3}{2}}}\int e^{ip(x-y)}\sum_{s_3}v(p,s_3)\bar{v}(p,s_3)\frac{ze^{-\beta p\cdot e}}{1\mp ze^{-\beta p\cdot e}}\frac{d^3p}{2\omega} \end{split}$$

Here e is a time-like vector which characterizes the rest frame of the heat bath and the \mp sign corresponds to Boson/Fermion statistics. Mistakes in the combinations of signs in front of the integrals can be easily corrected by remembering that the thermal correlation functions must have the same (anti)commutator functions as the standard free field correlation functions (in addition to the KMS property), i.e. the thermal aspect is an attribute of the state and not of the algebra. These relativistic correlation functions have rather interesting analytic properties [51]; they are analytic in $x - y =: \xi \to z$, $z \in \mathcal{T}_{\beta e}$ where $\mathcal{T}_{\beta e}$ is the tube $\mathcal{T}_{\beta e} = \{z \in C^d : Imz \in V_+ \cap (\beta e + V_-)\}$. The boundary values at the two edges fulfill as expected the KMS condition:

$$\exists F(z) \text{ analytic in } \mathcal{T}_{\beta e} \text{ s.t.}$$

$$\langle \psi(x)\bar{\psi}(y) \rangle = \lim_{z \to \xi} F(z), \ \langle \bar{\psi}(y)\psi(x) \rangle = \lim_{z \to \xi + i\beta e} F(z)$$

$$(3.184)$$

where the boundary values are taken from inside the analytic tube region. All the statements are easily checked by explicit computations. Although the boundary KMS condition is the standard one which relates the boundary values on the two sides of the temperature strip, the relativistic aspect generates a larger analytic tube in x-space which contains the strip in the *e*-direction. The temperature can be directly introduced as an extension of the Wigner theory. It should be interesting to combine the modular localization aspect with the heat bath temperature within the Wigner setting. It is very interesting to note that the net of modular localization subspaces and its modular automorphisms can be constructed from a finite (with the size dependent on the space-time dimensions d) generating skeleton of appropriately positioned ("half-sided modular") real subspaces. This is the analogue of a similar structure of nets of observables which developed from its beginning in chiral conformal QFT in the form of the "quarter circle situation" of the author [60], via the more general discussion of half-sided modular inclusions [85], to the mathematically rigorous work of Araki and Zsido [86]. In the present modular subspace version which is adapted to the Wigner representation theory, the mathematics is much simpler and very illustrative. We will refrain here from a presentation. In particular the nonexistence of a generalization of the CCR and CAR functors to d=2+1 anyonic Wigner spin and to d=3+1 "continuous spin" Wigner representations is already visible in the attempt to go beyond the net of modular wedge localized Wigner subspaces (chapter 6).

It is very important to distinguish between the localized subspaces of the Wigner representation space and localizes subspaces in Fock space. The latter are not the image of the former under the CCR (CAR) functor $H(\mathcal{O}) \to e^{H(\mathcal{O})} = \mathcal{H}(H(\mathcal{O}))$ but one rather finds a genuine inclusion $\mathcal{H}(\mathcal{O}) \subset \mathcal{H}(H(\mathcal{O}))$ where $\mathcal{H}(\mathcal{O})$ is the modular localization subspace of Fock space which is identical to the domain of $S(\mathcal{O})$.

3.7 Exotic Spin and Localization

It is well known that in d = 1 + 2 the Wigner spin is a priori not quantized. This is the reason why in the generic case $s \neq$ (half)integer, one uses often the terminology "any-on". It is remarkable that these representations reveal the nonexistence of a functor from a net of modular localized subspaces to von Neumann subalgebras in the process of refining the wedge localization.

First we note that the construction of $H_R(W_{stan})$ as an eigenspace of the unbounded Tomita involution parallels that for (half)integer spin with the only difference that instead of finding a phase factor *i* (as for halfinteger spin) which accounts for the difference between the quantum and the geometric opposite we now find:

$$H_R(W_{stan})' = e^{is\pi} H_R(W'_{stan}) \tag{3.185}$$

where the space $H_R(W'_{stan})$ is defined by applying a geometric rotation $W_{stan} \rightarrow W'_{stan}$ to $H_R(W_{stan})$. The net of wedges results from application of the Poincaré transformations to $H_R(W_{stan})$ and as in the previous cases, the positive energy is equivalent to the isotony properties of the wedge net. Surprisingly these geometric properties become lost if we try to refine the net by forming intersections. Whereas the triviality of the compactly localized subspaces obtained by intersecting at least 3 wedges is more or less expected on the physical grounds that anyonic spin should require exotic statistics which needs noncompactly localized operators (see our treatment of braid group statistics in a later chapter), the nonexistence of a isotonic net structure for noncompact localizations as e.g. spacelike cones (intersections of two wedges) is somewhat unexpected. From the

(doubled for antiparticle \neq particle) Wigner theory for the full (with reflections) Poincaré group we obtain δ^{it} and the TCP related j:

$$\begin{aligned} (\delta^{it}\phi)_{\pm}(p) &= e^{is\varphi(\Lambda(-2\pi t),p)}\phi_{\pm}(\Lambda(2\pi t)p) \\ j(\phi)_{\pm}(p) &= e^{is\pi}\overline{\phi_{\pm}(-jp)} \\ \mathfrak{s} &= j\delta^{\frac{1}{2}}, \ \mathfrak{s}H_{R}(W_{stan}) = H_{R}(W_{stan}) \end{aligned}$$
(3.186)

where ϕ_{\pm} denotes the two component wave function, $\varphi(\Lambda, p)$ is the Wigner phase for the standard boost in x-direction and $e^{is\pi}$ the TCP phase. By covariance one obtains the net of wedge spaces. By intersecting wedges we find the following obstruction for $s \neq$ (half)integer:

$$\begin{aligned} H_R(W_{stan}) & \frown & H_R(W_{rot}) \varsubsetneq H_R(gW_{stan}) \\ \forall g & \in & \mathcal{P}, \ W_{rot} = \vartheta \cdot W_{stan} \end{aligned}$$
 (3.187)

where $\vartheta \in U(1)$ denotes a spatial rotation by the angle ϑ . So even if geometrically $W_{stan} \frown W_{rot} \subset gW_{stan}$, the spaces will not be included! For the following it will be convenient to have a representation of $H_R(W)$ in terms of wedge supported test functions f(x). We choose f(x) with $\operatorname{supp} f \subseteq W$ and construct the corresponding on-shell momentum space wave function (leaving out the doubling in order to save on notation):

$$\phi(p) \equiv v(p)\hat{f}(p)|_{h_{+}}$$
(3.188)
$$v(p) = (p_{0} - p_{1})^{s} \frac{l(p)^{s}}{l(p)^{s}},$$

$$l(p) = p_{0} - p_{1} + m + ip_{2}$$

As expected the function ϕ is strip-analytic in rapidity and solves the quantum modular localization equation $\mathfrak{s}\phi = -\phi$ i.e. $\phi \in H_R(W)$; a fact which is easily established. The only nontrivial relation used is the intertwining relation:

$$v(\Lambda(-2\pi t) \cdot p)e^{is\varphi(\Lambda(2\pi t,p))} = e^{2\pi st}v(p)$$
(3.189)

which follows by straightforward calculation (from which one also obtains a very compact formula for the Wigner phase). For $t \to i\pi$ the exponential factor on the right hand side compensates the modular conjugation phase (equal to the TCP phase). In fact all of $H_R(W)$ is obtained in this way from the space of wedge supported test functions by closure.

The proof now consists in exhibiting a family of wave functions, which by construction belong to the left hand side of (3.187), but for no g are contained in the right hand Hilbert space. Since $H_R(W_{stan} \frown \vartheta \cdot W_{stan}) \subseteq H_R(W_{stan}) \frown H_R(\vartheta \cdot W_{stan})$, it follows that if we can prove that the left hand Hilbert space contrary to the geometric inclusion is not contained in $H_R(g^{-1} \cdot (W_{stan} \frown \vartheta \cdot W_{stan}))$ then this is a fortiori true for the right hand side. For this purpose we must get a better understanding of the latter space of the g-transformed

region. The relation (3.188) tells us how the standard boost acts on the rotated wave function. A somewhat lengthy but simple calculation gives:

$$\exists \phi \ s.t. \ (u(\Lambda(2\pi t))u(g^{-1})\phi)(p)$$
(3.190)
is nonanalytic in $t - strip$

For the construction of these functions it is convenient to parametrize the g with $g^{-1}(W_{stan} \frown \vartheta \cdot W_{stan}) \subset W_{stan}$ in the form $g^{-1} = \vartheta' \cdot \Lambda(-2\pi t') \cdot \frac{\pi}{2}$. By a rather lengthy calculation one shows that there are wave functions ϕ on which the application uf the above transformations (3.190) yield a function whose analytic continuation of t into the strip develops a cut for. We will spare the reader the details.

This result means that within the Wigner theory for $2s \neq integer$ one cannot refine the net of wedge localized subspaces by using intersections to e.g. a net of spacelike cones. So the situation is very different from the standard case 2s = integer where intersections even lead to nets which are indexed by compact regions as double cones. The case of compact localization is the only one which permits pointlike field generators. In the first case the extended spin statistics theorem would demand braid group statistics which is irreconcilable with a tensor product structure for multiparticle state. This explains the nonexistence of a functor. Nevertheless it is surprising that already the Wigner representation indicates a related obstruction. In a later chapter we will use scattering theory in order to determine the inner product structure of incoming free plektons and indicate how we can construct an affiliated x-space "free" field theory.

We expect similar localization problems with the d=1+3 finite energy "continuous spin" representation. Our modular localization method makes it possible for the first time to explore the physical consequences of these Wigner representations.

3.8 Localization and Hawking Temperature

In this section we present two physical interpretations of the Tomita-Takesaki formalism for the wedge localization of free field algebras which are a special case of the more general Bisognano-Wichmann property of interacting Wightman fields.

The first is a relation to Hawking-Unruh effect which was observed by Sewell [9]. The modular group for the wedge is according to Bisognano-Wichmann apart from a factor 2π equal to the wedge associated Lorentz-boost $\Delta^{it} = U\left[\Lambda(\chi = \frac{t}{2\pi})\right]$. In the integer spin case we have seen that there is complete harmony between the geometric and the quantum (in terms of von Neumann commutants) notion of localization and in the non-bosonic cases one only needs additional Klein factors. Furthermore thanks to the free field functors, the explicit construction of the modular operators and the wedge localized algebras can be delegated to the construction of an involutive unbounded antilinear soperator and its real closed subspaces H_R of -1 or +1 eigenvalue. according to

the previous section, the dense set of wedge-localized wave function, is simply $H_W^{loc} = H_R + iH_R$ and H_R consists of all momentum-space Wigner wave functions which are analytic in a strip of the wedge rapidity χ and fulfill the s-reality condition on the boundary. We know from the previous section:

$$H_R(W) + iH_R(W) = dom(\mathfrak{s}) \subset H_{Wigner} \tag{3.191}$$

$$\mathcal{H}_R(W) + i\mathcal{H}_R(W) = dom(S) \subset \mathcal{H}_{Fock} \tag{3.192}$$

These dense subspaces become Hilbert spaces in their own right if we use the graph norm of the Tomita operators. For the \mathfrak{s} -operators in Wigner space we have:

$$(f,g)_{Wigner} \rightarrow (f,g)_G = (f,g)_{Wig} + \overline{(\mathfrak{s}f,\mathfrak{s}g)}_{Wig}$$

$$= (f,g)_{Wig} + (f,\delta g)_{Wig}$$

$$(3.193)$$

The graph topology insures that the wave functions are strip-analytic in the wedge rapidity θ :

$$p_{0} = m(p_{\perp}) \cosh \theta, \quad p_{1} = m(p_{\perp}) \sinh \theta, \quad m(p_{\perp}) = \sqrt{m^{2} + p_{\perp}^{2}} (3.194)$$

strip: $0 < Imz < \pi, \quad z = \theta_{1} + i\theta_{2}$

where this "G-finiteness" is precisely the analyticity prerequisite for the validity of the KMS property for the two-point function. For scalar Bosons we have for the Wigner inner product restricted to the wedge :

$$(f,g)_{Wig}^{W} = \left\langle A(\overline{\hat{f}})A^{*}(\hat{g}) \right\rangle_{0} \stackrel{KMS}{=} \left\langle A^{*}(\hat{g})\Delta A(\overline{\hat{f}}) \right\rangle_{0}$$
(3.195)
$$\stackrel{CCR}{=} \left[A^{*}(\hat{g}) A(\overline{\hat{f}}) \right] + \left(f, \xi \right) W = \xi = \frac{2\pi K}{2\pi K}$$
(2.106)

$$\stackrel{CR}{=} \left[A^*(\hat{g}) A(\delta \overline{\hat{f}}) \right] + (f, \delta g)^W_{Wig}, \quad \delta = e^{2\pi K} \quad (3.196)$$

$$\sim (f,g)_{Wig}^W \equiv (f,g)_{K,T=2\pi} = \left[A^*(\hat{g})A(\frac{\delta}{1-\delta}\overline{\hat{f}})\right]$$
(3.197)

Here we used a field theoretic notation $(A^*(\hat{g}) \text{ is a smeared scalar complex field}$ of the type (3.197) linear in \hat{g} with $supp.\hat{g} \in W$) in order to emphasize that the temperature dependence on the right hand side is explicit via the δ acting on the complex-valued x-space smearing functions in the c-number commutator and not implicit, as the restriction of the wave functions to the wedge region on the left hand side. Of course the c-number commutator (without the state brackets) may be rewritten in terms of p-space Wigner wave functions for particles and $(\delta^{\frac{1}{2}}$ -transformed) antiparticles in such a way that the localization restriction is guarantied by the property that the resulting expression is finite if the wave functions are G-finite. We mention for experts that the difference between localization temperatures and heat bath temperatures on the level of field algebras in Fock space corresponds to the difference between hyperfinite type III_1 and type I von Neumann algebras. In this way one obtains a thermal representation of the wedge restricted Wigner inner product. The fact that the boost K appears instead of the Hamiltonian H for the heat bath temperature reveals one significant difference between the two situations. For the heat bath temperature of a Hamiltonian dynamics the modular operator $\delta = e^{-2\beta \mathbf{H}}$ is bounded on one particle wave functions whereas the unboundedness of $\delta = e^{2\pi K}$ in (3.196) enforces the localization (strip analyticity) of the Wigner wave functions.

This difference results from the two-sided spectrum of K as compared to the boundedness from below of H. In fact localization temperatures are inexorably linked with unbounded symmetry operators.

The generalization to fermions as well as to particles of arbitrary spin is easily carried out with the result that the localized thermal representation involves the anticommutator. The differences between K and H also leads to somewhat different energy distribution functions for small energies so that Boson K-energy distributions may appear as those of H heat bath Fermions. In this context one is advised to discuss matters of statistics not in Fourier space, but rather in spacetime where they have their unequivocal physical interpretation.

One may of course consider KMS state on the same C^* -algebra with a different K-temperature than 2π , however such a situation cannot be obtained by a localizing restriction. Mathematically C^* -algebras to different K-temperatures are known to belong to different folii (in this case after von Neumann closure to unitarily inequivalent III₁-algebras) of the same C^* -algebra. Or equivalently: a scaled modular operator $\Delta^{\alpha i\tau}$ cannot be the modular operator of the same theory at a different temperature as it would be the case for type I algebras.

As in the Bisognano-Wichmann situation, the modular wedge localization in the Wigner theory has led us to the Hawking-Unruh thermal situation. For those readers who are familiar with Unruh's work we mention that the Unruh Hamiltonian is different from K by a factor $\frac{1}{a}$ where a is the acceleration (see below). Any modular localization (not necessarily the wedge) leads to horizons and a thermal state. Only in very special cases one has a geometrical picture in terms of Killing vectors in space-time.. In the present setting it is the arena of Wigner space where one finds the isometries. In order to remove any doubt that these thermal properties are not typical for curved space-time QFT but constitute a general property of QFT and as such are very relevant for ordinary QFT. In order to illustrate this we look at the relation with crossing symmetry of formfactors.

More generally we may now consider matrix elements of wedge-localized operators between wedge localized multiparticle states. Then the KMS property allows to move the wedge localized particle state as an antiparticle at the analytically continued rapidity $\theta + i\pi$ from the ket to the bra. The simplest illustration is the two-particle matrix element of a free current of a charged scalar field $j_{\mu}(x) =: \phi^* \stackrel{\leftrightarrow}{\partial}_{\mu} \phi$: smeared with the

wedge supported function \hat{h} :

$$\left\langle 0 \left| \int j_{\mu}(x) \hat{h}(x) d^{4}x \right| f, \delta^{\frac{1}{2}} g^{c} \right\rangle$$

$$\overset{KMS}{=} \left\langle 0 \left| \left(\Delta \phi^*(\delta^{\frac{1}{2}} \widehat{g}) \Delta^{-1} \right)^* \int j_{\mu}(x) \hat{h}(x) d^4x \right| f \right\rangle$$

$$= \left\langle \bar{g} \left| \int j_{\mu}(x) \hat{h}(x) d^4x \right| f \right\rangle$$

$$(3.198)$$

Here $\delta^{\frac{1}{2}}g^c$ is the charged transformed antiparticle in the Wigner wave function g at the analytically continued rapidity $\theta + i\pi$ whereas \hat{g} denotes as before the wedge-localized space-time smearing function whose mass shell restricted Fourier transform corresponds to the boundary value of the analytically continuable Wigner wave function g. Moving the left hand operator to the left vacuum changes the antiparticle charge to the particle charge. Since the $H_R(W) + iH_R(W)$ complex localization spaces are dense in the Wigner space, the momentum space kernel for both sides of (3.198) takes the familiar form:

$$\langle p' | j_{\mu}(0) | p \rangle = anal.cont. \left\langle 0 | j_{\mu}(0) | p, p'(z) \right\rangle$$
 (3.199)

where p'(z) is the rapidity parametrization of above (3.194). This famous crossing symmetry, which is known to hold also in each perturbative order of renormalizable interacting theories, has never been derived in sufficient generality within any nonperturbative framework of QFT. It is to be thought of as a kind of on-shell momentum space substitute for Einstein causality and locality (and its strengthened form called Haag duality).

There are two slightly different interpretations of the thermal wedge situation.

The first physical interpretation is in terms of the Unruh effect i.e. the temperature experienced in the vacuum by a uniformly accelerated observer [55]. Such an observer moves on a world line: $t = \xi cosh\tau$, $x = \xi sinh\tau$ in an appropriately chosen Lorentz frame. In natural units (c = 1, $\hbar = 1$) his acceleration is $a = \frac{1}{\xi}$ and his proper time τ_a is related to the wedge "rapidity" τ by $\tau_a = \frac{\tau}{a}$ and the corresponding Hamiltonian is aK where K is the boost generator. The inside of the wedge (the Rindler world) as well as its boundary (the horizon of the Rindler world) are invariant under the boosts in the wedge direction $U(\Lambda_W)$. On the positive part of the wedge boundary bdW_+ , the action of the positive spectral light-like translations together with the boosts form an interesting two-parametric group which has a deep relation to so called half-sided modular inclusions [85], but here we will confine ourselves to more pedestrian methods.

Intuitively speaking, we expect that the global vacuum state appears similar to a heat bath with respect to an uniformly accelerated observer. After all, the wedge horizon generated by the acceleration signifies a loss of information (the opposite wedge suffers a causal blackout for the accelerated observer) which is also a characteristic feature of a heat bath temperature state. The above formula shows indeed that the restriction of the vacuum state to the wedge algebra (or the modular localization Wigner subspace belonging to W) satisfies the KMS condition with respect to the L_W -Lorentz boost with the Hawking-Unruh temperature $\beta_a \equiv kT_a = \frac{a}{2\pi}$. Actually Unruh did all his calculations explicitly (without recourse to modular localization theory) on a simple model of a two-level detector coupled to a free field. The reader finds a beautiful pedestrian review in⁷ [52].

Sewell [9] has shown that the Unruh effect may be generalized in such a way, that it serves to understand the Hawking effect in a quite general setting which includes black holes.

We now direct our attention to a second physical aspect of the wedge situation which is the Hawking effect: of pair creation and the antiparticle aspect, as well as its refined version namely crossing symmetry. For this reason we study the complex d = 1 + 1 Klein-Gordon field in a homogeneous external electric field which points into the wedge direction. We pick a gauge e.g. the axial gauge in time direction $A_t = Ex$, $A_x = 0$. With the Ansatz $\Phi = e^{-i\omega t}\varphi_{\omega}(x)$ the Klein-Gordon operator takes the form:

$$\left[\partial_x^2 + (\omega + Ex)^2\right]\varphi_\omega(x) = m^2\varphi_\omega(x) \tag{3.200}$$

By further canonical transformations $x \to \xi = \sqrt{E}(x + \omega/E), -i\frac{\partial}{\partial x} \to i\frac{\partial}{\partial \xi}$ and $u = \frac{1}{\sqrt{2}}(-i\frac{\partial}{\partial \xi} - \xi)$, the equation looks like a eigenvalue equation of an auxiliary Schrödinger Hamiltonian which is isomorphic to the boost or scaling generator:

$$u\frac{\partial}{\partial u}\varphi(u) = (i\varepsilon - \frac{1}{2})\varphi(u), \quad \varepsilon = \frac{m^2}{2E}$$
(3.201)

That this boost had to eventually appear is clear from the classical picture. The charged massive particles are uniformly accelerated and their trajectory is identical to that of the Unruh accelerator. Classically there is a perfect wedge horizon, but quantum field theoretically the vacuum state does not factorize into product states on the wedge and its causal opposite. Rather we expect particle-antiparticle pairs to appear near the horizon which as a result of the action of the constant electric field will be split into say right wedge particles and left (opposite) wedge antiparticles. A pedestrian treatment of this external field problem would approximate the constant field by a sequence of external fields which are different from zero only during a time T and inside a box of length L around the tip of the wedge with $T, L \to \infty$ at the end of the calculation. The relation between the free field creation and annihilation operators before (in) and after (out) is given in terms of a Bogoliubov automorphism:

$$a_{\omega}^{out} = \bar{\alpha} a_{\omega}^{in} + \beta b_{\omega}^{in*}$$

$$b_{\omega}^{out*} = \alpha b_{\omega}^{in*} + \bar{\beta} a_{\omega}^{in}$$
(3.202)

However this global automorphism is only unitarily implemented as long as T, L are kept finite (the spectrum of $\omega's$ is discrete and the density of ω states is proportional to T, L and E)

⁷But beware of certain pitfalls, e.g. KMS states are not Gibbs states (the latter need a quantization box which would wreck the wedge geometry) and massless free fields in d=1+1 have infrared divergent two-point functions. This requires an easy repair without change of conclusions.

The vacuum persistency probability follows the well known Schwinger formula:

$$|\langle 0_{out} | 0_{in} \rangle|^2 = exp\left(-\frac{ELT}{2\pi}ln(1+|\beta|^2)\right), \quad |\beta|^2 = e^{-m^2\pi/E}$$
(3.203)

which in turn is a consequence of the pair creation term in the formally unitary implementer U of the Bogoliubov transformation:

$$0_{out}\rangle = U |0_{in}\rangle = e^{bil(a^*, b^*)} |0_{in}\rangle$$
(3.204)

where *bil*. indicates the characteristic exponential bilinear dependence of implementers of Bogoliubov transformations on the $a^{\#'s}$ and $b^{\#'s}$. From the population ratio of:

$$\langle n_{\omega} \rangle = \left\langle 0_{in} \left| a_{\omega}^{out*} a_{\omega}^{out} \right| 0_{in} \right\rangle = \left| \beta \right|^2 \tag{3.205}$$

for two charged free fields with slightly different masses m and $m + \Delta m$ one obtains:

$$\frac{\langle n \rangle_{m+\Delta m}}{\langle n \rangle_m} = e^{-2\pi\Delta m/a}, \quad a = \frac{E}{m}$$
(3.206)

i.e. the ratio of created pairs obeys a thermal distribution with temperature $kT = 2\pi/a$ as in the Unruh interpretation of the wedge localization.

This last somewhat clumsy and conceptually controversial (the box-quantization is too much of a brute force computational device in scattering theory) derivation of the radiation aspects of the constant external electric field model (the same which in many textbooks serves to illustrate the Klein paradoxon) should be replaced by a more elegant method using appropriate representation concepts which do lead to exponentials free of TL-volume factors, in particular one expects a finite entropy density ratios per unit horizon surface for say two models with different matter contents. The afore-mentioned considerations suffer from the fact the type III_1 nature of the localized algebras together with the fact that there is no causally disjoint space-time between the wedge region and its opposite, lead to uncontrollable fluctuations which strictly speaking wreck the existence of Fock-space Bogoliubov operators and cause serious problems with an intrinsic notion of entropy. Here the split property (with the split distance approaching zero at the end of the calculation) is expected to have beneficial consequences, but the calculation has not been carried out. This split property which is related to the correct counting of degrees of freedom in QFT "phase space will be explained in a later chapter and in the mathematical appendix.

As a curious side result we mention that the low energy distribution of a system with a Hawking-Unruh temperature resulting from modular localization can be significantly different from that of a standard heat bath temperature. For free bosons and fermions, the energy distribution is given in terms of the spectral decomposition of:

$$\frac{1}{1 \mp e^{\beta K}} \tag{3.207}$$

with K being proportional to the infinitesimal generator of the modular group.

Only in the case of the heat bath situation is K the positive energy Hamiltonian in the rest frame of the heat bath. For the Rindler localization K is the wedgeaffiliated Lorentz-boost. The most dramatic difference between these two kinds of thermalizations show up for zero mass in odd space-time dimensions d (where the "reverberation" leads to a breakdown of Huygens principle). A convenient way to obtain the spectral distribution of (3.207) is to Fourier transform the zero-mass two-point function W in the rapidity (boost) variable τ :

$$\tilde{W}(\omega) = \int_{-\infty}^{+\infty} e^{-i\omega\tau} W(\xi(\tau))$$
(3.208)
$$W(\xi) = \frac{\Gamma(\frac{d}{2} - 1)}{4\pi^{\frac{d}{2}}} \frac{1}{\left(\sqrt{-(\xi_0 + i\varepsilon)^2 + \vec{\xi}^2}\right)^{d-2}}$$

$$\xi(\tau) = \frac{1}{a} (sh\tau a, ch\tau a, 0, ..0)$$

For d = 2 + 1 we obtain the well-known result[94] :

$$\tilde{W}(\omega) = \frac{1}{2} \frac{1}{1 + e^{\beta \omega}} \tag{3.209}$$

which is spectacularly different from the heat bath result $\frac{1}{2} \frac{1}{1 - e^{\beta \omega}}$.

This shows in addition, that one should avoid to analyze particle statistics in momentum space.

The zero mass model is conformally invariant. This means in particular that there are other situations with horizons which are conformally equivalent: the forward light-cone (modular group = dilatations) and double cones (modular group= radial conformal subgroup). They lead to identical energy distributions but their physical interpretation is more involved. One very surprising feature of modular wedge localization is that to compute thermal expectation values with the KMS condition for given commutation structure for low dimensional fields is often simpler than to handle ground state problems with that algebraic structure.

There is another important lesson to be drawn from this section. The modular localization aspects agree with the geometry of Killing isometries in the Rindler situation (and its conformal transforms for zero mass matter) with the Killing time being exponentially related to the (affine) geodesic time on the horizon. However the modular automorphism group, which exists for any region with a nontrivial causal complement, vastly generalizes these concepts to situations where the quantum aspects of localization do not allow for a geometric interpretation e.g. for general space-time regions as double cones. In our free field case at hand, this generalization of the Unruh picture is not describable in terms of finite dimensional geometric data, but rather in terms of localized subspaces of the Wigner space and their modular properties i.e. they depend more on the material content and not only on geometric properties. This means that the notion of "horizon" becomes more "quantum" and inherits its properties not so much from the geometry in space-time, but more from the position of modular localized quantum subspaces inside Wigner resp. Fock-space. Even those algebras localized close to the horizon are expected to become increasingly "fuzzy" for increasing "modular" times in the modular group action. Presently these general modular aspects are ill-understood. This remark also applies to the closely related concept of entropy and the dependence on the "split" property. The latter leads to a factorizing Rindler state which belongs to the same folium as the global vacuum. Of course paradoxes of quantum theory in the presence of evaporating black holes cannot be convincingly resolved without an entropy concept for nets.

Note that restriction of pure global states to local algebras always produce impure (thermal) states on $\mathcal{A}(\mathcal{O})$. This is a manifestation of the type III₁ nature of the local versus the type I global algebras. Overlooking this effect would lead to fake causality violations.

3.9 Examples of Dis(order)-Fields

In the following we will give an example for a field whose Borchers class is associated with (but not equal to) that of a free field. This construction is part of the d=1+1 duality (order- disorder) construction.

Let us start from a complex free Dirac field in d=1+1:

$$\psi(x) = \int \frac{dp}{2\omega} (e^{-ipx} u(p)a(p) + e^{ipx} v(p)b^*(p))$$
(3.210)
$$u(p) = \sqrt{\frac{m}{2}} \begin{pmatrix} e^{\frac{\theta}{2}} \\ -e^{-\frac{\theta}{2}} \end{pmatrix}, \quad v(p) = \sqrt{\frac{m}{2}} \begin{pmatrix} -e^{\frac{\theta}{2}} \\ -e^{-\frac{\theta}{2}} \end{pmatrix}$$
$$v(p) = u^C(p) = Ci\gamma_0 u(p), \quad p = m(\cosh\theta, \sinh\theta)$$

Here we took the following realization of the Dirac equation:

$$\begin{pmatrix} i\gamma_{\mu}\partial^{\mu} - m \end{pmatrix} \psi = 0$$

$$\gamma_{0} = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \qquad \gamma_{1} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$

$$C = i\gamma_{1} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

$$(3.211)$$

This field is U(1) covariant and the local generator is the conserved current $j_{\mu} =: \bar{\psi} \gamma_{\mu} \psi$:. This d = 1 + 1 current has (relatively to ψ) nonlocal pseudo-potential:

$$j_{\mu}(x) = \varepsilon_{\mu\nu}\partial^{\nu}\phi(x), \quad \phi(x) =: F_{x}(a^{\#}, b^{\#}) := \int_{-\infty}^{x} \varepsilon^{\mu\nu} j_{\nu}d\xi_{\mu} = (3.212)$$

$$= \int \left\{ \begin{array}{cc} \frac{-1}{\sinh\frac{1}{2}(\theta_{p}-\theta_{q}+i\varepsilon)}e^{i(p-q)x}\left[a_{p}^{*}a_{q}-b_{p}^{*}b_{q}\right] + \\ +\frac{1}{\cosh\frac{1}{2}(\theta_{p}-\theta_{q})}\left[e^{i(p+q)}a_{p}^{*}b_{q}^{*}+e^{-i(p+q)}a_{q}b_{p}\right] \end{array} \right\} d\theta_{p}d\theta_{q}$$

As naively expected, the ϕ is a localized field which, although relatively local with respect to the observables (generated by the current), fails to be local relative to the field ψ but instead fulfills:

$$\begin{aligned}
\phi(x), \psi(y) &= \theta(x_0 - y_0)\psi(y)\phi(x) \quad (3.213) \\
\lim_{x^2 \to -\infty, x^1 \to \infty} \phi(x) &= \sqrt{\pi}Q,
\end{aligned}$$

where Q is the global charge. Formally we may write

$$U(2\pi\lambda)_{<} = \exp{-2\pi i \int_{-\infty}^{x_1} j_0(x_0, y_1) dy_1} = \exp{-2\sqrt{\pi i \lambda}\phi(x)}$$
(3.214)

is the representation for "half-space" rotation i.e. $\mathrm{U}_{<}$ implements:

$$\psi(x) \to \begin{cases} e^{-2\pi i\lambda}\psi(x) & x_1 < 0\\ \psi(x) & x_1 > 0 \end{cases}$$
(3.215)

Such half-space transformations may be viewed as the point limit of Bogoliubov transformations. The correct normal product which is necessary in order to convert $U_{<}$ into a well defined expression, is the "triple" ordering. This is also recursively defined but different from the ϕ -Wick product; one subtracts all connected correlation functions and not just the two-point function. Formally we have the following simple exponential formula:

$$:e^{ia\phi(x)} := \frac{e^{ia\phi(x)}}{\langle e^{ia\phi(x)} \rangle}$$
 (3.216)

re-expressed in terms of the original ψ -Wick product we obtain a nonlocal looking expression, which is best written in terms of rapidities:

$$\mu(x) = \exp -2i\sqrt{\pi}\lambda\phi(x) = \exp L_{\lambda}(x) : \qquad (3.217)$$

$$L_{\lambda}(x) = \frac{\sin \pi \lambda}{2\pi} \int \left\{ \begin{array}{c} \frac{e^{-\lambda(\theta_{p}-\theta_{q})}}{\cosh \frac{1}{2}(\theta_{p}-\theta_{q})} \left[e^{i(p+q)x} a_{p}^{*} b_{q}^{*} + h.c. \right] + \\ \left[-\frac{e^{-\lambda(\theta_{p}-\theta_{q})}}{\sinh \frac{1}{2}(\theta_{p}-\theta_{q}+i\varepsilon)} \right] e^{i(p-q)x-i\pi\lambda} a_{p}^{*} a_{q} + \\ \left[\frac{e^{-\lambda(\theta_{p}-\theta_{q})}}{\sinh \frac{1}{2}(\theta_{p}-\theta_{q}-i\varepsilon)} \right] e^{-i(p-q)+i\pi\lambda} b_{q}^{*} b_{p}. \end{array} \right\} d\theta_{p} d\theta_{q} \quad (3.218)$$

Although L_{λ} is represented in terms of nonlocal kernels in rapidity space and is itself nonlocal, μ is a bosonic local field in the quantum sense which is however nonlocal relative to ψ i.e. outside the ψ -Borchers-class. It is easy to see that our special solution μ of the half-space commutation relation with ψ belongs to a whole family of solutions. We may modify the μ by any bosonic local function of the $\psi^{\#}$ without change in the relative commutation relations. Within our construction method this is made manifest by the "quasi-periodicity" (up to local operators) in λ mod 1. With the help of μ one can now construct another

3.9. EXAMPLES OF DIS(ORDER)-FIELDS

field $\sigma(x)$ which carries the same charge as ψ , but has quite different spacelike commutation relations. Through the definition:

$$\sigma(x) = N\left[\mu\psi\right](x) = \lim_{y \to x} \mu(x)\psi(y) \qquad (3.219)$$
$$= \frac{1}{\sqrt{4\pi}} : \int (e^{-ipx}a_p + e^{ipx}b_p^*)\mu(x)d\theta_p :$$

one obtains the same soliton like relative commutation relations with μ as those between ψ and μ :

$$\mu(x)\sigma(y) = \begin{cases} e^{i\alpha}\sigma(y)\mu(x) & x > y\\ \sigma(y)\mu(x) & y > x \end{cases}$$
(3.220)

However the σ carries a fractional spin and "statistics" (see later comments). Instead of dual commutation relations one finds symmetric commutation relations associated with abelian representations of the braid group i.e.

$$\sigma(x)\sigma(y) = e^{-2\pi i\lambda sign(x-y)}\sigma(y)\sigma(x) \tag{3.221}$$

These commutation relations appear as a interpolating continuous generalization of Fermions and Bosons and are called "anyonic". Their relation to particle statistics will be discussed later. The bosonic field as well as the anyonic field are living in the same Fock space generated by the free field ψ , but they are not members of the ψ -Borchers class. As a local field μ generates its own Borchersclass (it is an irreducible field in its own Hilbert space cyclically generated from the vacuum). The question of whether the notion of equivalence classes of fields can be generalized to anyonic fields will not be pursued here.

A physically more relevant illustration of duality and non-free Borchers classes is obtained by starting from a Majorana (selfconjugate) spinor field (a = b). In this case the symmetry is the discrete \mathbb{Z}_2 and the previous method based on a conserved Noether current is not applicable. There are however several alternative methods which lead to the following result ($c \equiv a = b$):

$$\mu(x) = :e^{\Lambda(x)}:, \quad \sigma(x) = s.d.l.: \hat{\psi}(x)\mu(x): \qquad (3.222)$$

$$\Lambda(x) = \frac{i}{4\pi} \int d\theta_p d\theta_q \left\{ \begin{array}{l} 2 \coth \frac{1}{2}(\theta_p - \theta_q + i\varepsilon) \cdot e^{i(p-q)}c_p^* c_q \\ + \tanh \frac{1}{2}(\theta_p - \theta_q) \cdot (e^{i(p+q)x}c_p^* c_q^* - h.a.) \end{array} \right\}$$

$$\hat{\psi}(x) = \frac{1}{2\pi} \int d\theta_p (e^{-ipx}c_p + e^{ipx}c_p^*) \neq \psi(x)$$

Here s.d.l. stands for the operator contribution which carries the leading short distance singularity). Whereas μ and σ fulfill the relative Z_2 -duality relation, both fields are bosonic. Hence σ generates a new Borchers class in \mathcal{H}_F which is inequivalent to the Fermion Borchers class. It is quite straightforward to show that the Ising lattice model can be described in terms of lattice Fermions which in the scaling limit (for fixed correlation length) become Majorana Fermions. In addition the lattice (dis)order variables go over into (μ) σ if one takes that

limit from the disorder side $(T \to T_c + \varepsilon)$. So we are justified to call our σ -fields the (order) "Ising fields". Let us compare the free Majorana field with the Ising field Borchers class. Consider the modular objects for the wedge algebras of the two classes. The modular operators Δ^{it} are identical and equal to the wedge-based Lorentz transformations. However the modular reflections J are different. For the free Fermion algebra the Wigner theory gave $J_F = K\mathcal{F}(j)$ with j being the antiunitary wedge reflection, \mathcal{F} the CAR-functor and K the Klein transformation in Fock space. The Boson algebra generated by σ on the other hand has $J_B = KJ_F$ because its commuting structure for space-like distances requires the absence of the twist. This can also be read off directly from the TCP transformation property of σ under the TCP in Fock space. Note that the Klein factor is a global operator whose half-space version is the disorder field μ (the Jordan-Wigner transformation in lattice theory). In our Ising example $(\mathcal{N}_F: \text{ fermion}\#)$:

$$K = \frac{1+iU}{1-iU}, \quad U = e^{i\pi\mathbf{N}_F} \tag{3.223}$$

Since the TCP symmetry θ of σ is related to the free field θ_0 TCP of the Majorana Fermion ψ in the same way as the above J's, namely by:

$$\theta = K\theta_0 \tag{3.224}$$

and since (as will be shown in the section on scattering theory) the unitary S-matrix is related to the antiunitary $\theta' s$ by $S = \theta \theta_0$, we conclude K = S. this means that the S of σ is energy independent and $S^{(2)} = -1$ for two particles. On a somewhat formal level we can understand this through:

$$\lim_{t \to \infty} \mu(x) = \begin{cases} U \\ 1 \end{cases}, \quad \lim_{t \to \pm \infty} \sigma(x) = \begin{cases} U\psi \\ \psi \end{cases}$$
(3.225)

Writing $U\psi = K\psi K^*$, we read off: S = K i.e. the Jordan-Wigner transformation approaches the global symmetry whose square root is the Klein transformation (which in this model coalesces with the S-matrix).

Returning for a moment to the λ -half-space rotation in the previous complex free Dirac field, we find by the same method in case of rational $\lambda = \frac{1}{N} (Z_N \text{ symmetry})$:

$$U = \sum_{n} e^{-2\pi i \frac{n^2}{N}} E_n \qquad (3.226)$$
$$K = \sum_{n} e^{-i\pi \frac{n^2}{N}} E_n$$

The quadratic n-dependence of the spin-statistic phase on the charge eigenvalues $\sim n^2$ is characteristic for anyonic commutation relations. The "exotic" nature of the commutation relation of this anyonic σ with itself shows up in the deviation of its modular reflection J from its expected geometric action J_0 . Such a twist factor $K = J \cdot J_0$ should be distinguished from the S-matrix in $S = J \cdot J_0$ in chapter 6. In the latter case J_0 has the interpretation of the incoming modular

3.9. EXAMPLES OF DIS(ORDER)-FIELDS

involution and J belongs to the interacting field algebra. Both involutions are geometric and belong to fields with the same local commutation relations. In this case there is no need for a twist which repairs the lost connection between spin and statistics. It should however be emphasized that in massive d=1+1 theories (different from chiral conformal or massive d=1+2 theories), the distinction between a scattering and a twist interpretation of $J \cdot J_0$ remains a delicate one. In some sense statistics in d = 1 + 1 massive QFT is an ambiguous notion. Consider the $a^{\#}(p)$ of a free Dirac field and write:

$$c(p) = e^{i\pi\lambda \int_{-\infty}^{p} n(p)dp} a(p) \qquad (3.227)$$

$$n(p) = a^{*}(p)a(p)$$

The commutation relation of the c's is now "anyonic":

$$c(p)c^*(q) = -e^{i\pi\lambda}c^*(q)c(p) + 2\omega\delta(p-q)$$

$$c(p)c(q) = -e^{i\pi\lambda}c(q)c(p), \quad etc.$$
(3.228)

The c's live in the same Fock space, are covariant with respect to the same representation of the Poincaré-group and (as it should be) create the same one particle states, even though the c's have anyonic (for $\lambda = 1$ bosonic) commutation relations. This is a special instance of a general phenomenon: particles in massive d = 1 + 1 theories are really statistical "schizons" [32] i.e. the nature of their charges (fusion laws etc.) does not determine their statistics (i.e. a Mendeleev periodic table in a d = 1 + 1 world allows for a bosonic description in terms of long range interactions). This is very different from all other situations, including chiral conformal QFT for which the field commutation relations (the "exchange algebra") is uniquely determined in terms of the charges carried by the fields. Warning: the statistical schizon phenomenon should not be confused with "bosonization" in chiral conformal QFT (see next section). In the latter case the commutation relations of the charge carrying fields are uniquely determined by the characteristics of these charges and cannot be changed. To the extend that the name "bosonization" indicates that a conformal fermion can live in the Fock space of bosonic creation and annihilation operators, this is a misunderstanding.

An exponential bose field cannot be interpreted as just an exponential function of an object in bosonic Fock space, rather it must be considered as a short hand notation for an field which defines (through its correlation functions) its own Hilbert space, which is a charged representation of the neutral current algebra. In this sense "bosonization" is a generic structure independent of spacetime dimensions and conformal invariance: any charged field (fermionic, for d<1+3 also plektonic) can be interpreted from arising from representing another sector on the (necessarily) bosonic observable algebra, i.e. in this sense algebraic QFT is "bosonization" par excellence. The only special feature is that the process of creating a charge by dumping an anticharge "behind the moon" can be expressed in terms of a very simple formula e.g.

$$\psi(x) = \lim_{y \to \infty} e^{i\pi \int_x^y jd\xi}$$
(3.229)

in which the limit leaves Fock space and produces the new representation space.

The schizon phenomenon is related to the fact that the natural framework for massive d = 1 + 1 theories is the "soliton" framework in which the concept of braid-group statistics is replaced by the more general "exotic" (or solitonic) commutation relations which can be changed at will (within certain limitations) without effecting the superselection structure. The most interesting new phenomenon in the algebraic QFT of solitons is that the problem of multiple vacuum states, even in situations where this cannot be blamed on spontaneous symmetry-breaking, becomes related in a profound way with the superselection structure. We will return to these problems in the sections on algebraic QFT (the net approach).

We conclude this section by some comments on the algebraic description [53] (independent of fields) of (dis)order. If one assumes that the theory is given in terms of a field net $\mathcal{F}(\mathcal{O}), \mathcal{O} \in \mathcal{K}$ (family of double cones). As usual the observable algebra is related to the field algebra by the invariance principle with respect to a symmetry group G:

$$\mathcal{F}(\mathcal{O})^G \mid_{H_{vac}} = \mathcal{A}(\mathcal{O}) \tag{3.230}$$

Whereas for $d \neq 1 + 1$ the so defined $\mathcal{A}(\mathcal{O})$ generically (apart from spontaneous symmetry breaking presented in a later section) is Haag dual if \mathcal{F} had this property (for fermionic \mathcal{F} the duality must be twisted):

$$\mathcal{F}^{tw}(\mathcal{O})' = \mathcal{F}(\mathcal{O}') \curvearrowright \mathcal{A}'(\mathcal{O}) = \mathcal{A}(\mathcal{O}') \tag{3.231}$$

However in massive d=1+1 theories this conclusion is incorrect for double cones but remains correct for wedges $\mathcal{O} = W$. The alternative definition in terms of $\mathcal{A}(W)$:

$$\mathcal{A}^{d}(\mathcal{O}) = \bigcap_{W \supset \mathcal{O}} \mathcal{A}(W), \quad \mathcal{O} \in \mathcal{K}$$
(3.232)

gives a bigger algebra (equal to the dual net) which is Haag dual. It comes as a bit of a pleasant surprise that the issue of \mathcal{A} versus \mathcal{A}^d is inexorably linked with the (dis)order structure. In any massive two-dimensional QFT with an internal group symmetry G (i.e. not just for free fields) there is a canonical way to introduce half-space transformations $U_l^{\mathcal{O}}(g)$ which implement the full g-transformation on the spacelike left of the double cone \mathcal{O} and is equal to the identity on its right. This construction uses the "split-property" (equivalent to the "nuclearity" i.e. a good phase-space behavior of QFT), a concept which will be explained in the mathematical appendix. Assume for the moment that $G=Z_N$ i.e. an abelian group which leads to one half-space generator $U_l^{\mathcal{O}}$, $(U_l^{\mathcal{O}})^N \sim \mathbf{1}$. We then extend the field algebra $\mathcal{F}(\mathcal{O})^{-8}$ by the disorder operators $U_l^{\mathcal{O}}$

$$\mathcal{F}_{ext}(\mathcal{O}) = \mathcal{F}(\mathcal{O}) \lor U_l^{\mathcal{O}}$$
(3.233)

 $^{^{8}}$ Here the reader is asked to be content with an intuitive understanding of local algebras and nets. The rigorous mathematics (see the appendix) is not really necessary for a first glimpse without proofs.

The map $\mathcal{O} \to \mathcal{F}(\mathcal{O})$ is still an isotonous net but it lost locality. The application of the invariance principle yields:

$$\mathcal{F}_{ext}(\mathcal{O})^{Z_n} = \mathcal{A}^d(\mathcal{O}) = \mathcal{A}(\mathcal{O}) \lor U_l^{\mathcal{O}} = \mathcal{A}_{ext}(\mathcal{O})$$

and we may arrange our result in form of the following "commuting square" of inclusions [53]:

$$\begin{array}{cccc}
\mathcal{A}^{d}(\mathcal{O}) & \subset & \mathcal{F}_{ext}(\mathcal{O}) \\ & & & \cup \\
\mathcal{A}(\mathcal{O}) & \subset & \mathcal{F}(\mathcal{O})
\end{array}$$
(3.234)

Is there an invariance principle which describes the entire commuting square, in particular of $\mathcal{A}(\mathcal{O}) \subset \mathcal{F}_{ext}(\mathcal{O})$? In the above abelian Z_N - illustration the halfspace transformations U commute with $G = Z_N$ but they suffer a nontrivial action of the dual group \hat{Z}_N ($\sim Z_N$):

$$\alpha_{\chi}(U(g)) = \chi(g)U(g), \quad \mathcal{A}^{d}(\mathcal{O})^{Z_{N}} = \mathcal{A}(\mathcal{O})$$

$$g \in Z_{N}, \quad \chi \in \hat{Z}_{N}$$

$$(3.235)$$

Whereas on \mathcal{F} only G acts, \mathcal{F}_{ext} is a natural domain for the action of the "double" $G \times \hat{G}$:

$$\mathcal{A}(\mathcal{O}) = (\mathcal{F}_{ext}(\mathcal{O}))^{G \times \hat{G}}, \quad \mathcal{F}(\mathcal{O}) = \mathcal{F}_{ext}(\mathcal{O})^{\hat{G}}$$
(3.236)

It turns out that this has an interesting counterpart for nonabelian G's. In that case the "double" has to be taken in the sense of Drinfeld which is the cross product Hopf algebra which was introduced in the third section of the first chapter:

$$D(G) = C(G) \bowtie_{adj} G \tag{3.237}$$

However in contradistinction to ordinary group symmetry, the double is always spontaneously broken and maximally only G survives as an unbroken symmetry. This mathematical manifestation of the (dis)order structure is presently the only case for which Hopf algebras emerge naturally from physical principles.

The expert reader recognizes the close relation with the global Kramers-Wannier Duality of statistical mechanics for lattice systems. In fact the formal scaling limit near a critical point towards a continuous QFT maintains the local (the Kadanoff-)form of the K-W symmetry⁹, although the Kramers-Wannier Duality and the notion of dual temperature T^* gets lost. The presentation of this section shows the close relation of the statistical mechanics duality concepts to the Haag duality of algebraic QFT. But whereas the former (in its relation to charge sectors) is limited to d = 1 + 1, Haag duality (and its controlled breaking) does not suffer such a limitation.

⁹The temperature becomes traded for the mass, but there is no "dual mass" which could substitute for the dual K-W temperature. Rather the dual symmetry becomes a pure algebraic concept in the sense of Kadanoff. It is always spontaneously broken, except in the conformal scale invariant zero mass limit.

As already mentioned, all this rich structure (including the statistical "schizon" aspect of d=1+1 particles) may be subsumed into the algebraic QFT framework for soliton sectors[87] which will not be presented in these notes.

Finally it is worthwhile to remark that all of the known d = 1 + 1 (dis)order models allow for a euclidean functional (Feynman-Kac)representation involving an external Aharonov-Bohm potential or (in the case of additive symmetries i.e. translations in field space) external "electric and magnetic" sources. Some more details can be found in the section on functional methods.

3.10 Special Features of m=0, d=1+1 Fields

It is well-known that the zero mass limit of massive free fields enhances the spacetime symmetry to the conformal group symmetry. In addition to the general well understood peculiarities of such an extension (Einstein causality "paradox" as the result of a continuous link through infinity of the space- and time-like), there is a very surprising phenomenon which only happens for d = 1 + 1: there are continuously many local quantum theories in the holomorphy region between the real and imaginary time boundary values. Let us verify this for the massless Dirac field which results from the formula (3.210):

$$\psi_r(v) = \frac{1}{\sqrt{2\pi}} \int_0^\infty dp (e^{-ipv} a_l(p) + e^{ipv} b_l^*(p)), \quad v = t - x \quad (3.238)$$

$$\psi_l(u) = \frac{1}{\sqrt{2\pi}} \int_0^\infty dp (e^{-ipu} a_r(p) + e^{ipu} b_r^*(p)), \quad u = t + x$$

where the right and left movers $a_r^{\#}(p)$, $b_r^{\#}(p)$ and $a_l^{\#}(p)$, $b_l^{\#}(p)$ anticommute with each other i.e. the chiral fields $\psi_{r,l}$ define independent chiral theories. Therefore from now on we will select one chirality and omit the r,l subscript.

As expected the two-point function can be rewritten in compact circular coordinates $z = e^{i\varphi}$ by introducing an appropriately normalized field ψ_c via stereographic projection (3.247):

$$\left\langle \psi(u)\psi^*(u')\right\rangle = \frac{1}{u-u'+i\varepsilon} \curvearrowright \left\langle \psi_c(z)\psi^*_c(z')\right\rangle = \frac{1}{z-z'(1+\varepsilon)}$$
(3.239)

where the ε -prescription in the last formula expresses the radial ordering. From this one reads off the spatial invariance group. It is the 3-parametric Moebiusgroup SL(2R) in u or SU(1,1) in z. The analytic continuation of this two-point function has a positive definite restriction onto any (Jordan) curve C circulating around zero i. e.

$$(f,g) = \oint_C \oint_C \frac{1}{z(\gamma(t)) - z'(\gamma(t'))} \bar{f}(t)g(t')dtdt'$$
(3.240)

is a (positive) scalar product, leading to a Hilbert space. Since the higher point correlation functions are products of the two-point functions, we obtain positivity. The algebra is still a CAR-algebra, but the quasifree state defined through

C differs from the vacuum state. The unitary equivalence of the representations is easily checked with the Hilbert-Schmidt criterion of the first chapter. This means that the transformation of the circle theory (the "living space" of the original real time theory) to the parametrized curve C theory is unitarily implemented. This emergence of the spatial automorphism associated to the $Diff(S^1)$ group and their unitary implementation from the existence of a family of noncommutative Wightman theories between the original angular circle and the euclidean (radial) cycle is the special property which fails in any other dimension. The infinitesimal manifestation is the well-known Lie-field structure of the energy-momentum tensor, in our case ¹⁰:

$$T(z) = :j^2(z):, \quad j(z) =: \psi^*(z)\psi(z):$$
 (3.241)

$$[j(z_1), j(z_2)] = i\delta'(z_{12}) \tag{3.242}$$

$$[T(z_1), T(z_2)] = -i\frac{1}{12}\delta'''(z_{12}) + i(T(z_1) + T(z_2))\delta'(z_{12})$$

The generic T-algebra is obtained from this special case by replacing the factor $\frac{1}{12}$ by $\frac{c}{12}$ with a positive c which turns out to take on any value above c = 1 and is quantized below this free value. The (nonlocal) Fourier components with respect to the rotation group lead to Kac-Moody and Virasoro algebras. The latter have commutation relations:

$$[T_n, T_m] = (n-m)T_{n+m} + \frac{c}{12}n(n^2 - 1)\delta_{n+m,0}$$
(3.243)

In this form it has been discovered by Virasoro while studying the dual Smatrix model of Veneziano. The local field theoretic version (3.242) on the other hand was found in 1973 (for the early history on low-dimensional models see [5] within two-dimensional QFT while persuing the construction of "Lie Fields", which were introduced in J. H. Lowenstein's thesis in 1968. The latter should be considered as the early version of what nowadays is called W-algebra.

Since all irreducible L-representations are one dimensional, the Lorentz spin s (halfinteger) fields can all be represented in one bosonic (or fermionic) Fock space generated by bosonic (or fermionic) $a^{\#}, b^{\#}$ which are independent of s. It is also easy to see that it is not possible to generalize this to arbitrary L-spin s i.e. to construct fields $\phi(u)$ with anyonic commutation relations:

$$\phi(u)\phi(u') = e^{2\pi i\lambda}\phi(u')\phi(u), \quad u > u'$$

$$\phi(u)\phi^*(u') = e^{-2\pi i\lambda}\phi^*(u')\phi(u), \quad u > u'$$
(3.244)

within the setting of Fourier transforms of creation/annihilation operators. Only anyonic momentum space operators as in (3.227) can be constructed in this way.

The rich world of general chiral QFT begins to open if one realizes the peculiar role of the scalar free field:

$$\varphi(x) = \frac{1}{\sqrt{2\pi}} \int (e^{-ikx}a(k) + h.a.) \frac{dk}{2\omega} = \varphi(t+x) + \varphi(t-x)$$
(3.245)

 $^{^{10}{\}rm The}$ triple dot denotes Wick ordering according to the frequency decomposition of j whereas the double dot refers to Fermion Wick ordering.

Due to the infrared divergence in this representation, the pointlike $\varphi(x)$ does not exist, only $\varphi(f)$ with $\tilde{f}(0) = 0$ defines an operator in Fock space. In order to maintain well defined local generators in Fock space, we consider the infrared finite first derivative $j(u) = \partial_u \varphi(u)$, u=t+x. A simple calculation shows that j can also be obtained as the chiral current of a free fermion field (3.238). Its commutation relations define the abelian current algebra (3.242). We now use the Weyl functor (2.6):

$$W(f) = e^{i \int j(x)f(x)}, \quad W(f)W(g) = e^{i\frac{1}{2}\sigma(f,g)}W(f+g)$$

$$\sigma(f,g) = \frac{1}{2} \int \{f(x)\partial g(x) - g(x)\partial f(x)\} dx \qquad (3.246)$$

In order to make the Möbius-covariance of this algebra manifest, one uses the angular parametrization for the compactified line:

$$u \to z = \frac{i-u}{i+u}$$
 $z = e^{i\varphi}$ (3.247)

In terms of this compact description, the above symplectic form σ becomes:

$$\sigma(f,g) = \int \frac{dz}{2\pi i} f'(z)g(z) = \sum n f_n g_{-n}$$
(3.248)
$$f(z) = \sum_n f_n z^n, \quad f_n^* = f_{-n}$$

Thanks to the aforementioned infrared property which forced us to define the Weyl algebra in terms of currents instead of fields, the symplectic inner product (3.248) is degenerate since it vanishes on constant functions ("zero modes"). These are carried into the center of the abstract C*-algebra which is generated freely from the W's, subject to the Weyl relation (3.246). The center defines an abelian charge algebra and there are continuously many superselected charge sectors obtained by diagonalization of the center.

In order to come to a more interesting situation one must extend the Weyl algebra by a lattice so that the extended algebra is not only characterized by the linear space of functions, but in addition has an underlying lattice. In mathematical terms the linear spaces are extended by "noncommutative tori". In order to allow for sufficient generality, we start from a multi-component abelian current algebra:

$$\left[J^{i}(z_{1}), J^{j}(z_{2})\right] = -g^{ij}\delta'(z_{1}-z_{2}), \quad i,j=1...N$$
(3.249)

where $\delta'(z_1 - z_2)$ is the appropriate circular δ -function:

$$\int \frac{dz'}{2\pi i} f(z')\delta'(z'-z) = -f'(z)$$
(3.250)

The symplectic form which now lives in LV i.e. smooth loops in the Ndimensional vector space V is given by:

$$\sigma(f,g) = \int \frac{dz}{2\pi i} \left\langle f'(z), g(z) \right\rangle \tag{3.251}$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product in V given in terms of the positive definite metric g^{ij} . The Möbius-group acts on LV as:

$$(u(g)f)(z) \equiv f_g(z) := f(g^{-1}z), \quad g \in PSU(1,1)$$
 (3.252)

and leaves σ invariant.

We are interested to classify all positive energy representations.

Theorem 10 Every covariant positive energy representation (π, H_{π}) of the C^{*} algebra \mathcal{U} generated by the Weyl-operators W(f) associated with σ is a direct sum of irreducible ground state representations i.e. $H = \sum_{i} H_{i}, H_{i} = \overline{U\Omega_{i}}, \Omega_{i} = ground$ state in H_{i}

We may recover the current algebra fields (3.249) if we restrict to regular representations i.e. those which are related to states ω fulfilling continuity of the function $\lambda \to \omega(W(\lambda f))$. This suggests the question whether the irreducible components can be created by applying (smeared out) covariant fields to the vacuum i.e. if the net point of view and the more standard field point of view are not only based on the same physical pictures but are even mathematical equivalent. Here we will only quote the result and leave the proof up to the last part on algebraic QFT.

Theorem 11 Every regular ground state representation of the (abelian or nonabelian) current- or the energy momentum tensor- algebra is generated by chargecarrying localized fields. The currents and the energy momentum tensor commute with the charge-carrying fields for noncoalescing points whereas the latter can be chosen in such a way that they obey braid group commutation relations (special case: permutation group) among themselves.

The representation theory of the above multicomponent Weyl-algebra is not very interesting since it processes a continuous set of representations labeled by additive (multicomponent) charges. They are generated by the following localized automorphisms γ_{ρ} [54]:

$$\gamma_{\rho}(W(f)) = e^{i\rho(f)}W(f) \qquad (3.253)$$
$$\rho(f) = \oint \frac{dz}{2\pi i}\rho(z)f(z)$$

Here the N-component function ρ is local with support $\subset S^1$ so that γ_{ρ} acts as the identity if f and ρ have disjoint supports. The total charge $q = \rho(1)$ labels automorphism classes which are "inner equivalent" i.e. for which there are unitaries $u(\sigma, \rho) \in \mathcal{U}$ which intertwine between the two automorphisms:

$$\gamma_{\sigma}(W) = u(\sigma, \rho)\gamma_{\rho}(W)u^{*}(\sigma, \rho) \quad W \in \mathcal{U}$$

$$u(\sigma, \rho) = W(f_{\sigma, \rho}) \quad if'_{\sigma, \rho} = \sigma - \rho$$
(3.254)

These equivalence classes of automorphisms are also referred to as abelian (superselection) sectors of \mathcal{U} . They exhaust the locally generated sectors of \mathcal{U} .

The properties of the automorphism immediately translate into properties of the associated representations $\pi_{\rho}(W) := \pi_0 \cdot \gamma_{\rho}(W)$ where π_0 denotes the vacuum representation. So the charge distribution $\rho(z)$ "measures" the local deviation from the vacuum. The representation formalism is more close to the standard formulation of QFT. Vice versa any representation of \mathcal{U} which deviates only locally from the vacuum (suitably defined) can be shown to allow a representation of the above form in terms of a local automorphism. This is a special case of a general theory, the so-called DHR (Doplicher, Haag, Roberts) theory, which we will meet in the section on algebraic QFT. These representations of the abelian current algebra was essentially known (though in a more conventional field theoretic language which is less precise) already by the author in collaboration with Swieca [56][57].

The C^* -Weyl algebra \mathcal{U} may be used as building blocks of structurally richer and more interesting C^* -algebras. The first step in this direction is the process of extensions by incorporating local sectors into the algebra. The naturalness of the so-called lattice-(or noncommutative torus-)extensions is best understood by looking first at subalgebras of $\pi_0(\mathcal{U})$ belonging to disconnected localization regions:

$$\mathcal{A}_{L}(I_{1} \cup I_{3}) = Alg \{\pi_{0}(W(f)) \mid f \in \mathcal{S}_{L}(I_{1} \cup I_{3})\}$$

$$\mathcal{S}_{L}(I_{1} \cup I_{3}) = \{f \in \mathcal{S} \mid f = const. in I_{2}, I_{4}; f(z_{2}) - f(z_{4}) \in 2\pi L\}$$
(3.255)

Here Alg stands for the generated von Neumann algebra, S is the Schwartz space of smooth test functions on the circle, $z_{2,4}$ are two arbitrary points from $I_{2,4}$ and L is an even lattice in V. The commutant of this operator algebra acting on the vacuum Hilbert space is not, as one could expect by a naive application of Haag duality equal to $A_L(I_2 \cup I_4)$, but it rather equals the bigger algebra [98]:

$$A_L(I_1 \cup I_3)' = A_{L^*}(I_2 \cup I_4)$$

$$L^* = dual \ of \ L$$
(3.256)

The reason for this state of affairs becomes clearer if one looks at the physical interpretation of these algebras. The $I_1 \cup I_3$ localized algebra contains, in addition to the naively expected operators which are separately neutral in I_1 and I_3 (zero values of f in $I_{2,4}$), also operators which are only globally neutral but locally charged with I_3 containing the compensating (anti-)charge to that in I_1 . The dual charge (being described by the dual lattice $L^* = V/L$) consists precisely of those values which lead to relative local commutativity:

$$W(f)W(g) = e^{i\sigma(f,g)}W(g)W(f) \quad f \in \mathcal{S}(I_1 \cup I_3), g \in \mathcal{S}(I_2 \cup I_4)(3.257)$$

$$\sigma(f,g) = 2\pi \langle l \cdot .l^* \rangle = 2\pi \cdot integer$$

The existence of these dual subalgebras of the vacuum representation of \mathcal{U} suggests to look for extensions of \mathcal{U} by lattices in V. For this purpose it is convenient to introduce homogeneous charge-transfer operators $\Gamma_{\alpha}, \alpha \in V$ in a subspace

 $H_L \in H_{uni}$ defined in the following. H_{uni} is the (nonseparable) universal representation space which is simply the direct sum of all charged q representations for all real values of q. $H_L \in H_{uni}$ contains only those charges lying on the L-lattice and Γ_{α} creates a charge $\alpha \in L$. As the charge q representation we simply take for H_q a copy of the vacuum representation Hilbert space H_0 but with \mathcal{U} acting through $\pi_0(\gamma_a(W))$. Hence the universal representation is:

$$(\pi_{uni}(W)\phi)_{\alpha} = \pi_0(\gamma_{\rho}(W))\phi_{\alpha} \quad \phi_{\alpha} \in H_0$$

$$(\Gamma_{\alpha}\phi)_{\beta} = \phi_{\beta-\alpha}$$

$$(3.258)$$

and the restriction to L means that this formula is restricted to $H_L = \sum_{\alpha \in L} H_\alpha$ i.e. all charge indices $\alpha, \beta \in L$. In particular the vacuum considered as a vector in $H_{\beta-\alpha}$ is mapped into the vacuum but this time considered as a vector in H_β . We will denote the universal representation restricted to H_L as $\hat{\pi}$. In order to speak about the ground state in each charge sector we need a hamiltonian. In conformal field theory there are two: the time translation and the rigid S^1 rotation generator L_0 . For the present discussion we only need the action of the rigid rotations:

$$(R(\tau)\phi)_{\alpha} := e^{\frac{i}{2}\langle\alpha,\alpha\rangle\tau} e^{iL_0\tau}\phi_{\alpha} \tag{3.259}$$

Then ground states are mapped into ground states and Γ_{α} commutes with $R(\tau)$ (rotational homogeneity) and the ground state energy in the sector α is $\frac{1}{2} \langle \alpha, \alpha \rangle \cdot \Gamma_{\alpha}$ implements a nonlocalized automorphism:

$$\Gamma^* \hat{\pi}(W) \Gamma = \hat{\pi}(\gamma_{\alpha}(W)) \quad \gamma_a(W(f)) = e^{i\langle\alpha, f_0\rangle} W(f) \qquad (3.260)$$
$$f_0 = \oint \frac{dz}{2\pi i z} f(z) \ i.e. \ \rho_{\alpha}(z) = \frac{1}{z} \alpha$$

Localized charge carrying operators in the same charge class with prescribed support properties for $\rho_{\alpha}(z)$ may be obtained by modifying Γ_{α} with a Weyl operator:

$$\psi_{\rho_{\alpha}}^{\zeta} = \eta_{\xi}(\rho_{\alpha})\hat{\pi}(W(\bar{\rho}_{\alpha})\Gamma_{\alpha}$$
(3.261)

One easily checks that the necessary test function $\bar{\rho}_{\alpha}$ solves the first order differential equation:

$$\frac{d}{dz}\bar{\rho}_{\alpha}(z) = i(\rho_{\alpha}(z) - \frac{\alpha}{z})$$

$$\bar{\rho}_{\alpha}(z) = i\sum_{n\neq 0} (\rho_{\alpha})_{-n} \frac{z^{n}}{n} - i\int \frac{dz}{2\pi i} \rho_{\alpha}(z) ln_{\zeta}(z)$$
(3.262)

Here $\zeta \in S^1$ denotes the direction of the cut along the line $\{\lambda \zeta \mid \lambda \geq 0\}$ which is necessary in order to define the branches of the logarithm. Remember that $\zeta = -1$ corresponds to infinity in the original **R**-parametrization. Here and in the following statement the lattice restriction is not yet needed, everything holds within the universal representation for an arbitrary charge α . The numerical factor η will be determined later. One finds: **Theorem 12** If $\zeta \notin supp \rho$ the charged operators $\psi_{\rho_{\alpha}}^{\zeta}(3.261)$ are local with respect to the observables U and fulfill the following relations:

$$\begin{array}{lll} (i) \quad \psi_{\rho_{\alpha}}^{\zeta} \psi_{\rho_{\beta}}^{\zeta} &= e^{\pm i\pi \langle \alpha, \beta \rangle} \psi_{\rho_{\beta}}^{\zeta} \psi_{\rho_{\alpha}}^{\zeta}, \quad if \; supp\rho_{\alpha} \cap supp\rho_{\beta} = \emptyset(3.263) \\ (ii) \quad \psi_{\rho}^{\zeta_{1}} (\psi_{\rho}^{\zeta_{2}})^{*} &= e^{-\sigma i\pi \langle \alpha_{\rho}, \alpha_{\rho} \rangle} e^{2\pi i \langle Q, \alpha \rangle}, \quad if \; supp\rho \subset S^{1} \setminus \{\zeta_{1}, \zeta_{2}\} \\ (iii) \quad R(\tau) \psi_{\rho}^{\zeta} R^{*}(\tau) &= \psi_{r(\tau)\rho}^{\zeta}, \quad if \; \eta_{\zeta} = e^{\frac{i}{2} \int \frac{dz}{2\pi i} \langle \alpha_{\rho}, \rho \rangle ln_{\zeta}(z)} \end{array}$$

The sign in (i) is coupled to the orientation of the path going from I_1 to I_2 through ζ . The σ in (ii) denotes $0, \pm 1$ according to whether the path which connects ζ_1 with ζ_2 and runs through $\operatorname{supp}\rho$ in addition runs through -1 ($\sigma = 0$) or not ($\sigma = \pm 1$, depending on the \pm orientation). The charge measuring operator Q is defined by:

$$(\langle Q, \beta \rangle \Phi)_{\alpha} := \langle \alpha, \beta \rangle \Phi_{\alpha} \tag{3.264}$$

The theorem is easily verified by explicit calculation.

Property (i) tells us that bosonic local fields correspond precisely to even lattices:

$$\langle \alpha, \beta \rangle = 2n, \quad n = 0, \pm 1, \pm 2, \dots$$
 (3.265)

Restricting to such lattices L, the right hand side of (iii) applied to H_L is equal to one and therefore independent of ζ i.e. those fields live on S^1 (and not a covering thereof).

Now we change our standpoint by and consider the von Neumann algebra \mathcal{A}_L generated by the extended operators together with its new (neutral) vacuum and the representation space H_L . It turns out that the new net \mathcal{A}_L has only finitely many additional positive energy representations. They are labelled by points on the dual lattice $L^* \supset L$ modulo L i.e. in L^*/L . Lattices with $L^* = L$ are called selfdual. They only have one sector (the vacuum sector) and they fulfill the "split" Haag duality:

$$\mathcal{A}(I_1 \cup I_3) = \mathcal{A}(I_2 \cup I_4)' \tag{3.266}$$

A famous illustration is the root lattice of E_8 as well as the Leech lattice Λ_{24} .

The charge sectors of \mathcal{A}_L corresponding to the abelian group L^*/L can again be described in a manner similar to the previous formula:

$$\phi_{\rho_{\alpha}}^{\zeta} := e^{i\pi(Q,\nu_{\alpha})}\psi_{\rho_{\alpha}}^{\zeta}|_{H_{L^{*}}} \quad \nu_{\alpha} := \lambda_{\alpha} + \sum_{\beta} \langle \lambda_{\beta}, \mu_{\alpha} \rangle \lambda_{\beta} \quad (3.267)$$
$$\mu_{\alpha} := \sum_{i=1}^{N-1} \langle \alpha, \alpha_{i} \rangle \lambda_{i} \quad i = 2...N, \quad \mu_{1} = 0$$

i.e. ν_a is a linear combination of the dual lattice basis vectors $\lambda_j : \langle \alpha_i, \lambda_j \rangle = \delta_{ij}$ i, j = 1...N. The first factor in ϕ is a Klein factor which plays a similar role as previously namely it adjusts certain commutation relations to standard form, in this case relative commutation with the observables \mathcal{A}_L for disjoint

localization. Again the unitary $\phi's$ implement localized automorphisms. It is easy to see that these sectors exhaust the possibilities of finite energy sectors. The restriction to L^* -charges results from the requirement that the action of L^* exhaust the possibility of leaving the set of L- charges invariant. The phenomenon of charge quantization by charge extension is a special case of the very general phenomenon of decrease in the number of superselection sectors with increasing size of algebraic extension. Note that the generators of \mathcal{A}_L are in physical terms loops which close modulo $2\pi L$ (and hence lead to univalued phase factors) instead of the "Weyl loops" f in W(f) for which f is strictly periodic. Mathematically they consist of "affine Hilbert spaces" i.e. multicomponent functions on the interval $[0, 2\pi]$ which fulfill lattice boundary conditions i.e. a combination of two well studied objects: Weyl algebras over vector spaces with a (possibly degenerate) symplectic form and Weyl-like algebras over (not necessarily even) lattices. The noncommutative tori of the mathematicians as well as the external magnetic field problems of Hofstedter are illustration of the latter. Whereas the von Neumann uniqueness applies to regular representations of the Weyl algebras over finite dimensional space with a nondegenerate σ , the tori algebras are never simple and therefore have several representations.

Having constructed all the charge sectors of the extended observable algebra \mathcal{A}_L one may consider the field algebra \mathcal{F}_{L^*} generated by all the charge carrying fields (3.267). It is easy to establish the following theorem [60].

Theorem 13 $\mathcal{F}_{L^*}(I) = \mathcal{F}_{L^*}(I')^{tw}$

Here the twist tw is a generalization of the fermionic twisted commutant. As in that case one must "twist" the von Neumann commutant with a Klein transformation which also in this case is a "square root" of the unitary operator which represents the 2π rotation $e^{-i2\pi L_o}$ [60]. This deviation of quantum physics and geometry increases with increasing amount of non-commutativity (from Fermions to Plektons) and naturally also makes the modular theory of e.g. anyonic field algebras for the wedge regions (in chiral conformal theories just intervals) less geometric than that of the observable algebras.

Besides the above extensions there is one other mechanism namely that of factorizing the observable net \mathcal{U} by a subgroup of its symmetry group. In the case of one current there is just the charge conjugation: $j \to -j$, whereas in the multicomponent case there are more possibilities. One finds new representations for the fixed point algebras. Some of these representations are not extendable to representations of the original $\mathcal{U}(S^1)$ but only to the noncompact $\mathcal{U}(R)$. These are called soliton representation because their charge distributions behave differently for $x \to \pm \infty$. If one prefers vague analogies to differential geometry to concrete expressions from physics, one may also call them "orbifolds".

The crucial remaining question is whether there exists a purely field theoretical systematics which also leads to the more interesting representations of algebras in which the charge sectors have branched fusion laws (as current algebras associated to nonabelian groups and W-algebras). The characteristic feature of those algebras is that they have representations π with nontrivial statistical (or quantum) dimensions $d_{\pi} > 1$ and endomorphisms instead of automorphisms. Since both concepts are far removed from standard QFT (Lagrangians etc.), their explanation will be postponed to the last chapter. Here we will only sketch how by "amplification" and "reduction" one may get away from the lattice-extended Weyl algebras.

By amplification we mean tensor products and in particular our interest is to study nets formed by diagonal tensor products of extended observable algebras:

$$\Psi^{(k)}(\rho) = \psi(\rho) \times \psi(\rho) \times \dots \psi(\rho), \quad \alpha_{\rho} \in L$$
(3.268)

where the tensor factors are of the form (3.261) which we now write as $\psi(\rho)$. If we would follow the logic of loop-groups, we would chose L = root lattice of e.g. SU(n) and $exp\rho \in \text{loop-group}$. Technically speaking one is dealing with a tensor product of k level one loop-group representations. It is well known that by reduction one obtains the higher level representations (with nontrivial branching laws) of the loop-group. There are also arguments by which Walgebras are related with current algebras through an invariance principle.

On the other hand a classification of admissible statistics by methods of algebraic QFT (exchange algebras with braid-group commutation relations) leads to 4-point functions which in simple cases exactly match the two families of current- and W-algebras [82]. This strongly suggests that the two families and possibly additional models with higher Virasoro c-values modulo 8 (see chapter 7 section 4) exhaust the possible plektonic (nonabelian braid group) commutation relations with finitely many sectors ("rational"). A direct proof that the amplification and reduction procedure leads to a family of irreducible nets, among which the nets with a finite number of plektonic charge sectors (rational theories) are exhausted by nonabelian current algebras and W-algebras, is still missing.

3.11 Counting of Localized Degrees of Freedom

In QM one finds that one degree of freedom occupies a phase space volume of $(2\pi)^3$. This result simply follows from the discretization of momentum space by enclosing the system in a box. Interactions modify this result somewhat but the number of degrees in a finite phase space cell remains always finite. As a consequence of the correct relativistic modular localization, the number of degrees of freedom in a e.g. double cone region \mathcal{O} with energy below E turns out to be infinite but "almost finite". Mathematically the following map $\Theta_{\Omega,E}$: $\mathcal{A}(\mathcal{O}) \to \mathcal{H}$ is compact:

$$\Theta_{\Omega,E}(A) = P_E A\Omega, \ A \in \mathcal{A}(\mathcal{O}) \tag{3.269}$$

Here P_E denotes the projector onto the Fock space \mathcal{H} below E and compact means that the unit ball of the algebra $\mathcal{A}(\mathcal{O})$ is mapped into a compact set of vectors, i.e. one for which each infinite sequence contains a convergent subsequence. Without loss of physical insight we may take $\mathcal{A}(\mathcal{O})$ to be the previously constructed free field algebra with $\mathcal{A}(\mathcal{O})\Omega$ the dense modular localization space. This statement we owe to the far sightedness of Haag and Swieca [3], who started the issue of counting of degrees of freedom in phase space of local quantum physics way back in the early 60^{ies} . The reason why the number was not finite but rather compact was precisely the mentioned denseness (the Reeh-Schlieder property of local quantum physics) of localized states. The Newton-Wigner localization (which is the localization of QM translated into QFT) would give finiteness, but it is inappropriate because it is not Einstein-causal. Actually the estimate of Haag and Swieca is unnecessarily conservative; by using better estimates one finds that the above sets of vectors are not only compact but also nuclear [3].

It turns out that there is an easier formulation of this physical property if one does not use a sharp energy cutoff but takes a hint from thermal aspects of either the heat bath-or the modular localization physics. The following maps $\Theta: \mathcal{A}(\mathcal{O}) \to \mathcal{H}$ turn out to be nuclear:

$$\Theta_{\Omega,\beta}(A) = e^{-\beta H} A\Omega, \ A \in \mathcal{A}(\mathcal{O})$$
(3.270)

137

$$\Theta_{\Omega,\Delta}(A)\Omega = \Delta_{\hat{O}}^{\frac{1}{4}}A\Omega, \ A \in \mathcal{A}(\mathcal{O})$$
(3.271)

Here $\hat{\mathcal{O}}$ is a region which contains \mathcal{O} and a collar around it; the damping modular factor $\Delta_{\hat{\mathcal{O}}}$ refers to the modular operator corresponding to $(\mathcal{A}(\hat{\mathcal{O}}), \Omega)$. Actually the two formulations are related to each other but for their conversion one needs the more detailed concept of index of nuclearity. Equivalently the following set of vectors is nuclear:

$$\mathcal{N}(O,\beta) = \left\{ e^{-\beta H} U\Omega : U \in \mathcal{A}(\mathcal{O}), U^*U = 1 \right\} \text{ is nuclear set}$$
(3.272)

A subset \mathcal{N} of a Hilbert space H is called a nuclear set if there exists a sequence of unit vectors $\phi_n \in H$ which span H and linear functionals l_n such that:

$$\sum_{n=1}^{\infty} \sup \left\{ |l_n(\psi)| : \psi \in \mathcal{N} \right\} < \infty$$

$$\sum_{n=1}^{\infty} l_n(\psi) \cdot \phi_n = \psi$$
(3.273)

The nuclear index is then defined as:

$$\nu(\mathcal{N}) = \inf \sum_{n=1}^{\infty} \sup \left\{ |l_n(\psi)| : \psi \in \mathcal{N} \right\}$$
(3.274)

On the basis of a naive picture which ignores the "fuzziness" generated by the Gibbs factor in front of the localized vectors one would identify these sets with those of a canonical ensemble occupying a box at temperature β^{-1} and hence expects the result:

$$\nu(\mathcal{N}(\mathcal{O}_r,\beta)) \le e^{const.(r/\beta)^3} \tag{3.275}$$

This naive expectation is indeed what is borne out by the explicit calculation below.. Remember that the zero temperature counting gave an almost finite (nuclear) instead of the naive finite degrees per phase space cell result in Schrödinger theory.

Using the Weyl functor, the nuclearity may be obtained from a corresponding property of suitably damped localization subspaces of the Wigner space. A slight complication is caused by the fact that the map from Wigner space to Fock space looses some of the functorial properties¹¹. The easy part is the nuclear estimate for the $e^{-\beta h}$ -damped localized wave functions:

$$e^{-\beta h} H_R^{(1)}(\mathcal{O}) \qquad is \ nuclear \ set \qquad (3.276)$$
$$H_R^{(1)}(\mathcal{O}) = unit \ ball \ in \ H_R(\mathcal{O})$$

An equivalent charaterization in terms of a trace class operator in Wigner space is:

$$\left\| E(\mathcal{O})\varepsilon^{-\beta h} \right\|_1 < \infty \tag{3.277}$$

Here $E(\mathcal{O})$ is the projector onto the real localization subspace $H_R(\mathcal{O})$ and the subscript denotes the trace norm. For the nuclearity property we have to prove the following two theorems:

Theorem 14 The previously defined operator has the following properties $(r \ge 1)$:

$$\left\| E(\mathcal{O})e^{-\beta h} \right\| < 1 \tag{3.278}$$

$$\left\| E(\mathcal{O})e^{-\beta h} \right\|_{1} < c(r/\beta)^{3}$$
 (3.279)

This is the input for the following theorem:

Theorem 15 Let $e^{it\gamma}$ be a unitary group in Wigner space which commutes with the TCP-operator and which in the standard way leads to a unitary operator e^{itG} on H_{Fock} which leaves the vacuum vector invariant. Then the above boundedness and trace class properties with $\gamma = \beta h$ are sufficient to establish the nuclearity of the set (3.272) with the nuclear index bounded by

$$v(\mathcal{O},\beta) \le \det(1-E(\mathcal{O})e^{-\gamma})^{-2}$$

Inserting the estimate (3.279) we obtain (3.275).

 $^{^{11}\}mathrm{e.g.}$ the intersection of localization spaces is much bigger than the localization space of the intersected region.

Similar estimates hold if one uses the modular damping with $\Delta_{\hat{O}}^{\frac{1}{4}}$. In fact the nuclearity property and the nuclear index of both versions are related. As in the case of the nonrelativistic counting of degrees of freedom nuclearity is expected to be a stable property in the presence of interactions with a possible change of the nuclear index. In these notes we require nuclearity as a restriction on the interaction. Grossly unphysical theories as those which posses exponentially increasing energy level densities and lead to pathological thermodynamic behavior (finite [Hagedorn] limiting temperature) are eliminated by the nuclearity requirement. One also expects this property to play an important role in the understanding of asymptotic completeness. We believe that the nonperturbative interactions based on modular localization of chapter 6 fulfill this requirement.

3.12 Split Property, Taming of Vacuum Fluctuations

The physical problems which gave rise to the split property reach back to the beginnings of QFT when Heisenberg observed that the QFT vacuum behaves very differently to nonrelativistic ground states with respect to local charges obtained by integrating conserved currents over a finite region. In contemporary terminology such a partial charge leads to infinite particle-antiparticle fluctuations at a sharp spatial boundary. Only if one allows a smoothened decrease inside a space-time collar around the localization region of the charge, one is able to control these vacuum fluctuations. In algebraic QFT this picture is the intuitive germ of a powerful concept, the "Split Property". We will come back to it in the later chapter on algebraic QFT. In its most practical version it states that for two spacelike separated regions \mathcal{O}_i such that they allow spacelike collars, the von Neumann algebra generated by $\mathcal{A}(\mathcal{O}_i), i = 1, 2$ is unitarily equivalent to the tensor product:

$$\operatorname{alg}(\mathcal{A}(\mathcal{O}_1) \lor \mathcal{A}(\mathcal{O}_2) \simeq \mathcal{A}(\mathcal{O}_1) \overline{\otimes} \mathcal{A}(\mathcal{O}_2)$$
(3.280)

If the two regions touch each other (examples: a wedge and its geometric opposite, a double cone and its outside spacelike complement), this property is definitely violated. Physically this is blamed on the lack of control of fluctuations near the common boundary. The statistical independence expressed by the right hand side of (3.280) is interpreted as the result of control of fluctuations thanks to the presence of a collar. It is well known that a boundary in Schrödinger QM (box-quantization) leads to a split in an inside and outside Hilbert space with the interaction causing cross contributions. The orthogonal sum of spaces in the conversion to the multiparticle Fock space (2^{nd} quantization) leads to ((anti)symmetrized) tensor products of two nonrelativistic Fock spaces together with an associated tensor product factorization of the nonrelativistic observable algebras \mathcal{A} :

$$H_F = H_F^{inside} \bar{\otimes} H_F^{outside} \tag{3.281}$$

$$\mathcal{L}(H_F) = \mathcal{A}^{inside} \bar{\otimes} \mathcal{A}^{outside}$$
$$\mathcal{A}^{outside} = (\mathcal{A}^{inside})'$$

This is not what happens in the relativistic case of the causal (modular) localization. Although in that case the algebra and its commutant still generate the algebra of all operators $\mathcal{L}(H_F)$, they are of hyperfinite type III₁ instead of the above quantum mechanical type I. Whereas type I factors are similar to $\mathcal{L}(H_F)$ in that they admit pure states or maximal measurements, type III only have impure states, a fact which is related to the thermal nature of modular localized states. One cannot ignore these subtle properties of local quantum physics without impunity. As an example of conceptual havoc, we cite some wrong statements claiming that Fermi's conclusion that the limiting velocity in QFT is c (as in its classical counterpart) is false [99] and that instead QFT allows for causality violations (and time machines). The mistake can be traced back to the incorrect implicit assumption that the localized algebras behave as type I factors [100]. Although experts have gotten tired to refute the never ending stories about superluminal velocities (with ot without tunelling), one may safely assume that they all suffer from the same conceptual flaw.

An equivalent but mathematically more natural definition of the split property is the following:

Definition 3 An inclusion $\mathcal{A} \subset \mathcal{B}$ is called split if there is a type I factor \mathcal{M} with:

$$\mathcal{A} \subset \mathcal{M} \subset \mathcal{B} \tag{3.282}$$

In that case $\mathcal{A} \subset \mathcal{M}$ and $\mathcal{B}' \subset \mathcal{M}'$ together with $\mathcal{L}(H_F) = \mathcal{M} \bar{\otimes} \mathcal{M}'$ results in the existence of an isomorphism Φ of $\mathcal{A} \vee \mathcal{B}'$ with $\mathcal{A} \bar{\otimes} \mathcal{B}'$:

$$\Phi(ab') = a\bar{\otimes}b' \in \mathcal{M}\bar{\otimes}\mathcal{M}' \tag{3.283}$$

In the case of interest $\mathcal{A} = \mathcal{A}(\mathcal{O}), \mathcal{B} = \mathcal{A}(\widehat{\mathcal{O}}), \mathcal{O} \subset \widehat{\mathcal{O}}$ we have more structure thanks to the fact that these two algebras as well as their collar algebra $\mathcal{A}' \cap \mathcal{B} \supset \mathcal{A}(\mathcal{O}' \cap \widehat{\mathcal{O}})$ are "standard" with respect to the vacuum vector Ω (cyclic+separating from the Reeh-Schlieder property). In this case the inclusion $\Lambda \equiv (\mathcal{A} \subset \mathcal{B}, \Omega)$ is called standard split. For such standard split inclusions the product state:

$$\phi_{\Lambda}(ab') \equiv (\Omega, a\Omega) \left(\Omega, b'\Omega\right) \tag{3.284}$$

has a normal extension to the v.N. algebra $\mathcal{A} \vee \mathcal{B}'$. Using the natural selfdual cone $\mathcal{P}_{\mathcal{A}'\cap\mathcal{B}}$ one obtains a representation in terms of a unique vector $\Omega_{\Lambda} \in \mathcal{P}_{\mathcal{A}'\cap\mathcal{B}} \subset H_F$ with:

$$\phi_{\Lambda}(a) = (\Omega_{\Lambda}, a\Omega_{\Lambda}), \ a \in \mathcal{A} \vee \mathcal{B}'$$
(3.285)

In this situation the above isomorphism has a unique unitary implementation $U_{\Lambda}: H_F \to H_F \bar{\otimes} H_F$

$$U_{\Lambda}ab'\Omega_{\Lambda} = a\Omega\bar{\otimes}b'\Omega \tag{3.286}$$

This leads to a canonical interpolating type I factor \mathcal{N}

$$\mathcal{N} \equiv U^*_{\Lambda}(L(H_F)\bar{\otimes}\underline{1})U_{\Lambda} \tag{3.287}$$

This quantum mechanical like subalgebra has of course pure states. The reader should be aware that the (im)purity of a state only has meaning with respect to an algebra. The unitary map U_{Λ} is called the *universal localizing map* and it has some interesting physical consequences. Among other things it leads to a purely intrinsic version of localized symmetry transformations [3] and represents a kind of local quantum physical version of Noether's theorem without obstructions from vacuum fluctuations. One simply takes the global symmetry operator $U(g), g \in G$ and defines:

$$U_{\Lambda}(g) = U_{\Lambda}^{*}(U(g)\bar{\otimes}\underline{1})U_{\Lambda}$$
(3.288)

It then follows that this operator is attenuated within the collar:

$$U_{\Lambda}(g) \in \mathcal{B}, \qquad (3.289)$$

i.e. $adU_{\Lambda}(g)b' = b'$
 $adU_{\Lambda}(g)a = adU(g)a$

This construction works for internal as well as for space-time symmetries.

In free field theories one can go beyond the mere existence and perform explicit computations of the localizing map. In case of zero mass, the geometrical nature of the modular theory of double cones simplifies the calculation. We defer such a calculation to the appendix.

3.13 Problems with Entropy

Let us add some speculative remarks on entropy. As the thermal aspect of modular localization has its characteristic properties which distinguish it somewhat from its standard heat bath setting, one expects also some peculiarities for the entropy of modular localization. By this we mean an entropy concept which carries the (quasi)classical observation of Bekenstein and its refinement by Hawking with its natural (God-given) black-hole horizons deeply in to the noncommutative world of modular structures of hyperfinite type III_1 localized factors (without Killing horizons). Whereas for type I factors we have von Neumann's definition of entropy and (in the thermodynamic limit) entropy-density (also relative entropy between different states on the same algebra), there is no direct definition [11] in the hyperfinite type III_1 case. Following the previous logic, we should start with e.g. two concentric double cones in order to create a collar for the intermediate type I factor \mathcal{N} (3.287). The global vacuum state restricted to \mathcal{N} is a thermal state Ω relative to the factorizing ground state ϕ_{Λ} . One expects Ω to have a restriction on \mathcal{N} which can be described by a well behaved density operator $D = e^{-K}$, where K is related to the modular theory of the pair (\mathcal{N}, Ω) and should be interpreted as a kind of regularized version of the modular theory of the smaller double cone $(\mathcal{A}(\mathcal{O}), \Omega)$ which in the massless case would be geometric and represented by a one parametric subgroup of the 16-parametric conformal group.

One expects that the K-Hamiltonian is in fact well enough behaved in order to allow for the existence of the von Neumann entropy:

$$S = -trDlnD \tag{3.290}$$

where $D = \frac{1}{tre^{-K}}e^{-K}$ is the density matrix defined in terms of the modular Hamiltonian K. This is a quantity which depends on the size of the collar ε and which diverges as $\varepsilon \to 0$ i.e. when the fuzzy type I factor becomes hyperfinite type III₁. If the result of the existing proposals [13] is compatible with this idea, we should expect a universal logarithmic divergence in the inverse size ε^{-1} of the collar which controls the vacuum fluctuations:

$$S = -trDlnD \sim Cln\varepsilon^{-1} \tag{3.291}$$

with C related to the longitudinal 2-dim. conformal theory which according to our previous discussion we expect to determine the geometric core of the fuzzy modular group of the double cone algebra $\mathcal{A}(\mathcal{O})$. Indeed the formula [13]

$$C \sim Area\sqrt{c}$$
 (3.292)

where Area denotes the area of the double cone and c the vacuum fluctuation strength of the energy momentum tensor. Although the limiting entropy is certainly infinite, we have not yet been able to confirm that this infinity is universal and behaves exactly as argued by Larsen and Wilszek [13].

In order to prove this one must do some new computations with modular methods and in particular on the "localizing map" which is the most convenient way to compute the distinguished type I factor [3]. The relevant degrees of freedom would "live", as we will argue later, inside the collar and the ratios of this "collar entropy" stay finite for vanishing collar size. This remains a fascinating program for the future.

Now we are able to formulate our two conjectures:

Conjecture 16 The modular group of the (nonconformal) massive double cone algebra $\mathcal{A}(\mathcal{O})$ with respect to the massive vacuum vector (i.e. the physical vacuum state restricted to $\mathcal{A}(\mathcal{O})$ is cocycle-related to the known geometric modular group of the associated conformally invariant situation belonging to the pair $(\mathcal{A}(\mathcal{O}), \omega_{m=0})$ where $\omega_{m=0}$ denotes the conformal invariant vacuum state. For the equivalence of the massive with the massless algebra one may either invoke the construction of the double cone algebra by canonical quantization or the fact that local algebras are always hyperfinite III₁ factors and the latter is unique modulo unitary equivalence. The cocycle accounts for the difference in the local propagation of massless (Huygens principle) and massive theories and its presence renders the action of the modular group "fuzzy". Only asymptotically near the horizon i.e. the boundary of the double cone, the fuzzyness decreases and the geometric conformal modular transformation reappears. Although a single algebra $\mathcal{A}(\mathcal{O})$ of the massive theory and its scale invariant limit may be identified, the two nets inside \mathcal{O} remain different. However the conjecture that the difference is due to the different propagation suggests that the massive net inside \mathcal{O} may be obtained from the massless $\mathcal{A}(\mathcal{O})$ by adjoining the action of the Poincaré covariances inside \mathcal{O} .

Remark 2 For a massive free Fermi field¹² in d=1+1 this can be shown. One notes that the restriction of such a free massive theory to the light rays which constitute the boundary of the d=1+1 double cone is simply the restriction of the corresponding massless theory and that by propagating the chiral conformal data on the one dimensional horizon inside with the massive propagator, one regains the massive free field net inside O. A general proof of this reduction of a d=1+1situation to its chiral conformal limit (+ possible covariance operators) would be extremely desirable because it would explain the association of the degree of freedoms of d=1+1 theories with the horizon, a phenomenon which has attracted considerable interest and has been termed "holographic behavior" [14].

Conjecture 17 The double cone algebras $\mathcal{A}(\mathcal{O})$ are identical to any of the twodimensional double cone algebras $\mathcal{A}(O^{(2)})$ obtained by cutting the double cone by a two-dimensional plane which contains the t-axis and one coordinate axis. The net inside $O^{(2)}$ may be obtained from the associated chiral conformal net on the one-dimensional horizon and a local representation of Poincaré covariances.

Remark 3 The first part is actually a consequence of Haag duality and the fact that the causal completion of $\mathcal{O}^{(2)}$ gives \mathcal{O} :

$$\mathcal{A}(\mathcal{O}^{(2)}) \equiv \cap_{\delta} \mathcal{A}(\mathcal{O}_{\delta}) \tag{3.293}$$

Where \mathcal{O}_{δ} denotes the middle slice of thickness δ by cutting the double cone parallel to the t-axis. Each \mathcal{O}_{δ} has \mathcal{O} as its causal completion and the property of "Haag Duality" demands the equality of $A(O_{\delta})$ with the algebra of the causal completion $\mathcal{A}(\mathcal{O})$. The essential step in the holographic reduction is the appearance of chiral conformal degrees of freedom after removal of the angular degrees of freedom due to angular symmetry (substituting the transversal symmetry in the case of the wedge). The envisaged entropy is therefore not proportional to area $(\mathcal{O}^{(2)}) \times$ angular volume but rather to horizon-length $(\mathcal{O}^{(2)}) \times$ angular volume = volume of horizon (\mathcal{O}) . Actually such a situation would also suggest that there may be an infinite hidden nongeometric (fuzzy) symmetry algebras in the nonperturbative structure of any QFT^{13} . Although they are local

 $^{^{12}\}mathrm{We}$ want to avoid the infrared problems of massless Bose fields

¹³In principle every modular automorphism has the interpretation of a (localized but fuzzy) physical symmetry. Some of these are "semi-geometric" i.e. they act geometric on subnets. The modular group of the previous modular intersection situation (which led to the transversal Galilei-transformation) is such a case of a "semi-hidden" symmetry [15].

144 CHAPTER 3. POINCARÉ SYMMETRY AND QUANTUM THEORY

in the sense of keeping things inside say \mathcal{O} , their action within \mathcal{O} is totally fuzzy. Such symmetries of nonperturbative local quantum physics would escape differential geometric methods. Note that the two conjectures cannot even be formulated in terms of properties of expectation values of fields; the use of the algebraic i.e. field coordinate independent concepts is indispensable for the formulation. If algebraic QFT did not already exist, one would have to invent it in order to understand the above thermal and entropic properties.

As we have seen, the thermal and entropic aspects which are erroneously attributed *exclusively* to black holes, are in fact a generic nonperturbative feature of the modular localization structure of QFT. They make their appearance e.g. in the formfactor bootstrap construction program (vis.. the KMS origin of crossing symmetry as a generalization of TCP) and also show up in CST QFT for the same (localization) reason. The latter case is only distinguished by the fact that these concepts allow for a classical (thermodynamic) interpretation which is of course the reason why they were first noticed there. This begs the question of the algebraic point of view about "Quantum Gravity". This time we put these words in quotation mark in order to indicate their precarious physical status, especially in the algebraic approach. It is agreed upon by most physicists that Quantum Gravity, whatever it is, does not fit into the framework of local quantum physics as say another spin=2 QFT. Therefore one may ask the more general question of physically consistent theories outside the framework local quantum physics. Surely there have been several attempts to imagine such possibilities, e.g. the pure S-matrix approach, the modifications of Lagrangians by formfactors or structure functions, SO(4) invariant cutoffs in euclidean QFT as candidates of real time relativistic nonlocal theories after analytic continuation and the peratization program (pairs of complex conjugate poles in Feynman rules) of A. Pais and T. D. Lee. If the proposal was not already defeated on the mathematical front, it turned out that the physical interpretation was either inconsistent (existence of precursors violating the indispensable macrocausality) or was not, as in local quantum physics part of the theory, but (as a consequence of the missing localization) had to be enforced from the outside. There is one recent proposal [97] which has survived recent years as a possible scenario (but its future survival is by no means guarantied) which roughly speaking consists in substituting the classical indexing of the algebras in a net by space-time regions in Minkowski space by noncommutative versions in the spirit of noncommutative geometry. Algebraic QFT would favor a situation in which no a priori space-time indexing (neither commutative or noncommutative) appears. Preferably one would like to have global algebra with an intrinsic substructure such that it would contain our physical world of localization and causality only in the germs of certain representations (states). Such an idea would be much more radical¹⁴ than say string theory, because going from pointlike to string extension does not mean that one abandons localization altogether.

¹⁴Note that this is the only radicalism in the algebraic approach since all other deviations from the standard QFT approach, as uncommon and revolutionary as they may appear at first sight, are all in accordance with the physical principles of QFT.
There is one intriguing property in chiral conformal QFT which has a certain quantum gravity "touch" to it. This is the fact that the "averaging over a Fuchsian group" of a chiral conformal QFT (assuming that the Poincaré series converge in some sense) converts the Moebius invariant vacuum expectation values into expectation values which loose this invariance but gain deformation parameters (generalizations of "compact temperatures"). Formally the positivity property holds as in Wightman theory, but the old localization region S^1 is now totally fuzzy. With other words there is no causal complement in the quantum sense. In such a scenario there would be no global concept (a priori knowledge of what is spacelike) of causality and hence of localization, and the place to find the lost net properties would be in certain states and even there they would only appear in their germs. Our second conjecture which led via modular theory to the speculative existence of a hidden "fuzzy" realization of the Moebius group would suggest that such a scenario may also be possible in d=1+3 theories.

We have made these extremely speculative remarks about quantum gravitation in connection with entropy, because we share with B. Kay [12] the believe that (holographic) entropy and quantum gravity belong to the same circle of problems. We are attracted by his idea of quantum gravitational degrees of freedom which have (apart from their classical long range tails) no observable aspect (i.e. no localization) in line with the previous conformal scenario, but we admit that we have not been able to reconcile the remaining different viewpoints.. It is interesting that both ideas are very removed from Weinberg's picture of quantum gravity as just a s=2 Lagrangian QFT.

Summing up our excursion on nonperturbative QFT we would like to stress again that the algebraic method allows for a completely intrinsic definition and understanding of QFT independent of its Lagrangian or non Lagrangian origin. Any quantum theory which fulfills the stability requirements of positive energy and allows for a net interpretation and the associated localization concepts is a QFT par excellence and enjoys all the general structural properties which feature in this article as TCP, spin & statistics, crossing symmetry & modular localization & thermality, wedge-localized fields without vacuum polarization, hidden modular symmetries, Haag duality (an abstract form of the 2-d Kramers-Wannier-Kadanoff Duality), nuclearity for the phase space degrees of freedom & the conjectured "Holographic Entropy" and all the other yet unraveled properties of nonperturbative local quantum physics. The main obstacle against progress is not so much the novel mathematics which these new physical concepts require, but rather (as always in the past) prejudices. One prejudice is that field theory has to be "Lagrangian". In view of the many existing lowdimensional non-Lagrangian models and the fact that they hardly rocked the Lagrangian boat, this appears to be the mightiest prejudice.

146 CHAPTER 3. POINCARÉ SYMMETRY AND QUANTUM THEORY

Chapter 4

Perturbative Interactions

4.1 Kinematical Decompositions

Before presenting an elementary approach to interactions and perturbation, it is helpful to have a closer look at those observable quantities which one wants to compute. Since among local "field coordinates" only currents have a distinguished physical meaning, one is naturally interested in matrix elements as:

$$\Gamma_{\mu}(p',p) = \langle p' | j_{\mu}(0) | p \rangle, \quad W_{\mu\nu}(p,x) = \left\langle p \left| j_{\mu}(\frac{x}{2}) j_{\nu}(\frac{-x}{2}) \right| p \right\rangle_{conn.}$$
(4.1)

The first quantity (where possible spin quantum numbers have been suppressed) is called the (electromagnetic) form factor of the p-particle and its static limit $(p - p')^2 \rightarrow 0$ can be probed by external magnetic fields and is related to the (anomalous) magnetic moment.

The second (diagonal) matrix element of two currents with the subscript connected ¹ gives rise to the notion of "structure function" of the p-particle and appears in the description of high-energy electron (more general: lepton) inclusive scattering on nucleons (scattering in which one does not observe the created outgoing hadrons).

Important energy shifts as the Lamb shift cannot be expressed in an elegant form in terms of such matrix elements between particle states (only if one defines "off-shell" extrapolations). These matrix elements between "ket" in-vectors and "bra" out vectors of localized (or multi-localized) fields O(x):

$$^{out} \langle p'_1, ..p'_m | O(0) | p_1, ..p_n \rangle^{in}$$
(4.2)

are referred to as "generalized formfactors". They are the most prominent measurable quantities, but turn out to be also the most important building

¹connected part means removal of the ill-defined vacuum contribution $\langle p' | p \rangle \langle 0 | j_{\mu} j_{\nu} | 0 \rangle$ before the limit p' \rightarrow p, which however does not influence the structure of the covariant decomposition.

blocks of the new constructive approach based on modular localization which will be presented in chapter 6.

The most important experimental observables are the generalized formfactors of the identity operator O(x) = 1 which are identical to the matrix elements of the *S*-matrix (or scattering operator):

$${}^{out}\left\langle p_{1}^{\prime},..p_{m}^{\prime}\left|\frac{1}{-}\right|p_{1},..p_{n}\right\rangle^{in}={}^{in}\left\langle p_{1}^{\prime},p_{2}^{\prime}...p_{m}^{\prime}\left|S\right|p_{1},p_{2}...p_{n}\right\rangle^{in}$$
(4.3)

from which via the momentum space kernel T the cross sections can be obtained:

$${}^{in} \langle p_1', p_2' \dots p_m' | S - 1 | p_1, p_2 \dots p_n \rangle^{in} = \delta(\sum_{i=1}^m p_i' - \sum_{k=1}^n p_k) T(p_1', \dots p_m', p_1, \dots p_n)$$

$$(4.4)$$

Here we used the fact that S is a Poincarė invariant operator in the Fock space of incoming particles (the energy-momentum conserving δ -function results from translation invariance).

Kinematical properties means the decomposition of covariant into invariant functions and the specification of the invariant variables on which the latter depend. For the formfactor of $s = \frac{1}{2}$ particles one finds the following decomposition (with k = p' - p):

$$\Gamma_{\mu}(p',p) = \frac{1}{(2\pi)^3} \bar{u}(p',s_3') \left(\gamma_{\mu}F(k^2) - \frac{i}{2m}(p'+p)G(k^2) + \frac{1}{2m}k_{\mu}H(k^2)\right)u(p,s_3) \quad (4.5)$$

The fastest way to see this is to first use the free field formalism to compute the matrix elements of the free current by "Wick-gymnastics":

$$\left\langle 0 \mid a(p', s_3') : \bar{\psi}(0)\gamma_{\mu}\psi(0) : a^*(p, s_3) \mid 0 \right\rangle = \frac{1}{(2\pi)^3} \bar{u}(p', s_3')\gamma_{\mu}u(p, s_3)$$
(4.6)

Then one has to construct the most general vector object from the γ -matrices and two mass shell momenta p and p' subject to the identity $\gamma_{\mu}p^{\mu} - m = 0$ which is valid on the intertwiner u(p). This leaves besides γ^{μ} itself, which appears already for a free current, only the above two momentum vectors (or linear combinations thereof). Current conservation $k_{\mu}\Gamma^{\mu} = 0$ gives $H \equiv 0$ (because H is a nonsingular function) and the value 1 of the total charge $Q = \int j_0(x) d^3x$ between the one-particle states requires F(0) = 1.

Due to kinematical identities of the u and v intertwiners, there are many different forms of covariant decompositions. For example the identity:

$$\bar{u}(p')i\sigma_{\mu\nu}q^{\nu}u(p) = \bar{u}(p')(2m\gamma_{\mu} + i(p'+p)_{\mu})u(p)$$
(4.7)

may be used to eliminate the γ_{μ} term in favor of the $(p' + p)_{\mu}$ and the $\sigma_{\mu\nu} = \frac{i}{2} [\gamma_{\mu}, \gamma_{\nu}]$ terms:

$$\bar{u}(p')\Gamma_{\mu}u(p) = \bar{u}(p')\{\frac{i}{2m}(p'+p)_{\mu}(F(q^2) + G(q^2)) + \sigma_{\mu\nu}q^{\nu}F(q^2)\}$$
(4.8)

148

4.1. KINEMATICAL DECOMPOSITIONS

In this form the leading contribution for small spatial momenta p and p' comes solely from the second term. The physical interpretation of F (which as G can only depend on k^2 since this is the only invariant which one can form two mass shell vectors) becomes clear if one rewrites the canonical coupling of the current to an external (classical) vector-potential as follows:

$$\left\langle p' \left| -e \int d^3 x \vec{j}(x) \vec{A}(x) \right| p \right\rangle = -e \frac{1}{(2\pi)^3} \int e^{i(\vec{p}-\vec{p}')\vec{x}} \vec{u}(p',s'_3) \vec{\Gamma}(p',p) u(p,s_3) \vec{A}(x) d^3 x d^$$

Here the last line is the static approximation of $\bar{u}\vec{\Gamma}u$ in first order of $\vec{p}\cdot\vec{p}'$ which brings in the angular momentum operator $\vec{J} = \frac{\vec{\sigma}}{2}$ (again just " γ -gymnastics" between *u*-intertwiner). The last step is to use $\vec{B} = \vec{\nabla} \times \vec{A}$ and to take Bconstant (static limit):

$$\left\langle p' \left| -e \int d^3 x \vec{j}(x) \vec{A}(x) \right| p \right\rangle \approx -\frac{e}{2m} F(0) 2p_0 \delta(\vec{p} - \vec{p}') \vec{B} \vec{\sigma}_{s'_3, s_3}$$
(4.10)

i.e. we obtain the magnetic moment interaction $-\vec{\mu} \cdot \vec{B}$ with $(\mu = |\vec{\mu}|)$

$$\mu = \frac{e}{2m} F(0) = \frac{e}{2m} + \mu_{anom.}$$

$$\mu_{anom.} = \frac{e}{2m} (F(0) - 1) = \frac{e}{2m} G(0)$$
(4.11)

In a similar fashion one decomposes the structure function (the form factor of two currents):

$$\int W^{\mu\nu}(p,x)e^{iqx}d^4x \doteq : W^{\mu\nu}(p,q) = -(\frac{q_{\mu}q_{\nu}}{q^2} - g^{\mu\nu})W_1(\nu,q^2) + \frac{1}{m^2}(p^{\mu} - \frac{p \cdot q}{q^2}q^{\mu})(p^{\nu} - \frac{p \cdot q}{q^2}q^{\nu})W_2(\nu,q^2)$$
(4.12)

The invariant structure functions W_i i = 1, 2 depend on two variables (q is offshell) q^2 and $\nu = \frac{p \cdot q}{m}$ (with m=target mass). Again the number of invariants has been reduced by using current conservation for the two currents. The W'sare measurable in processes involving deep inelastic lepton scattering.

The S-matrix is not measured directly, but rather through the ensuing scattering cross sections. The relevant formulas in most textbooks are derived by inventing a "box-quantization" (in order to solve the problem of "squaring the energy-momentum δ -function"). For scattering theory as well as for statistical mechanics of open systems such a trick goes against the basic concepts which require the infinitely extended Minkowski space. Therefore it is comforting to know that that there are suitable concepts and techniques which avoid such tricks in favor of working directly in the infinite volume e.g. the KMS condition in the case of statistical mechanics and techniques of wave packets in scattering theory.

4.2 Perturbative Realization of Interaction

In chapter 2.1 we solved the simple problem of a perturbation by an external source on a free bosonic system and found that there are two methods, one via unitary transformations (the so called "dressing" transformations), and the other by the use of the interaction picture in the form of a *time-ordered* exponential:

$$S(j) = Te^{i\int A(x)j(x)d^{4}x}, \quad A(x,j) = S^{*}(j)TA(x)e^{i\int A(y)j(y)d^{4}y}$$
(4.13)

By Wick-ordering these expressions, we saw that they agree with the dressing transformation method up to a phase (Feynman's famous "vacuum phase") and that phases show up in the form of cocycle factors in the composition law of the S(j).

In translation-invariant interaction problems (i.e. without external fields), there is a well-known obstruction against the existence of such a unitary dressing operator, the *Haag Theorem*. It says that in a globally translation-invariant theory the ground state of an interacting system cannot be described in the space of vector states of the free system [3], but it does not rule out the existence of a local dressing transformation which depends on the localizing region. The traditional way out is to overcome this No-Go theorem against Dirac's interaction picture in QFT is to start with a system which fulfills only "partial translational invariance" (similarly to the notion of "partial charges" in the free field theory in chapter 3.4). We begin by defining (A stands generically for the "would-be" Heisenberg field which corresponds to the free field A_0) :

$$S(g) = Te^{i \int g(y) \mathcal{W}(x) d^4 x}, \quad A(x,g) = S^*(g) TA_0(x) e^{i \int g(y) \mathcal{W}(y) d^4 y}$$
(4.14)
$$= S^*(g) \frac{\delta}{\delta h(x)} S(g,h) \mid_{h=0}, \quad S(g,h) = Te^{i \int \{g(y) \mathcal{W}(x) + A_0(x) h(x)\}} {}^{4}_{4:} 15)$$

 \mathcal{W} is an invariant Wick-ordered polynomial in terms of free fields which implements the notion of interaction via its use as a deformation of free fields. Mathematically it is a Lorentz scalar polynomial of at least trilinear degree in the free field Borchers class of the interacting free fields For g we choose a smooth function with compact support in Minkowski space which can be thought of as a smooth version of the characteristic function of a double cone with support in larger double cone and the constant value $g(x) \equiv g$ in a smaller cone placed inside the bigger one. If W contains several terms, there is one g(x) for each monomial. Before we show some remarkable properties of these formal operators in Fock space, some comments are in order.

• (i) Haag's theorem is not applicable to the S(g) formalism (no translation invariance), and we are allowed to do our calculations in Fock space. One of the remarkable properties is that the local observables localized within the smaller double cone fulfill *partial* invariances (including translations) in an algebraic sense explained later.

- (ii) The standard derivation of the above formula for A(x) (more precisely for the vacuum expectation values of time ordered products of A) goes through the canonical formalism and is known under the name of *Gell-Mann-Low formula*. Such derivations suffer from two conceptual weaknesses. On the one hand they give (physically unmotivated) preference to special field coordinates (only "Eulerian" free fields among the class of (m, s) Wigner fields are fitting into the canonical formalism) and on the other hand they rely on assumption that the fields A to be constructed are not more singular for short distances than the corresponding canonical free fields. These assumptions are only valid in certain very special low-dimensional models. The main culprit for the nonexistence of Fock space operators as above (even not after renormalization) is the use of the interaction picture with its time-ordered formalism². Physical concepts do not require the existence of objects in addition to the Heisenberg fields A and the asymptotic (in- or outgoing) free fields.
- (iii) The interaction density $\mathcal{W}(x)$ is a local function of free fields which (without the existence of a dressing transformation) has no direct (outside an interpretation in terms of infinitesimal deformations without global counterpart) physical interpretation. This means that there is (apart from external perturbations and some very special low dimensional models) no general physical reason to believe that after a certain necessary repair ("renormalization"), one obtains a mathematically well-defined theory. Perturbation theory generally has only a formal meaning as an *infinitesimal deformation theory.* There is no reason whatsoever why the "Bogoliubov axiomatics" i.e. the above scheme of operators S(g) in Fock space should have a solution (apart from the well-known superrenormalizable models as e.g. $P\phi_2$). It is very unfortunate that we use such a questionable framework for the baptization of interacting models. The nonperturbative approach of chapter 6 which aims at the global existence is based on quite different concepts.

Contrary to popular believes, it is not just the singular short distance behavior as such which endangers the existence of the theory, but rather the way interaction is introduced via the Ansatz in terms of a time ordered operator $S(g)^3$ which creates an obstruction against an intrinsic understanding of interactions. There is a very interesting lesson in this respect from the d=1+1 "bootstrap" constructions which show that short distance singularities can be worse than any given inverse power of the Minkowski distance, but without the existence of the theory being threatened. The simplest illustration of the fact that there is no obvious relation between the deviation of the short distance behavior from that of free fields is furnished by the Thirring model, for which

 $^{^{2}}$ Even the spatial cutoff g is generally not sufficient for the existence of the interaction picture or a dressing transformation.

³The so called Bogoliubov axiomatics may have no solution in higher dimensions and therefore the time-ordering method may be not appropriate for introducing interactions. In the nonperturbative approach to low dimensional QFT the time-ordering plays no role.

the singular short distance power can be made arbitrarily large by changing the coupling parameter. If there is at all a general relation between the existence of a QFT associated with $\mathcal{W}(x)$ and perturbative renormalization aspects, then it must be linked to the vanishing of the β -function i.e. the coupling parameter renormalization.

Let \underline{h} be a collection of test functions which couple to observable local fields (electro-magnetic. field strength, currents,..) and let us introduce the following relative Bogoliubov-Shirkov operators (see section 2 of this chapter):

$$V(g,h) = S^{-1}(g)S(g+\underline{h})$$
(4.16)

The causality of the local operators coupled to \underline{h} is then expressed as:

$$V(g, \underline{h_1} + \underline{h_2}) = \begin{cases} V(g, \underline{h_1}) V(g, \underline{h_2}), \ supph_1 \gtrsim supph_2\\ V(g, \underline{h_2}) V(g, \underline{h_1}), \ supph_2 \gtrsim supph_1 \end{cases}$$
(4.17)

i.e. the V factorizes in the second argument if there is no future directed causal curve from $supph_1$ to $supph_2$ or. vice versa. The spacelike consistency of this formula implies in particular Einstein causality. The double cone (=**C**) algebra may than be defined as

$$A_g(\mathbf{C}) = \overline{\{V(g,\underline{h}), \, supph \subset \mathbf{C}\}}$$

$$(4.18)$$

The next step consists of showing that a change of g(x) which maintains the constant value g inside W_{st} leads maximally (if the past "collar" is changed) to a h-independent unitary transformation. However such a common transformation does not change the net of wedge algebras inside W_{st} (by the very definition of isomorphy of nets). By Lorentz transformations one obtains the net of all wedge algebras and by intersecting wedges one finally obtains the full net of observables (which includes the double cones).

Let us now show that on a formal level the Fock space operator A(x,g) fulfills some remarkable formal properties. Suppose that we restrict the x to the double cone K in which $g \equiv 1$ i.e. we consider A(f,g)

$$A(f,g) = \int d^4x f(x) S^*(g) T A_0(x) S(g)$$
(4.19)

with $supp. f \subset K$. Then as a generalization of the composition operator S(j) in our old source model we find:

$$S(g_2 + g_1) = S(g_2)S(g_1), \quad \text{supp}g_2 \ge \text{supp}g_1 A(f,g) = A(f,g'), \quad \text{if } \text{supp}(g - g') \subset V_-(K)^{\perp}$$
(4.20)

where the notation means that the points in supp g_2 are either spacelike or timelike from those in supp g_1 and $V_-(K)^{\perp}$ is the complement of the smallest backward light cone which contains the double cone K. Furthermore any change of g to g' localized in $V_-(K)^{\perp} \setminus K$ can be implemented by a unitary ("partial dressing") transformation U(g) which is independent of f [35], i.e. the same for all operators in the algebra A(K):

$$A(f,g') = U(g',g)A(f,g)U^*(g',g)$$
(4.21)

152

Formally this unitary has the same form as S(h) where the smooth function h is compactly supported in the intersection of $V_{-}(K)^{\perp} \setminus K$ with a double cone K which contains the support of both g's. For the study of the net of double cone algebras localized in \hat{K} inside K the common unitary U is irrelevant since nets which are related by one common unitary are identical (isomorphic families define identical nets by definition) i.e. it is only the relative positions of these algebras and not the absolute position in the ambient space which counts. Hence even the limit $K \to \infty$, the net of algebras may be described within Fock space. Hence this Fock space is purely auxiliary. Physical states strictly speaking are to be obtained as states on the net of operator algebras with suitable localization properties. This would be the scenario for the construction of interacting theories within the setting of time ordered exponential of free field "interaction densities" $\mathcal{W}(x)$.

Before we look at the lowest nontrivial perturbative evaluation of these formal operators, let us briefly notice that A(x, g) fulfills Einstein causality within K:

$$[A(x,g), A(y,g)] = 0 \quad (x-y)^2 < 0 \quad and \quad x, y \in K$$
(4.22)

The formal reason is that for spacelike separations the product can be written in terms of one (cancellations between S's!) time-ordered free field expression:

$$S^{*}(g)T(A_{0}(x)A_{0}(y)S(g)) = S^{*}(g)T(A_{0}(x)S(g)) \cdot S^{*}(g)T(A(y_{0})S(g))$$

holds for $(x - y)^{2} < 0$ (4.23)

(remember: the T only acts on all the A_0 's to the right). The (Bogoliubov-Shirkov, Glaser-Epstein) renormalization approach allows to show that these formal relations are valid at least in every order of perturbation theory (expansion in W). The idea is to reduce the iterative determination of the operator T-products to a (Hahn-Banach type) extension problem of time-ordered vacuum expectations. In the case of renormalizable models this is possible with a finite number of parameters (counter-terms). The issue of whether this latter requirement is more than formal is still unsettled, although beyond the mere fact of computability of higher orders without new parameters there seems to be considerable physical success in this restriction.

Another remark, whose importance can only be fully appreciated later, is the statement that the local algebras of a net are all unitarily equivalent and there is (outside of perturbation theory) no relation between the particle structure of the ambient Fock space and the physical content of local algebras. The interaction generically speaking wrecks the one to one correspondence between particles and fields which existed in the free theory⁴. For local observables described in terms of local nets of algebras, the Hilbert space description allows great flexibility and the chosen massive Fock space of the above formalism is not to be interpreted as a commitment about physical parameters. This picture is unfortunately

 $^{^{4}}$ We are ignoring here the use of interaction polynomials as a mere placeholder for combining phenomenological correlation functions as in "phenomenological" Lagrangians. In those second derivative momentum space Taylor coefficients the particle and field contents are identical by fiat.

somewhat blurred by perturbation theory which maintains an unrealistic rigid correspondence between fields and particles (apart from the mentioned flexibility of choosing the Fock space mass parameter different from the physical mass). This (among other things) has created the misleading impression that QFT is nothing more than a relativistic made form of quantum theory of particles. Although it is a *quantum theory* and it is *relativistic* and its principle physical aim is to describe *particles*, it is primarily a *new physical realm* whose deep and unexpected concepts (despite its 70 years of existence) still await exploration. This will become much more evident in the later chapter on modular localization and the bootstrap-formfactor approach than on the present level of perturbation theory.

In these notes we only address some conceptual points of renormalized perturbation theory. The n^{th} order renormalization technology goes much beyond and should be studied in appropriate review articles by Zimmermann, Lowenstein, Piguet-Sorella [63] and others.

4.3 Perturbation and Adiabatic Parametrization

The naive expectation (i.e. by analogy to the external source problem in chapter 1) about S(g) would be that the limit of the theory for $K \to R^4$ exists and describes the physical S-matrix. Even in perturbative evaluation as an infinitesimal deformation theory with suitably adapted causality and positivity requirements, this picture needs two corrections. One is related to the infrared divergence problem in certain theories involving zero mass as QED, a somewhat special phenomenon whose physical basis will be reserved for a later discussion. The other is of a completely general nature related to the phenomenon of selfinteraction, well-known already from classical field theory where it leads to the famous problems of constructing consistent particle models within a classical field theory, as studied by Poincaré and Lorentz at the beginning of this century. As a result of selfinteraction, parameters with a physical name as mass, charge etc. which entered the construction of S(q) and A(x,q), do not represent the true measured value. Whereas for fields A and their correlation functions this does not matter⁵ (the true physical values can be recovered from asymptotic properties of correlation functions, see later), the large volume limit of S(g)

⁵The reason why in the classical theory the divergencies are not so easily disposed, is that unlike QFT the particle concept does not follow from the Poincaré transformations of the fields but has to be imposed on the classical field theory, a procedure which according to our best knowledge is inconsistent. Hence although Kramers renormalization idea was historically essential, it looses its importance as soon as one realizes, that unlike classical fields, poinlike quantum fields are intrinsically singular objects in the sense of Bohr and Rosenfeld. For those one needs Schwinger's physically motivated point-split form of field equation. Knowing this one, may safely deal with Feynman's simpler rules, as long as one confronts the intermediate infinities created by his formalism without confusing them with those of the classical theory studied by Lorentz and Poincaré. The latter are a physical consequence of forcing classical particles upon a classical field theory.

for $|K| \to \infty$ represents the physical S-matrix for the scattering of A-particles only if the true physical mass is used. The same applies to any quantity which is partially "on-shell" i.e. contains particles states as e.g. the electromagnetic formfactor. The reason is that the adiabatic switching on and off by multiplying $\mathcal{W}(x)$ with $e^{-\alpha|t|}$ and then $\lim t \to \infty$ is physically harmless only if the interaction includes the effect of persistent selfinteraction "counter-terms" which maintain the masses used in the Fock space in every order of \mathcal{W} at their physical value. In case of a neutral scalar $\mathcal{W} = g : A_0(x)^4 :$ model the modification is:

$$\mathcal{W}_{adiab}(x) = \mathcal{W}(x) + \frac{1}{2}\delta m^2 Z : A_0(x)A_0(x) : + \frac{1}{2}(Z-1) : (\partial_\mu A_0(x)\partial^\mu A_0(x) - m^2 A_0(x)A_0(x)) :$$
(4.24)

The "selfmass" δm^2 is chosen in every order to maintain m as the physical mass and $m_0^2 = m^2 - \delta m^2$ is an auxiliary unphysical mass (which loosely speaking corresponds to the mass without the stabilizing counterterm which changes in every order of W). The second Z-counterterm has been added in order to obtain a nicer form of the adiabatic principle which is the following requirement:

$$\lim_{a \to 1} \langle 0 | A(x,g) | p \rangle = \langle 0 | A_0(x) | p \rangle$$
(4.25)

By adjusting δm^2 and Z in every order such that this identity holds we took all selfinteractions into account. A subsequent adiabatic change of \mathcal{W}_{adiab} i.e.

$$\mathcal{W}_{adiab} \to e^{-\alpha|t|} \mathcal{W}_{adiab}, \quad \alpha \to 0 \text{ at end of calculation}$$
(4.26)

will not cause any harm i.e. does not change the one particle characteristics. In theories without selfinteraction e.g. in Schrödinger theory, this is automatically fulfilled. Using our formal time-ordered expressions we may rewrite the above requirement in second order:

$$\frac{(\delta m^2 Z)^{(2)} + (Z-1)^{(2)} (p^2 - m^2)}{p^2 - m^2 + i\varepsilon} = \frac{1}{2} \int \int \langle 0 | TA_0(0) \mathcal{W}(x_1) \mathcal{W}(x_2) | p \rangle \quad (4.27)$$

since the zero order terms agree and the second order term of the above requirement:

$$\lim_{g \to 1} \left\langle 0 \left| A(0,g) \right| p \right\rangle^{(2)} = 0 \tag{4.28}$$

consists of a $W \cdot W$ contribution and the lowest counterterm contribution (which we wrote on the left hand side). The evaluation of the right hand side (omitting combinatorial factors) gives:

$$g^{2} \int \int \Delta_{F}(0-x_{1}) \Delta_{F}^{3}(x_{1}-x_{2}) \frac{e^{ipx_{2}}}{(2\pi)^{\frac{3}{2}}} \simeq \frac{1}{p^{2}-m^{2}+i\varepsilon} \int e^{ip\xi} \Delta_{F}^{3}(\xi) d^{4}\xi \quad (4.29)$$

Therefore $\delta m^{2(2)} g^2 \int e^{ip\xi} \Delta_F^3(\xi) d^4\xi$ and $Z^{(2)}$ is the second Taylor coefficient in the expansion of the integral around $p^2 = m^2$ (note that $Z = 1 + Z^{(2)} + \dots$).

Later we will (via time dependent scattering theory) meet a formalism which relates off- shell quantities (correlation functions of interacting fields) in a natural way with distinguished free fields which have the correct physical mass (the "incoming" (outgoing fields). This relation is independent of the mass of the Fock space which we use for their perturbative construction. Scattering theory may be viewed as an extension of the adiabatic principle to multiparticle states. It is applied to off-shell correlation functions and requires the introduction of the physical mass i.e. a reparametrization which relates the mass of the auxiliary Fock space to that of the in and out Fock space of scattering theory. Since (Haag-Ruelle or LSZ-) scattering theory and (a fortiori) the above adiabatic principle are consequences of the general framework of QFT, a more systematic and conceptually clearer approach would have consisted in presenting the perturbative treatment of interaction after an introduction to the general framework. However a physical minded reader prefers to see some important results and understand them at least partially with a modest amount of mathematics and concepts before making a large formal investment.

Even though there are no physical reasons to introduce counterterms for off-shell quantities, the fact that the time-ordered products of $\mathcal{W}'s$ via Wick's theorem yield ill-defined (formally infinite) expressions as e.g. $i\Delta_F^3(x_1-x_2)$ forces "renormalization" for mathematical reasons. With other words our starting formula in Fock space although mathematically incorrect, was not beyond physical redemption. The integrand in S(g) i.e. $T\mathcal{W}(x_1)...\mathcal{W}(x_n)$, though not defined on all (Schwartz) testfunctions, is well-defined on the big class of testfunctions $f(x_1 \ x_n)$ which vanish of sufficient high order for coalescent points. If \mathcal{W} is a polynomial with $\dim \mathcal{W} \leq 4$, the order does not increase in n. This means that the (Hahn-Banach) extension to all testfunctions will lead to timeordered distributions which, although lacking uniqueness, have well-controlled ambiguities (possibly additional parameters) whose space-time dependence is given in terms of δ -functions (which spread in the next order where new δ functions arise) and derivatives up to a maximal finite order. With other words, different extensions differ by finite local counterterms. These counterterms may be used in order to achieve certain normalization conditions (as in the case of the adiabatic principle), but there is no mathematical necessity to take the ambient Fock space with a mass equal to the physical mass. The mass of a particle does not belong to the set of observables which can be measured locally. Globally however e.g. two free fields with different masses cannot be unitarily equivalent.

In the following we will use the more pedestrian regularization methods rather than the mentioned extension method (the latter will be used in the later *Curved Space Time* problems where it seems to be the *only renormalization method*). Whereas for structural arguments we mostly use the A^4 model, the explicit second order calculations will be done in Quantum Electrodynamics. We now specialize to the \mathcal{W} describing QED⁶ (first without the counterterms):

$$\mathcal{W}(x) = -e j_{0\mu}(x) A^{\mu}(x), \quad j_{0\mu}(x) =: \bar{\psi}_0 \gamma_{\mu} \psi_0(x) : \quad \psi_0, A_{0\mu} \text{ free fields} \quad (4.30)$$

and consider S(g) in second order:

$$S^{(2)}(g) = \frac{e^2}{2!} \int \int g(x)g(y)Tj_{0\mu}(x)j_{0\nu}(y)A_0^{\mu}(x)A_0^{\nu}(y)d^4xd^4y$$
(4.31)

with the Wick-reordering from the previous chapter, we obtain for the formfactor:

$$\langle p' | j_{\mu}(0) | p \rangle^{(2)} = \langle 0 | a(p', s'_{3}) S^{*} T(j_{0\mu}(0)S) a^{*}(p, s_{3}) | 0 \rangle^{(2)}$$

= $\langle 0 | a(p', s'_{3}) T(j_{0\mu}(0)S) a^{*}(p, s_{3}) | 0 \rangle^{(2)}_{v.c.}$ (4.32)

"vacuum-connected" (v.c.) has the same meaning as before: leave out the S^* in front of the T and ignore the vacuum "bubble" contributions. The evaluation of the right hand side amounts to look for the $\bar{\psi} - \psi$ contribution in the Wick reordering of:

$$T j_{0\mu}(0) j_{0\nu}(x_1) j_{0\kappa}(x_2) A_0^{\nu}(x_1) A_0^{\kappa}(x_2)$$
(4.33)

In order to keep track of the combinatorial possibilities, it is customary to draw graphs with vertices and edges (connecting lines). In our case their are three interaction points 0, x_1 and x_2 , one connecting photon line (one photon contraction) and two e-lines so that one uncontracted ψ_0 and $\bar{\psi}_0$ remains. One easily sees that there are three combinatorial distinct contributions according to whether the remaining pair may come from $j_{0\mu}(0)$, from the j(x)'s or if it is of mixed origin i.e. one from j(0) and the other from a j(x). The first case only contributes in zero order since :

$$\langle p' | j_{0\mu}(0) | p \rangle \langle 0 | S^* S | 0 \rangle^{(2)} = 0$$
 (4.34)

This cancellation is a general feature of all "vacuum bubble" contributions (which only contribute a phase factor to S and the opposite to S^*). More interesting are the terms in which both of the ψ - $\bar{\psi}$ "legs" (the graphical representation of the fields which remain after contractions) are contracted with legs in $S^{(2)}$. This constitutes the famous "vacuum polarization" contribution $\Gamma_{\mu,pol}$ to the form factor and the so called "one particle irreducible" form factor $\Gamma_{\mu,loop}$. The vacuum polarization contribution contains the fluctuation of the zero order current:

$$i\Pi_{\mu\nu}(x) = e^2 \left\langle 0 \left| T j_{0\mu}(x) j_{0\mu}(0) \right| 0 \right\rangle$$
(4.35)

The $\Gamma_{\mu,loop}$ -contribution originates from a contraction in which one leg goes to one W-vertex and the other to the second. The remaining "electron selfenergy

 $^{^{6}}$ The reader should follow the more detailed calculations of Weinberg ([61]) or Itzykson-Zuber ([62]). We only sketch those computations which we need in order to make some additional comments in line with our aims.

contribution" arises from the mixed contraction. It contain the electron selfenergy Σ and therefore is called $\Gamma_{\mu,e.s.}$. The three types of terms are conveniently pictured in terms of Feynman diagrams.(Fig.)

Inserting now the Fourier representation of the time ordered electron and photon propagators we obtain:

$$\langle p' | j_{\mu}(0) | p \rangle = \bar{u}(p', s'_{3})(\Gamma_{\mu, pol} + \Gamma_{\mu, loop} + \Gamma_{\mu, e.s.})u(p, s_{3})$$
 (4.36)

where the vacuum polarization, vertex-loop and electron-self energy contributions to the matrix-valued Γ_{μ} is

$$\Gamma_{\mu,pol}(p',p) = \frac{-1}{(p'-p)^2 + i\varepsilon} \Pi_{\mu\nu}(k) \gamma^{\nu}$$
(4.37)

$$\Pi_{\mu\nu}(k) = \frac{-ie^2}{(2\pi)^4} \int d^4q \frac{Tr([-i\not\!\!\!/ p+m]\gamma_{\mu}[-i(\not\!\!/ -k)+m]\gamma^{\nu})}{(q^2-m^2+i\varepsilon)((q-k)^2-m^2+i\varepsilon)}$$
(4.38)

$$\Gamma_{\mu,loop}(p',p) = \frac{ie^2}{(2\pi)^4} \int d^4q \gamma^{\rho} \frac{-i(\not\!p'-\not\!q) + m}{(p'-q)^2 - m^2 + i\varepsilon} \gamma_{\mu} \frac{-i(\not\!p-\not\!q) + m}{(p-q)^2 - m^2 + i\varepsilon} \gamma_{\rho}$$
(4.39)

$$\Gamma_{\mu,e.s.} = \left\{ i\Sigma(p')S_F(p')\gamma_{\mu} + \gamma_{\mu}S_F(p)i\Sigma(p) \right\}$$
(4.40)

The electron selfenergy $i\Sigma$ is not an observable quantity and fortunately its contribution drops out by the adiabatic principle. This happens because the two momentum variables p' and p are on the physical (mass m) mass shell and therefore the adiabatic principle forces us to work with \mathcal{W}_{adia} instead of \mathcal{W} and we have to fix the counterterm in such a way that the one particle matrix element of ψ equals that of ψ_0 . This is easily seen to be identical (in second order) to: $i\Sigma(p)_{adia}u(p,s_3)|_{p=m}=0$ with $i\Sigma_{adia}=i\Sigma+$ counter terms. The on-shell vanishing of the selfenergy is just the mathematical expression that the persistent selfenergy contribution to the large time asymptotics (equivalent to the momentum space mass shell limit) has been correctly taken into account and not "switched off". On the other hand as already mentioned, if we were to compute the off-shell 3-point function $\langle T\psi\bar{\psi}A_{\mu}\rangle$, we have the option to either use free fields with the "bare" mass m_0 and \mathcal{W} , or we can use free fields with the physical mass and \mathcal{W}_{adia} . This observation is well known from the Gell-Mann-Low representation or the Feynman-Kac representation of correlation functions (naturally off-shell).

In passing we mention that the Gell-Mann- Low representation for the correlation function of Heisenberg fields (in case of a scalar neutral selfinteracting field) has the form:

$$\langle TA(x_1)...A(x_n) \rangle = Z^{-1} \left\langle TA_0(x_1)...A_0(x_n)e^{i\int W(A_0(x))d^4x} \right\rangle_{Fock}$$
(4.42)
with $Z = \left\langle e^{i\int W(A_0(x))d^4x} \right\rangle$

158

We note that this is a special case of the previous formalism if we extend it to products of fields and consider formally the adiabatic limit $g \rightarrow 1$. The denominator Z is a phase factor and represents the (volume-dependent) "vacuumbubbles" which cancel against similar contributions from the numerator. The volume structure is completely analogous to the thermodynamical limit of the Gibbs representation of thermal correlation functions (at the end of chapter 1), with the only structural difference that there is no equivalent to the KMS condition. For physical reasons the correctly normalized ground state expectations should result from the KMS theory for $\beta \rightarrow \infty$.

Looking formally at the momentum space representation for the second order vacuum representation one would expect (by power counting in p) a quadratic divergence. Invoking current conservation (or gauge invariance), only a logarithmic divergence remains in $\pi_{\mu\nu}$. A closer look at the electron selfenergy term $i\Sigma$ reveals that the divergence is also logarithmic and the same is obviously true (by power counting) for $\Gamma_{\mu,loop}$. In the remainder of this section we present and explain the result of the renormalization on the second order formfactor. The presentation of the techniques and the actual calculation will be deferred to the next section. We collect the results (omitting tildes in Fourier transforms):

$$\Pi_{\mu\nu}(k) = i(k_{\mu}k_{\nu} - g_{\mu\nu}k^{2})\pi(k) \text{ with:}$$
(4.43)

$$\pi^{(2)}(k) = -\frac{\alpha}{3\pi} \left\{ +\frac{1}{3} + 2(1 + \frac{2m^{2}}{k^{2}}) \left[xarccotx - 1\right] \right\} - Z_{3}^{(2)}$$
where $x = (\frac{4m^{2}}{k^{2}} - 1)^{\frac{1}{2}} \text{ for } k^{2} < 4m^{2} \text{ and anal.cont.},$

$$Z_{3}^{(2)} = \frac{\alpha}{3\pi} \ln \frac{\Lambda^{2}}{m^{2}}$$

$$\Sigma^{(2)}(p) = \frac{\alpha}{2\pi} \left\{ \begin{array}{c} (\not p - m)(1 + \ln \frac{\mu^{2}}{m^{2}}) + 2m\frac{m^{2} - p^{2}}{p^{2}} \ln(1 - \frac{p^{2}}{m^{2}}) \\ - \not p \left[\frac{3}{2} \frac{m^{4} - (p^{2})^{2}}{(p^{2})^{2}} \ln(1 - \frac{p^{2}}{m^{2}}) + \frac{m^{2} - p^{2}}{p^{2}} \right] \right\}$$

$$+ \delta m + Z_{2}^{(2)}(\not p - m)$$
(4.44)

with
$$\delta m = \frac{3\alpha}{4\pi}m(\ln\frac{\Lambda^2}{m^2} + \frac{1}{2}), \ Z_2^{(2)} = \frac{\alpha}{2\pi}(\frac{1}{2}\ln\frac{\Lambda^2}{m^2} + \ln\frac{\mu^2}{m^2} + \frac{9}{4} + O(\frac{\mu}{m}))$$

$$\begin{split} \Gamma^{(2)}_{\mu,loop}(q^2) &= \gamma_{\mu} F^{(2)}_{loop}(\theta) + \frac{i}{2m} \sigma_{\mu\nu} q^{\nu} G^{(2)}_{loop}(\theta) + \gamma_{\mu} B \end{split}$$

$$\begin{aligned} F^{(2)}_{loop}(\theta) &= \frac{\alpha}{\pi} \left\{ (\ln \frac{\mu}{m} + 1)(\theta \coth \theta - 1) - 2 \coth \theta \int_{0}^{\frac{\theta}{2}} \chi \tanh \chi d\chi - \frac{\theta}{4} \tanh \frac{\theta}{2} \right\} \\ G^{(2)}_{loop}(\theta) &= \frac{\alpha}{2\pi} \frac{\theta}{\sinh \theta} \quad \text{with } q^2 = -4m^2 \sinh^2 \frac{\theta}{2}, \quad \theta:"rapidity" \end{aligned}$$

$$\begin{aligned} (4.45) &= \frac{\alpha}{2\pi} \left\{ (\ln \frac{\mu}{m} + 1)(\theta \coth \theta - 1) - 2 \coth \theta \int_{0}^{\frac{\theta}{2}} \chi \tanh \chi d\chi - \frac{\theta}{4} \tanh \frac{\theta}{2} \right\} \\ \end{aligned}$$

Here Λ is a cutoff i.e. a formal device which cuts off certain divergent momentum space integrals in a Lorentz-invariant manner. Although Λ carries no direct

physical significance (and will be removed shortly), it is important that the Λ -dependent terms have at most a polynomial p-dependence i.e. they are of the form of δ -functions and derivatives thereof. The infrared cutoff μ on the other hand has a physical origin. The interaction of charged particles with "soft" ($\omega \rightarrow 0$) photons is very strong and changes the character of the one particle states. Its projection on Wigner particles is zero and strictly speaking we must abandon our formulation of the adiabatic principle (and the standard forms of scattering theory) and think about "infraparticles". In order to not drift too far away from elementary treatments we followed standard practice and introduced a photon-mass μ into the A_{μ} propagator retaining at the end only the leading contribution for small μ . Note that the Λ -dependent (unrenormalized) Σ is infrared-finite (B contains a compensating contribution). In Γ there is no such compensation. We have separated the Λ -cutoff dependent δm^2 and B terms in Σ because the adiabatic principle fixes the counter terms in W_{adiab} to be:

$$\mathcal{W}_{c.t.} = \delta m^{(2)} \bar{\psi} \psi - B \bar{\psi} \, \partial \psi \tag{4.46}$$

This leads to a modification (renormalization) of Σ . We already noted that the resulting Σ_{adia} is the Λ -independent content of the curly bracket. Insertion into the formula (4.40) gives $\bar{u}(p')\Gamma_{\mu,e,s}^r u(p) = 0$ i.e. the renormalized contribution vanishes on the mass shell $p = m^2$. Enforcing the charge normalization for the diagonal matrix element:

$$\bar{u}(p,s_3)\Gamma^r_{\mu}u(p,s_3) = \bar{u}(p,s_3)\gamma_{\mu}u(p,s_3)$$
(4.47)

we also eliminate the Λ -dependence in $\Gamma_{\mu,pol}$ and $\Gamma_{\mu,loop}$. The result is:

$$\Gamma^{r}_{\mu} = \gamma_{\mu} F^{(2)}(\theta) + \frac{i}{2m} \sigma_{\mu\nu} q^{\nu} G^{(2)}(\theta)$$
(4.48)

with F and G given by the previous formulae in the regime $q^2 < 0$ and everywhere by analytic continuation (as $\pi(q^2)$ and $\Sigma(q)$ they can be represented by analytic functions with a cut on the real axis). According to the previous section we obtain for the anomalous contribution to the magnetic moment:

$$\mu_{an} = \frac{e}{2m}(F(0) + G(0)) = \frac{e}{2m}\frac{\alpha}{2\pi}$$
(4.49)

Note that only the infrared-finite G contributes to the zero Taylor term.

The calculation of the Lamb-shift is more complicated, computationally (since atomic physics enters) as well as conceptually. Here one is interested in the $2s_{\frac{1}{2}} - 2p_{\frac{1}{2}}$ energy split which the Dirac theory with external fields cannot explain. The idea is to study the electron selfenergy term Σ in the presence of an external field. Whereas it is not difficult to represent the resulting energy shift in a stationary A^{ex}_{μ} as:

$$\delta E_N = -\int \int d^3 p' d^3 p \bar{u}_N(p') \Sigma^r(p', p, E_N; A_\mu^{ext}) u_N(p), \qquad (4.50)$$

the evaluation of the low energy part of the renormalized external field dependent selfenergy part Σ^r requires significant skills and intuition. Since the atomic wave functions u_N prevent the electron to become free, one expects δE_N to be a "good" variable i.e. μ -independent. In distinction to the previous case, this independence is not manifest and can be used as a check on the approximation methods needed. The only manageable approximation method involves a different treatment of high and low energy contributions [61]. For the low energy part one uses the above formula. It turns out that the main modification consists in replacing the $S_F(x-y)$ function by $S_F(x, y, A_{\mu}^{ext})$ as well as a so-called "tadpole" term involving:

$$\left\langle 0\left|j_{0\mu}(x,A_{\mu}^{ext})\right|0\right\rangle = Tr\gamma_{\mu}S_{F}(x,x,A_{\mu}^{ext})$$

The latter occurs because the charge conjugation invariance, which prohibits any vacuum expectation with an odd number of j's, is broken by the external field. This new term requires a "tadpole" counter term. The $S_F(x, y, A_{\mu}^{ext})$ only involves Dirac's external field theory. The high energy part is calculated in first order of A_{μ}^{ext} and has the form:

$$\delta E_N \mid_{h.e.} = ie \int \int d^3p d^3p' \bar{u}(\mathbf{p}') \Gamma_{\mu}(\mathbf{p}, E_N; \mathbf{p}' E_N) u(\mathbf{p}) A_{\mu}^{ext}(\mathbf{p} - \mathbf{p}')$$

The Γ_{μ} is almost the previous form factor Γ , except for the fact that the hydrogen wave functions pushes it slightly off-shell. One replaces this by the on-shell Γ , assuming a smooth continuation. The infrared dependence of the latter causes a problem. One solution [62] consists in converting the unphysical photon mass μ into an infrared photon energy cutoff κ via use of the soft bremsstrahlung. K is then used to define an upper integration limit in $\delta E \mid_{l.e.}$, whereas the high frequency contribution $\delta E \mid_{h.e.}$ is calculated from the previous formula with μ in the formfactor Γ_{μ} being replaced in favor of κ . The result for the s-p splitting in hydrogen is:

$$\delta E_{2s} - \delta E_{2p_{\frac{1}{2}}} = \frac{\alpha^5 m}{6\pi} \left\{ \ln \frac{\Delta E_{2p}}{\alpha^2 \Delta E_{2s}} + \frac{19}{30} + \frac{1}{8} \right\}$$
(4.51)

Here the $\Delta E's$ are suitably averaged energies of the hydrogen atom (only numerically accessible). This result corresponds to the famous value 1052,19 MHZ. instead of a photon energy cutoff one may also base the division on the decomposition for the photon propagator:

$$\frac{1}{k^2 + i\varepsilon} = \frac{1}{k^2 - \mu^2 + i\varepsilon} + \left(\frac{1}{k^2 + i\varepsilon} - \frac{1}{k^2 - \mu^2 + i\varepsilon}\right)$$

The first part leads to the Γ contribution whereas the second faster decreasing part enters the atomic physics calculation. Again the infrared singular terms cancel. This somewhat more attractive calculation (invariant cutoff) can be found in [61]. Other physical problems related to the formfactor are the radiative corrections to the Coulomb scattering i.e. the second order correction to the Mott-formula and the bremsstrahlung correction to the Mott formula. Both are separately infrared divergent for $\mu \to 0$, but their joint cross section (for fixed photon infrared resolution) approaches a finite limit. This is a special case of good infrared behavior of photon inclusive cross sections which are the principle observables of QFT's involving photons.

In this section we met two slightly different reasons for renormalization. One is entirely physical: if we describe matrix elements between particle states (i.e. on-shell quantities) we must use the \mathcal{W}_{adiab} as our interaction, independent of whether the counter terms in the formal treatment have infinite coefficients or not. There is no other operator description for such quantities than the one in a Fock space with the correct mass. In the next section we will study offshell quantities which do not require \mathcal{W}_{adiab} . Any auxiliary mass Fock space may be used for their perturbative evaluation. It will be shown later, that scattering theory reconciles the description of on- and off-shell quantities. The Hilbert space for scattering theory requires a reparametrization from the offshell auxiliary mass to the physical mass as well as a multiplicative adjustment. In momentum space the difference between these two type of quantities looks deceivingly simple: one just sends certain p-variables to the physical mass shell. In x-space however this distinction looks more dramatic: it is the difference between global (particles involve asymptotic limits) and local quantities. In algebraic QFT the fact that the local spacetime description does not "perceive" the mass, corresponds to the local equivalence of algebras which belong globally to inequivalent (different charges) representations (see later sections).

We end this section with some formal remarks on how to use the above timeordered formalism to obtain perturbative correlation functions. As a prototype theory which is free of infrared problems, tensor-and spinor-indices etc., we take the model with $W(x)=:A_0(x)^4$:

Previously we have seen that the Gell-Mann Low representation for time ordered n-point functions has the following form:

$$\langle TA(x_1)...A(x_n)\rangle = \lim_{g(x)\to g} \left\langle TA_0(x_1)...A_0(x_n)e^{i\int g(x)\mathcal{W}(x)d^4x} \right\rangle_0^{v.c.}$$
(4.52)

The subscript 0 on the right hand side is a reminder that the free field expressions are to be evaluated in the A_0 -Fock space and the superscript v.c. (vacuum connected) indicates that "vacuum bubbles" in the Wick-ordering must be omitted. We also mentioned the extension method of distributions which succeeds to give an iterative definition of the expanded right hand side. More popular with physicists (but not necessarily more physical!) are the various regularization methods which we will discuss in the next section. Let us consider the purely formal aspects of the A^4 model. This time we introduce counterterms W_c solely for the elimination of the divergencies which arise from the removal of the unphysical regularizations:

$$\mathcal{W} = g : A_0^4 : + \mathcal{W}_c$$

$$\mathcal{W}_c = \delta m^2 Z : A_0^2 : + Z(: \partial_\mu A_0 \partial^\mu A_0 - m^2 A_0^2 :) + g(Z_g - 1) : A^4 :$$
(4.53)

The claim (proven partially later) of renormalization theory is that δm^2 , Z and Z_g can be chosen such that the correlation function stay finite in the limit of removed regularization. This time the mass appearing in Fock space does not have to be the physical one and the normalization of A is not required to be standard. However in order to have a simple form for scattering formulas it is convenient to implement the physical parametrization and the standard field normalization already in every order of perturbation theory.

4.4 Invariant Parametrizations, Regularization

The x-or p-space integrations of perturbation theory extend over noncompact regions and are difficult to perform in their original form. An efficient formalism which also allows to maintain the manifest Lorentz covariance in the presence of regularizations and cutoffs is based on Schwinger's α -parametrization (another one is due to Feynman):

$$\frac{1}{p^2 - m^2 + i\varepsilon} = \int_0^\infty e^{ia(p^2 - m^2 + i\varepsilon)} da \tag{4.54}$$

163

$$\frac{1}{\not p - m + i\varepsilon} = \frac{\not p + m}{p^2 - m^2 + i\varepsilon} = (\not p + m) \int_0^\infty e^{ia(p^2 - m^2 + i\varepsilon)} da$$

where the $i\varepsilon$ provides a damping factor for the upper integration limit. Applying this representation to the second order vacuum polarization we obtain:

$$\begin{aligned} \pi_{\mu\nu}(k) &= -e^2 \int \frac{d^4p}{(2\pi)^4} \left\{ \frac{Tr\gamma_{\mu}(\not p + m)\gamma_{\nu}(\not p - \not k + m)}{(p^2 - m^2 + i\varepsilon)((p - k)^2 - m^2 + i\varepsilon)} \right\} \tag{4.55} \\ &= -4e^2 \int \frac{d^4p}{(2\pi)^4} \left\{ \frac{p_{\mu}(p - k)_{\nu} + \{\mu \Leftrightarrow \nu\} - g_{\mu\nu}(p^2 - p \cdot k - m^2)}{(p^2 - m^2 + i\varepsilon)((p - k)^2 - m^2 + i\varepsilon)} \right\} \\ &= -4e^2 \int \frac{d^4p}{(2\pi)^4} \left\{ \frac{\partial}{\partial x^{\mu}} \frac{\partial}{\partial y^{\nu}} + \{\mu \Leftrightarrow \nu\} - g_{\mu\nu}(\frac{\partial}{\partial x} \cdot \frac{\partial}{\partial y} + m^2) \right\} \\ &\times \int \int_0^\infty d\alpha_1 d\alpha_2 \exp i \left\{ \begin{array}{c} \alpha_1(p^2 - m^2 + i\varepsilon) + \alpha_2((p - k)^2 - m^2 + i\varepsilon) \\ + x \cdot p + y \cdot (p - k) \end{array} \right\}_{x=0=y} \end{aligned}$$

where in the last step we used the α -parametrization and eliminated the polynomial in the numerator by differentiation and setting the auxiliary variables zero at the end. In this form the p-integration involves easy to do oscillatory Gaussian integrals and the original divergence has been shifted into the α -integrals as divergencies at $\alpha = 0$:

$$\pi_{\mu\nu}(k) = \frac{i\alpha}{\pi} \int \int_{0}^{\infty} \frac{d\alpha_1 d\alpha_2 \alpha_1 \alpha_2}{(\alpha_1 + \alpha_2)^4} \left\{ \begin{array}{c} 2(k_\mu k_\nu - g_{\mu\nu} k^2) \\ -g_{\mu\nu} \left(k^2 + \left[m^2 - \frac{i}{\alpha_1 + \alpha_2}\right] \frac{(\alpha_1 + \alpha_2)^2}{\alpha_1 \alpha_2}\right) \end{array} \right\} \times \\ \times \exp i \left\{ \frac{\alpha_1 \alpha_2}{\alpha_1 + \alpha_2} k^2 - (\alpha_1 + \alpha_2) m^2 \right\}$$
(4.56)

Here we split the polarization into a transversal and longitudinal part. Note that the transversality property:

$$k^{\mu}\pi_{\mu\nu}(k) = -e^{2} \int \frac{d^{4}p}{(2\pi)^{4}} Tr(k\frac{1}{\not p - m + i\varepsilon}\gamma_{\nu}\frac{1}{\not p - k - m + i\varepsilon}) \quad (4.57)$$
$$= -e^{2} \int \frac{d^{4}p}{(2\pi)^{4}} Tr\gamma_{\nu}(\frac{1}{\not p - k - m + i\varepsilon} - \frac{1}{\not p - m + i\varepsilon}) \stackrel{?}{=} 0$$

does not follow because the translation of integration variables is not allowed. Instead of enforcing the transversality condition by "brute force" (vanishing of the longitudinal term), we may also use regularizations which maintain transversality (gauge invariance). There are two gauge invariant methods: the Pauli-Villars method of auxiliary fields and the more recent dimensional regularization method. The P-V method adds fictitious spinor fields with masses $m_i = \lambda_i m$ and strength C_i :

$$\pi_{\mu\nu}^{PV}(k, m_1, m_2...) = \pi_{\mu\nu}(k, m) + \sum_{i=1}^n C_i \pi_{\mu\nu}(k, m_i)$$
(4.58)

The power counting of the integrand indicates convergence for:

$$1 + \sum_{i=1}^{n} C_i = 0 \tag{4.59}$$

For sufficiently convergent integrals one can shift integration variables and obtain the transversality of $\pi^{PV}_{\mu\nu}$. The transversal $\pi^{PV}_{\mu\nu}$ has the following α -representation:

$$\pi_{\mu\nu}^{PV}(k) = -i(g_{\mu\nu}k^2 - k_{\mu}k_{\nu})\pi(k)$$

$$\pi(k) = \frac{2\alpha}{\pi} \int_{0}^{\infty} \frac{d\alpha_1 d\alpha_2 \alpha_1 \alpha_2}{(\alpha_1 + \alpha_2)^4} \sum_{i=0}^{n} C_i \exp i \left\{ k^2 \frac{\alpha_1 \alpha_2}{\alpha_1 + \alpha_2} - (m_i^2 - i\varepsilon)(\alpha_1 + \alpha_2) \right\}$$

$$= \frac{2\alpha}{\pi} \int_{0}^{\infty} \int_{0}^{\infty} d\alpha_1 d\alpha_2 \alpha_1 \alpha_2 \delta(1 - \alpha_1 - \alpha_2) \times$$

$$\times \int_{0}^{\infty} \frac{d\rho}{\rho} \sum_{i=0}^{n} C_i \exp i\rho \left\{ k^2 \alpha_1 \alpha_2 - m_i^2 + i\varepsilon \right\}$$

$$(4.60)$$

where in the last line the identity $\int d\rho \delta(\rho - \alpha_1 - \alpha_2) = 1$ was used to introduce a radial variable ρ . By appropriate choice of the C_i one improves the small ρ behavior.. For our purpose it is sufficient that the above relation $\sum_{i=1}^{n} C_i = 1 + \sum_{i=1}^{n} C_i = 0$ leads to the vanishing of the sum in the integrand. For this n = 1 suffices. Higher order zeros could be obtained by requiring higher moments to vanish as well i.e. $\sum_{i=1}^{n} m_i^s C_i = 0$. The ρ integration gives (considering only leading contributions for large λ_i):

$$\lim_{r \to 0} \int_{r}^{\infty} \frac{d\rho}{\rho} \sum_{i=0}^{n} C_{i} \dots = \lim_{r \to 0} \sum_{0}^{n} C_{i} (-e^{i\sigma} \ln \sigma \mid_{\sigma = r(k^{2}\alpha_{1}\alpha_{2} - m_{i}^{2})} (4.61)$$

164

$$\begin{split} &+ \int_0^\infty d\sigma e^{i\sigma(1+i\varepsilon)} \ln \sigma) \\ &= -\left\{ \ln(1 - \frac{\alpha_1 \alpha_2 k^2}{m^2}) - \ln \frac{\Lambda^2}{m^2} \right\}, \\ \text{with } \sum_1^n C_i \ln \lambda_i^2 &= -\ln \frac{\Lambda^2}{m^2} \end{split}$$

This yields the Pauli-Villars regularized vacuum polarization:

$$\begin{aligned} \pi^{PV}(k) &= -\frac{2\alpha}{\pi} \int_0^1 dx x (1-x) \left\{ \ln(1-x(1-x)\frac{k^2}{m^2}) - \ln\frac{\Lambda^2}{m^2} \right\} \\ &= -\frac{\alpha}{3\pi} \left\{ 2(1+\frac{2m^2}{k^2}) \left[yarccoty - 1 \right] + \frac{1}{3} - \ln\frac{\Lambda^2}{m^2} \right\} \end{aligned} (4.62) \\ \text{with } y &= (\frac{4m^2}{k^2} - 1)^{\frac{1}{2}} \end{aligned}$$

Another more recent regularization scheme which also maintains gauge invariance of $\pi_{\mu\nu}$ is the dimensional regularization. This method only works in the euclidean formulation of perturbation theory which will be discussed in a later section. The regularized expressions for $i\Sigma$ and Γ_{μ} are also conveniently derived in the α -parametrization. since the integration over the loop momentum is always a simple (oscillatory) Gaussian, we write directly:

$$\Sigma(p) = \frac{\alpha}{2\pi} \int \int \frac{d\alpha_1 d\alpha_2}{(\alpha_1 + \alpha_2)^2} (2m - \frac{\alpha_1}{\alpha_1 + \alpha_2} \not p) \times \\ \times \exp\left\{i(\frac{\alpha_1}{\alpha_1 + \alpha_2} p^2 - \alpha_1 \mu^2 - \alpha_2 m^2)\right\}$$
(4.63)
$$= \frac{\alpha}{2\pi} \int_0^\infty \frac{d\rho}{\rho} \int \int d\alpha_1 d\alpha_2 \delta(1 - \alpha_1 - \alpha_2) (2m - \alpha_1 \not p) e^{i\rho(\alpha_1 \alpha_2 p^2 - \alpha_1 \mu^2 - \alpha_2 m^2)}$$

As in the previous case we add a PV regularization term but this time through an auxiliary "heavy photon field" of mass Λ^2 . Again only retaining the leading term, we obtain the answer by the substitution($C_0 = 1, C_1 = -1$):

$$\int \frac{d\rho}{\rho} e^{i\rho(above)} \to \int \frac{d\rho}{\rho} (e^{i\rho(above)} - e^{-i\rho\alpha_1\Lambda^2})$$
(4.64)

The ρ -integration and one of the α -integrations say α_2 can be done and we are left with the following integral representation:

$$\Sigma(p,\Lambda) = \frac{\alpha}{2\pi} \int_0^1 dx (2m - x \not p) \ln \frac{x\Lambda^2}{(1-x)m^2 - x(1-x)p^2 + x\mu^2 - i\varepsilon} \quad (4.65)$$

If we stay within $p^2 < m^2$ we may set $\mu = 0$ and obtain the explicit results

$$\Sigma = \frac{\alpha}{2\pi} \left\{ \begin{array}{c} \ln\frac{\Lambda^2}{m^2} (2m - \frac{1}{2}\not p) + 2m(1 + \frac{m^2 - p^2}{p^2}\ln(1 - \frac{p^2}{m^2}) \\ -\frac{1}{2}\not p[\frac{m^4 - (p^2)^2}{(p^2)^2}\ln(1 - \frac{p^2}{m^2}) + \frac{m^2}{p^2} + \frac{3}{2}] \end{array} \right\}$$
(4.66)

This is a matrix-valued analytic function in the cut p^2 -plane which has a diverging derivative on the mass shell $p = m^2$ as a reminder of the infrared problem. By keeping μ finite, the mass shell limit has finite derivatives.

• The regularization of the one particle irreducible second order contribution to the vertex function (4.39) is more involved, since as a result of the presence of three propagators one has to introduce $3 \alpha' s$. The α -representation for the three denominators reads as:

$$\int \frac{d^4q}{(2\pi)^4} \frac{e^{iqx}}{(q^2 - \mu^2 + i\varepsilon)(q^2 - 2p' \cdot q + i\varepsilon)(q^2 - 2p \cdot q + i\varepsilon)} \tag{4.67}$$

$$= \frac{1}{(4\pi)^2} \int \int \int \frac{d\alpha_1 d\alpha_2 d\alpha_3}{(\alpha_1 + \alpha_2 + \alpha_3)^2} \exp -i \left\{ \alpha_1 \mu^2 + \frac{(\frac{x}{2} - \alpha_2 p' - \alpha_3 p)^2}{\alpha_1 + \alpha_2 + \alpha_3} \right\}$$

where the exponential was added for the same reasons as in the previous case of $\pi_{\mu\nu}$ namely to convert the polynomial q-dependent numerator into a differentiation acting on the variable x. The (PV type) regularization can be again implemented through the photon propagator. Since there is no essential new idea involved but (even if one passes from the off-shell vertex to the on shell formfactor) only some lengthy calculations, we skip the details in the derivation of formula (4.43)The divergent Λ -dependent parts are evidently local (polynomial in the external momentum variables) ant therefore can be compensated by counter terms of the following kind:

$$\mathcal{W}_{c.t.} = -\frac{1}{4}(Z_3 - 1)F^2 + (Z_2 - 1)(\frac{i}{2}\bar{\psi}\overleftrightarrow{\partial}\psi - m\bar{\psi}\psi) + Z_2\delta m\bar{\psi}\psi - e(Z_1 - 1)\bar{\psi} \notA\psi$$
(4.68)

Finite parts in counterterms would remain unspecified unless one imposes normalization conditions. Natural normalization conditions are the conditions which result from the adiabatic requirement (a must for on shell quantities) augmented by the physical charge parametrization of the formfactor. The regularized formulas of the previous section (4.43) have been written in such a way that the natural normalization means omission of the δm and Z-terms. It is easily seen that the remaining second order π, Σ and Γ terms have the correct zeros required by the adiabatic principle and the physical charge (coupling constant) parametrization. The proof of n^{th} order renormalizability, i.e. the statement that the old local counterterms iterated together with the original \mathcal{W} in the S(q) expansion lead to higher order correlation functions which in turn may be liberated from their infinities by new higher order counterterms of the same local structure, requires a significant extension of the regularization formalism. These notes are not intended as a substitute for the rigorous treatment of n^{th} order renormalized pertubation theory. We only want to explain the physical concepts behind as well as to present some of the famous second order QED radiative corrections.

We already stressed the fact that the close connection between particles and fields is an artifact of perturbation theory and not a result of the use of the

166

=

choice of a particular Fock space for the definition of local operator algebras. In some sense the infrared singularities of Maxwell like (gauge) theories can be interpreted as a perturbative indication that the theory is not compatible with the imposed zero order particle content.

There exists a widespread misconception that a Lagrangian quantization viewpoint is important for the intrinsic physical understanding of interactions in QFT. From such a point of view the ultraviolet divergencies appear as serious flaw of perturbation theory. The history of renormalization [64]. After all, renormalization arose from some remarks of Kramers who suggested to use similar distinctions between bare and physical masses (and other parameters) in QED as they were used by Lorentz and Poincaré in their attempt to understand the selfenergy problem of particles grafted upon classical field theory. Additional support came from the observation that for a few models (e.g. A^4 in d=1+1) which were "well-behaving" in the perturbative treatment, it was possible to prove their mathematical existence by extending the perturbative method. In this connection also the functional integral method (based on the euclidean Feynman-Kac representation) which furnishes a rather direct relation between QFT and classical physics (more elegant than canonical quantization) ought to be mentioned. However the recent progress in a nonperturbative understanding of interactions from a different starting point (sometimes called the "bootstrap approach") has cast grave doubts on the universal correctness (apart from those few exceptions) of such quantization approaches to interactions (beyond an infinitesimal deformation picture). We will return to this important point at a more appropriate place.

It also should be stressed that renormalized perturbation theory does not lend credibility to the idea of a physical cutoff i.e. a Λ which cannot be absorbed into the renormalization constants but rather enters the physics. A physically interpretable nonlocal theory with an elementary length in form of a cutoff Λ does presently not exist⁷. If it ever comes into existence, it probably will be quantum gravity par excellence. For large families of lattice models many concepts of QFT as particle and scattering notions exist, even though they are much harder to prove [95] (as a result of the missing knowledge about commutativity properties at different times i.e. the substitute for spacelike commutativity). However even though the lattice cutoff does not wreck the interpretation in terms of particle excitations, the scarce rigorous results on scaling limits do not justify to attribute a physical meaning in the QFT use of lattice model cutoffs.

⁷The path of QFT is littered with proposals of theories which claimed to be nonlocal and physical: cut-off in Feynman integrals, formfactors in Lagrangians, "peratization" (pairs of complex poles in Feynman rules) etc. The only attempt which is still not disproved (although the causality issue has not yet been completely settled) is one in which the Einstein causality in Minkowski spacetime is replaced by another structure which originates from a model of noncommutative spacetime [97].

4.5 Specialities of Perturbative Gauge Theories

The calculations of QED radiative corrections in the previous section made a very naive use of covariant vectorpotentials in the spirit of classical gauge theory and relied on the hope that classically gauge invariant quantities will only involve the quantum physical degrees of freedom of photons. The incorporation of this classical gauge principle into the Heisenberg-Pauli-Fermi canonical field quantization then leads to the quantized abelian gauge theory. In order to achieve something similar for the nonabelian case, it is helpful to change from canonical to functional integral quantization, in which case one encounters the presence of the more complicated nonabelian Faddeev-Poppov ghosts which require the somewhat involved BRS formalism for the extraction of physical quantities.

In the spirit of a more intrinsic approach, we should however avoid quantization arguments altogether and start directly from the Wigner theory applied to free fields in Fock space.

The gauge concept originated in the form of "minimal coupling" in QM with external electro-magnetic interactions. As with many rules and recipes in QM (e.g. spin-statistics), one expects that this gauge principle is a consequence of the more fundamental spin=1 LQP together with perturbative renormalizability. Indeed as we show below, the consistency requirements on spin=1 perturbative interactions leave much less freedom than those between lower spins, in stark distinction to classical physics, where it is the other way around (and where one needs the gauge principle as a selection criteria. Faithful with Bohr's correspondence principle, it is the more basic LQP which explains the gauge principle of classical or external field theory of vectorfields rather than the other way around, as the quantization approach of the textbooks make believe. This leads not only to an alternative viewpoint e.g. on the Schwinger Higgs mechanism, but relates such disconnected issues as the decoupling of the alias Higgs field (in the terminology of the standard approach) with the appearacce of semiinfinite string-like localized charged fields in the zero mass limit for the mass of the vectormesons.

In this section we first we remind ourselves that the general relation between the (m,s) Wigner canonical creation and annihilation operators of (anti)particles and the covariant pointlike free fields is (see previous chapter):

$$\psi^{[n_+,n_-]}(x) = \int \sum_{s_3=-s}^{s} \left\{ e^{-ipx} u(p,s_3) a(p,s_3) + e^{ipx} v(p,s_3) b^*(p,s_3) \right\} \frac{d^3p}{2\omega}$$
(4.69)

where $u(p, s_3)$ are the columns of the rectangular $(2n_+ + 1)(2n_- + 1) \times (2s + 1)$ intertwiner matrix U(p) which intertwines the canonical Wigner representation with the covariant representation of the Poincaré group built on the calculus of (un)dotted spinors:

$$D^{[n_+,n_-]}(\Lambda)U(\Lambda^{-1}p) = U(p)D^{[s]}(R(\Lambda,p))$$
(4.70)

Independent of the choice of covariant field coordinates, for $s \ge 1$ one always

encounters a polynomial of degree $m \ge 2$ in p-space for the two-point function:

$$\langle \psi_{\cdot}(x)\psi_{\cdot}^{*}(y)\rangle = P_{\cdot}^{(m)}(i\partial)i\Delta^{(+)}(x-y)$$
(4.71)

In the massless case the Δ -functions have to be replaced by *D*-functions and in addition is has to be kept in mind that the family of possible intertwiners is more restrictive as a result of the helicity rule $n_+ - n_- = h$. This increasing operator dimension is the cause of a quantum obstruction against renormalizability: all spin (or helicity) ≥ 1 field operator have scale dimension ≥ 2 and hence in the free field Borchers class of the fields which enter in the invariant interaction W, we must have dim $W \geq 5$ (a W describing interactions must be at least trilinear!), whereas renormalizability demands dim $W \leq 4$. This means that even if one takes an algebraic point of view locality does not interacting generators W whose dimension is below 5. Note that their are certain derivatives and composite fields which retain their classical dimensions as e.g. $F_{\mu\nu}, F^{\mu\nu}F_{\mu\nu}, \psi$ etc. In fact the only fields associated with an (m,s) Wigner representation in case of $s \leq 1$ which (even for an optimal choice of field coordinates) must have an operator dimension beyond the classical one, is s=1 i.e. the vectorpotential.

Let us first look at the massive case which turns out to be conceptually simpler. The minimal description is in terms of a transverse vector potential with $\dim A_{\mu} = 2$ (this high operator dimension is resulting from the transversality inherent in the Wigner theory. There is no known way to use the milder nonlocal string-localized vector potentials in a perturbative renormalization scheme, and the dream of a renormalizable higher spin deformation theory could have ended right here. Fortunately one can outwit the above No Go argument by a magic trick which produces interaction-deformed observables (a power series *-algebra) generated by the subset of physical (composite) operators in Fock space. The physical generating fields in this process retained their classical dimensions modulo logarithms, and the unphysical fields involving vectorpotentials attain their classical value $\dim A=1 \pmod{\log(1+1)}$ and become formally local and covariant. In this approach the deformation theory only depends on a finite number of parameters as expected for a renormalizable theory. This magic is achieved by a cohomological representation of the Wigner theory for which the application of the Weyl (pseudo)functor commutes with the cohomological descend:

The horizontal arrows represent the (pseudo-)Weyl functors from wave function spaces to (pseudo-)von Neumann algebras and the vertical arrows denote the BRS descend for wave functions or algebras respectively. There is a corresponding cohomological descend from a pseudo Fock space to a physical (factor) Fock space. Here "pseudo" refers to the fact that the star operation with respect to which the Lorentz group fulfils a unitarity-like relation $U^{-1}(\Lambda) = U(\Lambda)^*$ in H^{ghost} is not the Hilbert space star related to the positive inner product. As in

169

the interaction free case of (m=0,h=1) in chapter3, one can (re)introduce physical vector potentials in H_{Wig}^{phys} , but they will be necessarily nonlocal. The magic trick spares one the conceptual pain to think about how a theory manages to be strictly local, even if some of the objects as vector potentials are in a physical sense mildly (semiinfinite stringlike) nonlocal.

The simplest cohomological extension of the Wigner wave function space which allows a nilpotent operation s with $s^2 = 0$, (such that the physical transversality condition $p^{\mu}A(p) = 0$ follows from the application of s) needs besides two scalar ghosts wave functions ω and $\bar{\omega}$ another scalar field φ_a :

$$(sA^{a}_{\mu})(p) = p_{\mu}\omega_{a}(p)$$

$$(s\omega_{a})(p) = 0$$

$$(s\bar{\omega}_{a})(p) = p^{\mu}A^{a}_{\mu}(p) - im_{a}\varphi_{a}(p)$$

$$(s\varphi_{a})(p) = -im_{a}\omega_{a}(p)$$

$$(4.73)$$

Here we have already added a multiplett index since we have a selfcoupling involving several vector mesons in mind with $m_a = m$ a common mass (the interesting question of whether the stability of the deformation via interaction can be maintained in the presence of unequal masses has not been systematically investigated). One immediately realizes that $s^2 = 0$ and that $s(\cdot) = 0$ enforces the vanishing of ω_a and relates φ_a to $p^{\mu}A_{\mu}$. At this point there is no grading in the formalism, i.e. the ω and φ are simply ungraded wave functions. The guiding principle for the cohomological extension of the Wigner theory is that the associated two point function (or propagation kernel) of the vectorpotential has a milder (renormalizable in suitable interactions) high momentum behavior with the physically required behavior only arising after the cohomolog*ical descend.* This is close to the particle theory spirit of Lewellyn-Smith [70]. In the geometrically motivated Faddeev-Poppov method on the other hand, it was the gauge fixing in the Lagrangian and the (euclidean) measure theoretical repair which assured the existence of a renormalizable deformation of free spin=1 fields. On the level of the Wigner theory it makes no sense to to impose a grading. However the functorial transition from Wigner theory to QFT requires the introduction of a grading with deg $\omega = 1$, deg $\bar{\omega} = -1$, and $\deg A_{\mu} = 0$, with s transferring degree 1. The reason is that only with this grading assignment, the s allows a natural tensor extension to multiparticle spaces with stable nilpotency, thus insuring the commutativity of the above diagram (which represents the cohomological ascend and descend) through the relation: $s(a \otimes b) = sa \otimes b + (-1)^{\deg a} a \otimes sb.$

This suggests to view the Fock space version δ of s as the image of a (pseudo)Weyl functor Γ as $\delta = \Gamma(s)$ and to write the δ in the spirit of a formal Noether symmetry charge Q associated with the free field (linear) version of the BRS charge [65] [66]:

$$Q = \int (\partial_{\mu}A^{\mu}(x) + m_{a}\phi(x)) \overleftrightarrow{\partial}_{0}\omega(x)d^{3}x = Q^{\dagger}$$
(4.74)

$$\delta F \equiv i \left\{ QF - (-1)^{\deg(F)} FQ \right\}, \ F \in \mathcal{F}$$

which acts formally like an abelian gauge transformation (i.e. adds a longitudinal contribution) on the vector potential. Here we keep the same symbols for the (pseudo)quantum fields as for the Wigner (pseudo)wave functions. As a result of the anticommutation relations of ω , the operator Q and the Z_2 graded δ are nilpotent i.e. $\delta^2 = 0$ and therefore defines a cohomology theory in the formal polynomial free field algebra \mathcal{F} . This is a formal *- algebra with a Z_2 -graded antilinear *-derivation. It is not difficult to show that:

$$\mathcal{A} = \frac{\ker \delta}{\delta(\mathcal{F})}$$

$$\ker \delta = \{B \in \mathcal{F} \mid \delta(B) = 0\}$$

$$(4.75)$$

where ker δ is the nullideal (coboundary) associated with s, is the algebra of physical photon operators. The *-operation and the associated notion of pseudo-hermiticity has an associated non-positive definite sesquilinear form on $H_{Fock}(A, \omega, \bar{\omega})$ with:

$$\langle \phi, F\psi \rangle = \langle F^*\phi, \psi \rangle \tag{4.76}$$

In fact the indefinite nature of this sesquilinear form is a structural consequence of the nilpotency and the pseudo-hermiticity of Q:

$$\langle Q\phi, Q\psi \rangle_{ps} = \langle \phi, Q^2\psi \rangle_{ps} = 0$$

$$(4.77)$$

$$\langle Q \equiv 0 \quad unless \quad \langle \phi, \psi \rangle_{ps} \text{ is indef.}$$

i.e. any cohomological construction based on a nilpotent charge requires indefinite metric. As expected, it is problem of Lorentz-covariance of vectorpotentials which is the origin of this structure. In the (necessarily indefinite) metric in which Q is (pseudo)hermitian, also the representation of the L-group comes out (pseudo)unitary. So in Wigner space as well as in the cohomological representation it is *always the Lorentz-group representation* in which the deviation from the standard situation shows up; in the Wigner space there was an additive term in the transformation law, and the vectorpotential was nonlocal. whereas now these aspects have been traded for the "pseudoness" of the transformation laws. This trading is formally advantageous, because the mathematics of the deformation does not care about correct physical behavior but only requires the formal locality and covariance properties.

It is worthwhile to note that in our dichotomic division of QFT into algebraic aspects and properties of states (and representations), the "pseudoness" is solely in the states and the GNS representation and not in the abstract algebra. As already for standard spin ≤ 1 renormalizable couplings, the algebras of the formal deformation theory are only *-algebras and not C*-algebras.

Now we come to the deformation of free fields via *W*-interaction polynomials. If one stays within the framework of completely massive spin=1 theories, the formal validity of (LSZ, Haag-Ruelle) scattering theory defines a reference space in which one can do all calculations: the incoming (pseudo)Fock space referring to the incoming scattering situation. This brings significant simplifications as compared to massless vector sons: in this reference space the conserved Q'shave a bilinear representation in terms of free fields (with significant simplifications as compared to the standard nonlinear BRS formalism). The bilinear structure of conserved charges (including the Poincaré-generators) in terms of the free incoming fields is absolutely crucial for the following. It gives a fixed position of the physical cohomological space inside the pseudo Fock space and it makes all basic fields as $F_{\mu\nu}, \psi$ etc. "physical" (except the vector potential and the ghosts). Note that in the standard approach based on quantization of gauge theories, the present massive spin1 situation would be termed "a completely broken gauge theory". The mentioned elementary physical fields would then be identical with the composite fields $F_{\mu\nu} \cdot \Phi, \psi \cdot \Phi$, where Φ is the Higgs field. This immediately poses the question of where and in what form the Higgs field enters the present formalism. This has been partially answered in the work of Scharf and collaborators [67]. In the following we will interpret and comment their findings. Take the simplest case of selfinteracting first order massive spin one fields and ghosts:

$$W^{(1)} = W^{(1)}_A + W^{(1)}_\omega + W^{(1)}_\phi \tag{4.78}$$

where the first term is the most general trilinear coupling $\sim \tilde{f}_{abc}A^{\mu}_{a}A^{\nu}_{b}\partial^{\nu}A^{\mu}_{c}$ and the two other terms contain the most general coupling to the w and ϕ -ghosts. Physical consistency now leads to [67]

1. In first order one obtains a list of constraints for the couplings. The result may be written in the form:

$$W^{(1)} = igf_{abc} (: \frac{1}{2} A^{\mu}_{a} A^{\nu}_{b} F^{\nu\mu}_{c} : - : A^{\mu}_{a} \omega_{b} \partial_{\mu} \bar{\omega}_{c} :) \qquad (4.79)$$

+ $igf'_{abc} : \partial_{\nu} A^{\nu}_{a} \omega_{b} \tilde{\omega}_{c} :$

where f and f' are totally antisymmetric.

2. In second order physical consistency leads above all to the Jacobi-identity for the f's i.e. the famous gauge group structure comes from just physical consistency of the cohomologically extended spin 1 theory. The consistency for the remaining term demands that it be a total divergence (surface term). But in the presence of the spin 1 mass, there is the necessity of yet another compensation; this time (if the compensatory field is taken as scalar field as the simplest possibility), it must be a physical field and not a ghost. The result looks like that of the spontaneous broken Yang Mills theory after the subtraction of the vacuum expectation of the Higgs field, except that in our case there was no gauge theory and the original $W^{(1)}$ did not contain a Higgs field to start with but it rather enters in order to maintain the consistency of the theory in higher order. What does this apparently very powerful "physical consistency"⁸ mean in mathematical terms? I is nothing but the requirement that the adiabatic limit:

$$S = \lim_{g \to const} S(g) \tag{4.80}$$

defines the physical S-matrix in a smooth way from the off-shell S(g). The natural mechanism for providing an S-matrix which commutes with Q (the physical consistency), is that the noncommuting contributions with S(g) for arbitrary g's which have a constant value in a large double cone and fall off to zero in a collar outside is:

$$[S(g),Q] = terms \ in \ \partial g \tag{4.81}$$

In the Bogoliubov-Shirkov deformation theory this leads to a restriction for the T-products:

$$S(g) = Te^{i \int W(x)g(x)d^{4}x} = = \sum_{n} \frac{1}{n!} \int T_{n}(x_{1}, ..., x_{n})g(x_{1})....g(x_{n}) with T_{1}(x) = W^{(1)}(x)$$

d. terms involving only ∂g , or in terms of the T_n :

$$[Q, T_n(x_1, ..., x_n)] = \sum \frac{\partial}{\partial x_i^{\mu}} T_{n,i}^{\mu}(x_1, ..., x_n)$$
(4.82)

Where the terms $T_{1,1}^{\mu}$ as well as all n > 1 terms should be iteratively determined and the Hahn-Banach extension problem for the $T'_n s$ solved (the latter step being the renormalization aspect in the Epstein-Glaser approach). It comes as somewhat of a surprise that this requirement has a retroactive action even on the first order trilinear input. For the computations up to second order which bring out the above results we refer to [67].

It is worthwhile to reflect about this result. The physical consistency requirement converted the apparent freedom of spin 1 couplings (which classically seemed to be much larger than that for interactions between spin<1 fields) into a very tight situation! The perturbative treatment does not work without introducing another (this time physical) scalar field. One may call this the Higgs field, however it should be clear that its role is strictly perturbative and nothing is said about the existence of a Higgs particle in a nonperturbative massive selfinteracting massive spin=1 theory. Note that in this LQP consistency approach, there is no intrinsic meaning to think of the Higgs particle as a "fattened" Goldstone mode. This underlines an old observation made in connection with the Schwinger-Higgs mechanism in the d=1+1 Schwinger model, since there is no

 $^{^{8}}$ In the work of [67], the surface term property in (4.81) is called "quantum gauge principle", but since this could be misunderstood as if an additional structure besides physical consistency of spin=1 interactions is needed, we avoid this terminology.

Goldstone mechanism in d=1+1). The result means that there is no other massive selfinteracting spin 1 theory other than that computed by the alias broken Yang-Mills gauge theory i.e. the result is unique but this terminology has no intrinsic physical meaning; it is at best a mnemotechnical concept. It is interesting to note that the present approach is natural for the massive theories which in the original Yang-Mills work was the desired (but with those concepts unattainable) dream.

A direct application of the cohomological method to zero mass vector mesons is necessarily more complicated as a result of the absence of a global reference space. In that case there is no possibility of working with a bilinear expression for Q, rather one has to face the presence of interacting trilinear contributions. As a consequence the physical cohomology space changes its position inside the big (pseudo)Fock space with the order of perturbation. Let us briefly look at the stability problem of this more complicated situation. Start from the cohomological representation:

$$H_{Fock} = \frac{\ker Q}{rangeQ} \tag{4.83}$$

$$\mathcal{A} = \frac{\ker \delta}{\delta(\mathcal{F})}, \ \delta(F) = QF - (-1)^{\deg F} FQ \qquad (4.84)$$
$$\ker \delta = \{B \in F \mid \delta(B) = 0\}$$

Check that ker Q is a positive semidefinite subspace and that range Q agrees with the space of nullvectors in ker Q. The descend from \mathcal{F} and H_{Fock}^{pseudo} of \mathcal{A} to H_{Fock} uses also a fairly standard argument: an element of the form $B+\delta(F)$ with $B \in \ker Q$ applied to a vector of the form $\phi + Q\psi$ with $\phi \in \ker Q$ (representative of the A-class applied to representative of the H_{Fock} -class) leads to $A\phi \in \ker Q$ and to:

$$(A + \delta(F))(\phi + Q\psi) - A\phi \in QH_{Fock}^{ps}$$

$$(4.85)$$

As mentioned before, for all gauge theories which allow an adiabatic limit (i.e. for which all perturbative contributions to S(g) in the limit $g(x) \to 1$ on all Minkowski space exist), one may use the incoming free Q. This excludes the (on-shell) infrared divergent theories as QED or gauge theories without a complete Schwinger Higgs (-symmetry breaking, better: -charge screening) mechanism.

Now we indicate of how to proceed, if such a free incoming reference situation is not available. In that case we have to expand Q into a power series. The main problem of the interacting theory is then to show that the cohomological descend is stable against interactions. Of course intuitively one expects the extended structure (which brings theories with photons on the same formal level of locality and covariance in a physical space as nongauge theories) to be stable under deformations as theories of spin zero and $spin = \frac{1}{2}$ are. To prove this is not entirely trivial and only possible if one works with a notion of positivity which is adjusted to formal power series. We call a formal power series $s = \sum g^n s_n$ positive if there exists another power series $t = \sum g^n t_n$ with:

$$t^{*}t = s, \ t_{n} \in IR$$

$$\exists k \in IN_{0} \smile \{\infty\} \ s.t. \ t_{n} = 0, \ if \ n < 2k$$

$$and \ b_{2k} > 0 \ if \ k < \infty$$

$$(4.86)$$

We need this weak sense of positivity because in general the interaction will change the position of ker Q and range Q inside the interaction independent fixed pseudo Fock space H^{sp}_{Fock} . The perturbative interaction deforms the free field operators into formal power series:

$$F = \sum g^{n} F_{n}$$

$$Q = \sum g^{n} Q_{n}$$

$$\delta = \sum g^{n} \delta_{n}$$

$$(4.87)$$

where the previous free operators appear in zero order:

$$Q_0 = Q_B, \ \delta = \delta_{previous}, \ F_0 = F_{previous}, \ etc.$$
(4.88)

with:

$$Q^{2} = 0, \ \delta^{2} = 0$$

$$\delta = AdQ$$

$$(4.89)$$

The new \mathcal{A} and $H = \frac{rangeQ}{\ker Q}$ are now formal power series in g. We have to prove the following three properties:

- 1. $\langle \phi, \phi \rangle \ge 0$, for $\phi \in \ker Q$
- 2. $\langle \phi, \phi \rangle = 0, \ \phi \in \ker Q \quad \curvearrowright \phi \in \operatorname{range} Q$
- 3. \mathcal{A} is faithfully represented on H

Here the positivity is meant in the previously explained weak sense of power series. So the position of the ker Q inside H_{Fock}^{ps} keeps changing with the perturbative order and fulfills is in general only the above weak positive semidefiniteness. An exception is the abelian case for which the ghost field remains decoupled from the physical fields i.e. it preserves its freeness, similar to the Gupta-Bleuler approach, even in the presence of interactions. Therefore perturbative QED has a better positivity status than QCD.

The proof of the three properties above is by induction. We refer the reader to the work of Duetsch and Fredenhagen [16].

The conceptual simplicity of the massive theory versus the complex zero mass situation means that the physical picture in LQP is opposite to that of gauge quantization. It suggests another more indirect but conceptually more interesting way than the above deformation theory in which the physical space has a position inside the big (pseudo)Fock space which keeps changing with the perturbative order. Whereas in the latter approach the massive theory is interpreted as a broken "symmetry" (we use the parenthesis because the gauge symmetry is really not a symmetry in any physical sense), the LQP picture (with no Lie group structure put in!) is that of "charge liberation" in the massless limit of the massive theory.

Charge liberation via the appearance [69] of new superselection sector in the scaling limit is a well studied physical (in contradistinction to gauge breaking) phenomenon in LQP. There are many illustrations in case of more standard models, e.g. the two sectors of the massive Ising field theory mutate into 3 chiral sectors for both chiral components (with the identification of the two vacuum sectors). Actually the LQP picture is much closer to Schwinger's little known contribution on the charge screening phase [68] which led him to propose the Schwinger model. The physical picture of $m \rightarrow 0$ based on the study of the Schwinger model [69] as well as on physical intuition reveals that there exist spacelike semiinfinite string localized operators (whose introduction into the massive theory would ply no important physical role, as a result of the excellent localization properties of a massive spin=1 theory), which in the massless limit carry a string localized Maxwell charge. In other words the charge liberation is just the opposite of Schwinger's screening mechanism. The Higgs "condensates" are nonintrinsic description dependent quantities which do not appear in the present spin 1 consistency approach. The corresponding structural theorem of Swieca on charge screening/charge liberation in Maxwell-like theories becomes, with the present perturbative hindsight, a rigorous statement on consistent spin=1 interactions. The main point of the present approach, which really merits strong emphasis, is that it places the relation classical-quantum physics there were it should be (in the spirit of Bohr): the consistent (in this case perturbative) local quantum theory in semiclassical approximation selects the classical gauge theory and not the other way around as in the gauge quantization approach. The correspondence principle of Bohr analyses the classical situation by semiclassical approximations of the more fundamental quantum theory and leaves the quantization parallelism as a piece of artistry outside mathematical physics⁹.

To strengthen the present LQP approach to spin one problems one needs more studies. In addition to the systematic n^{th} order investigation of the massive situation and a more profound understanding of the problem of uniqueness of the physical particle (alias Higgs), there are the problems of zero mass limits of this massive perturbation theory. Here one wants to understand which modifications (adjustments) in the massive theory must be done in order to have a smooth offshell limits for the expected semiinfinite string-localized charge-carrying fields. Another problem is to understand the expected decoupling of the two scalar field (the ϕ - and alias Higgs-field). The change from the gauge QFT formulation to the present one therefore causes an interesting change of paradigm in that the

 $^{^{9}}$ Apart from the case of Weyl and CAR algebras, which can be rigorously generated by applying an appropriate functor to classical function spaces.

emphasis is changed from the formal Higgs mechanism point of view to the more physical problem of how the charge generating string like fields arise from the massive theory in the limit. Another open question is whether the method of cohomological representation for the sake of maintaining renormalizability for spin=1 is the tip of an iceberg, i.e. if there are higher spin cases for which this magic works. In that case the present reformulation of the problem would have achieved much more than a clearer conceptual setting of (broken) Gauge QFT.

Needless to mention that the underlying idea of the original BRS work was precisely to incorporate the observables of the massive case into the class of perturbative renormalizable theories. The spirit of this early work on renormalizable spin1 problems [65] was more in the pragmatic vein of Lewellyn-Smith[70] and did not yet have to carry the present burden of the predominance of differential geometry over local quantum physics.

In this section we have traded one miracle (the gauge principle) with another one (removing renormalizability obstruction via a cohomological trick of pseudo-structures). In the physical philosophy of Bohr and Heisenberg one should always try to remove nonobservable (pseudo-physical) vestiges even in intermediate steps of the calculation. One may hope that ideas of modular localization, which already led to nonperturbative progress in factorizing d=1+1 models, will give new concepts by which one could be able to use directly string localized operators in a deformation approach, since a theory which explains the origin of weaker localization of free vectorpotentials should also give hints as to how to incorporate this property into interactions. Although this has not been achieved in the present LQP cohomology approach, we hope that our ideas may help to prepare the ground for such a step. The cohomological magic trick creates a lesser danger to be confused with a permanent achievement in QFT than gauge theory; it is manifestly of a transitional nature and asks for a future more fundamental physical approach in terms of quantum localization concepts.

4.6 Interactions with External Fields, CST-Problems

Interactions of quantum fields with external (classical) fields played an important role in the development of full QFT. The simplest situation of this kind one meets in case the quantum fields are free. In fact free Dirac or Schrödinger fields interacting with external electromagnetic fields preceded QED and led, with some hindsight concerning interpretations, (see the introduction in [61]) to many correct formulas. If we look at these external field problems from the point of view of Poincaré-invariant QFT, we notice a conceptual problem. Since the vacuum and also the particle states are defined in terms of Poincarécovariance properties, it is not immediately clear how one should define such reference states if Poincaré-covariance is broken by external fields[55]. In an elegant formalism like Schwinger's (referring to his treatment of the astrophysically relevant $\mu\bar{\mu}$ —pair creation in electro magnetic. fields), the formalism itself takes care of this problem without the user being aware. A closer look reveals that the reference state built into the Schwinger formalism is the "adiabatic vacuum", which in a more mundane formalism corresponds to the approximation of the actual external interaction by a sequence of softly switched on and off external interactions. Whereas this is eminently reasonable for external electromagnetic interactions, this is generically speaking unreasonable for problems in curved space time (CST) i.e. with external gravitational fields. Since the Hawking radiation effect belongs to this class of problems, these structural questions are not without (astro)physical interest and relevance.

In the following we will briefly sketch some ideas [71] [35] which not only led to an answer for the correct reference states, but gave a framework for the renormalized perturbation theory of *interacting* quantum matter in CST. Even if, as in the case of the present author, one is not an actively working specialist in this area, one should take notice of these developments for the following reason. General QFT as it stands, is not quite that perfect quantum counterpart of the classical Faraday-Maxwell theory with its "action at the neighborhood principle". Whereas the algebraic part (the net theory) is local, the energy positivity and the vacuum homogeneity are very nonlocal stability requirements. This is the cause of the above mentioned difficulty. Therefore if QFT in CST requires to think about a substitute, this may be beneficial even for Minkowski space QFT.

Since all of the renormalization schemes use either euclidean space or momentum space and none of these methods is meaningful in generic curved space time situations, one is forced to re-think the renormalization formalism. Looking at the literature, one notes that almost all the papers (before the above work) on the subject are about euclidean CST QFT and nobody ever tried to define a Wick-polynomial (needed e.g. for the real time energy-momentum tensor etc.)

Since the answer to both questions requires the use of somewhat unfamiliar concepts, let me make some qualitative comments on the "microlocal"- (or Fourier integral operator-) analysis developed by the mathematicians Hörmander and Duistermaat around 1971. It is this analysis which gives rise to the formulation of a "microlocal spectrum condition" in QFT [71][35].

The basic idea is to refine the local analysis of singularity structure Schwartz distribution u which deals with singular supports $supp_{sing}u$ (the $supp_{sing}u$ is the complement of the largest open smoothness region for u) and more generally of distribution densities on a manifold, by shifting it from the base space to the cotangent bundle. In brief, one first zooms in on a u-singularity and then one uses a directional Fourier "telescope". If ϕ is a localizing test function, one studies:

$$\widetilde{\phi u}(\xi) = \left\langle u, e^{-i\langle \cdot, \xi \rangle} \phi \right\rangle \tag{4.90}$$

where $\langle \cdot, \cdot \rangle$ denotes dual pairing. This is a fast decreasing function in ξ as long as supp ϕ does not touch the singularity points. If supp ϕ on the other hand does extend into the singular region of u, the singularity may be directional dependent and in certain ξ -directions one may still encounter a fast decrease. Therefore one uses the following mathematical definition (V denotes the euclidean base space):

Definition 4 The wave front set, WF(u), of u is the complement in $V \times$

 $\mathbf{R}^n \setminus \{0\}$ of the points (x_0, ξ_0) in cotangent space $V \times \mathbf{R}^n \setminus \{0\}$ s. t., for each ϕ there exists a neighborhood. $U \times \Gamma$, with Γ conic (directional) neighborhood. of ξ_0 , and an $N \ge 0$ with:

$$\left|\left\langle u, e^{-i\langle \cdot, \xi \rangle} \phi \right\rangle\right| \le C_{\phi, N} \left(1 + |\xi|\right)^{-N}, \quad \forall \xi \in \Gamma$$

$$(4.91)$$

Returning to physics, we recall that as the result of the positive energy property, (unordered) correlation functions belong to a class of distributions which can be freely multiplied. For example the product of two Minkowski space twopoint functions $w_i(x, y) = W_i(x - y)$ i = 1, 2 is again a well-defined distribution in the same class, because the convolution of their Fourier transforms with the V^{\uparrow} forward light cone spectral support amounts to an integration over a finite (phase space) region. Using the Kallén-Lehmann spectral representation:

$$W_i(\xi) = \int_0^\infty i\Delta^+(\xi,\kappa)\rho_i(\kappa)d\kappa, \quad i = 1,2$$

the convolution of the $\rho'_i s$ extends over a finite mass region. This property does not hold for time-ordered or retarded distributions since they do not posses a spectral support in momentum space.

The main property of the wave front sets of distributions is that they allow a simple criterion for the existence of the product: the conic neighborhoods Γ_i must add up to a resulting conic neighborhood. in $V \times \mathbf{R}^n \setminus \{0\}$. The coordinate free adaptation to densities (distribution valued differential forms) on manifolds M is easy. The wave front sets are now cones in T^*M and they behave additively under multiplication in the following sense:

$$WF(u_2u_1) \subseteq \{WF(u_2) \oplus WF(u_1)\} \cup WF(u_2) \cup WF(u_1) \tag{4.92}$$

The product exists, if the zero section in T^*M does not intersect $WF(u_2u_1)$. This is the analogue of the product structure of Wightman functions.

Let us now test this idea for free QFT in CST. We start with the structure of the algebra generated by a Klein-Gordon field ϕ in a globally hyperbolic space time[55]:

$$\begin{pmatrix} g^{\mu\nu}\partial_{\mu}\partial_{\nu} - m^2 \end{pmatrix} \phi(x) = 0, \quad [\phi(f), \phi(g)] = E(f \otimes g)$$

$$\forall f, g \in C_0^{\infty}(M), \qquad E(x, y) = \Delta^{av}(x, y) - \Delta^{ret}(x, y)$$

$$(4.93)$$

It is defined uniquely in terms of the manifold data i.e. the retarded (advanced) functions belong to the algebraic characterization and are uniquely determined by the geometry, whereas the unordered and time ordered correlation functions are determined by the states. One now defines a wave front set for the (yet unknown) two-point functions ω_2 :

$$WF(\omega_2) = \left\{ (x,k;x',-k') \in T^*M^2 \setminus \{0\} \mid (x,k) \sim (x',k'), \ k \in \bar{V}_+ \right\}$$
(4.94)

where the equivalence relation \sim means that there exists a light-like geodesic from x to x' s. t.. k is co-parallel to the tangent vector to the geodesic, and k' is

its parallel transport from x to x'. This physically appealing local requirement for the selection of physical states was shown by Radzikowski to be mathematically equivalent to the more global Hadamard condition (an older recipe to obtain physical states). Free field structure, i.e. the Wick combinatorics means that the higher point functions are products of ω_2 i.e. that the states ω on the algebra (4.93) are so called quasi-free states on a CCR algebra. Using the previous product structure (4.92) of distributions with a known wave front, one then proves the existence of the n-point functions and a formula for their WFset. In a similar vein one shows the existence of Wick-products as e.g. : ϕ^n : . As in the Minkowski case, the time-ordered propagator:

$$iE_F(x,y) = \omega_2(x,y) + E_{ret}(x,y)$$
 (4.95)

does not have the one-sided spectral structure in order to allow for pointwise multiplication. Its wave front set is:

$$WF(E_F) = \{ \text{same, but } x \neq x', k \in \overline{V}_{\pm} \text{ if } x \in \mathcal{T}_{\pm}(x') \} \qquad (4.96)$$
$$\cup \{ (x, k; x, -k), x \in M, k \in T_x^* M \setminus \{0\} \}$$

where $\mathcal{T}_{\pm}(x')$ are the future/past of x'.

Actually the formula (4.94) turns out to be not general enough in order to incorporate theories beyond free fields. The more general formula which does not contain the restriction to light like geodesics and its stable (under multiplication of n-point functions) generalization to ω_n (which is most conveniently expressed in terms of graphs with vertices x_i and directed geodetic edges between them), can be found in the work of Brunetti et. al..[35].

We still have to understand why the microlocal Spectrum Condition (μSC) is not capable of a unique selection of a state and what kind of family it selects. A local spectral condition is not able to single out states with global symmetry, in fact for generic CST they do not exist. With one particular state in this family, all other states in the same folium (vector or density matrix in the same GNS Hilbert space) turn out to share the same WF set. In fact the states with coalescing wave front sets form exactly one folium of the set of all states on the C^* -algebra \mathcal{A} . That folium contains of course states with different superselection charges, a situation which is vaguely reminiscent of the "no hair" property of black holes (in the sense that the standard global characterizations of states with their detailed assignments of quantum numbers loose their meaning in local folia). There are two physical questions which enter ones mind. One is whether physical properties (corrections to electro-weak effects as anomalous moments, Lambshift etc.) change significantly in one folium. For this one has to understand renormalization theory and the implementation of physical normalization conditions (the adiabatic separation of interaction discussed at the beginning of this chapter. The other is whether the μSC can perceive interactions, i.e. whether the wave front set of an interacting theory can be distinguished from that of a free theory. An affirmative answer to this question would be extremely interesting even for Minkowski QFT (since the global vacuum condition is not
capable of such a distinction. Both questions are presently under investigation [35]. The perturbative calculations of wave front sets is not an easy matter.

The last issue in this section is how to do renormalization theory for interacting QFT in CST (i.e. how to avoid euclidean- and momentum-space). A framework which stays in x-space is that of Epstein and Glaser [35]. The main problem in its adaptation to the present case, is how to avoid translational invariance on which the E-G approach relies heavily. Let us look at their starting formula for the nth order coefficients of the Bogoliubov Shirkov operator S(g)after Wick-ordering:

$$T_n^{k_1...k_n}(x_1,...,x_n) = \sum_{i=1}^{k_1...l_n} t^{l_1...l_n}(x_1,...,x_n) \times (4.97)$$

: $\phi^{k_1-l_1}(x_1)....\phi^{k_n-l_n}(x_n)$:

Where $t^{l_1...l_n}(x_1, ..., x_n)$ are numerical time ordered distributions. In the E-G approach it is crucial that Wick-products (i.e. operator-valued distributions) can be multiplied with translational invariant numerical distributions. The CST substitute is:

$$WF(t_n) \in \Gamma_n^{to}, \quad to: time \ ordered$$
 (4.98)

where Γ_n^{to} is a subset of T^*M^n with a certain graphical characterization. The construction of the time ordered operators $T_n^{k_1...,k_n}$ is achieved by induction in two steps. First one shows that T_n for $(x_1...,x_n) \notin \Delta_n$ (the total diagonal) can be patched together from all lower n T's. Let us denote this T_n on $M^n \setminus \Delta_n$ as T_n^0 . The second step (the more difficult one) is the extension to the diagonal. For a detailed presentation we refer to the recent literature [35]. The main point to be gathered from these highly technical investigations is that the E-G approach allows a uniform treatment of renormalization as a distributive extension problem with all the operators living in Fock space. As in the Minkowski case in chapter 4.2 the E-G formalism defines a formal *-algebra net with a folium of states on all local algebras. No general principles are known which select a distinguished global state as the Poincaré invariant vacuum or one-particle states.

The rigorous CST renormalization theory [35] contains of course the proof for the renormalizability of the standard theory as a special case.

It would have been too nice if QFT in CST could furnish a gateway into "Quantum Gravity". After all, QFT in external electro-magnetic fields was an essential step towards QED. But unfortunately presently this does not seem to be the case and Quantum Gravity continues to exist as only an enigmatic idea, which for the time-being lacks concrete physical content. With these remarks we conclude our short excursion into QFT on CST.

Literature to chapter 4:

S.Weinberg, "The Quantum Theory of Fields", Vol. I, Cambridge University Press 1995

C.Itzykson and J-B.Zuber, "Quantum Field Theory", McGraw-Hill 1980

R.Haag, "Local Quantum Theory" Springer Verlag 1992

R. M. Wald, "Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics", University of Chicago Press 1994.

Chapter 5

The General Framework of QFT

5.1 Model-independent Properties of pointlike Fields

The conceptual situation of QFT after the discovery of renormalized perturbation theory was at first somewhat confused. Despite the impressive agreement of low order radiative corrections, the precise relations between particles and fields as well as the mathematical consistency of QFT beyond perturbation theory were ill-understood. Most of the post renormalization progress was in the area of structural understanding about the particle-field dichotomy and on mathematical formulations of well posed physical requirements. These developments are often linked with the names of the principle protagonists of those problems: Lehmann, Symanzik and Zimmermann (LSZ) in the first case and Wightman in the second case (the Wightman framework).

The strong return of perturbative methods in the 70^{ies} via the Standard Model, only led to a temporary lull in the ongoing research on general structural properties of QFT, especially in view of the fact that those nonabelian gauge theories, after some initial success (notably in the area of small distance behavior off mass shell) run into tough nonperturbative problems which appeared unsolvable in the standard approach. Up to this date fundamental insights, as an intrinsic understanding of gauge theory on the basis of its gauge invariant correlation functions (i.e. without reference to the quantization method by which it has been constructed) are still missing (see also chapter 4.5).

We already have explained the relation between free fields and the net of local algebras generated by them in terms of an analogy to differential geometry: the fields are like coordinates and the net corresponds to the coordinate-free (intrinsic) approach to QFT. Many of the properties of fields appear in a clearer physical light, if one thinks about them in terms of local generators of nets, in analogy to the enveloping algebras of noncompact Lie algebras. Therefore let us list some properties (the main properties of the Haag-Kastler net theory) of nets before we write down the (model-independent) postulates for fields.

• (i) There is a map of double cones \mathcal{O} in Minkowski space into von Neumann operator algebras $\mathcal{A}(\mathcal{O})$ which are subalgebras of all operators $\mathcal{B}(\mathcal{H})$ in some Hilbert space \mathcal{H} :

$$\mathcal{A}:\mathcal{O}\to\mathcal{A}(\mathcal{O})$$

The C^{*}-completion of this family yields the global C^{*}-algebra \mathcal{A}_{quasi} :

$$\mathcal{A}_{quasi} = \bigcup_{\mathcal{O} \in \mathcal{M}} \mathcal{A}(\mathcal{O})$$

• (ii) The family \mathcal{A} forms a "net" i.e. a coherent (isotonic) family of local algebras:

if $O_1 \subset O_2$ then $\mathcal{A}(\mathcal{O}_1) \subset \mathcal{A}(\mathcal{O}_2)$

In case the local algebras represent observables one requires another physically motivated coherence property namely Einstein causality:

$$\mathcal{A}(\mathcal{O}) \subset \mathcal{A}(\mathcal{O}')'$$

• (iii) covariance with respect to the Poincare group. For observable nets:

$$\alpha_{(a,\Lambda)}(\mathcal{A}) = \mathcal{A}(\Lambda \mathcal{O} + a)$$

As already mentioned, the subsequent properties of fields and their physical interpretation is facilitated by thinking about them as coordinatizations of generators for local nets. The main difference of the field approach versus the net approach is that properties of charge-carrying fields are put in, and not derived from those of observable neutral fields. In the net approach charges (and their field carriers) are constructed via the superselection theory. The latter approach is more fundamental and more suitable in situations which are far away from quantization prescriptions and Lagrangians (e.g. low dim.QFT with braid group statistics). In the sequel we explain the properties of fields in the setting of Wightman. Here and in the following we use the symbol A as a generic notation for collection of generating fields but the standard situation underlying illustrations and proofs is mostly that of one generating scalar field.

Properties of Fields:

- <u>A</u> \mathcal{H} -space and \mathcal{P} -group
- 1. Unitary representation $U(a, \alpha)$ of $\widetilde{\mathcal{P}}$ in $\mathcal{H}, \widetilde{\mathcal{P}}$:covering of $\widetilde{\mathcal{P}}$
- 2. Uniqueness of the vacuum Ω , $U(a, \Lambda)\Omega = \Omega$
- 3. Spectrum condition: $specU \in \overline{V}_+$, the forward light cone.

• $\underline{\mathbf{B}}$ Fields

1. Operator-valued distributions: $A(f) = \int A(x) f(x) d^4x$, $f \in \mathcal{S}$ (the Schwartz space of "tempered" testfunctions) is an unbounded operator with a dense domain \mathcal{D} such that the function $\langle \psi_2 | A(x) | \psi_1 \rangle$ exists as a sesquilinear form for $\psi_i \subset \mathcal{D}$

- 2. Hermiticity: with A, also A^* belongs to the family of fields and the affiliated sesquilinear forms are as follows related: $\langle \psi_2 | A^*(x) | \psi_1 \rangle = \langle \psi_1 | A(x) | \psi_2 \rangle$
- <u>C</u>. $\widetilde{\mathcal{P}}$ -covariance of fields: $U(a, \alpha)A(x)U^*(a, \alpha) = D(\alpha^{-1})A(\Lambda(\alpha)x + a)$
- 1. For observable fields only integer spin representations (i.e. representations of ${\mathcal P}$) occur.
- <u>D</u>. Locality: $[A^{\#}(f), A^{\#}(g)]_{\mp} = 0$ for suppf×suppg (supports are spacelike separated).
- <u>E</u>. Stability of local algebras under causal completion: $Alg(A, \mathcal{O}) = Alg(A, \mathcal{O}'')$, where (for \mathcal{O} convex) the causal completion is the smallest double cone which contains \mathcal{O} . A weaker form of this requirement is the so called "time-slice" property.

Comments:

The domain requirements on (unbounded) smeared-out fields A(f) are reminiscent of properties which are required of generators of noncompact groups. The motivation for this Wightman postulates are entirely pragmatic; they insure that the standard calculational methods of physicists are applicable. These more technical domain requirements will be absent in the net approach. The only domain properties of the algebraic approach are the very fundamental (and physical) domain properties of the Tomita-Takesaki modular theory. But it turns out that as a result of the new concept of modular localization, the Wightman domain properties, far from being merely technical postulates, become carriers of important physical informations about properties of quantum versus geometric localization. The Wightman domain turns out to be simply the intersection of all the dense modular localization spaces.

The existence of the sesquilinear forms for pointlike fields is the substitute for the classical notion of field strength. The $\tilde{\mathcal{P}}$ -transformation property of the hermitian adjoint field is that of the complex conjugate transformation which is isomorphic to the antiparticle field: $A^{(\bar{\lambda})}(x) = CA^{(\lambda)*}(x)$, C=charge conjugation matrix.

Strictly speaking the causality requirement applies to locally observable fields only (example: electromagnetic field strength and currents, but not to vector potentials and charged matter fields). The restriction to local fields, which by definition obey the \mp commutation relations, is too strong in $d \leq 2 + 1$ (see last chapter). In d=3+1 all compactly localizable charge carrying fields are equivalent (by Klein-transformations) to local Fermi- or Bose-fields. We will use the term "localizable" instead of local for fields with noncompact localization properties.

The strong causal completion property is the substitute for an hyperbolic equation of motion (which, due to ill-defined nonlinear terms, is a priori meaningless in QFT). Its formulation and exploration is more natural in the algebraic setting where it simply means that $\mathcal{A}(\mathcal{O}) = \mathcal{A}(\mathcal{O}'')$.

Another physically important property which has been omitted here, but makes its appearance in the net theory later on, is the nuclearity or compactness property. This is the QFT counterpart of the statement that a finite cell in phase space can only accommodate a finite (in QFT a nuclear instead of finite set of vectors) number of degrees of freedom.

The most useful objects which one can form in such a Wightman setting of fields are the vacuum expectation values or (terminology of condensed matter physics) correlation functions:

$$w_{n}^{(\lambda_{1}...\lambda_{n})}(x_{1}....x_{n}) = \left\langle 0 \left| A^{(\lambda_{1})}(x_{1})....A^{(\lambda_{n})}(x_{n}) \right| 0 \right\rangle$$
(5.1)
$$= W_{n}^{(\lambda_{1}...\lambda_{n})}(\xi_{1},...,\xi_{n-1}) \quad \xi_{i} = x_{i} - x_{i+1}$$

$$= \left\langle 0 \left| A^{(\lambda_{1})}(0)e^{-i\mathbf{P}\xi_{1}}....A^{(\lambda_{n-1})}(0)e^{-i\mathbf{P}\xi_{n-1}}A^{(\lambda_{n})}(0) \right| 0 \right\rangle$$

$$= \int \cdots \int \tilde{W}_{n}^{(\lambda_{1}...\lambda_{n})}(q_{1}....q_{n-1})e^{-i\sum_{k}q_{k}\xi_{k}}d^{4}q_{1}....d^{4}q_{n-1}$$

The spectrum property: $spec(\mathbf{P}) \subset \overline{V}_+$ evidently implies that

$$suppW^{(..)}(q_1..q_{n-1}) \subset \otimes^n \overline{V}_+$$

and, as a property of a Fourier-Laplace transform of a cone supported distribution we encounter the "tube analyticity", namely W is boundary value of a function $W^{(..)}(z_1....z_{n-1})$ analytic in the tube $T^{(n-1)} z_i = \xi_i - i\eta_i$ with $\eta_i \in \overline{V}_+$ fulfilling the "tempered" bound (assuring the temperedness of the singular boundary values):

$$\left| W_n^{(..)}(z_1....z_{n-1}) \right| \le C \frac{\left(1 + \sum_i^{n-1} |z_i|^2 \right)^{\kappa}}{\left(\min_j(\eta_j^2) \right)^l}, \quad |z|^2 := \sum_{\mu} |z_{\mu}|^2 \tag{5.2}$$

This tube analyticity together with the Lorentz-invariance of the W's (a consequence of the invariance of the vacuum and the covariant transformation properties of the fields) yields the invariance under the complex Lorentz-group L^c :

$$L^{c} = \{A, B\} \quad A, B \in SL(2, C)$$

$$\underline{z} = \sigma_{\mu} z^{\mu} \to A \underline{z} B^{*}, \quad z^{\mu} z_{\mu} = inv.$$
(5.3)

This complex extension is a rather direct consequence of the previous analyticity and the fact that the finite dimensional representations $D^{(A,\dot{B})}$ permit an extension to a transformation in which the undotted and dotted spinors transform independently. The details can be looked up in the literature. L^c has

different from L, only two instead of four connected components: det= \pm . It takes some additional calculations to prove that the extended tube is of the form $T_{ext}^{(n-1)} = L^c T^{(n-1)}$. This is a natural analyticity region and the W are univalued. It is remarkable that T_{ext} contains real points. It is easy to see that the convex real set:

$$\xi_k^2 < 0, \quad (\sum_k \lambda_k \xi_k)^2 < 0 \quad \lambda_k \ge 0 \quad \sum_k \lambda_k > 0 \tag{5.4}$$

(the so called Jost points) is contained in T_{ext} . The locality binds all the n! different $w^{(n)}(x_{i_1}...x_{i_n})$ together to one (anti-)symmetric holomorphic "master function" with T_{ext}^{perm} the extended permuted tube being the enlarged analyticity region on which the permutations act:

$$w_n^{(\lambda_{i_1}\dots\lambda_{i_n})}(z_{i_1}\dots z_{i_n}) = \begin{cases} w_n^{(\lambda_1\dots\lambda_n)}(z_1\dots z_n)\\ signPw_n^{(\lambda_1\dots\lambda_n)}(z_1\dots z_n) \end{cases}$$
(5.5)

The mathematical structure behind this extension is the so called "edge of the wedge" theorem which generalizes the well known Schwarz reflection principle from one to several complex variables. The resulting "permuted extended tube" is not a natural holomorphy domain but its holomorphic completion is difficult to understand (and fortunately physically not as relevant as it appeared during the 60^{ies}). For a discussion of this and related matters we refer to the literature. One physically relevant fact is the univaluedness of the master function in d=3+1 theories. In d=1+1 the possibility of richer spacelike commutation relations (e.g. braid group statistics) which have multivalued master functions. Naturally its restriction to the real analytic (Jost) points is always univalued (the branching happens in the euclidean region), since otherwise the Hilbert space setting of quantum physics would get lost [18] [73].

The crucial question is now wether the family of w's with those properties following from the operator postulates suffice in order to reconstruct uniquely (up to isomorphism) the quantum field theory. From our experience with the GNS construction we would expect a positive answer. However the "field algebras" are not C^* -algebras of bounded operators and therefore a special construction which is more adapted to this problem is necessary. One uses the polynomial algebra $\mathcal{P}(M)$:

$$\left\{f_{0}\underline{1} + \sum_{n=1}^{N} \int \cdots \int f_n(x_1...x_n) A(x_1)...A(x_n) \mid f_n \in \mathcal{S}(\mathbf{R}^{dn}), \forall N\right\}$$
(5.6)

Here we have again supressed all Lorentz- and charge-indices i.e. we used our standard neutral scalar illustration. As in the case of CCR and CAR we can interpret the expectation values $w^{(n)}$ as affiliating a positive linear functional on a *- algebra of test functions:

$$f = \{f_0 \dots f_N\} \in \bigoplus_n \mathcal{S}(R^{dn}) \equiv T\mathcal{S}$$
(5.7)

$$(f \cdot g)_n(x_1...x_n) = \sum_k f_k(x_1..x_k)g_{n-k}(x_{k+1}..x_n)$$
(5.8)

$$(f^*)_n(x_1....x_n) = \overline{f_n(x_n....x_1)}$$
 (5.9)

Note that different from the CCR and CAR case this is not a Hilbert space of "one particle" functions but a tensor algebra T(M) on sequences of functions. Here M indicates that the testfunction space consists of functions on Minkowski-space. The localized polynomial algebra $\mathcal{P}(\mathcal{O})$ is a subalgebra of $\mathcal{P}(\mathcal{M})$. The vacuum expectation values w_n just define a state (positive definite normalized functional) on T:

$$w(f) = \sum_{n} W_n(f_n), \quad w(f^*f) \ge 0$$
 (5.10)

In the operator way of writing, this is just the positivity of the norm squared:

$$\left\| \left(f_0 \underline{1} + \sum_{n=1}^N \int \cdots \int f_n^{(\lambda_1 \dots \lambda_n)}(x_1 \dots x_n) A^{(\lambda_1)}(x_1) \dots A^{(\lambda_n)}(x_n) \right) |0\rangle \right\|^2 \ge 0$$
(5.11)

With an appropriately defined action α of $\mathcal P$ on the tensor algebra, W is covariant:

$$w(\alpha_{a,\Lambda}(f)) = w(f) \tag{5.12}$$

The reconstruction is completely analogous to the GNS situation. One obtains a triple (H, π, Ω) i.e. a *-representation (of the so-called Borchers-Uhlmann tensor-algebra) which is covariant with positive spectrum and a unique vacuum vector Ω . Certain properties as the time-slice requirement and its local version have no known equivalent in terms of correlation function; they need the reconstructed operator theory for their formulation.

5.2 Simple Structural Properties

1. The Cluster decomposition property. Its weak form is defined:

$$\lim_{\lambda \to \infty} W(f \alpha_{\lambda x}(g)) = \lim_{\lambda \to \infty} \left\langle 0 \left| A_f^* U(\lambda x) B_g \right| 0 \right\rangle = W(f) W(g) \quad (5.13)$$
$$A_f, B_g \in \mathcal{P}(M)$$

and results from the fact that only the discrete part of the energy momentum spectrum (i.e. the assumed unique vacuum contribution) survives, whereas the continuum oscillates to zero (Riemann-Lesbegue Lemma). The strong form is conveniently formulated in terms of the connected correlation functions:

$$\langle 0 | A_{f_1}(x_1) \dots A_{f_n}(x_n) | 0 \rangle_{con} \xrightarrow{clustering} 0$$

$$A_f(x) = U(x) A_f U^*(x), \quad A_f \in \mathcal{P}(M)$$

$$(5.14)$$

It uses locality (in order to disentangle overlapping clusters) and needs more mathematical effort for its derivation from the postulates. Note that a vacuum

degeneracy would show up in form of a very specific violation (containing information about the dimension of the vacuum projector) of the cluster property.

2.The Reeh-Schlieder Theorem. The localized polynomial algebra $\mathcal{P}(\mathcal{O})$ is cyclic and separating on Ω , i.e.

$$\overline{[\mathcal{P}(\mathcal{O})\Omega]} = H, \quad A\Omega = 0 \curvearrowright A = 0, \quad A \in \mathcal{P}(\mathcal{O})$$
(5.15)

For the cyclicity assume that $\psi \in [P(O)\Omega]^{\perp}$, $\psi \neq 0$. Then for $A_i \in \mathcal{P}(\tilde{O})$, $\tilde{O} \ll O$ (no boundary touching) define:

$$F(x_1....x_n) = \langle \psi | \alpha_{x_1}(A_1)....\alpha_{x_n}(A_n) | 0 \rangle \quad \psi \in \mathcal{D}$$

$$= 0 \quad on \left\{ (x_1....x_n) | x_i \in V, \ \tilde{O} + V \in O \right\}$$
(5.16)

The Fourier transform $\tilde{F}(p_1...,p_n)$ vanishes outside the support: $\cap_l \left\{ \sum_l^n p_i \in \bar{V}^+ \right\}$, as follows from the spectrum condition. Therefore also the matrix element F enjoys tube analyticity in $z_1...,z_n$ (instead of n-1 z's as the W's). They agree with the (obviously holomorphic) zero function in the above real neighborhood.. The already mentioned multi-dimensional generalization of the Schwarz reflection principle termed "edge of the wedge theorem" will then lead to the identical vanishing. But this contradicts the assumption of nontriviality of ψ , q.e.d.

The proof of separability of Ω with respect to $\mathcal{P}(\mathcal{O})$ can be reduced to cyclicity by using locality. We have:

$$AA'\Omega = A'A\Omega, \quad A \in \mathcal{P}(\mathcal{O}), \quad A' \in \mathcal{P}(\mathcal{O}')$$
 (5.17)

Since \mathcal{O}' is non-void, $\mathcal{P}(\mathcal{O}')$ acts cyclically on Ω and therefore $A'\Omega = 0 \curvearrowright A' = 0$ on the dense set $\mathcal{P}(\mathcal{O})\Omega$ and hence $A \equiv 0$.

With the Reeh-Schlieder theorem we have met the first characteristic property of local quantum physics. It has no counterpart in Schrödinger theory and general quantum theory. Indeed the idea that one can emulate vacuum excitations "behind the moon" by operating with hardware localized on an earthly laboratory with increasing accuracy, sounds somewhat exotic. It has led to many misunderstandings especially with respect to causality. One of the more spectacular conceptional mistakes even casts doubt on Fermi's conclusion that Einstein's causality statements about classical relativistic field theory are also valid in QFT [74]. On the positive side this property led to deeper thoughts about long range correlation and the proper operational formulation of causality as well as phase space localization of degrees of freedom (nuclearity).

3. Irreducibility of $\mathcal{P}(\mathbf{M})$ Starting from the time-development automorphism which (according to the positive energy assumption) is implemented by a positive Hamiltonian:

$$\alpha_t(A) = e^{iHt} A e^{-iHt}, \quad H \ge 0, \quad A \in \mathcal{P}(\mathcal{M}), \tag{5.18}$$

we study the analytic properties of matrix elements of time translated operators from the commutant:

$$f(t) \quad : \quad = \left\langle A_1 \Omega \left| e^{-iHt} A' e^{iHt} \right| A_2 \Omega \right\rangle = \left\langle \Omega \left| A_1^* \alpha_{-t}(A') A_2 \right| \Omega \right\rangle \quad (5.19)$$

$$A_i \in \mathcal{P}(\mathcal{M}), \quad A' \in \mathcal{P}(\mathcal{M})',$$
$$\mathcal{P}(\mathcal{M})' \equiv \{C \mid \langle A^*\phi, C\psi \rangle = \langle \phi, CA\psi \rangle \ \forall A \in \mathcal{P}(\mathcal{M}), \ \phi, \psi \in \mathcal{D} \}$$

One computes:

$$f(t) = \langle A'^*\Omega, \alpha_t(A_1^*A_2)\Omega \rangle = \langle A'^*\Omega, e^{itH}A_1^*A_2\Omega \rangle$$

$$= \langle \alpha_t(A_2^*A_1)\Omega, A'\Omega \rangle = \langle A_2^*A_1\Omega, e^{-itH}A'\Omega \rangle$$
(5.20)

The first line represents f(t) as a matrix element of e^{itH} , and the second of e^{-itH} . Therefore f is a bounded function which is simultaneously analytic in the upper and lower halfplane. According to a theorem of Liouville this forces f to be a constant i.e.

$$\begin{split} f(t) &= \langle A'^*\Omega, E_0 A_1^* A_2 \Omega \rangle = \langle \Omega, A'\Omega \rangle \langle A_1\Omega, A_2\Omega \rangle, E_0 = \text{proj on vac.} \\ &\curvearrowright \quad A' = \langle \Omega, A'\Omega \rangle \cdot \mathbf{1} \end{split}$$

4. TCP Symmetry. We first remind ourselves of the TCP-transformation property of free particles:

$$\Theta |p, \lambda, i\rangle = \sum_{\lambda'} |p, \lambda', \bar{i}\rangle D_{\lambda', \lambda}(i\sigma_2) \text{ in } H_1, \text{ antilinear}$$
(5.21)
$$\bar{i} : \text{ antiparticles of type i, } \Theta^2 = (-1)^{2s} \mathbf{1}$$

$$\Theta \Phi^{[A,\dot{B}]}(x)\Theta^{-1} = (-i)^F (-1)^{|\dot{B}|} \Phi^{[A,\dot{B}]*}(-x) \equiv \Phi^{\theta}(-x)$$
(5.22)

$$F \quad \# \text{ of fermions, } \begin{vmatrix} \dot{B} \end{vmatrix} = \# \text{ of dotted spinor indices}$$

We will show that this formula holds in general (for local interacting fields). If $\Phi = A$ =scalar field, the proof starts from first rewriting the content of TCP symmetry in terms of correlation functions:

$$\langle \Theta A(x_m) \dots A(x_1) \Omega, \Theta A(x_{m+1}) \dots A(x_n) \Omega \rangle$$

$$= \langle A(x_{m+1}) \dots A(x_n) \Omega, A(x_m) \dots A(x_1) \Omega \rangle$$

$$\Leftrightarrow w(-x_1, \dots - x_n) = w(x_n \dots x_1)$$

$$(5.23)$$

where in the last line we used the above Θ -action. We now take notice of the fact that by combining the symmetry relation from locality in T_{ext}^{perm} with the $L_+(C)$ invariance (which included the total reflection) we have:

$$w(x_n...x_1) = w(x_1...x_n) \stackrel{L_+(C)}{=}$$
 (5.24)

$$w(-x_1....-x_n)$$
 (5.25)

This means that we obtain the above relation in T, and hence also on the physical boundary (the boundary $i\varepsilon$ -prescription in the above relation is the same on both sides) which is the desired relation for the operators.

5. Spin & Statistics. If we require the wrong local commutation relations for $\Phi^{[A,\dot{B}]}$:

$$\{\Phi(x)\Phi(y)\} = 0, \quad (x-y)^2 < 0, \quad A+\dot{B} = \frac{n}{2}, n \text{ odd}$$
(5.26)
$$[\Phi(x)\Phi(y)] = 0 \quad (x-y)^2 < 0, \quad n \text{ even}$$

then $\curvearrowright \Phi \equiv 0$. With other words within the framework of local fields, the standard relation between spin and statistics is a consequence of the postulates. The proof (again for neutral scalar fields) only employs the two-point function W(z)which is analytic in T_{ext} and fulfills (as a consequence of $L_+(\mathbf{C})$ -invariance) W(z)=W(-z):

$$\langle 0 | \{ \Phi(x), \Phi(y) \} | 0 \rangle = W(\xi) + W(-\xi) = 0, \ (x - y)^2 < 0$$
 (5.27)
 $\sim 2W(z) = 0, \ \sim \Phi(x)\Omega = 0$

Finally the Reeh-Schlieder theorem gives $\Phi \equiv 0$. The general case with dotted and undotted spinors is left to the reader.

The TCP and Spin&Statistics theorem are considered to represent the deeper parts of structural QFT. They even gave the title for the first book on the subject [18]. Later we will see that they continue that role in algebraic QFT with an additional gain in profoundness (in a particular evident form in low dimensional QFT)..

6. Normal C.R. and Klein transformations. The previous theorem left open the commutation relations between different Lorentz-multiplets. One defines as "normal" the spacelike commutativity of two local fields of which at least one is bosonic, as well as the spacelike anticommutativity between two fermionic $(A + \dot{B} = \text{halfinteger})$ fields $\psi^{(A,B)}$. As a preparatory step towards proving that normal commutation relation can always be achieved, let us prove that commutation relations remain stable under transition to the hermitian adjoint field:

$$[\Phi_1(x), \Phi_2^*(y)]_{\pm} = 0 \quad (x-y)^2 < 0 \tag{5.28}$$

$$\curvearrowright \quad [\Phi_1(x), \Phi_2(y)]_{\pm} = 0 \quad (x-y)^2 < 0$$

The proof uses the cluster decomposition property (i.e. the uniqueness of the vacuum):

$$\langle \Omega, \Phi_1^*(f) \Phi_2^*(g) \Phi_2(g) \Phi_1(f) \Omega \rangle$$

$$= \| \Phi_2(g) \Phi_1(f) \Omega \|^2 \ge 0$$

$$\stackrel{loc.}{=} \sigma \langle \Omega, \Phi_1^*(f) \Phi_1(f) \Phi_2^*(g) \Phi_2(g) \Omega \rangle$$

$$\stackrel{cluster}{\Rightarrow} \sigma \| \Phi_1(f) \Omega \|^2 \| \Phi_2(g) \Omega \|^2$$

$$(5.29)$$

Here $\sigma = \pm$. Consistency requires that σ agrees with the ± 1 in the original C.R.. between Φ_1 and Φ_2^* . Now we are prepared to construct the Klein transformation which carries anomalous into normal commutation relations. For the typical anomalous situation assume that:

$$[\varphi(x), \psi(y)] = 0, \quad (x - y)^2 < 0$$

$$\varphi : \text{ bosonic, } \psi : \text{fermionic}$$

$$(5.30)$$

Define:

$$\begin{aligned}
\varphi'(x)H_{even} &= \varphi(x)H_{even} \\
\varphi'(x)H_{odd} &= -\varphi(x)H_{odd} \\
\psi'(x)H_{even} &= \psi(x)H_{even} \\
\psi'(x)H_{odd} &= \psi(x)H_{odd}
\end{aligned}$$
(5.31)

or briefly: $\varphi'(x) = (-1)^{\mathbf{F}} \varphi(x), \quad \psi'(x) = \psi(x).$ One obtains:

$$\{\varphi'(x), \psi'(y)\} = 0, \quad (x-y)^2 < 0$$
 (5.32)

The general situation is analogous.

=

7. Characterizations of free fields. Assume first that the two-point function agrees with that of a free field, i.e.

$$\langle \Omega, \varphi(x)\varphi(y)\Omega \rangle = i\Delta^{(+)}(x-y) \curvearrowright$$

$$\varphi(x) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int (e^{-ipx}a(p) + h.c.)\frac{d^3p}{2\omega}$$

$$(5.33)$$

in case of a neutral scalar field. The first step in the proof consists in deriving the Klein-Gordon equation $(\partial^{\mu}\partial_{\mu} + m^2)\varphi(x) = 0$. From the two-point function one obtains

$$\langle \Omega, j(x)j(y)\Omega \rangle = 0, \quad \text{where } j(x) := (\partial^{\mu}\partial_{\mu} + m^2)\varphi(x)$$
 (5.34)
 $\sim j(x)\Omega = 0$

The analytic properties in the tube T of the mixed j- φ correlation functions together with the relative spacelike commutativity bring about an edge of the wedge situation with:

$$\langle \Omega, \varphi(x_1)..., j(x_i)\varphi(x_{i+1})...,\Omega \rangle$$

$$= \langle \Omega, \varphi(x_1)..., \varphi(x_{i+1})..., j(x_i)\Omega \rangle = 0$$

$$(5.35)$$

on an open set of the boundary and hence the vanishing of all matrix elements of j on the dense domain \mathcal{D} i.e. $j(x) \equiv 0$. Therefore φ indeed fulfills the free field equation and hence permits a frequency decomposition:

$$\varphi(x) = \varphi^{(-)}(x) + \varphi^{(+)}(x), \quad \varphi^{(-)}(x)\Omega = 0$$
 (5.36)

A characterizing property of free fields is their c-number (anti)commutator, in our case:

$$[\varphi(x),\varphi(y)] = i\Delta(x-y)\mathbf{1}$$
(5.37)

But this follows by again using analyticity properties. First we use the spectrum condition to obtain:

$$\varphi^{(-)}(x)\varphi^{(+)}(y)\Omega = i\Delta^{(+)}(x-y)\Omega$$
 (5.38)

since $\varphi^{(\pm)}$ transfers momentum on the forward (backward) mass shell and hence the spectral transfer of the product is spacelike including zero i.e. the intersection with the physical spectrum consists of just p = 0 corresponding to the vacuum vector Ω . This is a much stronger statement than the assumed twopoint function structure. For the commutator applied to Ω we now have:

$$\left[\varphi(x),\varphi(y)\right]\Omega = i\Delta(x-y)\Omega + \left[\varphi^{(+)}(x),\varphi^{(+)}(y)\right]\Omega \tag{5.39}$$

Let $\psi \in \mathcal{D}$ and consider the analytic properties of:

$$F(x,y) := \left\langle \psi, \left[\varphi^{(+)}(x), \varphi^{(+)}(y)\right] \Omega \right\rangle$$
(5.40)

The momentum transfer of each $\varphi^{(+)}$ is on the forward mass shell, and hence this distribution is the boundary value of a function $F(z_1, z_2)$ analytic in $z_i = x_i - iy_i$, $y_i \in V^+$. Since this function vanishes in a neighborhood of the real boundary, $\sim F \equiv 0$. But:

$$B(x,y) := [\varphi(x),\varphi(y)] - i\Delta(x-y)\mathbf{1}$$
(5.41)

is a bilocal operator-valued distribution ($B(f,g)\in \mathcal{P}(\mathcal{O})$) for which Ω is a separating vector, i.e.

$$B(x,y)\Omega = 0 \frown B(x,y) \equiv 0 \quad q.e.d. \tag{5.42}$$

This property has no analogue in quantum mechanics i.e. the pair interation does not show up in the nonrelativistic two-point function (related to the absent selfinteraction). This conclusion continues to hold for semiinfinite string-localized operator-valued solutions of free field equations[91].

Remembering that a free field has vanishing connected n-point correlation functions for n>2, the question arises whether this property is typical. The affirmative answer is:

if
$$\exists n > 2 \quad w_n(x_1, \dots, x_n)^{conn} = 0$$
 (5.43)
 $\curvearrowright \quad \varphi(x)$ is a generalized free field

A generalized free field shares with the free field the property of having a cnumber commutator. But this commutator is a (continuous) superposition of free field commutators:

$$[\varphi(x),\varphi(y)] = \int d\rho(\kappa^2) i\Delta(x-y,\kappa^2)$$
(5.44)

We will not prove the above statement here.

Another characterization of (generalized) free fields not presented here is in terms of gaps in the spacelike momentum transfer of fields. Reductions to generalized free fields are effectively reductions to free fields in view of the time slice property and in particular of the phase space nuclearity property presented in a later section.

8. Shape of Energy Momentum Spectrum. The asymptotic factorization or clustering of correlation functions suggests that the energy-momentum spectrum specP is an additive set i.e. with $p_1, p_2 \in \text{specP}$ also $p_1 + p_2 \in \text{specP}$. To see this consider the vector:

$$\psi_{21}(a) = U(a)A_2U^*(a)A_1\Omega, \quad A_i \in \mathcal{P}(\mathcal{M})$$
(5.45)

Assume that the energy-momentum transfer is limited to regions $\Delta_i \in \text{specP}$. Then the Fourier-transform of ψ has its support in $\text{supp}\tilde{\psi} \in \Delta_1 + \Delta_2$. The clustering:

$$\lim_{n \to \infty} \left\| \psi_{21}(a) \right\|^2 = \langle \Omega, A_2^*(a) A_2(a) \Omega \rangle \langle \Omega, A_1^* A_1 \Omega \rangle$$

$$= \left\| \psi_2 \right\|^2 \left\| \psi_1 \right\|^2, \quad \psi_i = A_i \Omega$$
(5.46)

serves to show that $\|\psi_{21}(a)\| \neq 0$ i.e. does not vanish identically thus assuring the nontriviality of the vector carrying the sum of the momenta.

Classically the hyperbolic causal propagation in classical field theory is inexorably linked with Lorentz-covariance. By analogy one would expect that causality, even if it does not extend translational covariance to full Poincaré covariance, at least forces the energy-momentum spectrum to have a Lorentzinvariant shape. Indeed, the implementer of the translation can always be chosen in such a way:

$$\exists U(a)s.t.\alpha_a(A) = U(a)AU^*(a)$$

$$U(a) = e^{i\mathbf{P}a}, \quad spec\mathbf{P} \text{ inv. shape}$$
(5.47)

This theorem is easier to prove in an algebraic setting and hence will be deferred.

5.3 Euclidean Fields

C

Analytic continuations through Wick-rotation have been useful in perturbation theory, because certain regularization techniques (e.g. the dimensional regularization) only work if noncompact L-invariance can be replaced by compact euclidean invariance. Therefore it is interesting to know if a euclidean formulation is also possible outside perturbation theory and whether it is useful there. Schwinger and later Symanzik and Nelson were the first to realize that a euclidean framework indeed opens a useful connection with statistical mechanics and classical probability theory.

The starting point for a nonperturbative euclidean approach is the analyticity and univaluedness of the analytic extension of correlation functions

into the extended permuted tube T_{perm}^{ext} . It is obvious that the non coinciding $(\hat{x}_i \neq \hat{x}_j \forall i, j)$ euclidean points are inside this domain. The Wick-rotation $(\vec{x}, x^0) \rightarrow (\hat{x} = \vec{x}, \hat{x}^0 = ix_4)$ relates the Minkowski inner product with the Euclidean one and the group $\mathbf{O}_+(4)$ with a subgroup of $\mathbf{L}_+(\mathbf{C})$. Here and in most of what follows we present the euclidean formulation for integer spin fields, the adaptation to halfinteger spin will be commented on later. The restriction of the analytically continued correlation functions to the euclidean points $(\vec{x}, x_4) \in \mathbf{E} = \mathbb{R}^d$ are called Schwinger functions:

$$s(x_1....x_n) = w(\hat{x}_1....\hat{x}_2)$$
(5.48)
= $S(\xi_1....\xi_{n-1})$

where we used translation invariance in the last line. As for time-ordered functions, there is no spectrum condition which assures that they are distributions on the Schwartz-space S, rather their natural domain of definition are those test-functions which vanish at coinciding points of sufficiently high order. If the dimension of the fields is canonical i. e. for scalar fields dim $\varphi = \frac{d}{2} - 1 = \dim(\text{free}$ field) then S is naturally (i.e. without Hahn-Banach extension) integrable and hence a $S(\mathbf{E}^n)$ distribution. We now collect those properties of the Schwinger functions which allow to reconstruct a local Poincaré-invariant QFT. These properties are called the Osterwalder-Schrader axioms. In the following we present these axioms for the illustrative case of scalar neutral fields.

• **S1** The Schwinger functions are translation invariant real analytic function for non coinciding euclidean variables. They are distributions in $\mathcal{S}'(\mathcal{E}^{n-1}_{-})$ with $\mathcal{E}^{n-1}_{-} = \{\xi \in \mathbf{E}^{n-1} \mid \xi_1^4 < \xi_2^4 \dots < \xi_n^4\}$ where $\mathcal{S}(\mathcal{E}^{n-1}_{-})$ is given a weaker topology which is defined by the following system of seminorms:

$$\|f_{<}\|_{l,m} = \left\|\tilde{f}\right\|_{l,m} \quad f_{<} \in \mathcal{S}(\mathcal{E}_{-}^{n-1})$$
 (5.49)
$$\tilde{f}(q_{1}....q_{n-1}) = \int \dots \int e^{\sum_{1}^{n-1} (\xi_{j}^{4} q_{j}^{0} + i\boldsymbol{\xi}_{j} \mathbf{q}_{j})} f_{<}(\xi_{1}....\xi_{n-1}) d^{d}\xi_{1}....d^{d}\xi_{n-1}$$

Here we used the property of the Laplace-Fourier transforms of mapping continuously $\mathcal{S}(\mathcal{E}^{n-1}_{-})$ onto a dense set in $\mathcal{S}(\mathcal{M}^{n-1}_{+})$ which are Minkowskispace test functions \tilde{f} with

$$supp\tilde{f} \in \left\{ q \in \mathbf{M}^{n-1} \mid q_i^0 \ge 0 \ \forall i \right\} \equiv \mathcal{M}_+^{n-1} \tag{5.50}$$

The above topology is the one which $\mathcal{S}(\mathcal{E}^{n-1}_{-})$ inherits from $\mathcal{S}(\mathcal{M}^{n-1}_{+})$ through this map. The Schwinger distributions are just the continuous linear functionals on $\mathcal{S}(\mathcal{E}^{n-1}_{-})$ in this topology. It is the analog of the growth condition on the holomorphically extended correlation functions W which insured the temperedness of their distributional boundary values and often called the Osterwalder-Schrader growth condition.

• S2. Hermiticity. For the Schwinger functions of a scalar neutral field:

$$s(x_1....x_n) = \overline{s(Tx_1....Tx_n)}$$
(5.51)
$$Tx = (\vec{x}, -x_4) \quad \text{euclidean time reversal}$$

• **S3.** Reflection-Positivity:

$$\sum_{n,m} \int s(Tx_m, ..., Tx_1, y_1, ...y_n) \overline{f_m(x_1, ...x_m)} f_n(y_1, ...y_n) d^d x_1 ... d^d y_n \ge 0$$
(5.52)

where the sum only involves a finite sequence of test functions

 $(f_0, f_1 \dots f_n \dots f_N)$ with their support on the time simplex

$$\mathbf{E}_{<}^{n} = \left\{ x \in E^{n} \mid 0 < x_{1}^{4} < \dots < x_{n}^{4} \right\}$$
(5.53)

Clearly this property is the analogue of the Wightman-positivity for the W's. In fact it results from the positivity of "euclidean states":

$$\psi(x_1....x_2) = \varphi(\vec{x}_1, ix_1^4)....\varphi(\vec{x}_n, ix_n^4)\Omega, \quad x \in \mathbf{E}_{<}^n$$
(5.54)

Note that the spectrum condition allows to interpret the analytic continuation as a smearing with a an exponential damping factor (fast decreasing test function in time).

• **S4.** Euclidean covariance:

$$s(Rx_1....Rx_n) = s(x_1....x_n)$$
(5.55)

• **S5.** Permutation symmetry:

$$s(x_{P(1)}, \dots, x_{P(n)}) = s(x_1, \dots, x_n)$$
(5.56)

• **S6.** Cluster property:

$$\lim_{a \to \infty} \int s_n(x_1, \dots, x_m, x_{m+1} + a, \dots, x_n + a) f(x_1, \dots, x_m) g(x_{m+1}, \dots, x_n) (5.57)$$
$$= \int s_m(x_1, \dots, x_m) f(x_1, \dots, x_m) \times \int s_{n-m}(x_{m+1}, \dots, x_n) g(x_{m+1}, \dots, x_n)$$

The generalization to charged fields with arbitrary finite spin is obvious: the covariance law involves the representations of the $SU(2) \times SU(2)$ covering of O(4) and the permutation symmetry carries an additional sign(P)

It is fairly obvious that a theory in terms of correlation functions fulfilling positivity, hermiticity, P-covariance and locality leads to Schwinger functions

fulfilling S1-S6. One just defines euclidean vectors $\psi(x_1, ..., x_n)$ as above. The reflection positivity allows to equip the linear vector space:

$$\left\{\sum_{n=1}^{N} \int \dots \int f_n(x_1, \dots, x_n) \psi(x_1, \dots, x_n) \mid f_n \in \mathcal{S}(\mathbf{E}^n_{<})\right\}$$
(5.58)
$$\mathcal{S}(\mathbf{E}^n_{<}) \text{ with } \|f\|_{l,m} - \text{topology}$$

with a positive semidefinite inner product. Factoring out the null-space and forming the closure one obtains a euclidean Hilbert space which thanks to the Reeh-Schlieder theorem is equal to the GNS space of the real time correlation functions. The short-distance growth condition of the W's in the tube (controlling the temperedness of the distributional boundary values) are equivalent to the $\|\cdot\|_{lm}$ topology of the Schwinger functions:

$$s(x...x) = \langle \Omega, \psi(x...x) \rangle \tag{5.59}$$

The permutation symmetry of s is a result of that symmetry for the analytic w's (from locality). Actually already the $\psi's$ are symmetric as real analytic functions in the euclidean domain for $x_i \neq x_j, i \neq j, x_i^4 > 0$, as can be shown by he application of the edge of the wedge theorem. Note that the Osterwalder-Schrader (euclidean) reflection positivity S3 cannot be interpreted as a state on a *-algebra (the Borchers-Uhlmann tensor algebra of functions) but only serves to define a scalar product onon a linear space (finite sequences of test functions $f_n \in S(E_{\leq}^n)$. The reconstruction of the real time theory can then be carried out in two different ways. Either one uses functional analysis (contractive properties of space-time semigroups) or the analytic properties of the previous Laplace-Fourier transforms in S1 which relate the Schwinger distributions $\in S(\mathcal{E}^{n-1})$ to the spectral supported correlation functions $\tilde{W} \in \mathcal{S}(\mathcal{M}^{n-1}_+)$ and carries the reflection positivity into the W positivity. The latter method is more appropriate in the present context whereas the first method also works in situations without space-time analyticity e.g. the derivation of the transfer matrix formalism in classical statistical mechanics on a lattice (see a later section). We collect the result:

Theorem 18 (Osterwalder-Schrader) Every set of Schwinger functions with S1-S6 comes from a real time QFT by analytic continuation and restriction to the euclidean points.

The euclidean framework described here is primarily a structural reformulation, it does not really solve any problem of the real time theory which the latter is unable to solve by itself. In fact even from a mathematical viewpoint some of the axioms look somewhat mocked up, since the topology we used on $S(E_{<}^{n})$ is not natural. Only under very special circumstances it becomes a powerful constructive tool of QFT. This happens e.g. if the Schwinger functions allow an interpretation in terms of a continuous classical statistical mechanics. Mathematically this amounts to the Feynman-Kac representability of the Schwinger functions in terms of a (infinite dimensional) functional measure theory e.g. (the φ^4 -theory):

$$s(x_1, \dots, x_n) = \frac{1}{Z} \int [d\varphi] e^{-A[\varphi]} \varphi(x_1) \dots \varphi(x_n)$$

$$A[\varphi] = \frac{1}{2} (\partial \varphi \partial \varphi + m^2 \varphi^2) + g \varphi^4$$
(5.60)

A physically fruitful formal interpretation is in terms of a continuous version of a Gibbs formula for classical statistical mechanics on a lattice:

$$\langle \varphi(x_1)....\varphi(x_n) \rangle_{Gbbs} = \lim_{\Lambda \to \infty} \frac{1}{Z_\Lambda} \sum_{conf,\Lambda} e^{-\beta H_\Lambda[\varphi]} \varphi(x_1)....\varphi(x_n) \quad (5.61)$$
$$Z_\Lambda = \sum_{conf,\Lambda} e^{-\beta H_\Lambda[\varphi]}$$

Here the dynamical variables φ over each lattice point take on either values in a discrete (e.g. \mathbf{Z}_n) or continuous manifold (e.g. \mathbf{C} , SU(2) etc.) in which case the sum over configurations represents an integral over the field values at each lattice point within the volume Λ . There are two questions to be asked:

- (i) can one work out a measure theory for stochastic variables such that the above functional integrals have mathematical meaning?
- (ii) can one control "critical limits" (second order phase transitions) of classical statistical mechanics precisely enough in order to obtain possibly existing local QFT?

Deferring the second problem to a later section, we comment here only on the first one, namely the relation between a Nelson-Symanzik stochastic euclidean theory and real time QFT. Euclidean fields are continuous linear maps ϕ from test function spaces $S(E^d)$ into random variables over a probability space (Q, Σ, μ) with μ a normalized measure on Q and Σ the μ -measurable subsets. Let us define a generating functional W for the euclidean correlation functions of ϕ in a reference state (the euclidean "vacuum") which has the following properties:

$$S(f) = \int_{Q} e^{i\phi(f)} d\mu, \quad i.e.S(0) = 1, \ S(f) = \overline{S(-f)}, \tag{5.62}$$

S(f) is of positive type (5.63) and invariant under euclidean time reflections.

Here we may declare any axis to be the time axis. According to a famous theorem of Minlos, this measure-theoretical setting is equivalent to the following (Nelson-Symanzik) positivity and covariance properties of the functional S(f) (the functional Fourier-transform of μ):

$$\sum_{i,j=1}^{n} \bar{c}_i c_j S(f_i - f_j) \geq 0, \quad S(0) = 1$$

$$S(f) = S(\vartheta f), \quad S(f) = S(\alpha_{a,R} f)$$
(5.63)

the last line expressing the time reflection ϑ (the choice of the time axis is arbitrary) and euclidean invariance. In addition S(f) is continuous on \mathcal{S} in the Schwartz topology.

This setting of euclidean fields is obviously appropriate for the Feynman-Kac representation which assumes that the measure μ on the space of field configurations is given by an invariant statistical mechanics-like local "Hamiltonian" which consists of a quadratic free and a polynomial interacting part. We already know that the validity of the reflection positivity is a prerequisite for obtaining real time local quantum physics. It is not difficult to prove that such a stochastic euclidean theory with reflection positivity is equivalent to a special class of real time QFT namely the so called stochastic positive QFT.

Definition 5 A QFT is said to fulfill stochastic positivity if its associated von Neumann algebra \mathcal{A} contains an abelian subalgebra \mathcal{B} ("fields at one time") and an automorphism α_t ("time translation") such that: $\bigcup_t \alpha_t(\mathcal{B}) = \mathcal{A}$

Theorem 19 [37] A reflection positive stochastic euclidean theory is equivalent to a stochastic positive real time QFT.

Hence the equivalence requires the stochastic theory to have an additional QFT positivity property (reflection positivity) and the QFT to posses an additional stochastic (Nelson-Symanzik) positivity. We will not prove this theorem since our main motivation here is pedagogical namely to counteract the erroneous but widespread belief that QFT can be always be defined in terms of measure theory or Feynman-Kac Formulas. Only theories which "stay close" to the d=1+1 ϕ^4 -theory (the standard relativistic illustrative example of the above theorem) allow for a Feynman-Kac representation. Whatever the intuitive appeal of Lagrangian quantization and functional integrals may be worth, one of its conceptual and mathematical limitation is set by the above theorem. In quantum mechanics involving charges coupled to vectorpotentials, it is possible to go somewhat beyond the above standard setting of euclidean functional integrals at the expense of loosing the tight physical relation of the euclidean theory to statistical mechanics. But no such framework is known for e.g. the Chern-Simons Feynman-Kac representations.

Note that we are here not concerned with mathematically fine points caused by renormalization (e.g. ϕ^4 in d=1+2 or d=1+3) wrecking the canonical (equal time) structure. Rather we mean that certain theories are *structurally incompatible* with the stochastic Feynman-Kac representations; they simply do not even posses a formal Feynman-Kac like representation as the ϕ_4^4 theory. Examples are chiral conformal theories and, as mentioned before, d = 1 + 2 theories with braid group statistics (Chern-Simons actions). They are easily shown to fail on the stochastic positivity property. The reason is the nonexistence of an abelian subalgebra with the required density property.

On the other hand, if there is any quantum theory at all associated with the Chern-Simons Lagrangian, then the combinatorial theory [75] defined by the Markov trace on the ribbon braid group RB_{∞} i.e. the theory behind the knot

invariant and the associated 3-manifold invariants of V. Jones [26] is the only reasonable candidate. Witten introduced rules for Wilson loops which indeed give this result [77]. Additional support comes from algebraic QFT which finds these invariants in the type II_1 intertwiner algebras within the DHR theory of superselection sectors (see last chapter). The system of intertwiners between different sectors together with the Markov trace forms a "combinatorial QFT" par excellence [89]. In this form the algebras underlying the topological theories was known already in the famous 1969 DHR work [3]. Of course the DHR analysis becomes richer and more interesting for the more recent braid group statistics in low dimensional field theory in which case one obtains the Jones knot invariants as well as the invariants of 3-mf. of Witten's approach (see appendix of [89]). Mathematically very close related to the algebraic QFT approach to intertwiner algebras is the combinatorial theory which does not use physical principles but rather quantum group methods. This is due to Reshetikin, Turaev and Viro [75] and was extended by Karowski and Schrader [76]. Neither this method nor the Witten method furnishes a physical interpretation .Only the intertwiner approach of algebraic QFT which places the type II combinatorical intertwiner algebra into a localizable (and hence interpretable) QFT carries such information. Witten's rules for extracting the knot and 3 mf. invariants from the Chern-Simons action is however somewhat surprising and poses the question, whether this observation can also be understood in the spirit of algebraic QFT by putting suitable states on e.g. appropriately extended Weyl-like algebras. It is has been my firm belief that the use of singular state will remove the "meat" of these Weyl-like algebras and just leave the combinatorial "bones" [74][5]. This, if true, would bridge the gap between the approach of Witten, which looks like continuous QFT and the combinatorial approach and also answer some questions concerning the relation between real time and euclidean time. As it stands, the situation presents a very interesting paradox.

Singular states harmonizes very well with the formal idea of integrating over infinitely many *gauge copies* in euclidean path integrals, except that singular states are more noncommutative and in this way reconcile the difference between real time and imaginary time theories: the time development automorphism of the Weyl like algebras is wiped out by the singular nature of states. Whereas in the standard formulation of gauge theories there is no mathematical veto against considering non gauge-invariant formally space-time dependent correlation functions, singular states create such a mathematical veto. This aspect of singular states is very desirable, because something which is unphysical, should also be unmathematical. The idea is that whereas in "full" gauge theories with physical photons and matter the singular states become regular on a huge translation covariant subalgebra [96] (the gauge invariant algebra of the quantization picture), in Chern-Simons theories the regular part is so small [74] that it can only support combinatorial type II_1 data. In fact singular states are the only states which are capable of distinguishing a subalgebra. In quantum mechanics these states are usually excluded by the regularity assumption, i.e. one is only interested in those representations of the Weyl algebra (the von Neumann uniqueness theorem) for which the translations $U(\alpha)$ are continuous in α . The

only known situation in QM where singular states are apparently needed is the Hofstedter model of particles in a constant magnetic field (mathematically: the "noncommutative torus"). The abelian \mathbf{Z}_n -Chern Simons theory in the real time formulation is a maximally extended QFT of one-forms in d=2+1 where the extension is done by admitting semiinfinite one forms which are closed but not exact in the angular variable. Since I am not going to use such an approach in these lectures, I refer the interested reader to [74] where he finds references to important previous work on singular states. Unfortunately there exists presently no solution of this interesting problem of topological versus combinatorial theory.

Excepting the correctness of this picture, it is not reasonable to attempt to extract a theory of anyons from the pure Chern-Simons theories. In fact the only physical use of topological field theory should be the illustration of the working of singular states. In the last chapter we will use an extension of the d=2+1 Wigner particle classification method because we expect more concrete results on "free" anyons and plektons than with the Chern-Simons approach. In this way we also follow the historical route, since free Bosons and Fermions where first obtained by operator-methods and Wigner theory before this description was transcribed (in connection with perturbation theory) into functional language.

A closely related, conceptually more robust constructive idea is to try to define QFT as scaling limits of mathematically controllable lattice systems instead of working with formal Feynman-Kac representations. The guiding principle (going back to Kadanoff, Wilson and others) was to use the possible existence of second order phase transitions ("criticality") to loose the memory of the lattice and recover \mathcal{P} -covariance and locality. This approach always has a "light" start since the mathematical control of lattice systems is rather simple. But in the last step, the investigation of criticality and the execution of the scaling limit, one has to pay heavily for the easy life at the beginning. The mere control of existence via the various lattice inequalities is not enough, the last step requires a deep structural understanding of the lattice theory. Whereas it is true that most of the QFT concepts as conserved charges, particles, multiparticle scattering, antiparticles etc. can be transferred to the lattice (albeit with much more sweat, since the helpful causality structure is absent), a sufficiently detailed structural control is only possible under special circumstances as integrability (meaning the Yang-Baxter structure for 2-dim. lattice systems, as in the transition from the discrete Ising model to the continuous Ising QFT). This kind of temporary practical restriction is quite different from the above restriction through lack of Feynman-Kac representability. In particular there is no limitation on the short-distance behavior: the operator short-distance dimensions of e.g. the Ising-, Thirring-, RSOS- etc. models in the scaling limit is too far away from canonicity as to permit a euclidean F-K representation. Real time short distance singularities which go significantly beyond canonical behavior do not endanger the existence of real time QFT, but only limit certain methods as quantizations by functional integrals. We do not really pursue a lattice approach and refer the interested reader can find details on this subject in a later section. Our main constructive contribution (presented only after the chapter on algebraic QFT) will be based on the net approach.

5.4 Scattering Theory

Whereas scattering theory in e.g. Schrödinger QM is very important for the comparison of theory with experiments but less so for the formulation and construction of quantum mechanical models, the S-operator takes on a more fundamental significance in local quantum physics. The reason is threefold: in addition to its standard role of permitting experimental verification of the theory, S is an invariant of the net (i.e. S is attached to a Borchers class and should not be affiliated with individual fields) and finally S is related to the modular reflection J for the wedge algebra and the TCP-operator θ by $S = JJ_0 = \theta\theta_0$ where the subscript zero refers to the incoming fields (considered as a free theory). In this section we will present the scattering content and the class invariance property of S. The S-matrix adds an important aspect to the particle-field dichotomy. Whereas the renormalization in the sense of physical parametrization required the understanding of the relation of one-particle properties and fields (or local observables in the algebraic approach) the S-matrix deepens this connection by resolving continuum states into multiparticle scattering states. With other words, all the important aspects of the interpretation of the QFT formalism are determined by the basic causality and spectral concepts of the theory and nowhere does one have to add prescriptions from the outside. The completeness property separates QFT from other theoretical attempts about fundamental physics as e.g. string theory.

In the perturbative approach we already met the S-matrix as the adiabatic limit of S(g). But we also realized that from a conceptual point of view such limits should be avoided since that formalism is good for the local net properties, but becomes unnatural for the calculation of "on-shell" quantities, in particular for the scattering operator. The conceptually most satisfying method is to first calculate (the approximations for) the correlation function and then to use the scattering theory for on-shell quantities. Similar to the nonrelativistic theory, the main objective is to use the time dependent formulation because of its physical clarity, and to convert its content into analytically simple stationary formulas.

This aim is accomplished in the Lehmann-Symanzik-Zimmermann (LSZ) approach. As quantum mechanical time dependent scattering theory relates interacting wave functions for $t \to \infty$ with those of a free system, scattering theory in QFT should relate interacting (Heisenberg) fields with free fields. By checking with stationary external source models as well as with renormalized perturbation theory these authors proposed the following asymptotic condition (for the standard scalar situation):

$$\lim_{t \to \pm\infty} \langle \phi | A_f(t) | \psi \rangle = \langle \phi | A_f^{ex} | \psi \rangle \quad ex = out, in$$
(5.64)

$$A_f(t) = \int_{x^0 = t} f(x) \stackrel{\leftrightarrow}{\partial}_0 A(x) d^3x$$

Here f(x) is a solution of the Klein-Gordon equation, A_f^{ex} is defined by the same formula with A replaced by the free incoming or outgoing field (and therefore time-independent) and the state vectors ϕ, ψ are taken from a dense set of in states (with nonoverlapping wave functions in velocity space, as we know nowadays). Later the Haag-Ruelle formulation, which is based on strong convergence, was derived from the locality and spectral principles of QFT and it was shown that (5.64) follows. But before we discuss these refinements, we will derive the useful LSZ reduction formulas from (5.64).

Let us start with the reduction of an incoming particle in the following matrix-element:

$$\sup_{t \to -\infty}^{out} \langle f_{n+1}....f_{n+m} | A(x) | f_1....f_n \rangle^{in}$$

$$= \lim_{t \to -\infty}^{out} \langle f_{n+1}....f_{n+m} | A(x)A_f(t) | f_2....f_n \rangle^{in}$$

$$= \left\langle f_{n+1}....f_{n+m} \left| \int K_y T A(x)A(y)f(y)d^4y \right| f_2....f_n \rangle^{in} + c.t.$$

$$(5.65)$$

where T denotes the time ordering, K is the Klein-Gordon operator and c.t.(contraction terms) is the generic notation for terms in which f's in the in or out states have been contracted with resulting $(f_i, f_j) \times \text{lower terms}$ (example: the annihilation part of A_f^{in} may contract with f_i in the in state if the overlap is nonvanishing). In the third term the time ordering occurs since we want the outgoing boundary contribution in: $\lim_{t\to-\infty} \{A_f(t) - A_f(-t)\} = c.t.$ to appear on the left hand side of the local operators whose matrix elements we are reducing (then its contribution just produces outgoing contraction terms). The same statements apply verbatim to the reduction of outgoing states. The iterative application of this procedure therefore leads to the following reduction formula:

$$\int \dots \int \bar{f}_{n+1} \dots \bar{f}_{n+m} |A(x)| f_1 \dots f_n \rangle^{in}$$

$$= \int \dots \int \bar{f}_{n+1} (y_{n+1}) \dots \bar{f}_{n+m} (y_{n+m}) f_1(x_1) \dots f_n(y_n) \times$$

$$K_{n+1} \dots K_{n+m} K_1 \dots K_n \langle 0 | TA(x)A(y_1) \dots A(y_{n+m}) | 0 \rangle$$
(5.66)

Instead of A(x) we could have also started with any multilocal product of local fields. In the special case of $A \to \mathbf{1}$ we obtain the reduction formula for the S-matrix:

$$\int \frac{\partial u}{\partial t} \langle f_{n+1}....f_{n+m} | f_1....f_n \rangle^{in}$$

$$\int \int \bar{f}_{n+1}(y_{n+1})...\bar{f}_{n+m}(y_{n+m})f_1(x_1)...f_n(y_n) \times$$

$$K_{n+1}...K_{n+m}K_1...K_n \langle 0 | TA(y_1)....A(y_{n+m}) | 0 \rangle + c.t.$$

$$(5.67)$$

By going to the limit of plane waves one obtains for the connected part of the

momentum space kernel of the S-matrix:

$$S(p_{n+1}..p_{n+m}; p_1..p_n)^{conn.} = \lim_{p_i^2 \to m^2} \prod_i (p_i^2 - m^2) \times (5.68)$$

$$\tau(-p_{n+1}..-p_{n+m}, p_1..p_n)$$

i.e. we obtain the residua on mass shell of the Fourier transforms of the time ordered function τ . These reduction formulas are very suggestive of the so called crossing symmetry:

incoming particle
$$p \rightarrow$$
 outgoing antiparticle $-p$ (5.69)

for the generalized formfactores of local fields:

$$= {}^{out} \langle p_1 ... p_n | \mathcal{O}(0) | p_{n+1} ... p_{n+m} \rangle^{in} \\ = {}^{out} \langle p_1 ... p_n, -p_{n+1} | \mathcal{O}(0) | p_{n+2} ... p_{n+m} \rangle^{in}$$

where $-p_{n+1}$ stands for the momentum of the antiparticle on the backward mass shell which by on-shell analytic continuation is related to the process with the antiparticle momentum on the forward mass shell. With other words such a "symmetry", in order to be physically meaningful, must be interpretable as a relation between different boundary values of an on-shell meromorphic "master" function. Although in renormalized perturbation theory this turned out to be true in each checked case, a proof of the necessary analyticity derived from the principles of QFT is only known in special cases i.e. the reduction formula is only suggestive but does not establish the crossing symmetry. A related question is the existence of time-ordered functions outside perturbation theory. According to the best of my knowledge, this has not been demonstrated in the general setting of QFT¹. A closer look at the derivation of the reduction formula reveals that a pointlike covariant time ordering is not needed; any asymptotic ordering will lead to the same on-shell values i.e. the residua on mass shell are independent on the precise ordering prescription for finite space time separations. In a later section we will see that time ordered fields is not a natural concept in nonperturbative QFT. The more natural objects turn out to be certain sesquilinear forms of the fields, the so called "generalized formfactors".

In the following we will derive the Haag-Ruelle scattering theory in the general setting of QFT and then comment in the derivation of the LSZ theory.

In *n*-particle Schrödinger theory, the physical input for the existence of scattering state vectors as large time limits of suitably chosen time dependent vectors is the strong fall-off property of the two body potential. Although one can somewhat relax those properties, potentials as the Coulomb potential fall-off too weakly in order to belong to the standard situation (the large time wave functions oscillate with a logarithmic factor which does not contribute to the probabilities). In QFT the corresponding property is the strong cluster property

 $^{^{1}}$ In view of the fact that in the bootstrap-form factor construction the time ordering plays no role, it would be unreasonable to postulate its existence.

of correlation functions in spacelike directions. A sufficient condition for this property is the existence of a spectral gap in the mass operator.

An operator from the polynomial \mathcal{P} algebra (see section 2 of this chapter):

$$Q = \sum_{n} \int f_n(x_1, \dots, x_n) A(x_n) \dots A(x_n) d^d x_1 \dots d^d x_n \qquad (5.70)$$
$$f_n \in \mathcal{S}^{4d}$$

will be called "almost local" (if $supp f_n \in O$, O is local). We will be interested in the behavior of correlation functions of $Q(x) := U(x)QU(x)^{-1}$. The relevant theorem is

Theorem 20 (Ruelle, 1962, [3]) In a local QFT with a spectral mass gap (isolated one-particle mass shells) the quasilocal operators fulfill the strong cluster property:

$$\forall N \in \mathbf{N}, \exists C \ s.t. \ \langle Q_1(x_1)...,Q_n(x_n) \rangle_{con} < C_N R^{-N}$$
(5.71)

Here R denotes the maximal space like distance:

$$R = \max_{i,k} -(x_i - x_k)^2 \tag{5.72}$$

We will not prove the theorem, but rather try to understand how it can be used in order to understand the convergence for large times and the structure of the incoming and outgoing multi-particle states. We first pick $Q_i s$ which applied to the vacuum create one-particle states with given wave function $\tilde{\varphi}(p)$. By choice of $f_n \in S^{nd}$ with appropriate energy-momentum support this is always possible. Then we form the operators:

$$Q_i(h_i;t) := i \int_{x_0=t} Q_i(x) \overleftrightarrow{\partial_{x_0}} h_i(x) d^3x$$
(5.73)

where h_i is a positive energy solution of the Klein-Gordon equation and the derivative act with a minus sign to the left. Clearly:

$$Q_i(h_i;t)\Omega = |\psi_i\rangle, \quad \tilde{\psi}_i(p) = \tilde{\varphi}(p)\tilde{h}_i(p)$$
(5.74)

i.e. one obtains time independent one-particle states. On the other hand the multiple application (at least two) of these operators leads to time dependent states whose large time behavior is controlled by the following theorem:

Theorem 21 (Haag 1958, [3])

(i) The sequence of state vectors

$$\Psi(t) = \prod_{i}^{n} Q_i(h_i; t) \Omega$$
(5.75)

converge strongly for $t \to \pm \infty$. The limiting states have the physical interpretation of incoming and outgoing multiparticle scattering states:

$$\Psi^{in} = \lim_{t \to -\infty} \Psi(t) = |\psi_1, \dots, \psi_n\rangle^{in}$$

$$\Psi^{out} = \lim_{t \to +\infty} \Psi(t) = |\psi_1, \dots, \psi_n\rangle^{out}$$
(5.76)

(ii) The scalar product of these scattering states has the Fock space structure:

$$\sum_{out}^{in} \left\langle \psi_1' \dots \psi_n' \mid \psi_1 \dots \psi_m \right\rangle_{out}^{in} = \delta_{nm} \sum_{P \in S_n} \left\{ \begin{array}{c} + \\ sign(P) \end{array} \right\} \prod_k \left\langle \psi_{P(k)}' \mid \psi_k \right\rangle$$
(5.77)

according to bosonic or fermionic spacelike behavior of the Heisenberg fields A(x). One should add that the Poincaré transformation act naturally on the asymptotic Fock space structure, i.e. the in and out states do not remember the special frame in which the time direction was defined.

The idea of the proof consists in showing that $\Psi(t)$ is a Cauchy sequence i.e. that $\left\|\frac{d}{dt}\Psi(t)\right\| < Ct^{-\frac{3}{2}}$. What one needs in addition to the cluster properties of the Q(x)-correlations is a refined asymptotic estimate on the single particle wave functions which goes beyond the result of the well-known stationary phase method:

$$h(\mathbf{x},t) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int \tilde{h}(\mathbf{p}) e^{-i(\omega(\mathbf{p})t-\mathbf{p}\mathbf{x})} \underset{t \to \infty}{\to}$$
(5.78)
$$= const.t^{-\frac{3}{2}} \exp(-im\gamma^{-1}t)(\gamma^{\frac{3}{2}}\tilde{h}(m\gamma\mathbf{v}) + O(t^{-1}))$$

$$\gamma = \frac{1}{\sqrt{1-\mathbf{v}^{2}}}, \quad \mathbf{v} = \frac{\mathbf{x}}{t}$$

The refined version determines the "essential" x-space support of h in terms of the velocity support in momentum space $\Sigma = \left\{ \mathbf{v} = \frac{\mathbf{p}}{\omega} \mid p \in supp\tilde{h} \right\}$. one has:

Theorem 22 (Ruelle 62) Let h be a positive energy solution of the Klein-Gordon equation and Σ its velocity support. With \mathcal{U} an open set containing Σ we have:

we have: (i) for $v \in \mathcal{U} : |h(\mathbf{v}t, t)| < C |t|^{-\frac{3}{2}}$ as 5.78 (ii) for $v \notin U : |h(\mathbf{v}t, t)| < C_N (1 + |\mathbf{v}|)^{-N} |t|^{-N}$

If we now choose one-particle wave functions h' with nonoverlapping velocity supports relative to the unprimed h then

$$\left\langle \Omega \left| Q_1(h'_1;t)^* \dots Q_m(h'_m;t)^* Q(h_n;t) \dots Q(h_1;t) \right| \right\rangle$$

$$\xrightarrow{}_{t \to \infty} \delta_{nm} \left\{ \sum_{P \in S_n} \left\{ \begin{array}{c} + \\ sign(P) \end{array} \right\} \prod_{k=1}^n \left\langle \psi'_{P(k)} \mid \psi_k \right\rangle \right\}, \quad \begin{array}{c} bosonic \\ fermionic \end{array}$$

The connected part, upon integration with the dissipating wave packets, does not contribute at all to the limit, as follows from the elementary geometrical

(essential) support pictures in Minkowski space.. The same holds for any connected cluster with more than two operators Q_i . this fixes the structure of the in/out scalar products. The fall-off of $\left\|\frac{d}{dt}\Psi(t)\right\|^2$ is even simpler to understand, because each term which contributes to this norm square for large t contains one two-point function where one operator is a time derivative of Q(h; t) which vanishes upon acting on the vacuum.

The restriction to nonoverlapping wave packets has a physical origin: parallel flying particles lead to a weaker convergence. The best strategy is to prove formulas for the nonoverlapping situation and only at the end take the plane wave limit. The formalism does not only allow to derive the LSZ theory and the reduction formulas, but also gives higher order t corrections to LSZ ([3], chapter II section 4).

The above scattering formalism needs to be modified in an essential and interesting way if the fields have a spacelike commutation structure which leads to braid group statistics. In the physically interesting case of d=2+1 dimensions, these "plektonic" fields have really a string-like spatial extension i.e. they are not fields in the sense of Lagrangian QFT. Their construction falls into the realm of general or algebraic QFT. One still can prove the asymptotic convergence, but the asymptotic state vectors loose their tensor product structure and the cut between kinematics (in/out structure) and dynamics (genuine interactions) has to be essentially modified. The fact that such theories are outside the Lagrangian framework and even outside quantization ideas, does not make them any less physical or susceptible to explicit and perturbative constructions, but the perturbation around free "plektons" is expected to have more in common with ideas on perturbing around chiral conformal theories than with Feynman perturbation theory around bosonic/fermionic free fields. the structure of d = 2 + 1 plektons will be investigated in the last chapter.

Literature to chapter 5:.

R.F. Streater and A.S. Wightman, "PCT, Spin and Statistics and All That" New York, Benjamin 1964

R.Jost "The General Theory of Quantized Fields", American Math. Soc. 1965

N.N.Bogoliubov, A.A.Logunov, A.I.Oksak and I.T.Todorov "General Principles of Quantum Field Theory" Kluwer 1990

J. Glimm and A. Jaffe, "Quantum physics. A functional integral point of view". Springer 1987

R. Haag "Local Quantum Physics, Fields, Particles, Algebras" Springer 1992

Chapter 6

Nonperturbative Constructions

6.1 Introductory Comments

Presently QFT appears as being formed of several parts which seem to drift apart into different directions. On the one hand there is the standard approach presented in the previous chapters which is centered around renormalized perturbation theory and the various quantization methods (canonical, functional, the causal Bogoliubov-Shirkov approach). Enriched with geometrical ideas the standard formulation based on Lagrangians and actions has prepared the ground for many mathematical advances including the duality structure of Seiberg-Witten as well as to string theory. On the other hand there is the more algebra-based low dimensional approach which has led to the construction of rich families of chiral conformal and factorizing QFT without Lagrangians. This constructive approach, although being somewhat conservative in its use of physical principles, has nevertheless given many startling nonperturbative results of particle physics concerning e.g. fusion charges, in particular of antiparticles from particles, confined objects and solitons as being two opposite sides of the same subject, and other more general (and somewhat surprising) manifestations of the principle of "nuclear democracy". The third approach to QFT which forms the backbone of these notes, was carried out by a rather small group of theoretical physicist with a strong mathematical background on operator algebras without restrictions of space-time dimensionality and to integrability, with the aim to arrive at a general constructive nonperturbative approach. This will be of main concern in this chapter.

The most interesting message of the low dimensional constructive bootstrapformfactor program is that the emphasis on the scattering matrix advocated way back by Heisenberg and Wheeler and later by Chew, Stapp and others, was in a certain sense physically well founded. What went wrong in those early "bootstrap" attempts is mostly related to the enforced and artificial separation of S from local QFT and the (unfortunately cyclically recurrent) working hypothesis of a "TOE", a theory of everything (in this case of everything minus gravity). Looking at the old articles it is hard to understand the fervor with which the S-matrix concepts were cleansed from all field theoretic "impurities".

The main theme of this chapter is the realization that the S-matrix in algebraic QFT acquires a new, hitherto unknown pivotal role in the construction of local nets (whose generators are local fields). In this way it is regaining some of its importance before the gauge theory began to shift the emphasis from on-shell to off-shell properties. It turns out that the S-matrix belongs to the foundation of the local field theory (in its role as the net invariant which carries local modular information) as well as to its roof (in its role as describing scattering observables); a truly vexing "bootstrap" situation. The fact that in d=1+1 factorizable theories Chew's bootstrap ideas for the S-matrix work without mentioning fields (but with the help of "fusion" and "Yang-Baxter") is not due to the correctness of the underlying philosophy but rather to undeserved luck: the physical rapidity (scattering) variable is at the same time the uniformization parameter of the analytic properties¹. In higher dimensions or without the factorization, Chew's program would fail without the use of local fields (and it did fail). In that case one could hope hat an iterative procedure which corrects the trial input S-matrix together with a locality improvement of states and fields, may have a constructive chance, a situation which could be more vaguely reminiscent of the Hartree-Fock iteration in Schrödinger theory than of renormalized perturbation theory.

In this section we will apply the modular localization introduced in chapter3 to interacting theories [38]. In this way we will reconquer the lost unity in QFT. In particular, we will learn a new and very interesting lessons from the d=1+1 formfactor program. Far from being a special "exotic" construction, remote from any "real" QFT, this approach, if analyzed with general and deep concepts related to the TCP theorem and the S-matrix (interpreted as an invariant of a local net), reveals a surprising new and powerful nonperturbative construction principle which, so we hope, may turn out to be the basis of a future new iterative constructive approach in d=1+3 theories.

Locality of observables and localization of states (always relative to the vacuum or some other distinguished reference state) in QFT is a conditio sine qua non for its physical interpretation. Global topology (as in the combinatorial or so called "topological field" theories, or in the vacuum degeneracy structure beyond spontaneous symmetry), remains part of mathematics, as long as its connection with the local structure remains unclear.

The fastest way to get a glimpse at the "modular localization" [38] is to recall that concept in connection with the Wigner representation theory for positive energy representations of the Poincaré group and free fields, as it was explained in chapter 3. There we learned that there are infinitely many free fields in Fockspace and they constitute the linear part (in creation and annihilation operators)

 $^{^1\}mathrm{Even}$ in d=1+1 the situation is very far removed from the desired uniqueness of Chew's S-matrix approach.

6.1. INTRODUCTORY COMMENTS

of a huge local equivalence class of fields, the so called Borchers class $\mathcal{B}(m, s)$ [3] [18] Any cyclic (with respect to the vacuum) representative field from this class generates the same net of local von Neumann algebras in H_F :

$$\mathcal{O} \to \mathcal{A}(\mathcal{O}) \tag{6.1}$$

In fact the emerging picture of pointlike fields that behave similar to coordinates in differential geometry, was the prime historical motivation for formulating algebraic QFT in terms of nets of algebras [3]. For a detailed recent presentation of the physical motivations and aims of this net approach in algebraic QFT we refer to [5].

In the following it is important to understand the *direct* construction of this net in terms of the "modular localization" principle. Let us briefly review what we learned in chapter 3. We use the d=3+1 Wigner (m,s)-representations as an illustrative example. In case of charged particles (particles≠antiparticles) we double the Wigner representation space:

$$H = H^p_{Wig} \oplus H^p_{Wig} \tag{6.2}$$

in order to incorporate the charge conjugation operation as an (antilinear in the Wigner theory) operator involving the p- \bar{p} -flip. On this extended Wigner space one can act with the full Poincaré group (where those reflections which change the direction of time are antiunitarily represented). For the modular localization in a wedge we only need the standard L-boost $\Lambda(\chi)$ and the standard reflection r which (by definition) are associated with the t-x wedge:

$$\delta^{i\tau} \equiv \pi_{Wig}(\Lambda(\chi = 2\pi\tau)) \tag{6.3}$$

$$j \equiv \pi_{Wig}(r) \tag{6.4}$$

These operators have a simple action on the p-space (possibly) doubled Wigner wave functions, in particular:

$$(j\psi)(p) \simeq \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \bar{\psi}(p_0, p_1, -p_2, -p_3)$$
 (6.5)

By functional calculus we form $\delta^{\frac{1}{2}}$ and define:

$$s \equiv j\delta^{\frac{1}{2}} \tag{6.6}$$

This unbounded antilinear densely defined operator s is involutive on its domain: $s^2 = 1$. Its -1 eigenspace² is a real closed subspace H_R of H which allows the following characterization of the domain of s:

$$dom(s) = H_R + iH_R$$

$$s(h_1 + ih_2) = -h_1 + ih_2$$
(6.7)

²This is one of the few places where a sign mistake has no grave consequences since a multiplication by i transforms the + into the - real eigenspace.

Defining:

$$H_R(W) \equiv U(g)H_R, \ W = gW_{stand} \tag{6.8}$$

where g is an appropriate Poincaré transformation, we find the following theorem:

Theorem 23 $H_R(W)$ is a net of real Hilbert spaces i.e. $H_R(W_1) \subsetneq H_R(W_2)$ if $W_1 \subsetneq W_2$.

The proper containment is an easy consequence of a theorem by Borchers [7] which relates this property to the positivity of the energy (in fact the geometric property is equivalent to the energy positivity.

If we now define:

$$H_R(\mathcal{O}) \equiv \bigcap_{W \supset \mathcal{O}} H_R(W) \tag{6.9}$$

then it is easily seen (even without the use of the u,v-intertwiners) that the spaces $H_R(\mathcal{O}) + iH_R(\mathcal{O})$ are still dense in H_{Wig} and that the formula:

$$s(\mathcal{O})(h_1 + ih_2) \equiv -h_1 + ih_2$$
 (6.10)

defines a closed involutive operator with a polar decomposition:

$$s(\mathcal{O}) = j(\mathcal{O})\delta(\mathcal{O})^{\frac{1}{2}} \tag{6.11}$$

Although now j(O) and $\delta(\mathcal{O})^{i\tau}$ have no obvious geometric interpretation, there is still a bit of geometry left, as the following theorem shows:

Theorem 24 The $H_R(\mathcal{O})$ form an orthocomplemented net of closed real Hilbert spaces, i.e. the following "duality" holds:

$$H_R(\mathcal{O}') = H_R(\mathcal{O})' = iH_R^{\perp}(\mathcal{O}). \tag{6.12}$$

Here \mathcal{O}' denotes the causal complement, H_R^{\perp} the real orthogonal complement in the sense of the inner product $Re(\psi, \varphi)$ and H'_R is the symplectic complement in the sense of Im (ψ, φ) .

The direct construction of the interaction-free algebraic bosonic net for (m,s=integer) is now achieved by converting the "premodular" theory of real subspaces of the Wigner space into the Tomita-Takesaki modular theory for nets of von Neumann algebras using the Weyl functor:

The application of the Weyl functor - to the net of real spaces:

$$-: H_R(\mathcal{O}) \longrightarrow \mathcal{A}(\mathcal{O}) \equiv alg \{ W(f) | f \in H_R(\mathcal{O}) \}$$

$$(6.13)$$

leads to a net of von Neumann algebras in \mathcal{H}_{Fock} which are in "standard position" with respect to the vacuum state with a modular theory which, if restricted to the Fock vacuum Ω , is geometric:

$$\Gamma(s) = S, \ SA\Omega = A^*\Omega, \ A \in \mathcal{A}(W)$$

$$S = J\Delta^{\frac{1}{2}}, \ J = \Gamma(j), \ \Delta^{i\tau} = \Gamma(\delta^{i\tau})$$
(6.14)

The proof of this theorem uses the functorial formalism of section 2.6. It should be evident from the derivation that the wedge localization concept in Fock obtained in this functorial way from the Wigner theory only holds for interaction free situations. The Fock space is also important for interacting QFT, but in that case the wedge localization enters via scattering theory as in section 4, and not through Wigner's representation theory.

Clearly the W- or \mathcal{O} - indexing of the Hilbert spaces corresponds to a localization concept via modular theory. Specifically $H_R(\mathcal{O}) + iH_R(\mathcal{O})$ is a certain closure of the one particle component of the Reeh-Schlieder domain belonging to the localization region \mathcal{O} . Although for general localization region the modular operators are not geometric, there is one remaining geometric statement which presents itself in the form of an algebraic duality property [4]:

$$\mathcal{A}(\mathcal{O}') = \mathcal{A}(\mathcal{O})', \quad Haag \ Duality \tag{6.15}$$

Here the prime on the von Neumann algebra has the standard meaning of commutant. In the following we make some schematic additions and completions which highlight the modular localization concept for more general free cases [39].

In the case of s = halfinteger, the Wigner theory produces a mismatch between the "quantum" (in the sense of the commutant) and the "geometric" opposite of $H_R(W)$, which however is easily taken care of by an additional factor *i* (interchange of symplectic complement with real orthogonal complement). This requires, via the physical localization property, the application of the CARfunctor instead of the CCR-functor as well as the introduction of the well-known Klein transformation K which takes care of the above mismatch in Fock space:

$$J = K\mathcal{F}_{CAR}(ij)K^{-1}$$

$$\mathcal{A}(\mathcal{O}') = K\mathcal{A}(\mathcal{O})'K^{-1}$$
(6.16)

where the K is the twist operator of the "twisted" Haag Duality.

In section 3.8 we exposed the thermal properties of this modular localization by constructing via the KMS property a representation which is only defined in the thermal Hilbert space belonging to the wedge localization. We also commented on the relation between these thermal properties and the crossing symmetry of particle physics. This famous crossing symmetry, which is known to hold also in each perturbative order of renormalizable interacting theories, has never been derived in sufficient generality within any nonperturbative framework of QFT. It was thought of as a kind of on-shell momentum space substitute for Einstein causality and locality (and its strengthened form called Haag duality). As such it played an important role in finding a candidate for a nonperturbative S-matrix, an important contribution known under the name of the Veneziano dual model. Although it stood (in this indirect way) on the cradle of string theory, the recent string theoretic inventions of duality result from formal generalizations which appearantly have little do with the original physical concepts of nonperturbative relatiistic scattering theory.

If crossing symmetry is a general property of local QFT, a conjecture (proven in every order of perturbation theory) which nobody seems to doubt, then it should be the on-shell manifestation of the off-shell KMS property for modular wedge localization. In the construction of wedge localized thermal KMS states on the algebra of mass shell operators satisfying the Zamolodchikov-Faddeev algebraic relations³ on the momentum space rapidity axis [21], the derivation of crossing symmetry is similar (albeit more involved) to the previous free field derivation [2][39] and the argument can be found in section 4 of this chapter. Very recently arguments were proposed [40] which were bases on the idea that Haag-Ruelle scattering theory can be adapted to the wedge situation. I think that such ideas are untainable. In our approach it turns out that the general crossing symmetry (in any dimension) is indeed related to the wedge KMS condition but that this statement cannot be derived just from scattering theory alone. It rather follows from the existence and the modular intertwining property of the "modular Møller operator" U (next section) which, unlike the S-matrix, is not an object of scattering theory but is only defined in terms of modular wedge localization (for this reason we maintain the prefix "modular"). We intend to use this object in order to prove [31] the uniqueness of the main inverse problem of QFT: $S_{scat} \rightarrow QFT$.

In fact the previous free field formalism of the first two sections may be generalized into two directions (or into both):

- interacting fields
- curved space-time (without or with interacting quantum matter)

As mentioned before, low-dimensional interacting theories will be discussed in the next section. For the generalization to curved space time (e.g. the Schwarzschild black hole solution) it turns out that only the existence of a bifurcated horizon together with a certain behavior near that horizon ("surface gravitation") [9][10] is already sufficient in order to obtain the thermal Hawking aspect. In the standard treatment one needs isometries in space-time, i.e. Killing vectors. In chapter3 we have already presented the thermal aspects with the wedge situation. Here we address a more general situation. The idea of modular localization suggests to consider also e.g. double cones for which there is no space-time isometry but only an isometry in H_{Wigner} or \mathcal{H}_{Fock} . Of course such enlargements of spaces for obtaining a better formulation (or even a solution) of a problem are a commonplace in modern mathematics, particularly in noncommutative geometry. The idea here is that one trades the ill-defined Killing isometries by a geometrically "fuzzy" but well-defined symmetry transformations in quantum space, which only near the horizon loose their space-time fuzziness. The candidates for these nongeometric symmetries are the modular automorphisms of von Neumann algebras of arbitrary space time regions together with suitable faithful states from the local folium of admissable states.

 $^{^{3}}$ As will become clear in the next section, although these operators are nonlocal, they generate the wedge localized states and as a consequence the modular KMS formalism is applicable to them.

Although the restriction of the global vacuum state is always in that folium, it is often not the most convenient for the construction of the modular automorphism.

In this context one obtains a good illustration by the (nongeometric) modular theory of e.g. the double cone algebra of a massive free field. From the folium of states one may want to select that vector, with respect to which the algebra has a least fuzzy (most geometric) behavior under the action of the modular group. Appealing to the net subtended by spheres at time t=0 one realizes that algebras localized in these spheres are independent of the mass. Since m=0 leads to a geometric modular situation⁴ for the pair $(\mathcal{A}_{m=0}(S), |0\rangle_{m=0})$, and since the nonlocality of the massive theory in the subtended double cones is only the result of the fuzzy propagation inside the light cone (the breakdown of Huygens principle or the "reverberation" phenomenon), the fuzziness of the modular group for the pair $(\mathcal{A}_{m\neq 0}(C(S)), \Omega_{m\neq 0})$ is a pure propagation phenomenon i.e. can be understood in terms of the deviation from Huygens principle. In view of the recent micro-local spectrum condition, one expects such nonlocal cases to have modular groups whose generators are pseudo-differential instead of (local) differential operators [35]. In fact for free massive theories in d=1+1one can give a rigorous proof together with an explicit formula for the modular automorphism. In order to avoid the pathology of the d=1+1 scalar zero mass field, we start with a massive free spinor field whose massless limit gives a twocomponent field with the first component only depending on the left light cone and the second on the right hand light cone. The same zero mass theory results from Sewell's restriction to the light cone horizon (boundary) of the double cone. But in the latter case we know the modular group for the massless double cone algebra together with the massless vacuum vector. It is a one-parametric subgroup of the conformal group [3]. The massless theory on the horizon is then propagated inside with the massive causal propagator and this last step is responsible for the delocalization inside the double cone. This and similar subjects will be the content of a separate paper together with Wiesbrock. [20].

The Hilbert space setting of modular localization offers also a deeper physical understanding of the universal field domain \mathcal{D} which plays a rather technical role in the Wightman framework [18] In the modular localization approach the necessity for such a domain appears if one wants to come from the net of localization spaces which receive their natural topology from the (graphs) net of Tomita operators $\bar{S}(\mathcal{O})$ to a net of (unbounded) polynomial algebras $\mathcal{P}(\mathcal{O})$ such that:

$$dom \ S(\mathcal{O}) \cap \mathcal{D} = \mathcal{P}(\mathcal{O})\Omega = dom \ \mathcal{P}(\mathcal{O}) \tag{6.17}$$

this domain is of course also expected to be equal to $\mathcal{A}(\mathcal{O})\Omega$. Here we used a more precise notation which distinguishes between the operator S defined on the core $\mathcal{A}(0)\Omega$ and its closure \bar{S} which is defined on $\mathcal{H}_R(\mathcal{O}) + i\mathcal{H}_R(\mathcal{O})$. One may round off these new interpretations of old domain problems by Fredenhagen's speculative remark about the modular role of pointlike fields. This idea is based on modular observations in the algebraic approach to chiral conformal

⁴The modular group is a one-parametric subgroup of the conformal group.

field theory [101]. There it is possible to extract the pointlike covariant fields without additional technical assumptions directly from the net. This, together with the known modular structure of the algebras in the net gives a beautiful characterization of these fields: the "one field states" obtained by applying a field to the vacuum and smearing with test functions form a representation space for the universal modular group. The latter is defined as the group generated by all modular groups belonging to arbitrary double cones.

6.2 Modular Localization and Interaction

In order to obtain a clue of how to incorporate an intrinsic notion of interactions into this modular localization setting, we remind ourselves that if we do use pointlike fields, the modular localization for free fields agrees with what we get by applying the polynomial in the localization region supported smeared fields. In contrast to the conventional characterization of localization in terms of xspace pointlike fields, the modular characterization works in the *momentum*-(*Fock*)space of the (incoming) free particles. It attributes a physical significance to the precise position of the Reeh-Schlieder [3] dense set of localization region, i.e. it primarily deals with subspaces and subalgebras and only in second place with individual vectors and operators.

In order to formulate the modular localization principle in the case of interactions, one must take note of the fact that the scattering matrix S of local QFT is the product of the interacting TCP-operator Θ with the free (incoming) TCP operator Θ_0 and (since the rotation by which the Tomita reflection J differs from Θ is interaction-independent as all connected Poincaré transformations are interaction-independent) we have:

$$S = \Theta \cdot \Theta_0, \quad S = J \cdot J_0 \tag{6.18}$$

and as a result we obtain for the Tomita involution \check{S} :

$$\check{S} = J\Delta^{\frac{1}{2}} = SJ_0\Delta^{\frac{1}{2}} = S\check{S}_0 \tag{6.19}$$

Again we may use covariance in order to obtain $\hat{S}(W)$ and the localization domain of $\check{S}(W)$ as $\mathcal{D}(\check{S}(W)) = \mathcal{H}_R(W) + i\mathcal{H}_R(W)$ i.e. in terms of a net of closed real subspaces $H_R(W)$ of the incoming Fock space. However now the construction of an associated von Neumann algebra is not clear since an "interacting" functor from subspaces of the Fock space to von Neumann algebras is not known. In fact whereas the existence of a functor from the net of localized *Wigner subspaces* H_R^{Wig} to a net of von Neumann algebras is equivalent to the equality:

$$H_R^{Wig}(W_1 \cap W_2) = H_R^{Wig}(W_1) \cap H_R^{Wig}(W_2)$$
(6.20)

The equality becomes an inequality \subset for the above localized subspaces of Fock space \mathcal{H}_R . It also turns into an inequality for Wigner subspaces as soon as the Wigner spin becomes anyonic or Wigner's continuous spin [91] as in section
3.7. We will make some remarks (still short of a solution of this important problem) in the concluding section of this chapter and continue here with some more helpful comments on modular localization of interacting state vectors.

As in the free case, the modular wedge localization does not use the full Einstein causality but only the so-called "weak locality", which is just a reformulation of the TCP invariance [18] Weakly local fields form an equivalence class which is much bigger than the local Borchers class but they are still associated to the same S-matrix (or rather the same TCP operator). Actually the S in local quantum physics has two different interpretations: S in its role to provide modular localization in interacting theories, and S with the standard scattering interpretation in terms of (nonlocal!) large time limits. There is no parallel outside local quantum physics to this state of affairs. Whereas all concepts and properties which have been used hitherto in standard QFT (perturbation theory, canonical formalism and path integrals) as e.g. time ordering ⁵ and interaction picture formalism, are shared by nonrelativistic theories, modular localization is a new structural element in local quantum physics ⁶ and its characteristic property. No physically viable alternative (i.e. physically interpretable) to Einstein causality has ever been found in the long history of QFT.

The simplest kind of interacting theories are those in which the particle number is at least asymptotically ("on-shell") conserved i.e. $[\check{S}, \mathbf{N}_{in}] = 0$.

In the next section we will briefly review the d=1+1 bootstrap-form factor program in a manner which facilitates the later application of modular localization.

6.3 The Bootstrap-Formfactor Program

In this section we will meet a constructive approach for "integrable" d=1+1 QFT. Our first task is to obtain an intrinsic QFT understanding of integrability in a way which avoids classically inherited notions such as a complete sets of conservation laws etc. For this purpose we note an important d=1+1 peculiarity. Our generic expectation is that large spatial separation of the center of wave packet of two particles in the elastic two-particle scattering matrix leads to the weakening of scattering, or in momentum space:

$$\left\langle p_{1}^{'}p_{2}^{'}|S|p_{1}p_{2}\right\rangle = \left\langle p_{1}^{'}p_{2}^{'}|p_{1}p_{2}\right\rangle + \delta(p_{1}+p_{2}-p_{1}^{'}-p_{2}^{'})T(p_{1}p_{2}p_{1}^{'}p_{2}^{'})$$
 (6.21)

where the identity contribution is more singular (has more δ -factors) than the interaction *T*-term and therefore the second term drops out in x-space clustering. This argument fails precisely in d=1+1 and therefore the cluster property of the S-matrix is not suitable in order to obtain an intrinsic understanding of interaction. In the large distance clustering process, the two-particle S-matrix

⁵There is a conspicuous absence of the time-ordering operation in the bootstrap construction of factorizable field models. Instead the basic objects are generalized formfactors i.e. sesquilinear forms on a dense set of state vectors.

 $^{^{6}}$ This characteristic modular structure lifts local quantum physics to a new realm by its own which cannot be obtained by specialization from general quantum theory.

looses its higher particle threshold structure, but it remains nontrivial (in distinction to d=3+1). However for all higher particle scattering processes the behavior for d=1+1 is qualitatively the same as in higher dimensions: the decreasing threshold singularities (which decrease with increasing particle number) are responsible for the spatial decrease. Therefore any d=1+1 QFT is expected to have a limiting S_{lim}-matrix which is purely elastic and solely determined by the elastic two-particle $S^{(2)}$ -matrix. The Yang-Baxter relation results as a consistency relation for the elastic $3\rightarrow 3$ particle $S_{\text{lim}}^{(3)}$ -matrix. If this limiting S-matrix would again correspond to a localizable QFT, we would have a new class division of QFT, this time based on a long distance limit (which in some sense is opposite to the scale invariant short distance limit). It is this (long distance) class property 7 which makes these factorizing models so fascinating, as much as the fascination of chiral conformal QFT results from their role of representing short distance universality classes. In d=3+1 $S_{\text{lim}} = 1$ and therefore the limiting theory is expected to maintain the same superselection rules but in the "interaction-freest" possible way (literally free theories as we will argue later on). Hence in d=1+1 we are invited to speculate on the validity of the following commutative diagram:

$$\mathcal{F} \searrow \begin{array}{c} \mathcal{F}_{ld} \\ \downarrow \\ \mathcal{F}_{sd} \end{array} \tag{6.22}$$

Here ld(sd) labels the long (short) distance limits. There are also arguments [21] that with the help of a perturbative idea one may ascend from \mathcal{F}_{sd} to \mathcal{F}_{ld} . It is however presently not clear how one can use the known properties of the ld theories (i.e. the integrable models) in order to formulate a constructive program for the nonintegrable members of the ld equivalence class. We hope that our modular localization principle (which is not restricted to factorizable models) may turn out to be helpful for this purpose.

The constructive approach based on the bootstrap idea proceeds in two steps. One first classifies unitary and crossing symmetric solutions of the Yang-Baxter equations which fulfill certain minimal (or maximal, depending on the viewpoint) requirements. Afterwards we use these factorizing S-matrices together with the Watson equations (a notion from scattering theory relating formfactors with the S-matrix) and analytic properties for formfactors in order to compute the latter. One obtains the complete set of multi-particle matrix elements of "would be" local fields, i.e. one constructs the fields as sesquilinear forms. It is characteristic of this method, that one does not use the "axiomatic" properties of the beginning of this section, but rather less rigorously known momentum-space analytic properties which, although certainly related to causality and spectral properties, are more part of the LSZ+dispersion theoretic induced folklore (example: crossing symmetry) than of rigorous QFT. As long as one demonstrates at the end that the so obtained fields fulfill local commutativity, this is a legitimate procedure[30][27]. It leaves open the question of whether there exists a

⁷Although I know of no article in which this has been spelled out, its pervasive presence behind the scene is is recognizable in some publications.

more direct conceptual link between the S-matrix and the local fields or rather the field independent local nets. That this is indeed the case will be shown after the presentation of the formfactor program.

6.3.1 Properties of Factorizing S-Matrices

Consider first the analytic structure of an elastic S-matrix for a scalar neutral particle. In terms of the rapidity variable θ :

$$|p_{1},p_{2}\rangle^{out} = S |p_{1},p_{2}\rangle^{in} = S_{el}(p_{1},p_{2}) |p_{1},p_{2}\rangle^{in}$$
 (6.23)

$$S_{el}(p_1, p_2) = : S(\theta), \quad p_i = m(\cosh\theta_i, \sinh\theta_i), \quad \theta := |\theta_1 - (\theta_2 24)|$$

$${}^{in} \langle p'_1, p'_2 | S | p_1, p_2 \rangle^{in} = S(\theta)^{in} \langle p'_1, p'_2 | p_1, p_2 \rangle^{in}$$

Usually the elastic S-matrix is written in terms of the invariant energy $s = (p_1 + p_2)^2 = 2m^2(1 + cosh\theta)$ and the momentum transfer (not independent in d=1+1) $t = (p_1 - p_2)^2 = .2m^2(1 - cosh\theta)$. As a result of undeserved fortune, the rapidity θ turns out to be a uniformization variable for the real analytic S i.e. the complex s-plane with the elastic cut in $s \ge 4m^2$ is dumped into the strip $0 \le Im\theta \le \pi$ and the S-matrix becomes a meromorphic function $S(\theta)$ with $S(-\theta) = S^*(\theta) = S^{-1}(\theta)$. (unitarity). Hence the strip $-\pi \le \theta \le \pi$ is the physical strip for $S(\theta)$. Crossing symmetry in our special (neutral) case means a symmetry on the boundary of the strip: $\theta \to i\pi - \theta$. Note that the presence of inelastic thresholds would destroy the uniformization.

The factorization implies the operator relation:

$$S_{12}(p_1, p_2)S_{13}(p_1, p_3)S_{23}(p_2, p_3)$$

$$= S_{23}(p_2, p_3)S_{13}(p_1, p_3)S_{12}(p_1, p_2)$$
(6.25)

According to Liouville's theorem, the only minimal solution (minimal number of poles, smallest increase at ∞) for this scalar diagonal case is $S = \pm 1$. More general solutions are obtained by placing bound-state poles into the minimal solution. In order to maintain unitarity, the pole factor must be of the form:

$$P(\theta) = \frac{\sinh\theta + i\sin\lambda}{\sinh\theta - i\sin\lambda} \tag{6.26}$$

Transforming back this pole at $\theta = i\lambda$ into the original individual particle variables, we obtain the following parametrization in terms of a center of mass and relative rapidity:

$$p_{1} = m\left(\cosh(\chi + \frac{i\lambda}{2}), \sinh(\chi + \frac{i\lambda}{2})\right)$$

$$p_{2} = m\left(\cosh(\chi - \frac{i\lambda}{2}), \sinh(\chi - \frac{i\lambda}{2})\right)$$
(6.27)

Clearly the two-particle bound state has the momentum:

$$p_{1,2} = (p_1 + p_2)_{at \, bd.state} = 2mcos \frac{\lambda}{2} (cosh\chi, sinh\chi)$$

$$p_{1,2}^2 = m_2^2, \quad m_2 = \frac{m}{2sin\frac{\lambda}{2}}sin\lambda$$
(6.28)

The "fusion" of particles may be extended. For a 3-particle bound state we would look at the 3-particle S-matrix which, as a result of factorization has the form:

$$S^{(3)}(p_1, p_2, p_3) = S(\theta_{12})S(\theta_{13})S(\theta_{23})$$
(6.29)

We first fuse 1 with 2 and simultaneously 2 with 3 as before. The center of mass + relative rapidity parametrization yields:

$$p_{1} = m \left(\cosh(\chi + i\lambda), \sinh(\chi + i\lambda) \right)$$

$$p_{2} = m \left(\cosh\chi, \sinh\chi \right)$$

$$p_{3} = m \left(\cosh(\chi - i\lambda), \sinh(\chi - i\lambda) \right)$$
(6.30)

Again we get the mass of the 3-particle bound state by adding the zero components in the $\chi = 0$ frame:

$$m_3 = (p_1 + p_2 + p_3)_0 = m_2 \cos\frac{\lambda}{2} + m\cos\lambda = 2\frac{m}{2\sin\frac{\lambda}{2}}\sin\frac{3\lambda}{2}$$
(6.31)

Induction then gives the general fusion mass formula:

$$m_n = 2\mu sin\frac{n\lambda}{2}, \quad \mu = \frac{m}{2sin\frac{\lambda}{2}}$$
 (6.32)

We will meet such trigonometric fusion formulas later in algebraic QFT where they are related to the statistical dimensions of fused charge sectors. They became first known through the Dashen-Hasslacher-Neveu quasiclassical approach. The above fusion calculation was done as far back as 1976 [22] and consisted in a synthesis of the quasiclassical work of DHN with some suggetive ideas of Sushko using the factorization principle, but still without the ideas of Yang and Baxter (which are not needed for this scalar case). The decisive step towards a general factorizable bootstrap program was taken two years later [23][24].

The consistency of these particles as incoming and outgoing objects leads to additional structures. Consider the scattering of the mass m_2 bound state with a third m-particle. This S-matrix for the scattering of these two different particles is obtained from $S^{(3)}$ by:

$$S_{b.e.}(p_1 + p_2, p_3) \mid_{(p_1 + p_2)^2 = m_2^2} = \frac{1}{R} \frac{Res}{(p_1 + p_2)^2 \to m_2^2} S_{12} S_{13} S_{23}$$
(6.33)

where the projector P_{12} together with a numerical residue value R is defined by:

$$\underset{(p_1+p_2)^2 \to m_2^2}{Res} S(p_1, p_2) = RP_{12}$$
(6.34)

and we used the subscript elementary e and bound b. as labels on the new two-particle scattering operator: $S_{b.c.}$. The factorization insures that:

$$P_{12}S_{13}S_{23} = S_{23}S_{13}P_{12} \tag{6.35}$$

A prominent family of scalar S-matrices with N-1 bound state fulfilling all these requirements are the Z_N models [25]. Consistency requires that the bound state of N-1 mass *m*-particles is again a mass *m*-particle (the antiparticle). For N=2 this family contains the Ising field theory with $S_{Ising}^{(2)} = -1$ which we already met in the section on (dis)order variables.

Instead of elaborating this scalar factorization situation, we pass immediatly to the matrix case where we meet a new and interesting phenomenon. We assume that the particle from which we start has an internal "charge" which can take on a finite number of values i.e.

$$|p,\alpha\rangle \in H_1 \otimes V, \quad dimV < \infty$$
 (6.36)

The two-particle S-matrix is then written as a matrix acting on $V \otimes V$ whose entries are operator-valued (represented as in the previous case by momentumspace kernels):

$$S|p_{1},...,p_{n}\rangle_{\alpha_{1}...\alpha_{n}}^{in} = |p_{1},...,p_{n}\rangle_{\alpha'_{1}...\alpha'_{n}}^{in} S_{\alpha_{1}...\alpha'_{n}}^{\alpha'_{1}...\alpha'_{n}}(p_{1},...,p_{n})$$
(6.37)

$$S^{(n)}(p_1, \dots, p_n) = \prod_{i < j} S^{(2)}(p_i, p_j)$$
(6.38)

The factorization requires a specific order of the product of matrices. Consistency demands the validity of the Yang-Baxter relation:

$$S_{12}(p_1, p_2)S_{13}(p_1, p_3)S_{23}(p_2, p_3)$$

$$= S_{23}(p_2, p_3)S_{13}(p_1, p_3)S_{12}(p_1, p_2)$$
(6.39)

a relation which resembles the Artin (braid-group) relation, but upon closer inspection reveals conceptual and mathematical differences to the latter. The notation should be obvious: the subscript on S indicates on which of the tensor factors in the 3-fold tensor product of one-particle spaces the object acts. The relation with the Artin relations would be clear, if one could ignore the p-dependence and rewrites the Y-B relation in terms of $\tilde{S}_{\cdot} = PS$, where P is the permutation of two tensor factors.

We arrived via our S_{lim} arguments at the famous Yang-Baxter relation, since as a result of the (independent) discovery by Yang and Baxter it was clear, that such a mathematical structure had appeared before outside QFT in a quite different setting of statistical mechanics models and nonrelativistic scattering theory of δ -function interactions. In our context this identity permits to change the temporal order of individual re-scatterings so that the n-particle scattering $S^{(n)}$ is independent of those (graphically: invariance under parallel shifts of 2-momenta in graphical illustrations of scattering processes). The problem of finding the natural parametrization (e.g. Baxter's elliptic parametrization) for these Yang-Baxter relations does not arise in QFT; the uniformizing physical rapidity θ is already the natural Yang-Baxter variable:

$$S_{12}(\theta)S_{13}(\theta + \theta')S_{23}(\theta') = S_{23}(\theta')S_{13}(\theta + \theta')S_{12}(\theta)$$
(6.40)

If fermion-antifermion pairs can go into boson-antiboson pairs, the object which fulfills the Yang-Baxter relation is not S but σS where $\sigma = \pm 1$ with + for bosons.

As the braid group relation, this is an overdetermined system of equations. For the former one found a powerful mathematical framework within V. Jones subfactor theory [26]. Although the attempts to get an equally powerful mathematical framework for the latter was less than successful (the "Baxterization" of the subfactor representations of Artin braids), one was able to find many interesting families of nontrivial solutions of which some even allowed a comparison with Lagrangian perturbation theory.

The S-matrix bootstrap idea originated in the early 60^{ies} from dispersion theory. Its revival in connection with d=1+1 factorization in the late 70^{ies} showed that its premises were physically reasonable, except the idea that it could be seen as a "theory of everything" (TOE) which was wrong and even absurd (for the more recent TOE's one would be hard pressed to say friendly words about their physical content).

The basic new message [27][28] is that one should use these factorizing Smatrices as computational tools for the construction of local fields and local nets as explained in the following subsection

6.3.2 Generalized Formfactors

Now we will probe the idea that these S-matrices belong to localizable fields. Let A be any local field which belongs to a Borchers equivalence class of local fields. We write the generalized formfactor of A(x) as:

$$_{\alpha_{1}...\alpha_{m}}^{out} \langle p_{1},...,p_{m} | A(0) | p_{m+1},...,p_{n} \rangle_{\alpha_{m+1}...\alpha_{n}}^{in}$$
(6.41)

We are interested in its analytic p-space properties. "On shell" p-space analytic properties are more elusive than x-space analytic properties. For the latter the spectral support properties play the important role, whereas p-space analyticity relies heavily on causality. The above matrix element still contains energy-momentum δ -functions resulting from contracting incoming p's with outgoing. These are removed by taking the connected parts of the formfactors. Only for the distinguished formfactor:

$$\langle 0 | A(0) | p_1, ..., p_n \rangle_{\alpha_1...,\alpha_n}^{in} = \langle 0 | A(0) | p_1, ..., p_n \rangle_{\alpha_1...,\alpha_n}^{in, \, con}$$
(6.42)

we have coalescence with its connected part. Similar to x-space analyticity, one expects the existence of one analytic master-function whose different boundary values correspond to the different n-particle formfactors:

$$_{\alpha_1...\alpha_m}^{out} \langle p_1, ..., p_m | A(0) | p_{m+1}, ..., p_n \rangle_{\alpha_{m+1}...\alpha_n}^{in, con}$$
(6.43)

6.3. THE BOOTSTRAP-FORMFACTOR PROGRAM

$$= F_{\underline{\alpha}}^{A}(s_{ij} + i\varepsilon, t_{rs} - i\varepsilon, s_{kl} + i\varepsilon), \quad i < j \le m < k < l \le n$$

$$t_{rs} = (p_r - p_s)^2, \quad r \le m < s \le n$$

There are Watson relations between the S-matrix and the form factors. In the d=1+3 dispersion theory setting it is well known that the cuts below the inelastic threshold of $\langle 0 | A(0) | p_1, p_2 \rangle$ is related to the partial wave phase shifts in that elastic region. In a factorizing d=1+1 theory these Watson relations can be written down in general:

$$F^{A}_{\alpha_{1}...\alpha_{n}}(s_{ij}+i\varepsilon) = \langle 0 | A(0) | p_{1},...p_{n} \rangle^{in}_{\alpha_{1}...\alpha_{n}}$$

$$= \sum_{out} \langle 0 | A(0) | out \rangle \langle out | p_{1},...,p_{n} \rangle^{in}_{\alpha_{1}...\alpha_{n}}$$

$$\curvearrowright F^{A}_{\alpha_{1}...\alpha_{n}}(s_{ij}+i\varepsilon) = F^{A}_{\alpha'_{1}...\alpha'_{n}}(s_{ij}-i\varepsilon)S^{\alpha'_{1}...\alpha'_{n}}_{\alpha_{1}...\alpha_{n}}(s_{ij})$$

$$(6.44)$$

and for the mixed formfactors(6.41):

$$F_{\underline{\alpha}}^{A}(s_{ij}+i\varepsilon,t_{rs}-i\varepsilon,s_{kl}+i\varepsilon)$$

$$= S_{\alpha_{1}'...\alpha_{m}'}^{\alpha_{m}...\alpha_{1}}(s_{ij})F_{\underline{\alpha}'}^{A}(s_{ij}-i\varepsilon,t_{rs}+i\varepsilon,s_{kl}-i\varepsilon)S_{\alpha_{m+1}...\alpha_{n}}^{\alpha_{n}'...\alpha_{m+1}'}(s_{kl})$$

$$(6.46)$$

Using the uniformazing $\theta's$, this is like a generalized quasi periodicity property on θ -strips for the F's (instead of the periodicity of S). The first who considered formfactors beyond two-particles [30] and presented a system of axioms for their calculation was Smirnov [27] Following a recent presentation by Babujian, Fring and Karowski [28] in a more standard field theoretic setting (LSZ+ dispersion theory), the formfactor program for the construction of d=1+1 QFT is as follows. Introduce the ordered formfactors:

$$f_{\underline{\alpha}}^{A}(\theta_{1},...,\theta_{n}) := \langle 0 | A(0) | p_{1},...,p_{n} \rangle_{\underline{\alpha}}^{in}, \quad \theta_{1} > ... > \theta_{n}$$
(6.47)

and define the value for reordered $\theta's$ by analytic continuation (starting with this ordering in the physical region). Demand that the f's fulfill the following properties ("axioms"):

• (i)
$$f^A_{\dots ij\dots}(\dots,\theta_i,\theta_j,\dots) = f^A_{\dots ji\dots}(\dots,\theta_j,\theta_i,\dots)S_{ij}(\theta_i-\theta_j) \quad \forall \ \theta's$$

• (ii)
$$f_{12...n}^A(\theta_1 + i\pi, \theta_2, ..., \theta_n) = f_{2...n1}^A(\theta_2, ..., \theta_n, \theta_1 - i\pi)$$

• (iii)
$$f_{1...n}^A(\theta_1,...\theta_n) \underset{\theta_1 \to \theta_2 + i\pi}{\approx} \frac{2i}{\theta_1 - \theta_2 - i\pi} C_{12} f_{3...n}^A(\theta_3,...,\theta_n) (1 - S_{2n}...S_{23})$$

where $C_{\alpha\beta} = \delta_{\bar{\alpha}\beta}$ is the charge conjugation matrix.

Here we have not mentioned the poles from bound states (states which appear by the previous fusion) since they are automatically entering the formfactors via the S-matrix. The word "axiom" in the context of this paper has the significance of working hypothesis i.e. an assumption which receives its legitimation through its constructive success. Physical principles on the other hand,

as the spectral and causality properties of general QFT, will not be called axioms. Our main aim is to show how one can reduce the above axioms of the bootstrap-formfactor approach to the principles of QFT, and thereby recuperate the unity of this nonperturbative approach with the rest of QFT.

The conceptually somewhat unusual property is the "symmetry" property (i). Here one should bear in mind that from the point of view of the LSZ formulation f is an auxiliary object to which the statistics property under particle exchange does not apply (it would apply to the original matrix-element). The above exchange property for f is a statement about analytic continuation. The statistics of incoming particle is only used in order to get the charges (i.e. the tensor factors) into the same j - i order as the analytically interchanged $\theta's$. Following BFK [28] let us first remind ourselves of the standard argument for (i) in somewhat detail. For the special case $\langle 0 | A(0) | \theta_1 \theta_2 \rangle^{ex} ex = in, out$ it is evident that:

$$lim_{\varepsilon \to 0} F(s_{12} \pm i\varepsilon) = \begin{cases} \langle 0 | A(0) | \theta_1 \theta_2 \rangle^{in} \\ \langle 0 | A(0) | \theta_1 \theta_2 \rangle^{out} \end{cases}$$
(6.48)

i.e. there is one analytic masterfunction f(z) (assuming identical particles) with different boundary values on the $s \ge 4m^2$ cut, having the *in*, *out* interpretation. Assuming Bose statistics, the physical matrix elements on the right hand side are symmetric under the interchange of the $\theta's$. In terms of the uniformization variable θ_{12} in F the transition from $in \to out$ means a change of sign via analytic continuation i.e. without changing the charge quantum numbers α i.e. the position of the tensor factors. After accomplishing this last step by the bose commutation relation, the negative θ_{12} formfactor $F(\theta_{21})$ can according to the definition (6.47) be identified with $f_{21}(\theta_2, \theta_1)$ and the relation

$$\langle 0 | A(0) | \theta_2 \theta_1 \rangle^{out} S(|\theta_1 - \theta_2|) = \langle 0 | A(0) | \theta_1 \theta_2 \rangle^{in}$$
(6.49)

agrees with (6.47). The generalization to ^{out} $\langle \theta_3...\theta_n | A(0) | \theta_1 \theta_2 \rangle_{con}^{ex}$ has a problem, because replacing in by out means passing from time-ordering to anti-timeordering but the LSZ scattering theory produces boundary terms contributing to the connected part. Although they are absent for theories in which the number of in-particles are conserved (on-shell conservation), it is unclear what property of general QFT is bringing about (i) through specialization to factorizing d=1+1 models.

On the other hand (ii) and (iii) are consequences of the following standard crossing formula [28] which relate the connected part of the generalized form-factors to the analytic master function f:

$$\frac{1}{\langle 0 | A(0) | p_2, ..., p_n \rangle_{2...n}^{in} }{\sum_{j=2}^{n} \frac{1}{\langle p_1 | p_j \rangle_j f_{2..\hat{j}..n}^A S_{2j}...S_{j-1j} + f_{12...n}^A (\theta_1 + i\pi_-..., \theta_n) }{\sum_{j=2}^{n} \frac{1}{\langle p_1 | p_j \rangle_j f_{2..\hat{j}..n}^A S_{jn}...S_{jj+1} + f_{2...n1}^A (..., \theta_n \theta_1 - i\pi_-) }$$

$$(6.50)$$

The fastest way to understand this is to draw the corresponding graphs and remember that a positive energy particle crosses into a negative energy antiparticle. Successive application leads to a formula which expresses the formfactors in terms of the analytic auxiliary function f. The analytic part of this relation gives (ii) whereas the δ -function part is responsible for (iii). A argument that these properties do not only insure TCP-invariance (weak locality) but also Einstein causality, can be given formally[27]. Apart from the Ising model, the formfactor program has not been carried out to the end although all of the two particle formfactors associated with the computed S-matrices are known.

It would be nice to have a direct derivation of all the bootstrap-formfactor axioms from the principles of QFT but this is still an open problem. It is part of the complicated and incomplete momentum space analyticity problem. Even the derivation of forward dispersion relations in particle physics took several years, not to mention the derivation of the analytic aspects ⁸ of crossing symmetry which still remain quite incomplete. It is precisely at this point where our modular localization approach shows its strength. To anticipate one result, it shows that the crossing symmetry is a kind of strengthened TCP-property, and that the cyclicity [38] [40] it leads to is identical to the KMS-temperature (≡Hawking -Unruh temperature in this special case) characterization of the (Rindler-)wedge based Hawking-Unruh effect. From our point of view, the most valuable result is that it opens for the first time the way to a new constructive iterative (but non perturbative) approach to *non-quantization*, non-Lagrangian based QFT. My confidence that this may amount to more than just another fashion rests on the observation that the tool of modular localization comes from a refinement of TCP which, as anybody will immediately admit, the central structure of local QFT.

6.3.3 Modular Theory and the Formfactor Program

There is one more important idea which is borrowed from scattering theory namely the existence of a "modular Møller operator" [39] U which is related to the S-matrix as:

$$S_s = U J_0 U^* J_0 (6.51)$$

This correspond to the well-known standard formula $S_s = (\Omega^{out})^* \Omega^{in}$. This analogy with a scattering interpretation has to be taken with a grain of salt however. Note that the Haag-Ruelle scattering theory (as well as its more formal but better known LSZ predecessor) in local quantum physics does not provide a Møller isometry between Heisenberg states and incoming states because the scattering state space and the space for the interacting fields are identical (in local quantum physics, different from the nonrelativistic rearrangement scattering theory one does not introduce a separate space of fragments). The name Møller appears here only in a modular context.

 $^{^8 \}rm Only$ together with the (mass shell) analytic properties the crossing symmetry aquires a physical content.

The idea of introducing such an object into our modular approach comes from the unitary equivalence of the interacting and the free hyperfinite type III_1 wedge algebras:

$$\mathcal{A}(W) = U\mathcal{A}^{in}(W)U^* \tag{6.52}$$

together with the U-invariance of the vacuum $U\Omega = \Omega$. Note that this situation is very different from the problem of equivalence of the canonical equal time commutation relations in the free versus the interacting case. A unitary equivalence in this case (of an algebra belonging to a region with trivial spacelike complement) would be forbidden by Haag's theorem on the nonexistence of the interaction picture in QFT. The above characterization of U may be replaced by a slightly more convenient one in terms of an intertwining property between modular operators:

$$US_0 = SU \tag{6.53}$$

Such a Møller operator cannot commute with the Poincaré group (apart from the boost associated with the wedge). The latter serves to define a family of W-affiliated U's from the standard wedge. In terms of localized spaces, the Uhas the property:

$$U\mathcal{H}_R^{in}(W) = \mathcal{H}_R(W) \tag{6.54}$$

Let us first illustrate these concepts in an explicitly known (including pointlike fields) model. Between the two possibilities Ising field theory with $S_s^{(2)} = -1$ and the (non-parity invariant) Federbush model with $S_{I,II}^{(2)} = e^{i\pi g\varepsilon(\theta_1 - \theta_2)}$ we chose the latter because it allows also a Lagrangian interpretation (i.e. more old fashioned nostalgia than conceptual necessity). The model consists in coupling two species of Dirac fermions via a (parity violating) current-pseudocurrent coupling [19][20]:

$$\mathcal{L}_{int} = g : j^I_\mu j^{II}_\nu : \varepsilon^{\mu\nu}, \quad j_\mu =: \bar{\psi}\gamma_\mu\psi :$$
(6.55)

One easily verifies that:

$$\psi_I(x) = \psi_I^{(0)}(x) : e^{ig\Phi_{II}^{(l)}(x)} :$$
(6.56)

$$\psi_{II}(x) = \psi_{II}^{(0)}(x) := e^{ig\Phi_{I}^{(r)}(x)} :$$

$$\psi_{I,II}^{(0)}(x) = \frac{1}{\sqrt{2\pi}} \int \left(e^{-ipx} u(\theta) a_{I,II}(\theta) + e^{ipx} v(\theta) b_{I,II}^{*}(\theta) \right) d\theta \quad (6.57)$$

where $\Phi^{(l,r)} = \int_{x' \leq x} j_0 dx'$ is a potential of $j_{\mu 5}$ i.e. $\partial_{\mu} \Phi \sim \varepsilon_{\mu\nu} j^{\nu} = j_{\mu 5}$ and the superscript l, r refers to whether we choose the integration region for the line integral on the spacelike left or right of x. For the form of the u and v spinors we refer to (3.210). The triple ordering is needed in order to keep the closest possible connection with classical geometry and localization and in particular to maintain the validity of the field equation in the quantum theory; for its meaning we refer to the above papers. This conceptually simpler triple ordering

can be recast into the form of the analytically (computational) simpler standard Fermion Wick-ordering in terms of the (anti)particle creation/annihilation operators $a_{I,II}^{\#}(\theta), b_{I,II}^{\#}(\theta)$.

$$\psi_{I}(x) = \psi_{I}^{(0)}(x) : e^{L_{g}(x)} :$$

$$L_{g}(x) = \frac{\sin\pi g}{2\pi} \int \left\{ \begin{array}{c} \frac{e^{-g(\theta_{p}-\theta_{q})}}{\cosh\frac{1}{2}(\theta_{p}-\theta_{q})} \left[e^{i(p+q)x} a_{II}^{*}(\theta_{p}) b_{II}^{*}(\theta_{q}) + h.c. \right] + \\ \left[-\frac{e^{-g(\theta_{p}-\theta_{q})}}{\sinh\frac{1}{2}(\theta_{p}-\theta_{q}+i\varepsilon)} \right] e^{i(p-q)x - i\pi g} a_{II}^{*}(\theta_{p}) a_{II}(\theta_{q}) + \\ \left[\frac{e^{-g(\theta_{p}-\theta_{q})}}{\sinh\frac{1}{2}(\theta_{p}-\theta_{q}-i\varepsilon)} \right] e^{-i(p-q) + i\pi g} b_{II}^{*}(\theta_{q}) b_{II}(\theta_{p}). \end{array} \right\} d\theta_{I} d\theta_{I}$$

and similar formula for ψ_{II} . Although in this latter description the classical (manifest) locality is lost, the quantum exponential do still define local Fermifields[20]; in the case of relative commutation of ψ_I with ψ_{II} the contributions from the exponential (disorder fields) compensate. This model belongs to the simplest class of factorizing models (those with rapidity independent S-matrix) and its explicite construction via the formfactor program is almost identical to that of the massive Ising field theory [102]. The reason why it does not appear under this approach in the literature is that the bootstrap classification was limited to strictly parity conserving theories. For our present purposes it serves as the simplest nontrivial illustration of new concepts arising from modular localization.

Despite the involved looking local fields (6.56), the wedge algebras are easily shown of utmost simplicity:

$$\mathcal{A}(W) = alg \left\{ \psi_{I}^{(0)}(f) U_{II}(g), \psi_{II}^{(0)}(h); suppf, h \in W \right\}$$
(6.60)
$$\mathcal{A}(W') = \mathcal{A}(W)'_{Klein} = alg \left\{ \psi_{I}^{(0)}(f), \psi_{II}^{(0)}(h) U_{I}(g); suppf, h \in W' \right\}$$

i.e. the two wedge-localized algebras (W denotes the right wedge) are generated by free fields "twisted" by global U(1)-symmetry transformation of angle g(coupling constant)⁹ and the subscript "Klein" denotes the well-known Klein transformation associated with the 2π Fermion rotation. The right hand side follows from the observation that with x restricted to W, one may replace the exponential in ψ_I in (6.56) (which represents a left half space rotation) by the full rotation since the exponential of the right halfspace charge is already contained in the right free fermion algebra etc. The following unitarily equivent description of the pair A(W), A(W') has a more symmetric appearance under the extended parity symmetry $\psi_I(t, x) \leftrightarrow \psi_{II}(t, -x)$:

$$\mathcal{A}(W) = alg \left\{ \psi_{I}^{(0)}(f) U_{II}(\frac{g}{2}), \psi_{II}^{(0)}(h) U_{I}(-\frac{g}{2}); suppf, h \in W \right\}$$
(6.61)
$$\mathcal{A}(W') = alg \left\{ \psi_{I}^{(0)}(h) U_{II}(\frac{g}{2}), \psi_{II}^{(0)}(f) U_{I}(-\frac{g}{2}); suppf, h \in W' \right\}$$

⁹The equality of the $\mathcal{A}(W)$ net (6.60) to the net obtained by the subsequent modular method adapted to the Federbush model is not a very easy matter.

The computation [20] of the scattering matrix S_s from (6.56) is most conveniently done by Haag-Ruelle scattering theory [3]:

$$S_{s} \left| \theta_{1}^{I}, \theta_{2}^{II} \right\rangle = S_{s}^{(2)} \left| \theta_{1}^{I}, \theta_{2}^{II} \right\rangle = e^{i\pi g\varepsilon(\theta_{1}-\theta_{2})} \left| \theta_{1}^{I}, \theta_{2}^{II} \right\rangle$$

$$S_{s}^{(n)} = \prod_{pairings} S_{s}^{(2)}$$

$$(6.62)$$

These formulae (including antiparticles) can be collected into an operator expression [20] :

$$S_s = \exp i\pi g \int \rho_I(\theta_1) \rho_{II}(\theta_2) \varepsilon(\theta_1 - \theta_2) d\theta_1 d\theta_2$$
(6.63)

Where $\rho_{I,II}$ are the momentum space charge densities in the rapidity parametrization.

The surprising simplicity of the wedge algebra as compared to say double cone algebras consists in the fact that one can choose on-shell generators. We will show that modular wedge localization for factorizing models always leads to on-shell generators though for rapidity dependent S-matrices they are less simple than (6.61).

It would now be easy to solve the n-particle modular localization equation¹⁰:

$$S\mathcal{H}_{R}^{(n)}(W) = -\mathcal{H}_{R}^{(n)}(W)$$

$$\mathcal{H}_{R}(W) = \left\{ \int F(\theta_{1}, \theta_{2}, ..., \theta_{n}) |\theta_{1}, \theta_{2}, ..., \theta_{n}\rangle \, d\theta_{1} d\theta_{2} ... d\theta_{n} \mid F \in H_{strip} \right\}$$
(6.64)

Here H_{strip} denotes the space of square integrable function which allow an analytic continuation into simplices inside a multi-strip:

$$Imz_{i_1} > Imz_{i_2} > \dots > Imz_{i_n}$$

$$0 < Imz_i < \pi, \ i = 1...n$$

$$(6.65)$$

This is just the p-space on-shell analyticity which comes from the wedge localization. As for Wightman functions in x-space, the n! different boundary prescriptions $Imz_{i_1} > Imz_{i_2} > \dots > Imz_{i_n} \to 0$ yield the generally n! different boundary values $F(\theta_{i_1}, \theta_{i_2}, \dots, \theta_{i_n})$. Similar statements hold for the boundary values on the upper rim. This boundary prescription (which for Wightman functions is a consequence of the energy positivity) follows from the analytic aspects of the KMS properties and the remark that the group parameters of the modular automorphisms are the rapidities. In the free case i.e. for $S_s = 1$, there are no discontinuities (i.e. the F's just incorporate the Fermi statistics) but with the Federbush S-matrix the space consists of strip-analytic functions which are

¹⁰The Tomita operator S for Fermions is different from that of Bosons by a Klein transformation. For a special family of d=1+1 solitons the correct TCP operator has been computed by Rehren[17]. Since all the known families of factorizing models are described by Fermions and Bosons and since it is not clear whether this generalization is compatible with the factorization we will ignore this more general TCP-situation in the present context.

a solution of a Riemann-Hilbert boundary problem, i.e. the Tomita eigenspace equation for $\mathcal{H}_R(W)$ relates the boundary values (products of the two particle S-matrices) on the various boundaries obtained by placing each single rapidity z_i on the lower/upper edge of the strip in the various ordered manners of the (real parts fo the) rapidities. The general solution of this problem (i.e. the characterization of the subspace $\mathcal{H}_R(W)$ within the full multiparticle wave function space) may be presented as a special solution of the Riemann-Hilbert problem convoluted with the general solution of the interaction free problem in $\mathcal{H}_R(W)^{in}$. A physically more enlightening way consists in writing the localization subspace in a field theoretic manner as:

$$\int d^2 x_1 \dots d^2 x_n f_n(x_1, \dots, x_n) \quad : \quad Z_{I,II}(x_1) \dots Z_{I,II}(x_n) : \Omega, \qquad (6.66)$$
$$supp f_n \quad \in \quad W^{\otimes n}, \ f_n \ real$$

where the Z's are on-shell operators whose frequency positive and negative momentum space components have to fulfil commutation relations which must be compatible with the boundary relations governed by products of two particle Smatrices. On immediately realizes that this leads to the Zamolodchikov-Faddeev algebra relations for the Federbush S-matrix:

$$Z_{I,II}(x) = \frac{1}{\sqrt{2\pi}} \int \left(e^{-ipx} u(\theta) c_{I,II}(\theta) + e^{ipx} v(\theta) d^*_{I,II}(\theta) \right) d\theta \tag{6.67}$$

where the c and the corresponding anti d can be formally expressed in terms of the incoming (anti)particle creation and annihilation operators:

$$c_{I,II}(\theta) = a_{I,II}(\theta)e^{-i\pi g \int_{-\infty}^{\theta} \rho_{II,I}(\theta')d\theta'}$$

$$d_{I,II}(\theta) = b_{I,II}(\theta)e^{i\pi g \int_{-\infty}^{\theta} \rho_{II,I}(\theta')d\theta'}$$
(6.68)

with the Zamolodchikov-Faddeev relations¹¹:

$$\begin{aligned} c_{I,II}(\theta_{1})c_{II,I}(\theta_{2}) &= -S^{(2)}(\theta_{1} - \theta_{2})c_{II,I}(\theta_{2})c_{I,II}(\theta_{1}) & (6.69) \\ c_{I,II}(\theta_{1})c_{II,I}^{*}(\theta_{2}) &= -S^{(2)}(\theta_{1} - \theta_{2})^{-1}c_{II,I}^{*}(\theta_{2})c_{I,II}(\theta_{1}) + \delta(\theta_{1} - \theta_{2}) \\ d_{I,II}(\theta_{1})d_{II,I}(\theta_{2}) &= -S^{(2)}(\theta_{1} - \theta_{2})d_{II,I}(\theta_{2})d_{I,II}(\theta_{1}) \\ d_{I,II}(\theta_{1})d_{II,I}^{*}(\theta_{2}) &= -S^{(2)}(\theta_{1} - \theta_{2})^{-1}d_{II,I}^{*}(\theta_{2})d_{I,II}(\theta_{1}) + \delta(\theta_{1} - \theta_{2}) \\ d_{I,II}(\theta_{1})c_{II,I}(\theta_{2}) &= -S^{(2)}(\theta_{1} - \theta_{2})^{-1}c_{II,I}(\theta_{2})d_{I,II}(\theta_{1}) \\ d_{I,II}(\theta_{1})c_{II,I}^{*}(\theta_{2}) &= -S^{(2)}(\theta_{1} - \theta_{2})c_{II,I}^{*}(\theta_{2})d_{I,II}(\theta_{1}) \\ &\{\cdot,\cdot\} &= 0, \forall others \end{aligned}$$

¹¹The natural appearance of the (rapidity-dependent) Yang-Baxter structure in these on shell Z-F operator algebras contains the interesting mathematical message that whereas the Artin braid group, which represents statistics of plektons, can be naturally represented in combinatorical type II_1 algebras ("topological field theory"), the natural representation of the Yang-Baxter structure requires "bigger" algebras related to scattering theory with spacetime aspects. There seems to be no easy "Baxterization" od the V. Jones tracial representations of the infinite braid group.

The simplicity of the model is reflected in the fact that interactions only take place between species I and II and the independence of $S^{(2)}$ on θ . The interaction does make a distinction between left and right and parity is only conserved if one also interchanges the two species. The operators (6.69) inserted into (6.67) lead to the same commutation relations as those of the generators of $\mathcal{A}(W)$ in (6.61). We still have to check that the *J*-transformed opposite algebra commutes with $\mathcal{A}(W)$ and is equal to the geometric opposite. Since there is no proper Lorentz transformation which transforms $\mathcal{A}(W)$ into $\mathcal{A}(W')$ we must take the previously mentioned unitary parity transformation *P* which in addition changes $I \leftrightarrow II$ for the definition of the geometric opposite. This amounts to the following relation

$$\mathcal{A}(W)' = J\mathcal{A}(W)J = \mathcal{A}(W') \equiv PA(W)P \tag{6.70}$$

The first equality is a consequence of the following relation for the generators:

$$j(c_{I,II}(\theta)) \equiv Jc_{I,II}(\theta)J = SJ_0a_{I,II}(\theta)e^{-i\pi g \int_{-\infty}^{\theta} \rho_{II,I}(\theta')d\theta'}J_0S^*$$
(6.71)

$$= Ka_{I,II}(\theta)K^*e^{i\pi g \int_{\theta}^{\infty} \rho_{II,I}(\theta')d\theta'}$$
(6.72)

where the adJ_0 transformation only changed the sign in the exponential and the adjoint transformation with the S-matrix generates an exponential factor : $\exp -i\pi g \int_{-\infty}^{\infty} \rho_{II,I}(\theta') d\theta'$: of which part of it compensates against the signchanged previous exponential factor. K is the Fermion twist of the modular theory for free Fermi-fields (chapter 3). Similarly we have

$$j(d_{I,II}(\theta)) \equiv J d_{I,II}(\theta) J = b_{I,II}(\theta) e^{-i\pi g \int_{\theta}^{\infty} \rho_{II,I}(\theta') d\theta'}$$
(6.73)

Therefore the TCP (\equiv J in d=1+1) transformed Z-F fields are:

$$j(Z_{I,II}(x)) = \frac{1}{\sqrt{2\pi}} \int \left(e^{ipx} j(c_{I,II}(\theta)) + e^{-ipx} j(d_{I,II}^*(\theta)) \right) d\theta$$
(6.74)

The relative commutation relations of $Z_I^{\#}(x)$ for $x \in W$ with $j(Z_I^{\#}(y))$ for $y \in W'$ are precisely those of a free field since e.g. in the Z_{I} - $j(Z_I^*)$ commutator the exponentials add up to the θ - independent term $\exp -ig\pi \int_{-\infty}^{\infty} \rho_{II}(\theta')d\theta'$ which multiplies the free (anticommutator) function with support outside the spacelike region. The argument for the Z_{II} - $j(Z_{II}^*)$ commutator with $locZ \in W$ is identical, whereas all other $Z^{\#}$ - $j(Z^{\#})$ commutators vanish without giving rise to contraction terms. There is one more property of the wedge algebra which ought to be checked: the opposite wedge $\mathcal{A}(W)'$ is geometric i.e. the justification for the Haag duality $\mathcal{A}(W)' = \mathcal{A}(W')$. Since the only geometric interaction free transformation which links the two wedges in a unitary way is the generalized parity covariance:

$$a_I^{\#}(\theta) \longleftrightarrow a_{II}^{\#}(-\theta), \ b_I^{\#}(\theta) \longleftrightarrow b_{II}^{\#}(-\theta)$$
 (6.75)

One easily checks that this transformation indeed transforms the wedge generators $Z^{\#}$ into those of $\mathcal{A}(W)'$. To see this one uses

$$a_{I}(\theta)e^{-ig\pi\int_{-\infty}^{\theta}\rho_{II}(\theta')d\theta'} \rightarrow a_{II}(-\theta)e^{-ig\pi\int_{-\infty}^{\infty}\rho_{I}(-\theta')d\theta'} \qquad (6.76)$$
$$= a_{II}(-\theta)e^{-ig\pi\int_{-\theta}^{\infty}\rho_{I}(\theta')d\theta'}$$

and the analogous transformation law for the b's. With this relation we have completed the checks on the generators of the wedge algebras. The surprising fact is that the wedge algebra which fulfills the cyclic and seperating conditions of the Reeh-Schlieder theorem admits on-shell generators generators which applied to the vacuum create one-particle states without the vacuum polarization clouds. For compact localization regions such operators cannot exist. If one looks at the more general factorizing models one finds that such generating onshell operators always exist. As a matter of fact it is not even necessary to look for formulas which represent the Z-F algebras in the incoming Fock space. Rather the only important point is to start from the Ansatz of modular localized states in the form:

$$\int \hat{F}(x_1, \dots, x_n) Z^*(x_1) \dots Z^*(x_n) \Omega$$

$$= \int F(\theta_1, \dots, \theta_n) \tilde{Z}^*(\theta_1) \dots \tilde{Z}^*(\theta_n) \Omega$$
(6.77)

where we have surpressed all indices distinguishing the various Z-F fields. The multivariable strip θ -analyticity comes from the localization and the relation between the various boundary values of the on-shell momentum space functions in rapidity variables function $F(z_1, ..., z_n)$ for the different orderings $\theta_{i_1} \ge \theta_{i_2} \ge ... \ge \theta_{i_n}$ is dictated by the Z-F commutation relations among the $\tilde{Z}^{*'}s$. It is very helpful to write the wave functions which characterize the Hilbert space $H_R^{(n)}$ in terms of the free functions $H_R^{(n)in}$ by splitting off a reference wave function F^{ref} :

$$F(\theta_1, ..., \theta_n) = F_n^{ref}(\theta_1, ..., \theta_n) f(\theta_1, ..., \theta_n)$$

$$H_R^{(n)} = F^{ref} H_R^{(n)in}$$
(6.78)

The reference wave function is most conveniently obtained by defining a state on the $Z^{\#}$ -generated wedge algebra $\mathcal{A}(W)$ which fulfills the KMS condition with respect to the modular group generated by the boost operator:

$$\left\langle Z^{\#}(x_1)...Z^{\#}(x_n)Z^{\#}(y_1)...Z^{\#}(\rho_n,\chi-i\pi) \right\rangle_{T=2\pi}$$
 (6.79)

$$= \left\langle Z^{\#}(\rho_n, \chi + i\pi) Z^{\#}(x_1) ... Z^{\#}(x_n) Z^{\#}(y_1) ... Z^{\#}(y_{n-1}) \right\rangle_{T=2\pi} (6.80)$$

Here all the mass shell Z-F operators have to be placed inside the wedge i.e. $y_i, x_k \in W$, and the χ denotes the x-space rapidity and ρ_n the radial coordinate of y_n . Besides this KMS condition we have enough boundary conditions from the commutation relation between the positive and negative mass shell components of the $Z^{\#'s}$ which lead to a recursive system linking the 2n correlation to the 2n-2 etc. This is not only a very elegant way for finding a special solution of the above Riemann-Hilbert problem, but it also emphasizes the physical role of the auxiliary operators as being attached to the wedge algebra $\mathcal{A}(W)$ which does not have any pure state (it is hyperfinite type III_1). The Lorentz-invariance in the momentum space rapidity together with the indicated iterative pairing

enforces the following diagonal θ -structure:

$$(\tilde{Z}^{\#}(\theta_1')...\tilde{Z}^{\#}(\theta_n')\Omega_{2\pi},\tilde{Z}^{\#}(\theta_1)...\tilde{Z}^{\#}(\theta_n)\Omega_{2\pi}) = \prod_i \delta(\theta_i' - \theta_i)F_n^{ref}(\theta_1,...,\theta_n)$$
(6.81)

The KMS condition relates this function to another one in which a particle on one side is missing and an antiparticle on the other created. This KMS property of the auxiliary fields is the germ for the crossing symmetry of the local fields. The reference F^{ref} ((suitably normalized) defines one concrete realization of the n-particle component of the modular Møller operator. As a matter of fact, any wedge localized vector $\psi = B\Omega_{2\pi}$ could have been used to define via $(\psi, Z^{\#}(y_1)...Z^{\#}(y_n)\Omega_{2\pi})$ a modular Møller isomorphism $U^{(n)}$ from $H_R^{(n)in}$ to $H_R^{(n)}$; i.e. there are as many transformations as there are localized vectors with n-particle components. On the other hand if we would have a modular Møller operator U in Fock space, the on mass shell auxiliary fields are defined as in (6.52) $Z^{\#}(0) = U\psi^{\#}(0)U^*$, $Z^{\#}(x) = U(x)Z^{\#}(0)U^*(x)$, $x \in W$, where in the last formula U(x) denotes the translation.

The F^{ref} for different particle number are in principle all independent. What we are really interested in for the formfactor program is of course the case where B is either a pointlike field B(0) or an operator from a double cone algebra. In that case the function transformed to rapidity space is directly related to the B-formfactor between n-particles and the vacuum and the rapidity transformed KMS condition becomes the crossing relation since the vectors generated by the application of the $\tilde{Z}^{\#}$ are intimately related to the on mass shell matrix elements of B(0) between incoming particle states. The raison d'etre for the existence of the auxiliary wedge-localized fields becomes now clear: these operators are the mediators between on-shell analytic properties and the space-time localization properties. Despite their wedge localization properties (which are reponsible for avoiding the consequences of the Haag theorem), the algebra generated by them can resolve one particle states which would not be the case for smaller localization regions which are not left invariant by a boost subgroup of the Poincaré group. Their existence cannot be derived from scattering theory since the wave packets in scattering theory cannot be localized in a wedge. Note also that the crossing symmetry does not hold for the incoming on-shell matrix elements of the auxiliary fields, but it is the above KMS property which leads to the crossing symmetry for *local* fields. What is truly amazing is that these fields bring together three aspects which up to now had their separate places: The pure quantum aspects of the Bekenstein-Hawking-Unruh issue (whose semiclassical manifestation shows up in black hole physics, the on-shell crossing properties and the related nuclear democracy or duality (that one which was suggestive for the Veneziano dual model) and the uniqueness of the field theory related to an admissable S-matrix (the inverse problem of LQP) and the nonperturbative construction idea based on modular localization. This of course begs the question of whether the existence of the mass shell vacuum polarization free fields can also be guarantied outside of factorizing models, to which we have some comments in the next section.

The last and difficult question, for which we will again use the Federbush model for clarification is: how can we understand double cone algebras and local fields? From the formula (6.56) as well as from the formfactor rules (axiom III) in the previous section it is clear that for local $\psi's$ i.e. for formfactors F_B of pointlike fields B, the n-point components become related by a kind of cluster property which identifies the residuum of poles in the analytic continuation with lower formfactors. The mechanism behind this can be better understood by looking at the form of (6.56) and asking the question how does the local field manage to fulfill a pointlike TCP transformation $J\psi(0)J \sim \psi^{C}(0)$ whereas the $Z^{\#}$ -fields only insure the correct transformation (6.71) of the entire wedge algebra $\mathcal{A}(W)$. So what kind of magic in the formfactors $\left\langle \Omega \left| \psi^{\#}(0) \right| \theta_1, ..., \theta_n \right\rangle, \ \theta_1 \geq \theta_2 ... \geq \theta_n$ is responsible for the pointlike TCP-property? Obviously the pointwise (geometric) TCP property is most manifest in the representation (6.56) whereas the (more quantum) ordering which allows a clear-cut separation of n-particle components is the Fermion Wick-ordering. The ordering we use for our auxiliary operators (6.68) is different from the Fermion ordering by a very simple cumulant expression in which the two terms are only different by a simple c-number factor whereas in the presence of fluctuation (particle-antiparticle) terms the operator terms change and the Fermion ordered bilinear exponentials have a nontrivial $i\varepsilon$ pole structure in rapidity space. It is precisely this structure which, similar to the cluster decomposition property, relates the formfactors for different n. Therefore the formfactor rule axiom *III* which relates the pole structure with a lower formfactor (i.e. the only difficult part of bootstrap-formfactor program) corresponds to the step from the wedge localization to the compact or pointlike localization. This was to be expected on the basis of LQP physical intuition. For the wedge particle massive state can still be identified and the modular localized n-particle states can be chosen independently whereas for compact localization regions it is not possible to separate single particles from the "clouds" which accompany them and which regulate the relation of the formfactors (the components of local field vector states) for different particle number.

The computation of the above reference wave functions from the KMS structure (6.79) is particularly simple for the Federbush model. All thermal two-point functions are equal to the wedge restricted free field vacuum expectations which according to chapter 3.8 have a manifest thermal representation. This is a property of all local fields. Only for the nonlocal auxiliary fields this identity between restriction and thermal representation is violated and, as we have seen above, we must take the thermal formula for the calculation of the coefficient functions F which appear in the particle rapidity representation of the wedge localized states. Obviously the lowest nontrivial function for which our KMS formalism becomes relevant, is the 4-point function:

$$\left\langle Z_{II}(\theta_2') Z_I(\theta_1') Z_I^*(\theta_1) Z_{II}^*(\theta_2) \right\rangle_{2\pi} \tag{6.82}$$

$$\sim e^{-\frac{1}{2}i\pi g} \left\langle Z_{I}(\theta_{1}')Z_{I}^{*}(\theta_{1}) \right\rangle_{2\pi} \left\langle Z_{II}(\theta_{2}')Z_{II}^{*}(\theta_{2}) \right\rangle_{2\pi} + e^{\frac{1}{2}i\pi g} \left\langle Z_{II}(\theta_{2}')Z_{II}^{*}(\theta_{1}) \right\rangle_{2\pi} \left\langle Z_{I}(\theta_{1}')Z_{I}^{*}(\theta_{2}) \right\rangle_{2\pi}$$
(6.83)

where up to numerical factors the rapidity representation coalesces with the modular group (boost variable) labeling. The thermal two-point functions at $T = 2\pi$ are equal to the wedge restricted free expressions. The first nontrivial functions are the 6-point functions:

$$\langle Z_{II}^{*}(\theta_{3}')Z_{II}(\theta_{2}')Z_{I}(\theta_{1}')Z_{I}^{*}(\theta_{1})Z_{II}^{*}(\theta_{2})Z_{II}(\theta_{3})\rangle_{2\pi}$$

$$\sim e^{-\frac{1}{2}i\pi g} e^{-\frac{1}{2}i\pi g} \frac{e^{-g\pi(\theta_{2}-\theta_{3})}}{\cosh\frac{1}{2}(\theta_{2}-\theta_{3})} \cdot \\ \cdot \langle Z_{I}(\theta_{1}')Z_{I}^{*}(\theta_{1})\rangle_{2\pi} \langle Z_{II}(\theta_{2}')Z_{II}^{*}(\theta_{2})\rangle_{2\pi} \langle Z_{II}^{*}(\theta_{3}')Z_{II}(\theta_{3})\rangle_{2\pi}$$

$$+ \dots$$

$$(6.86)$$

where the dots stand for three other terms which result from permutations of θ_1 with θ_2 and θ_1 with θ_3 weighted with appropriate phase factors. These expressions should be consistent with $\langle \Omega, Z_{II}(\theta'_2)Z_I(\theta'_1)N(\psi^*_I(x)\psi^*_{II}(x))\Omega \rangle$ and $\langle \Omega, Z_{II}^*(\theta'_3)Z_{II}(\theta'_2)Z_I(\theta'_1)N(\psi^*_I(x)\psi^*_{II}(x)\psi_{II}(x))\Omega \rangle$ where N denotes the leading operator term in a short distance expansion. This is easily checked if one pays attention to the charge terms in the $i\varepsilon$ prescription which are contained in the exponential of (6.59). The reference functions for the description of the wedge localization spaces of the more general factorizable models are also constructed by these thermal method. The important new contributions are the minimal formfactors which. We will present the systematics together with non-trivial illustrations in a seperate paper.

Although Gibbs states also share the KMS property, it should be stressed that the thermality which originates from restriction in QFT, is described by KMS without a Gibbs representation. There seems to be a confusion in the literature on this point.

For the Federbush model it is very easy to solve all these problems since the S-matrix does not depend on the individual rapidities but only on their order. For more general factorizing models, the special wave functions obtained from the KMS formalism as well as the formfactors are (as the S-matrix) meromorphic functions in the rapidity variables which in this way is elevated to a uniformization variable of the problem. Although this simple Federbush model has no bound states, one expects that they show up in the thermal expectations as soon as they contain one local local field in addition to the wedge localized Z's. Note that they would occur slightly outside the strip region; if one θ is in the strip, another one must be at the reflected point below the strip; only in this pair-formation one encounters particle poles. Expressed in terms of the difference variables this of course agrees with the old findings of Karowski-Weisz [30]. We mention that in a previous $paper^{12}$ [39] we have discussed the relation between the eigenvalue equations for the vectors in the modular localization space $\mathcal{H}_{R}(W)$ and the Riemann-Hilbert problems resulting the formfactor "axiomatics" following [30][27][103] without the thermal use of the Zamolodchikov-Faddeev algebra. But it is only this thermal aspect which allows to incorporate

 $^{^{12}}$ Unfortunately there were some errors in the formulas which characterize the wedge localization spaces in [39].

the latter into the general framework of QFT. Whereas the representation (6.61) for $\mathcal{A}(W)$ is a peculiar property of models with constant S-matrices, the auxiliary thermal Z-fields exist for all factorizable models. The modular study of the Federbush model has supplied us with physically rich new nonperturbative concepts which are collected in the following remark:

Remark 4 For a given physically admissable S-matrix there exists a unique interacting wedge algebra which is generated by semilocal on-shell fields which for d=1+1 factorizing theories fulfill the Zamolodchikov-Faddeev algebra in momentum space. Such a field applied to the vacuum creates a one particle state. Although the generated algebra is TCP-covariant, these fields are not yet TCPcovariant. In order to achieve the latter property of covariant transformation property under TCP, one must find the generators for double cone algebras of arbitrary small size. These generators (more generally all compactly localized operators) have a rich virtual particle structure such that the vector they generate from the vacuum has its higher particle components determined in terms of the lower ones. In fact the coefficient functions of the local operators are nothing else then their generalized formfactors. Whereas the general analytic structure of an n-particle formfactor is already determined by the auxiliary fields (in particular the KMS property of the latter are responsible for the crossing symmetry of the local operators), the relation between the formfactors for different n as well as their pole-structure is only understood after having passed from wedge to compact (double cone) modular localization. The existence of the intermediate semilocal wedge algebra generating fields is related to the existence of a modular Møller operator; in fact the auxiliary semilocal field is nothing but the field T(x)(6.87). There are as many T's as there are free Wick-polynomials of the free fields related to the S-matrix (including those which describe bound states).

This situation for factorizable models is only the tip of an iceberg. Every local QFT has thermal variables T(x) without vacuum fluctuations which generate the wedge algebra and are related to the modular Møller operator U:

$$T(x) \equiv U(x)UA_{in}(0)U^*U^*(x), \ x \in W$$
(6.87)

where the translation U(x) is acting on $T(0) \equiv UA_{in}(0)U^*$. In spite of our simplified notation, the U depends on the wedge and we should rather talk about a family U(W).

We will defer all explicit calculations concerning the construction of the local model fields to a separate paper. The reason for this postponement is that we want to present the construction of the factorizable models fully in the spirit of algebraic QFT (where special local field coordinates are avoided), i.e. by constructing the net of double cone algebras from the wedge algebras¹³ and the relevant concepts for these calculations are still unfinished. In particular our

 $^{^{13}}$ The main step in the algebraic constructive program is really the calculation of the wedge algebra with a *geometric* commutant. Barring the possibility that all intersections of wedge algebras are empty, the existence of local double cone algebras (and thei generating local fields) is secured by the above main step.

knowledge of the interpretation of the modular objects (J, Δ^{it}) for double cones is extremely scarce. Apart from the special case of conformal theories and a conjecture about a geometrically "fuzzy" action of Δ^{it} in general, nothing is known. The wedge situation, for which all the interaction resides in J and Δ^{it} is given in terms of Lorentz boosts, is certainly no guide for the double cone localization. Even in the case of the simple Federbush model one can show that there are no generators which (if applied to the vacuum) create one particle states without clouds of virtual particle-antiparticle pairs. This is true, although the model has no real (on-shell) particle creation.

One can show that the modular construction of "free" anyons and plektons [39] in d=1+2 leads to similar mathematical problems. In this case the whole construction takes place in a scattering space which, in contradistinction to Fermions and Bosons, has no tensor product structure in terms of Wigner spaces. The braid group commutation relation leads to a Tomita J which again involves a constant matrix S, however in this case it does not carry scattering information but is identical to the braid group representation R-matrix. Formally such a situation has some resemblance with the Federbush model, apart from the fact that now the opposite of $\mathcal{A}(W)$ in the quantum sense of the von Neumann commutant is different from the geometric opposite $A(W)' \neq A(W')$. This difference is accounted for by a statistics "twist" S_{twist} . Again the model has a rich virtual particle structure even though no real particle is created. But since this time the rich virtuality is a result of the nontrivial statistics twist S_{twist} , there is no reason whatsoever to expect that this fades away in the nonrelativistic limit. The fact that the nonrelativistic limit of Fermions and Bosons leads to the Schrödinger QM is related to the existence of relativistic free fields in Fock space. But since the Fock space structure in d=1+2 cannot support anyons and plektons, there is all reason to expect a kind of nonrelativistic field theory which can incorporate the virtuality which is *necessary* to maintain the relation between spin and statistics in the nonrelativistic limit. Indeed all attempts to incorporate braid group statistics into QM ever since the time of Leinaas and Myrheim have only led to a deformation of (half)integer spin but not to nonrelativistic operators with the correct spin-statistics commutation structure. A consistent multiparticle QM with braid-group statistics i.e. a theory which leads to a multiparticle S-matrix fulfilling the clusterproperties does presently not exist. The investigation of two-particle Aharonov-Bohm scattering is not sufficient to settle the issue of braid group statistics in QM. The above discussion casts doubt on the quantum mechanical nature of braid group statistics i.e. on the possibility to have a nonrelativistic description without virtual particle-antiparticle creation. This could explain the negative results of all pure geometric attempts in terms of Schrödinger wave functions. The method of using the Wigner representation and the correct multiparticle structure from scattering theory [92] together with the present modular localization method looks well defined and promising, but still needs to be carried out. A somewhat easier problem is the use of the modular localization method in order to construct chiral conformal theories with given plektonic statistics. Some remarks are contained in the next section.

In the remainder of this section we would like to make some pedagogical remarks for readers with an insufficient knowledge of general properties of nonperturbative QFT. In connection with rapidity dependent Zamolodchikov-Faddeev algebra operators, as well as x-dependent chiral conformal operators, one often finds the erroneous concept of "analytic field operators" and "holomorphic algebras. Because their use is so widespread (the few articles where this misleading concept was not used are an exception), it is interesting to ask where such incorrect ideas are coming from. Since I am not an expert on string theory, I have limited my search to QFT. The oldest paper which could be interpreted as alluding to "analytic operators" $A(z), z \in \mathbf{C}$ seems to be the famous BPZ [59] paper¹⁴ on minimal models. Although the authors do not use such terminology in print, the notation used in that paper could be misunderstood (and has been misunderstood by many physicist whose first experience with QFT came through that famous work or was influenced by string theory). The truth is that field algebras never have holomorphic properties. The analytic properties of correlation functions and state vectors depend entirely on the nature of states one puts on those algebras. Whereas vacuum ground states lead to the famous BHW-domain (in chiral conformal QFT equal to a uniformormization region with poles for coalescing coordinates), temperature states will only yield strip analyticity. In the above thermal construction the problem is to compute a L-boost KMS state at temperature $T = 2\pi$ on the Zamolodchikov-Faddeev algebra and not to construct a regularized Gibbs state on an invented algebra of analytic operators $Z(\rho, z_{boost})$ or $Z(z_{rapidity})$.

It is a fact that the associated analytic Bargman-Hall-Wightman domain for conformal covariant QFT and factorizing models is larger than that of the corresponding Poincaré covariant massive theories) and as a result of braid group statistics one looses the unicity of these analytic continuations (but never the univaluedness in the real time physical localization points!) . But neither the old nor the new BHW domains have anything to do with the "living space" of fields in the sense of quantum localization of operators in this article. The analogy between x-space analytic properties of conformal QFT and momentum rapidity space properties of factorizing theories have been observed by many authors. They are related to the similarity of p-space spectral support properties with xspace localization properties. These analogies do however not justify the search for a common formalism; the conceptual situations remain essentially different.

In an attempt to attribute a constructive meaning to the above unfortunate but existing terminology, one could point at a property which has *no analogue in higher dimension* (not even in the conformal invariant limit of higher dimensional theories). Whereas generally with vacuum expectation values one can relate at most two physical theories: a noncommutative real time QFT and a commutative euclidean candidate for a statistical mechanics, a chiral conformal theory on one light cone has infinitely many noncommutative boundary values (and no commutative) each of which defines a set of positive definite correlation

 $^{^{14}}$ In an older paper on conformal blocks e.g. [56][57] (called nonlocal components in a conformal decomposition theory with respect to the center of the universal covering) such "holomorphic" terminology was never used.

functions and hence a theory. Namely the restriction of the analytically continued correlation function defines a positive QFT not only on the circle (the standard living space of chiral theories) but also on each boundary encircling the origin (with the right $i\varepsilon$ Wightman boundary prescription). Although this does not legitimize the notion of "holomorphic operators" in the literal sense, the existence of a operator conformal QFT for each chosen boundary within the BWH analyticity domain is a significant difference to the mentioned higher dimensional situation on which one may base this terminology. The reader recognizes easily that this operator structure is equivalent to the existence of the infinite dimensional diffeomorphism group which is related to the Virasoro algebra structure in fact it is the LQP version of the diffeomorphism covariance of the theory. The application of a symmetry, which does not leave the vacuum reference state invariant, defines another set of positive Wightman functions which at the case at hand belong to the deformed boundary. Its deep relation to the modular theory will be commented on in the next section.

6.3.4 Modular Construction in General Case, open Problems

One important general structural question is whether a physically admissable scattering matrix S_s belongs to only one QFT (or better LQP which is the field-coordinate independent content of QFT). This is the famous inverse problem of QFT. As we saw before, the modular Møller operators are highly nonunique, so the question is if $\mathcal{A}(W)$ defined by (6.52) is uniquely determined. Suppose their would be two different U's say U_1 and U_2 . Then $U = U_2^*U_1$ commutes with the free Tomita involution S_0

$$US_0 = S_0 U \tag{6.88}$$

so the problem has been reduced to the question of whether $S_s = 1$ implies a free field theory. The proof of this old folklore statement of QFT is surprisingly difficult. Here it helps to recall the physical picture that in order to identify the vacuum and the one-particle state of a QFT it is not necessary to know the theory globally, but it suffices to know the wedge algebra and the geometric action of the modular L-boosts. Only if one also in addition wants to identify scattering states, one of course needs the whole Minkowski space.

The modular localization spaces for wedges are equal and agree with that of a free field theory. From this we would like to conclude that the same is true for the associated algebras i.e. $\mathcal{A}(W) \equiv U\mathcal{A}_0(W)U^* = \mathcal{A}_0(W)$. Since J commutes with U, the commutant of $\mathcal{A}(W)$ is

$$\mathcal{A}(W)' = U\mathcal{A}_0(W)'U^* = U\mathcal{A}_0(W')U^*$$
(6.89)

In fact from the equality of the modular operators and real spaces one could try to prove a slightly stronger statement that the so called natural cones $\Delta^{\frac{1}{4}}\mathcal{A}_{+}(W)\Omega$ and $\Delta^{\frac{1}{4}}\mathcal{A}_{0,+}(W)\Omega$ are equal. According to a theory of Connes this implies the equality of the algebras. A more direct physicist's argument would consist in using the linear map L between the algebras defined by

$$L : \mathcal{A}(W) \to \mathcal{A}_0(W)$$

$$A \to A_0, \ A\Omega = A_0\Omega$$
(6.90)

This is a *-linear map with the following identity for the special matrix elements:

$$\langle \Omega, A\chi \rangle = \langle \Omega, A_0\chi \rangle, \ \chi \in \mathcal{A}_0(W)\Omega = \mathcal{A}(W)\Omega$$
 (6.91)

Taking for χ wedge localized n-particle states and using free field crossing symmetry, the equality generalizes to general bra-states χ' .

As far as the strategy of constructing interacting models in d>1+1 is concerned, we do not have the possibility to extract a bootstrap program for the S-matrix. Hence the idea to introduce interactions via known S-matrices is not a good one. A better object for the general case seems to be the operator U, which, as the formfactors, is partially off-shell (more "local" than S) and has a quadratic relation to the S-matrix. In that case the first step of the program namely to find an "interacting pair" $\mathcal{A}(W)$ and $\mathcal{A}(W)'$ which is U-related to corresponding free algebras such that $SU = US_0$, $[U, \Delta] = 0$ is easily carried out with the result that the candidate for the modular scattering matrix would be $S_s = JJ_0 = UJ_0U^*J_0$. Certainly one would not expect that an interaction introduced this way and used first for the construction of a net of modular localized spaces and then for a net of algebras, will lead all the way to nontrivial local nets of algebras of a LQP. It would however be very interesting to find the obstructions and understand them.

The analogy with the miraculous "corner transfer matrix" in 2-dim. lattice problems is very tempting. In a relativistic theory the standard Hamiltonian picture of implementing interactions (which distinguishes a time direction) is not very natural, although one has accumulated a lot of experience and intuition. The modular aspects of the formfactor program for factorizing theories suggests to take the wedge proposal serious and to develop the necessary intuition for "good U's". A guiding idea should be the existence of mass shell auxiliary fields which create vacuum polarization free states. Formally one would imagine to write for $x \in W$:

$$\psi_{aux}(x) = U(x)U\psi_0(0)U^*U^*(x) =$$

$$= \int \sum \left\{ e^{-ipx}u(\theta,..)a_{aux}(\theta,..) + e^{ipx}v(\theta,..)b_{aux}^*(\theta,..) \right\} d\theta..$$
(6.92)

where θ is the wedge affiliated rapidity and the dots stand for the transversal variables. For the auxiliary fields one again expects a form

$$a_{aux}(\theta,..) = a(\theta,..)e^{i\eta_{-}(\theta)}, \ b^{*}_{aux}(\theta,..) = ...$$
(6.93)

with the operator $\eta_{-}(\theta)$ of the form:

$$\eta_{-}(\theta) = \sum_{\theta}' \int_{\theta}^{\infty} \eta(\theta_{1}, ..., ..., \theta_{n}, ..) : \tilde{\psi}(\theta_{1}, ..) \tilde{\psi}(\theta_{n}, ..) :$$
(6.94)

Here ψ denotes the Fourier transform of the free field, i.e. the creation/annihilation operators of particles/antiparticles, and the double dots represent the free field Wick-ordering. The integration extends over all transverse variables and over the interval $[\theta, \infty]$ in the longitudinal rapidities θ_i . The η -operators do not contain transitions between vacuum as well as one-particle states to other states. This is expressed by the dash on the sum and is nothing but the formulation of absence of vacuum polarization in the ψ_{aux} operators. In analogy with the integrable case, this operator should be thought of as the $[\theta, \infty]$ part of the operator η in a representation:

$$Sa(\theta, ..)S^* = a(\theta, ..)e^{i\eta} \tag{6.95}$$

An approach via interacting models of modular Møller operators is of course not expected to lead to explicit 4-dimensional nets of operator algebras. Rather one expects to obtain a controllable iterative framework which is in analogy to QM more like the Hartree Fock theory than standard perturbation theory. There is of course always a chance that there exist simple and controllable LQP's which are not accessible from the perturbative point of view (e.g. because they do not contain a deformation parameter).

The close relationship of the modular localization approach with its thermal aspects leads to the question whether the wedge generating auxiliary fields allow for a lattice version which could bring in a quantum mechanical aspect. Lattice theories in infinite lattices with short range interactions are conceptually somewhat similar to QFT in that they have a sharp particle concept an fulfill cluster properties which (in the absence of causality) are strong enough to allow for the derivation of scattering theory. The (vague) analogy of U with the corner transfer matrix also points into this direction. The modular localization approach would however not support an ad hoc momentum space cutoff since the modular structures are inexorably linked with Einstein causality and furthermore since such a cutoff wrecks the basis of time dependent scattering theory and more generally the physical interpretability altogether..

Note that we have used the word "nonperturbative" in an intrinsic sense, i.e. we do not expect that there are special field coordinates in terms of which the theory becomes long- or short-distance perturbative. The modular ideas are consistent with the construction of scale invariant limiting theories but do not support the wild scale-sliding which seems to be a hallmark of the quantization approach.

6.4 Old Ideas on Conformal Theories in new Setting

The desire of constructing nontrivial and dynamically interesting nonperturbative models of QFT is as old as the Wightman framework. The attempts in this direction can be divided into two groups. The main thrust was to follow the formalism of perturbation theory enriched with ideas from rigorous statistical mechanics. The constructive approach was subdivided into two steps of which the first one was to control the system in a finite space-time volume and the second step consisted in performing the thermodynamic limit [106]. Since the first step in most cases used the canonical formalism, it was severely limited to superrenormalizable couplings of the ϕ_2^4 -type. Even with all the subsequent improvements (partially due to better operator regularizations, in particular the lattice regularization), it was not possible to control a strictly renormalizable (i.e. non-superrenormalizable) model by these methods. The later success with low dimensional models which started in the late 60^{ies} with the massless Thirring-model was based on different construction methods.

Our nonperturbative methods in this chapter would suggest that any method, which follows the conceptual framework of the perturbative approach too closely, is liable to suffer from spurious short distance problems caused by perturbative methods. These problems are known to every field theorist, and they led to the questionable belief that there is (even outside of perturbation theory) a direct connection between the existence of a QFT and the short distance behavior of its fundamental fields. The algebraic framework of LQP, with its emphasis on local algebras as the carriers of the physical content, would cast doubt on this idea and the concrete construction of factorizing theories based on modular wedge localization with its emphasis on polarization free on-shell auxiliary fields has substantiated this doubts. It seems that the traditional constructive approach ("constructive QFT") is very limited indeed.

On the other hand there were always attempts to look for nontrivial models away from concepts related with perturbation theory. The first attempt to get away from free fields was to study so called generalized free fields; but unfortunately they turned out to be dynamically quite useless, although their pathological aspect had the beneficial effect of directing attention towards the important problem of "degree of freedom counting" i.e. the issue of phase space behavior in QFT.

A second idea was to look for "Lie-fields", i.e. a set of fields whose space-time commutation relation close on themselves (the predecessors of the *W*-algebras). This proposal apparently was made by W. Greenberg in the 60^{ies} . The special case of equal time current relations (current algebra) was known to restrict, but not to completely determine the models. More interesting was the idea of space-time Lie-fields, but in the first attempt to analyse this situation [107], the examples given were too trivial and somewhat discouraging.

The obvious place to look for was 2-dimensional conformal QFT. Already in 1970 I presented some results on the conformal invariant aspects of the massless Thirring model and their local generation by the conformal energy momentum tensor at a "Conference on Special Topics in Quantum Field" July 27-31, St. Louis. This material entered a joint paper with John Lowenstein [108]. It appears somewhat curious in retrospect that we did not use the Lie-field idea immediately in order to liberate ourselves from special models. Only in 1973 it became clear to me that a general structure analysis on conformal energymomentum tensors can be performed. The result I presented at the January 1974 Rio de Janeiro V Brazilian Symposium and it was published in the proceedings which the reader finds in their original form attached as an appendix to these notes. The reason why the result was not submitted to an international journal was that the topic was quite marginal to most of the theoreticians at that time. In that article there also appear for the first time differential identities (equ. 6.22) in addition to the equations of motion which are characteristic of chiral conformal theories and which many years later became known (together with the equation of motion and in a somewhat more general context) as the Knishnik-Zamolodchikov equations. The paper also shows that charge carrying fields for nonabelian conformal current algebras were an object of investigation already in the early 70^{ies} . In fact the main motivation for my contribution was the observation that the interchange of equations of motions with differential identities led to the erronous claim that an appearantly new nontrivial conformal fixed point of the generalized Thirring model had been found. In more recent times these models were called Wess -Zumino models by Witten; the more precise terminology would have been W-Z-W "field coordinates"¹⁵ in the old current algebra models. These field coordinates are helpful if one wants to separate these conformal invariant theories from the generic family of generalized (massive) Thirring models within a Lagrangian framework; they are of lesser help in the actual calculation of the correlation functions of the charge-carrying fields of the conformal current algebras.

Three years before the L_n -algebra of Fourier coefficients had appeared in a totally different context of the dual model S-matrix in a paper of Virasoro [110]. Apparantly Virasoro was not aware about the ongoing work on d=2conformal QFT, in particular the massless Thirring model, since otherwise he could hardly have missed the central term c which inevitably shows up in every conformal model. I certainly did not know Virasoro's findings since, apart from sociological reason for not being a member of any group close to one of the big laboratories, that global L_n on-shell (S-matrix) formalism of the dual model was physically too far away from my local (off-shell) mode of thinking which, as far as physical content is concerned, was closer to the conceptual framework of the present notes. The first physicists who also studied the conformal aspects of the Thirring model and observed the formal relation to Virasoros contribution on the algebraic structure of Veneziano's dual S-matrix model was an group of Italian physicist [109]. Looking back at its content, one observes the rare sociological curiosity that an author, whose prior printed work on the same subject is not referered to, is nevertheless mentioned in the acknowledgments. That very same paper introduced for the first time (I was not able to find an earlier reference) the terminology "Virasoro-algebra" for the algebra generated by the Fourier coefficients of energy-momentum tensor.

The above mentioned unfashionable aspect of algebraic structures of twodimensional QFT was related to their off-shell physics aspect, whereas the Zeitgeist in those times was definitely on mass shell before it went off shell with the renewed interest in conformal field theory of the 80^{ies} . This is another of the

 $^{^{15}}$ Group-valued fields in d=2 euclidean theories similar to those introduced by Witten also made their natural appearance in the older literature on QCD₂ Fermion determinants [111].

curiosities of those times.

These findings in 1973 were sufficiently encouraging for Jorge Andre Swieca and myself to collaborate on conformal QFT. In 1974/75 the results were published in two papers [56][57]. The conformal blocks (called "nonlocal component fields") resulted from the decomposition of the local fields with respect to the center of the conformal covering. Physically this amounted to a resolution of the "Einstein causality paradox" which consisted in the observation that any violation of Huygen's principle in massless theories means that the would be local conformal fields are, contrary to popular opinion reducible and decompose into irreducible components under the central decomposition of the conformal covering, i.e. Einstein causality of the original local fields requires the existence of nontrivial irreducible components under the conformal covering; the idea that the local field is conformally irreducible was too naive and responsible for the paradox.

The discovery of the nontrivial family of minimal models by BPZ [59] 10 years later was making use of the increased knowledge of special algebras and their Verma module representation theory. In that case one is working outside known quantum algebras and the problem of passing from a Verma module to a Hilbert space with a representation of a *-algebra belongs to the most subtle points. Here one uses special methods like factorizing out certain zero states and appealing to special results of V. Kac which unfortunately play no role in higher dimensional QFT. This always endangers the main physical motivation for doing these low-dimensional calculations, namely to develop a theoretical laboratory for getting some intuition about the conceptual framework of nonperturbative QFT (and not for the purpose of enriching mathematics). This was the motivation of Swieca and myself and also that of Mack and Luescher, who two years later understood the origin of the quantization of the cocycle constant c, the strength of the energy-momentum tensor two-point function. In [58] one finds an interesting historical account. In order to complete the historical review, I have attached a scanned version of my own paper which was published in a (hard to access) Brazilian journal as a second appendix.

The reason for this historical excursion in this section is not just historical piety or curiosity, but to point out that only now, with the arrival of the modular techniques, we have for the first time the means to approach the classification of chiral conformal QFT's with the help of generally valid concepts of QFT and in this way relate it more tightly with the rest of QFT. In this way one may hope to complete the old dream to use conformal QFT as a theoretical laboratory for the study of the principles which underlie nonperturbative QFT. In the case of conformal fields with braid group statistics, the idea was to compute the plektonic correlation functions in a similar spirit as in case of free Bosons/Fermions the algebraic (anti) commutation structure together with the causality and positivity of the energy spectrum determines the correlation functions (the Jost-Schroer theorem [18][91]). There one constructs charge-carrying free fields directly without the intervention of the representation theory of observable algebras. The guiding principle is that chiral conformal QFT's are like free fields in that the (plektonic) commutation relations and the internal symmetry determine the theory uniquely, i.e. chiral theories have no possibility of continuously varying coupling parameters. This turns out to be correct. In order to see this we must first adapt our modular localization approach to zero mass problems.

In the presence of zero mass fields, the previous wedge localization method is not suitable, because there is generally no in- Fock space from scattering theory which may serve as a reference space. Only double cones, whose modular theory is presently ill understood, would define compact localization regions where the interacting algebras would be unitarily equivalent to a free reference algebra of zero mass Fermions/Bosons. This observation has a natural relation to the vanishing of the LSZ limits (from the softening of the mass shell singularity through the infraparticle structure). If the theory involving zero mass is however conformal invariant, then the wedge region (via compactification) can be considered as a "double cone at infinity" or in the case of a half line or circle (chiral theories) as a special finite interval.

In plektonic chiral QFT one starts with the exchange algebra commutation relations in momentum space. The multiparticle spaces on which these operators act are represented by path spaces on a fusion graph for which the vertices are labeled by the possible nonabelian quantum charges. The n-particle Hilbert space is the direct sum over those charge sectors which can be reached by the nonabelian addition of n nonabelian fundamental charges which are represented by a path-space spanned by the iterated fusions:

$$H_{n} = \sum_{\rho(n)} H_{\rho(n)}$$

$$H_{\rho(n)} = \sum_{path(\rho(n),0)} \int f(\chi_{n},...,\chi_{1}) |(\chi_{n},\rho(n)),...,(\chi_{1},\rho(1)),0\rangle d\chi_{n}...d\chi_{1}$$
(6.96)

The $\chi's$ denote the exponential parametrization of the momenta on the light line of say the right line $p = e^{\chi} > 0$, whereas the $\rho(i)$ label the charge vertices on a fusion graph which one can reach by applying the fundamental charge i-times onto the vacuum. The plektonic statistics i.e. the R-matrices (see chapter 8.7) appears, if one compares two path which differ at one vertex. The path structure exactly agrees with that obtained by the iterated application of exchange algebra operators in momentum χ -space. The natural order is again $\chi_1 \geq ... \geq \chi_n$ and the other orders are obtained by the following commutation structure of annihilation and creation operators:

$$a_{\bar{e}_{1}}(\chi_{1})a_{e_{2}}^{*}(\chi_{2}) = \sum_{\substack{e_{1}',e_{2}'}} R_{\bar{e}_{1}e_{2}}^{e_{2}'\bar{e}_{1}'}a_{e_{2}'}^{*}(\chi_{2})a_{\bar{e}_{1}'}(\chi_{1}) + \delta(\chi_{1}-\chi_{2})\delta_{\bar{e}_{1}e_{2}} \quad (6.97)$$

$$a_{e_{1}}^{*}(\chi_{1})a_{e_{2}}^{*}(\chi_{2}) = \sum_{\substack{e_{1}'e_{2}'}} R_{e_{1}e_{2}}^{e_{2}'\bar{e}_{1}'}a_{e_{2}'}^{*}(\chi_{2})a_{e_{1}'}^{*}(\chi_{1})$$

Here the e's label charge edges which consists of a source charge, the charge c transferred by the creation operator and the range charge e = (r(e), c, s(e))

and \bar{e} is the cojugate charge edge which labels the annihilation operator. The new (non-Fock) feature is the source and range dependence. Related to this is the fact that the Fourier transforms are nonlocal fields in x-space. In order to obtain localized fields which fulfil the exchange algebra relations in x-space, we have to process these creation and annihilation operators through the modular localization machine for the half-line (the one-dimensional wedge). Despite the change of the n-particle spaces as compared to Fermions/Bosons, the discussion of modular localization is similar. The deviation of J from the geometrically defined J_0 is determined by the statistics operators which are piecewise constant in χ . For the Tomita involution we have the same structure as previously:

$$\check{S} = SJ_0\Delta^{\frac{1}{2}} \tag{6.98}$$

The mathematics of modular localization does not care whether S is (as in the Federbush model) a rapidity independent scattering matrix or a Klein factor or an R-matrix due to a change of statistics. The difference would only become visible if one compares the quantum opposite $\mathcal{A}(\mathbf{R}_+)'$ with the geometric opposite $\mathcal{A}(\mathbf{R}_-)$ where the latter must be constructed with a unita \mathbf{R}_+ ry geometric parity reflection. The first step constructs the \mathbf{R}_+ localized eigenspaces of \check{S} . Here one again uses the thermal KMS method. The generating semilocal operators are again non polynomial expressions in the $a^{\#'s}$ but yet without vacuum polarization. The last step from the semilocal polarization free auxilary operators to the local fields is as involved as in the integrable case¹⁶.

Although at the time of writing I had not yet finished the construction of the pointlike fields and their correlation functions, it is already evident that the two simplifications render the conformal problem much simpler than the construction of the massive factorizing models. The aim is to determine the theory in terms of its quantized statistics (R-matrix) structure and to obtain the FQS-quantization of the c-value as a consequence of the statistics rather than the other way around. We hope to present these conformal constructions in a separate paper.

Finally there is the question if the constant c_{\pm} in the energy-momentum commutation relation (or the central extension cocycle of the Virasoro algebra) have a more fundamental modular significance within the LQP setting based on observable nets. The Moebius symmetry group of the vacuum which allows a modular construction (in terms of modular half-sided inclusions) does not give any such information. The classical invariance group of an interval on the circle is a transformed dilation group. It acts fixed point free on the interior of the interval and has the two end points as its only fixed points. All these groups are modular groups of algebras localized in that interval. The generalization of this idea is a diffeomorphism of the circle with more than two fixed points. Of particular interest are those simple situations which are covering-equivalent to the old Moebius situation. In terms of circular coordinates $z = e^{i\varphi}$ these

 $^{^{16}}$ A more direct way of using the x-space exchange algebra is not possible, because in contradistinction to the momentum space situation the distributional character at coalescing points is not directly known from the algebraic structure.

diffeomorphisms are of the form:

$$z \to \sqrt{\frac{a+bz^2}{c+dz^2}}, \ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SU(1,1)$$
 (6.99)

The simplest illustration is in a somewhat symbolic short hand:

$$Dil(t)^{(2)} \equiv \sqrt{z} \circ Dil(t)^{(1)} \circ z^2$$
 (6.100)

Our notation for the half-circle is $z = e^{i\varphi}$, $0 \le \varphi \le \pi$, and $Dil(t)^{(1)}$ denotes the usual dilation $x \to e^t x$ rewritten into circular coordinates by stereographic projection. The complex poles and zeros of such transformations lie outside (poles) or. inside (zeros) the circle and may be joined by two cuts which do not cross the circle so that the map is well defined as a circular diffeomorphism. The point now is that the covering dilations have their own inner product in which they act unitarily:

$$(f,g)^{(2)} = \frac{1}{2} \int dx dy \frac{(1+x^2)(1+y^2)}{(x-y+i0)^2 (1+xy)^2} f(x)g(y)$$
(6.101)

The new inner product is only different from the old one by a multiplicative change of the measure. It is easy to verify that both inner products have the same symplectic form $\omega^{(1)}(f,g) = \omega^{(2)}(f,g) = \frac{1}{2} \int g(x) df(x) = \omega(f,g)$ hence belong to the same Weyl algebra

$$W(f)W(g) = e^{i\omega(f,g)}W(f+g)$$
 (6.102)

The SL(2,R) group acts unitarily in both spaces, but the action on the second space is different; it is the two-fold covering action

$$U^{(2)}(\alpha)f(x) = \Gamma f\left(\frac{a\frac{2x}{1-x^2}+b}{c\frac{2x}{1-x^2}+d}\right), \quad (\Gamma f)(x) \equiv f\left(-\frac{1}{x}+\sqrt{1+(\frac{1}{x})}\right) = 0$$

$$\alpha = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2,C)$$

In particular the above $Dil(t)^{(2)}$ is a special one parametric subgroup with 4 fixed points $(\mp)\infty, -1, 0, 1$ which acts as:

$$Dil(t)^{(2)}(f)(x) = f\left(-\frac{1-x^2}{2t^2x} + sign(x)\sqrt{1+\frac{1-x^2}{2t^2x}}\right)$$
(6.104)

The suspicion that this group is the Tomita-Takesaki modular group of a state on the two-fold algebra $\mathcal{A}[-\infty, -1] \vee \mathcal{A}[0, 1]$. $\subset Weyl(\mathbf{R})$, i.e. a von Neumann subalgebra of the the Weyl algebra on the line. The higher coverings lead to a similar situation. This can be confirmed by checking the KMS property for $Dil(t)^{(2)}$. We will defer all proofs to a forthcoming publication in collaboration of the author with H. W. Wiesbrock and limit ourselves to comment on some results.

- 1. The construction of the n-fold covering diffeomorphisms from the modular groups of algebras of n disjoint intervals in different positions follows the same pattern as the modular construction of the Moebius group.
- 2. The underlying states (associated with the above inner products) which lead to geometric modular situations are, with the exception of the vacuum, not cyclic with respect to the algebra of a single interval. This "defect" is related to the fact that a geometric situation for Haag duality for disjoint intervals overlooks the presence of higher superselection sectors as charge-anticharge operators in the commutant of disjoint interval algebras.

These are fascinating developments since the modular point of view liberates conformal QFT from its special dependence on structures (Virasoro-, affine-algebras) which have no place in the conceptual framework of general local quantum physics. By emphasizing multi-local algebras and their modular structure we have the chance to discover a whole world of "hidden" symmetries, i.e. symmetry transformations which unlike diffeomorphisms have no classical geometric aspect.

6.5 Comparison with Ideas from String Theory

One look at the content of theoretical publications under the heading of "high energy physics" reveals that what most physicist in this area are doing is very different from the content of these notes. A more detailed comparison shows that there are a few but perhaps significant concepts with an interesting overlap.. One idea which immediately comes to ones mind are certain structural properties which one affiliates with "light cone canonical quantization. This idea had a recent come back in relation with the so called M-theory, both in the sense of Matrix theory and M-theory proper [14]. The observation was that this quantization formally leads to simpler fields which resemble the nonrelativistic situation in that their application to the vacuum is free of particle-antiparticle polarization clouds. This of course has a resemblance with the structural properties of auxiliary fields which generate the wedge localized algebras. The relation becomes even more striking if on realizes that the transversal Galilei-group generators fit naturally into the subgroup G(8) of the Poincaré-group which arises in connection with the split into longitudinal, transverse and mixed generators. As with almost all standard quantization methods, the assumptions (canonical structure, functional Feynman-Kac representation) are not true properties of the results drawn from this assumptions, but rather play the role of a working hypothesis or better mental mark which suggests other structures which are true properties, despite the fact that the original assumptions become violated in the process of construction (e.g. through necessary renormalization repairs. There are some more interesting connections, and the best procedure seems to be to be to put these observations into a more systematic context.

The fastest way to understand the ideas underlying the present situation is to look at history of post Feynman QFT. The common cradle of all present frameworks of local quantum physics in general, and relativistic particle physics in particular, is certainly Feynman's renormalized perturbation theory and the ensuing understanding of QED. In the aftermath of this work, three schools of thought have emerged. Although their aim was the same, namely to abstract a nonperturbative framework (in order to incorporate strong interactions), the paths taken towards this goal were quite different. The first school of ideas which we will briefly refer to as Lagrangian field theory, took the covariant perturbative formalism of Lagrangian quantization as a starting point and interpreted it as a realization of an extension of the very successful quantization idea from quantum mechanics to the realm of infinite degrees of freedom. In this parallelism to classical physics, renormalization was understood (following a suggestion of Kramers) in analogy to the selfenergy problems of classical field theory at the beginning of this century. At that time Lorentz, Poincaré and others showed that these problems arise inevitably if one adds to classical field theory models of particles. Hence the occurrence of certain infinities was natural in a quantization approach and, apart from problems of mathematical consistency, their "dumping" into physical parameters was already partially anticipated in the classical theory. The Lagrangian quantization approach led to functional integral representations which, if it would not be for the necessity to perform infinite renormalization, could serve as nonperturbative definitions of physically relevant objects¹⁷. But since in local QFT, if analyzed from Wigner's particle picture, the one-particle aspects are already a consequence of the Poincaré covariance via the covariant fields which define the QFT, the infinities in a proper intrinsic formulation should never have appeared. With this remark we are already entering the territory of the second school : general QFT or in more recent terminology LQP. This school of thought, after some conceptual reformulations incorporated renormalized perturbation theory (see chapter 4) as a deformation on free fields but believed that a nonperturbative understanding would not come from generalizing Feynman's formalism but rather require a different conceptual framework where the physical intuition is mainly entering through local observables. The third school of thought, the S-matrix school, insisted that the nonperturbative understanding will come from an analysis of the scattering operator which is an important global observable. Although this approach in form of the bootstrap program came after 10 years to an end, certain of its notions survived up to this date in the form of effective actions and (as a result of its emphasis on crossing symmetry=duality) via Veneziano's S-matrix model in the present string theory. Through the constructive approach based on modular wedge localization as explained in this chapter, we also noticed that even in LQP the S-matrix has interesting modular aspects. But if one only

¹⁷In the lattice approach the discrete analogon to euclidean functional integrals is indeed taken as a definition and the renormalization is merged with the herculean task of controlling the scaling limit via second order phase transitions. This task was only accomplished in cases of existence of dynamical variables which are stable under scaling (e.g. the Lieb-Matthis-Schulz Ising model fermions).

considers the present sociological situation, the areas appear worlds apart. A comparison therefore is difficult and endangered by misunderstandings.

Let us first establish some rules for comparison. As we remarked above the quantization approach works with structures as canonical commutation relations, actions, functional integrals etc. which, although for themselves are almost never true properties of the would be theory, nevertheless lead to new properties which sometimes are at least perturbatively obeyed by the resulting theory. In fact the initial canonical assumptions tie the existence of the theory to a certain short distance behavior which even in renormalized perturbation theory is violated. To think about a cutoff does not help because a physically interpretable non Einstein-causal theory does not exist¹⁸. So in some way any QFT quantization approach has an artistic start. On the other hand all the properties on which LQP and in particular modular localization is based are true properties up to the very end. Hence the only reasonable way to compare LQP with Lagrangian QFT or string theory is to look at the results at the latter and not to pay attention to the words which enter the derivation. We first collect schematically those properties, results and conjectures of LQP which we want to compare. For a QFT with a mass gap with a complete scattering interpretation, one can derive the following facts (chapter 3 and 6).

• The modular theory of wedge algebras is geometric with the modular group Δ^{it} being the wedge associated Lorentz boost Λ_w of the incoming particles and the modular reflection J being related to that of the incoming free field theory J_{in} through the scattering operator (S-matrix) S_{scat} :

$$\Delta^{it} = U(\Lambda_w(2\pi t)), \quad J = S_{scat}J_{in} \tag{6.105}$$

The dense set of wedge localized state vectors can be represented in the form:

$$H_{loc}(W) = H_R(W) + iH_R(W)$$

$$H_R(W) \equiv \overline{\{\psi; S\psi = \psi\}}, S \equiv J\Delta^{\frac{1}{2}}$$
(6.106)

where $H_R(W)$ is the real closed subspace generated by the +1 eigenvectors of the antilinear unbounded Tomita operator which is involutive on its domain $S^2 = 1$. This brings the thermal Hawking Unruh aspects, which one usually relates with black holes, into ordinary QFT (chapter 3).

• As a standard reference wedge W_{stan} we may take the z-t wedge in which case we call z,t the longitudinal and x,y the transversal coordinates. This situation suggests to decompose the Poincaré group generators into longitudinal, transversal and mixed generators

$$P_{\pm} = \frac{1}{\sqrt{2}} (P_0 \pm P_z), \ M_{0z}; \ M_{12}, \ P_i; \ G_i^{(\pm)} = \frac{1}{\sqrt{2}} (M_{i0} \pm M_{iz}), \ i = 1, 2$$
(6.107)

¹⁸Such a cutoff would go against the modular localization ideas on which the only explicitely known nonperturbative results are based. One can regularize concrete Feynman integrals and even euclidean functional integrals, but those objects cannot be related to a real time physically interpretable (with time dependent scattering) QFT.

The generators $G_i^{(\pm)}$ are precisely the "translational" pieces of the euclidean stability groups $E^{(\pm)}(2)$ of the two light vectors $e^{(\pm)} = (1, 0, 0, \pm 1)$ which appeared for the first time in Wigner's representation theory for zero mass particles. More recently these "translations" inside the homogenous Lorentz group appeared in the structural analysis of "Modular Intersections" of two wedges [8]. Its role is analogous to that of the true translations P_{\pm} with respect to halfsided "Modular Inclusions" [8]

1. As one reads off from the C.R., $P_i, G_i^{(+)}, P_{\pm}$ have the interpretation of a central extension of a transversal "Galilei group"¹⁹ with the two "translations" $G_i^{(+)}$ representing the Galilei generators, P_+ the central "mass" and P_- the "nonrelativistic Hamiltonian". The longitudinal boost M_{0z} scales the Galilei generators $G_i^{(+)}$ and the "mass" P_+ . Geometrically the $G_i^{(+)}$ change the standard wedge (it tilts the logitudinal plane) and the corresponding finite transformations generate a family of wedges whose envelope is the halfspace $x_- \geq 0$. The Galilei group together with the boost M_{0z} generate an 8-parametric subgroup $G^{(+)}(8)$ of the 10-parametric Poincaré group:

$$G^{(+)}(8): P_{\pm}, M_{0z}; M_{12}, P_i; G_i^{(+)}$$
 (6.108)

The modular reflection J transforms this group into an isomorphic $G^{(-)}(8)$. All observation have interesting generalizations to the conformal group in massless theories in which case the associated natural space-time region is the double cone.

• The position of the subspace $H_R(W)$ within the incoming Fock space allows to define a modular Møller operator U(W) which intertwines the wedge affiliated Tomita involution S with that of the corresponding incoming involution:

$$SU(W) = U(W)S_{in} \tag{6.109}$$

and leaves the vacuum unchanged. The interacting wedge algebra $\mathcal{A}(W)$, which together with the vacuum vector has Δ^{it} and J as its modular data, is defined in terms of $\mathcal{A}_{in}(W)$ as:

$$\mathcal{A}(W) \equiv U(W)\mathcal{A}_{in}(W)U^*(W) \tag{6.110}$$

The Haag's theorem prevents the existence of analogous intertwining unitaries for the type I equal time canonical algebras:

$$A_{can} \equiv \cap_{\varepsilon} A(\varepsilon) \tag{6.111}$$

which in the algebraic approach are represented as the intersection of time slice algebras of thickness ε . This leads to the nonexistence of the interaction picture in local quantum physics and the necessity of the artificial

¹⁹This G's are only Galileian in the transverse sense; they tilt the wedge so that one of the light like directions is maintained but the longitudinal plane changes.

infinite volume limiting procedure involving a quantization box (which is unfortunately not related to modular localization). The above intertwining relation between the unique hyperfinite type III₁ algebras (all localized subalgebras in local quantum physics which have a nontrivial causal complement are of this kind) is protected against such No Go theorems. The existence of the modular Møller operator [113]U(W) (in QFT it does not seem to be possible to define a "scattering" Møller operator) leads to the existence of generators of $\mathcal{A}(W)$ which are localized in $\mathcal{A}(W)$ but allow no smaller localization inside W i.e. they are nonlocal inside W. They are "on shell" i.e. contain a negative frequency part which annihilates the vacuum. Formally they are given as $A_W(x) = U(x)U(W)A_0(0)U^*(W)U^*(x)$. It is essentially the absence of vacuum polarization pairs i.e. the mass shell support of their Fourier transform which makes these generators²⁰ of $\mathcal{A}(W)$ extremely helpful. In factorizable d=1+1 theories, the positive and negative frequency components of these semiglobal operators fulfill the Zamolodchikov Faddeev algebra [112]. Be aware that the x in the Utransformed fields has nothing to do with localization around x inside W. Rather localization has to be constructed via smaller algebras defined by intersecting wedge algebras:

$$\mathcal{A}(\mathcal{O}) = \bigcap_{W \supset \mathcal{O}} \mathcal{A}(W)$$

$$\mathcal{A}(W) = U(\Lambda)\mathcal{A}(W_{stand})U^*(\Lambda)$$

$$W = \Lambda \cdot W_{stand}$$
(6.112)

where W_{stand} is the standard x-t wedge and the net of W's is generated from W_{stand} by Poincaré transformations. It is very important for the interacting case to realize that U(W) depends on W i.e. $U(W_{stand})$ does not commute with the space-time transformations U(L) except with the W-associated Lorentz boost. It should be clear that ideas about how to construct such U's should not be viewed in the setting of the perturbative split $H = H_0 + H_{int}$ (the free incoming situation does not correspond to H_0). Whereas the representations of the Poincaré group of the interacting situation agrees with that of the incoming fields, this is not so for the unperturbed theory belonging to H_0 . In fact the latter theory does not even live in the same Hilbert space (only its local folium of states agrees with that of the interacting theory).

• The Møller operator U(W) can be explicitly computed for d=1+1 factorizable models and it is intimately related to the Riemann-Hilbert properties of the modular localized real subspace $H_R(W)$ (chapter 6). The mass shell components of the nonlocal generators $A_W(x)$ turn out to satisfy

 $^{^{20}}$ Strictly speaking these polarization free operators are only well-defined objects in the thermal Hilbert space associated with the modular wedge localization. However they are very valuable as a basis for the pointlike local fields which allow for a nonsingular extension outside the wedge [112].

[112] the Zamolodchikov-Faddeev algebra and the modular localization property defines a thermal KMS state on this algebra.

- The rich physical structure emerging from inclusions and intersections of local algebras in the net. Algebraic QFT interprets the external (space-time) and internal global symmetries from how one algebra is positioned with respect to another one (see mathematical appendix):
 - 1. "Shallow" inclusions. These are inclusions $\mathcal{N} \subset \mathcal{M}$ which posses a (noncommutative) conditional expectation i.e. $\mathcal{N} = E(\mathcal{M})$ of the kind studied by Vaughn Jones. The local endomorphism of algebraic QFT of the DHR theory and its extension to low dimensional QFT are typical examples. The relation between the algebraic QFT endomorphism and the Jones setting is well understood. Shallow inclusions are related to inner symmetries²¹. For d=1+3 dimensional theories this leads to Fermi-Bose Statistics and compact internal group symmetry whereas for low dimensional theories this yields the class of (physically) admissable unitary braid group representations through Markov traces on the braid group B_{∞} .
 - 2. "Deep" inclusions. By this one means inclusions which have no conditional expectation but obey a modular restriction which lead to space-time symmetries. There are two "modular inclusions" whose geometric consequences have been studied: halfsided modular inclusions and modular intersections. The first case is illustrated by two touching wedges, a situation resulting from a light like shift of a wedge into itself. A halfsided modular inclusion leads to noncompact group isomorphic to the longitudinal (d=1+1) Poincaré group. The second case of modular intersections is illustrated by two wedges which have one light ray in common. In that case the intersection data lead among other things to the above Galilei generators. The full Poincaré group is obtained by the relative "modular position" of a finite number of algebras (the minimal number depends on the space-time dimensionality). In this way one may generate the whole net from a finite "modular skeleton net".

Some comments are in order.

Let us add to these rigorous results two conjectures which are important for the later comparison. These conjectures are related to the Hawking-Unruh issue of horizon physics of quantum matter in black hole solutions. Even though our comments are only conjectures, we will try to stick to the conceptual rigor of the rest of this article. For this reason we will not use the word "quantum

 $^{^{21}}$ Note however that in low-dimensional QFT's there is no sharp distinction between inner and spacetime symmetries because the charge structure relates to the spatial covering aspects.
gravity" in this discussion and emphasize the fact that the physical origin of the Hawking temperature is the modular localization in Minkowski- or curvedspace-time and not primarily a black hole horizon. The latter mainly plays the role of enforcing a natural localization (by creating a bifurcated horizon via the e.g. black hole metric) and constitutes a special case of the former. The notion of a bifurcated horizon through modular localization is more abstract, since it is not related to metric Killing vectors but rather to isometries in the space of wave functions or the underlying Hilbert space of QFT [112]. It nevertheless leads to the same physical consequences of thermal behavior and Hawking radiation. For this reason the main concepts which are usually attributed to gravitation theory can be perfectly understood in terms of thermality through localization (instead of the standard heat bath thermal behavior). The main difference of the two thermalization concepts can be traced back to that of the two sided spectrum of the modular localization operators (e.g. the Lorentz boost) versus the one sided spectrum of the heat bath Hamiltonian which leads to the boundedness of $e^{-\beta H}$.

This raises the question whether modular localization also leads to a fundamental algebraic notion of entropy. Here it is helpful to mention the "degrees of freedom" counting in local quantum physics which deviates in an interesting and significant fashion from that in e.g. Schrödinger quantum mechanics [3]. In the latter case one learns, that the phase space cells (standard notion of localization and momentum restriction) leads to a finite number of degrees of freedom per $2\pi\hbar$ size phase space cell. The first attempt in algebraic QFT by Haag and Swieca led to the notion of compactness. Later this notion was sharpened to the "nuclearity criterion" of Buchholz and Wichmann [3] which does not use a sharp cutoff in Hamiltonian- or. in the "modular"-energy but rather an exponential fall off. Contrary to the nonrelativistic case and to popular opinion, the relativistic localization concept (as opposed to the standard box quantization) together with the finite energy projection does not lead to a finite number of quantum states ("bits") but rather only to a compact (Haag-Swieca) or nuclear (Buchholz-Wichmann) set. A computation for free fields reveals that this behavior is optimal i.e. local quantum physics cannot reproduce the finite degrees of freedom behavior of quantum mechanics, but comes pretty close:

$$H - S : \{ P_E A \Omega \mid A \in \mathcal{A}(\mathcal{O}), \|A\| \le 1 \} = compact set \quad (6.113)$$

$$B - W : e^{-\beta H} \mathcal{A}(\mathcal{O}) = nuclear set$$

The interrelation between these slightly different forms of relativistic "local degree's of freedom counting" has been discussed in [3]. This property forbids infinite towers of particles (as they occur e.g. in genus ≤ 1 string perturbation) and an associated limiting Hagedorn temperatures. The most valuable consequence is a very profound interpretation of the ancient issue of Heisenberg-Weisskopf vacuum fluctuations: if a spatial volume is not interpreted as a quantization box, but rather as a region for localization of a partial charge via a conserved current (say inside e.g. an already defined Minkowski space free field theory), then the vacuum fluctuation near the boundary are infinitely large. In order to control them, it is necessary to allow a smooth transition to zero charge density inside a "collar" around the localization region. The "split property" of algebraic QFT [3], which is a consequence of the above "nuclearity" property of degree of freedom counting, just provides the mathematical precision for this intuitive idea²²:

$\exists type \ I \ factor \ \mathcal{N} \ s.t. \ \mathcal{A}(\mathcal{O}_1) \subset \mathcal{N} \subset \mathcal{A}(\mathcal{O}_2) \tag{6.114}$

Here one should imagine two concentric double cones \mathcal{O}_i with their associated hyperfinite III₁ factors. The type I factor \mathcal{N} has a "fuzzy" localization inside the bigger double cone, and it is just this fuzziness which allows the definition of partial charges without infinite vacuum fluctuation and with a clear-cut split between the physics "inside and outside" [3]. Needless to add that the algebras underlying QM are always of type I, whereas the relativistic causality and associated localization structure always lead to hyperfinite III₁ factors at least if the regions allow for a nontrivial causal complement. So in order to find quantum mechanical structures inside local quantum physics, one needs type I factors inside local hyperfinite III₁ factors. The split property gives also a preferred candidate [114] for such an interpolating type I factor \mathcal{N} . The scenario for a definition of entropy from first principles is in terms of the modular group of \mathcal{N} . As a consequence of type I this modular group is inner, i.e. there exists a "Hamiltonian" described by a hermitian operator K associated to \mathcal{N} (this never happens for the $\mathcal{A}(\mathcal{O})$ factors).

The issue of entropy is then closely related to the problem of the modular Hamiltonian K of \mathcal{N} which in turn is determined in terms of the modular objects of the split data [3]:

$$\Delta_{\mathcal{O}_1}^{it}, \Delta_{\mathcal{O}_2}^{it}, \Delta_{\mathcal{O}_1' \cap \mathcal{O}_2}^{it}; J_{\mathcal{O}_1}, J_{\mathcal{O}_2}, J_{\mathcal{O}_1' \cap \mathcal{O}_2}$$
(6.115)

In zero mass conformal theories the double cone J and Δ^{it} relative to the massless vacuum are geometric transformations inside the full (including reflections) conformal group [3], in particular Δ^{it} transforms the r,t coordinates but not the two transversal angular variables of the double cone. On the other hand the massive double cone theory can be incorporated into the same Hilbert space or more precisely, the massless and the massive. This suggests that the modular object of the massive situation are nonlocal deformations of the conformal massless split situation. One expects that the K-Hamiltonian is well enough in order to allow for the existence of the von Neumann entropy:

$$S = -trDlnD \tag{6.116}$$

where $D = \frac{1}{tre^{-K}}e^{-K}$ is the density matrix defined in terms of the modular Hamiltonian K. This is a quantity which depends on the size of the collar ε and which diverges as $\varepsilon \to 0$ i.e. when the fuzzy type I factor becomes hyperfinite

 $^{^{22}}$ The idea to define entropy with the help of the split property is due to Heide Narnhofer (this was pointed out to me by H-W. Wiesbrock) [11]. It constitutes one important element in a joint unfinished project of Wiesbrock with the present author.

type III₁. If the result of the existing proposals [13] is compatible with this idea, we should expect a universal logarithmic divergence in the inverse size ε^{-1} of the collar which controls the vacuum fluctuations:

$$S = -trDlnD \sim Cln\varepsilon^{-1} \tag{6.117}$$

with C related to the longitudinal 2-dim. conformal theory which according to our previous discussion we expect to determine the geometric core of the fuzzy modular group of the double cone algebra $\mathcal{A}(\mathcal{O})$. Indeed the formula [13]

$$C \sim Area\sqrt{c}$$
 (6.118)

where Area denotes the area of the double cone and c the vacuum fluctuation strength of the energy momentum tensor. Although the limiting entropy is certainly infinite, we have not yet been able to confirm that this infinity is universal and behaves exactly as argued by Larsen and Wilszek.

In order to prove this one must do some new computations on the "localizing map" (math. appendix) which is the most convenient way to compute the distinguished type I factor [3]. The relevant degrees of freedom would "live", as we will argue later, inside the collar and the ratios of this "collar entropy" stay finite for vanishing collar size. This remains a fascinating program for the future.

Now we are able to formulate our two conjectures:

Conjecture 25 The modular group of the (nonconformal) massive double cone algebra $\mathcal{A}(\mathcal{O})$ with respect to the massive vacuum vector (i.e. the physical vacuum state restricted to $\mathcal{A}(\mathcal{O})$ is cocycle-related to the known geometric modular group of the associated conformally invariant situation belonging to the pair $(A(O), \omega_{m=0})$ where $\omega_{m=0}$ denotes the conformal invariant vacuum state. For the equivalence of the massive with the massless algebra one may either invoke the construction of the double cone algebra by canonical quantization or the fact that local algebras are always hyperfinite III₁ factors and the latter is unique modulo unitary equivalence. The cocycle accounts for the difference in the local propagation of massless (Huygens principle) and massive theories and its presence renders the action of the modular group "fuzzy". Only asymptotically near the horizon i.e. the boundary of the double cone, the fuzzyness decreases and the geometric conformal modular transformation reappears. Although a single algebra $\mathcal{A}(\mathcal{O})$ of the massive theory and its scale invariant limit may be identified, the two nets inside \mathcal{O} remain different. However the conjecture that the difference is due to the different propagation suggests that the massive net inside \mathcal{O} may be obtained from the massless $\mathcal{A}(\mathcal{O})$ by adjoining the action of the Poincaré covariances inside O.

Remark 5 For a massive free Fermi field²³ in d=1+1 this can be shown. One notes that the restriction of such a free massive theory to the light rays which

 $^{^{23}\}mathrm{We}$ want to avoid the infrared problems of massless Bose fields

constitute the boundary of the d=1+1 double cone is simply the restriction of the corresponding massless theory and that by propagating the chiral conformal data on the one dimensional horizon inside with the massive propagator, one regains the massive free field net inside O. A general proof of this reduction of a d=1+1 situation to its chiral conformal limit (+ possible covariance operators) would be extremely desirable because it would explain the association of the degree of freedoms of d=1+1 theories with the horizon.

Conjecture 26 The double cone algebras $\mathcal{A}(\mathcal{O})$ are identical to any of the twodimensional double cone algebras $\mathcal{A}(O^{(2)})$ obtained by cutting the double cone by a two-dimensional plane which contains the t-axis and one coordinate axis. The net inside $O^{(2)}$ may be obtained from the associated chiral conformal net on the one-dimensional horizon and a local representation of Poincaré covariances and this holographic modular picture is the local quantum physical property behind the Bekenstein-Hawking quasiclassical entropical observations.

Remark 6 The first part is actually a consequence of Haag duality and the fact that the causal completion of $\mathcal{O}^{(2)}$ gives \mathcal{O} :

$$\mathcal{A}(\mathcal{O}^{(2)}) \equiv \cap_{\delta} \mathcal{A}(\mathcal{O}_{\delta}) \tag{6.119}$$

Where \mathcal{O}_{δ} denotes the middle slice of thickness δ by cutting the double cone parallel to the t-axis. Each \mathcal{O}_{δ} has \mathcal{O} as its causal completion and the property of "Haag Duality" demands the equality of $A(O_{\delta})$ with the algebra of the causal completion $\mathcal{A}(\mathcal{O})$. The essential step in the holographic reduction is the appearance of chiral conformal degrees of freedom after removal of the angular degrees of freedom due to angular symmetry (substituting the transversal symmetry in the case of the wedge). The envisaged entropy is therefore not proportional to area $(\mathcal{O}^{(2)})$ × angular volume but rather to horizon-length $(\mathcal{O}^{(2)})$ × angular volume = volume of $horizon(\mathcal{O})$. Actually such a situation would also suggest that there may be an infinite hidden nongeometric (fuzzy) symmetry algebras in the nonperurbative structure of any QFT^{24} . Although they are local in the sense of keeping things inside say \mathcal{O} , their action within \mathcal{O} is totally fuzzy. Such symmetries of nonperturbative local quantum physics would escape differential geometric methods. Note that the two conjectures cannot even be formulated in terms of properties of expectation values of fields; the use of the algebraic *i.e.* field coordinate independent concepts is indispensable for the formulation. If algebraic QFT did not already exist, one would have to invent it in order to understand the above thermal and entropic properties.

Summing up our excursion on nonperturbative LQP we would like to stress again that the algebraic method allows for a completely intrinsic definition and

 $^{^{24}}$ In principle every modular automorphism has the interpretation of a (localized but fuzzy) physical symmetry. Some of these are "semi-geometric" i.e. they act geometric on subnets. The previous modular intersection situation which led to the transversal Galilei-transformation is such a case of a "semi-hidden" symmetry [15].

understanding of QFT, independent of its Lagrangian or non Lagrangian origin. Any quantum theory which fulfills the stability requirements of positive energy and allows for a net interpretation and the associated localization concepts is a QFT par excellence and enjoys all the general structural properties of these notes as TCP, spin & statistics, crossing symmetry & modular localization & thermality, wedge-localized fields without vacuum polarization, hidden modular symmetries, Haag duality (an abstract form of the 2-d Kramers-Wannier-Kadanoff Duality), nuclearity for the phase space degrees of freedom & the conjectured "Holographic Entropy" and all the other yet known properties of nonperturbative local quantum physics. The main obstacle against progress is not so much the novel mathematics which these new physical concepts require, but rather (as always in the past) prejudices. One prejudice is that field theory has to be "Lagrangian". In view of the many existing low-dimensional non-Lagrangian models and the fact that they hardly rocked the Lagrangian boat up to now (they are simply ignored, because their construction does not fit a quantization approach), this appears to be the mightiest prejudice.

Before we mention some recent results of string theory obtained with the help of light cone quantization and its discretization as well as finite truncation, we take a rapid look at its history since this allows a clearer conceptual understanding than the somewhat confused picture obtained from only looking at the recent hep-theory publications.

A good starting point is the dispersion theory which was the main nonperturbative attempt of the 50 and 60 to go beyond the Feynman approach. The main issue was to find sufficiently many "on-shell" properties of QFT such that an S-matrix theory or at least a phenomenologically successful scheme could emerge. Besides the obvious properties like unitarity and certain analytic properties, the on-shell property which apparently was most intimately and deeply related to the off-shell causality principle of QFT was crossing symmetry. The complete nonperturbative understanding of crossing symmetry including the requires on shell analytic behaviour and the precise relation to TCP invariance and causality was never achieved. Therefore it was considered a major achievement when Veneziano succeeded to construct a S-matrix model which fulfilled crossing symmetry exactly and allowed for a systematic unitarization which maintained crossing symmetry in each step and in some sense was reminiscent of the perturbative systematics, although it had little in common with ordinary perturbation theory. Later it was realized that the quantum mechanics of strings can successfully describe this model and its unitarization. The infinite tower of particles which, if they would remain stable under unitarization, would violate the principles of local quantum physics (the aforementioned degree of freedom behavior, leading to a Hagedorn temperature) could, as in Feynman's perturbation theory, become unstable particles i.e. poles in the second Riemann sheet of the S-matrix and in this way the model could be perfectly consistent with nonperturbative QFT. This was at least what I and many of my contemporary QFT colleagues thought when we got used to those nice pictures involving Regge trajectories.

But, as everybody knows, things happened differently. Instead we had to wit-

ness the "Bartholomew night massacre"²⁵ (a bit poetic, but not completely unrealistic since part of the story really happened in Paris [115]), also often referred to as the first string revolution, in which the old string theory, which served as a laboratory of certain aspects of nonperturbative QFT (notably strong interactions), was killed and the mathematical formalism (first without any change) was pushed upward in energy by more than 15 orders of magnitude and physically outed as "quantum gravity". Only later string theory obtained the modern differential geometric wrapping, which partially expressed the increasing mathematical sophistication of the string community. In fact in this modern fashion it became an impressive source of mathematical innovations. It was precisely the distance from any kind of laboratory physics, which protected these developments from usual fate of theories whose relevant energy scales stays close to the experimentally accessible region. Because of the involved fantastic "scale sliding" and its innovative differential geometric content, it is often referred to as the "second string revolution".

From the point of view of exhausting the scenarios offered by Lagrangian quantization, the sliding up the energy scale was very logical indeed. It is legitimate and even useful to stretch a framework (as the Lagrangian quantization, canonical or functional), which was so successful as Feynman's renormalized perturbation theory, to its physical limits set by the Planck scale. A successful formalism in the history of physics was always pushed to its limits. Such extensions only become somewhat counterproductive if, as a result of apparent lack of alternatives, one identifies its consequences (as the "big desert" region beyond the present laboratory energies up to the Planck mass, or as the omnipotence of supersymmetry in its underlying differential geometric mathematical formalism) with what should be expected from nature. The very existence of alternative concepts as in LQP which do not lead to such consequences shows that one is exploring the limits of quantization approaches rather than of local quantum physics. Let us first look at light cone quantization since most recent papers on string theory use this structure. We briefly remind the reader about its meaning in the simplest Lagrangian QFT context.

In analogy to canonical quantization but different from the discussion of the bifurcated wedge situation one distinguishes one x-t plane as the canonical quantization plane and the other for the definition of a propagating time. In the Lagrangian setting we have for a selfinteraction of a scalar field:

$$L = \int dx^{-} dx^{\perp} \left\{ \partial_{+} \varphi \partial_{-} \varphi - \frac{1}{2} (\partial_{\perp} \varphi)^{2} - \mathcal{L}_{int} \right\}$$

where $x^{\pm} = \frac{1}{\sqrt{2}}(t \pm x)$, $x^{\perp} =$ transversal coordinates. Using longitudinal mo-

²⁵The name some of my QFT collegues attributed to this event expressed their bewilderment after being told to forget the nice Regge trajectory interpretation of Veneziano's dual crossing symmetric S-matrix model which they already had gotten used to (as a proposal for structure of a strongly interacting S-matrix which could originate from a nonperturbative QFT).

menta k_{-} , the "Hamiltonian" becomes:

$$H = \int dk_{-} dx^{\perp} \left\{ \frac{\partial_{\perp} \varphi^{*}(k_{-}, x^{\perp}) \partial_{\perp} \varphi(k_{-}, x^{\perp})}{2k_{-}} + \mathcal{H}_{int} \right\}$$

The main difference to standard canonical quantization is the absence of vacuum polarization in the fields $\varphi(k_-, x^{\perp})$ [14]. The prize for this apparent simplification is a somewhat hidden nonlocal interpretation of these auxiliary fields which went unnoticed by the authors of (M)atrix theory²⁶. The light cone theory in the context of the Unruh Hawking mechanism was studied by Sewell [9]. His setting only applies to free field theories and to those interacting fields which have the same short distance behavior as e. g. φ^4 in d=1+1. There is one essential difference to the standard canonical commutation relations. The canonical rules in light cone coordinates are [9]:

$$[\varphi(x_+=0,\partial_-f),\varphi(x_+=0,\partial_-g)] = \int f\partial_-g dx_- dx_\perp$$
(6.120)

$$\varphi(x_{+},h) = \int \varphi(x_{+},x_{-},x_{\perp})h(x_{-},x_{\perp})dx_{-}dx_{\perp}, \ h \in \mathcal{S}(R^{3})_{0} \ (6.121)$$

where the subscript 0 denotes Schwartz functions with vanishing integral over all space $\int h = 0$. Hence the infrared behavior of light cone fields is analoge to that of zero mass fields in d=1+1. The characteristic initial value problem is easily shown to be uniquely soluble if the initial data are in $\mathcal{S}(R^3)_0$, and the causality condition which leads to localizability is easily reexpressed in terms of the light cone description. There are two caveats in the interacting case. On the one hand the restriction to a light front has the same limitations as the standard canonical formalism: any theory with stronger than free field short distance singularities does not permit such a restriction. This only leaves the superrenormalizable $P\varphi_2$ interactions. Here one could be open minded and argue that the light cone idea should not be taken literal, but that the important message is that there are field coordinates without vacuum polarizations [14]. However such degrees of freedom are necessarily nonlocal, i.e. the interaction of an originally local theory causes the appearance of nonlocal light cone degrees of freedom (even for the $P\varphi_2$. theories). There is nothing wrong with the use of nonlocal field coordinates in a local theory as long as one knows the relation to the local fields. Unfortunately there is no mentioning of this important problem in the literature on light cone quantization. Whereas these short-comings may be viewed of a technical nature (as most of the starting assumptions in the standard approach they must be repaired in the course of calculations), the third one is more physical and hence more serious. The light cone approach does not reveal how the interacting nonlocal polarization-free light cone fields are related to the local fields. The knowledge of this relation is crucially important for the interpretation of the light cone approach.

 $^{^{26}{\}rm The}$ LCQ field variables are very nonlocal with respect to the usual local canonical fields. Only in this way these degrees of freedom manage to surpress vacuum fluctuations.

The suggestion from our modular localization approach would be that the light front situation is closely related with the wedge; the horizon of the wedge consists of two light fronts. In this way the precise relation between wedge localized fields without vacuum polarization and local pointlike fields will be clear. In fact the division into wedge-affiliated longitudinal and transversal degrees of freedom led us to the existence of the 8-parametric "Galileian extension" $G^{(+)}(8)$ of the longitudinal modular group. Interestingly enough, the Galilean subgroup of $G^{(+)}(8)$ constitutes the starting point of the BFSS [116] light cone quantization framework. The standard obstacle against a quantum mechanical description, namely the presence of vacuum polarization, has been taken care of if one uses these semiloval fields and this is perhaps the main reason why these variables and their QM discretization is relevant for the exploration of ideas on M-theory. Here the terminology QM should be understood in sufficient generality. Galilei-invariant field theories without vacuum polarization, but with rich channel couplings between different multiparticle sectors (as the T.D. Lee model , just to mention one) can a priori not be excluded. In any case the analogy of the polarization cloud free state vectors $\varphi_{light-cone}\Omega$ with the wedge thermal space affiliated vectors $A_W \Omega$ of this chapter is very startling.

By compactification of the light cone time x^{-} one formally obtains a discretization of LCQ called DLCQ. As with lattice discretization, DLCQ allow a matrix approximation which, following BFSS, posses a natural interpretation in the modern string setting. In our attempt to translate these situations into the conceptual framework of algebraic QFT, we would draw the analogy to the compact double cone situation $\mathcal{A}(\mathcal{O})$ which is the modular substitute for the box quantization in QM. This is the spacetime localization by which one must substitute the wedge region if there is no LSZ scattering theory as a result of the presence of infrared photon clouds. In that case the electrically charged "infraparticles" have vanishing LSZ limits and one only can work with a Fock reference space for compact modular localization regions. The claimed proximity of the light cone quantization idea and string- and M-theory begs the question whether the latter theories are local in the presence of interactions. If they are, than string theory is a special case of LQP and all the algebraic results can be taken over. I expect that they are not local, i.e. that one is not dealing with nonlocal variables in a local theory, but the string framework itself is nonlocal. In QFT it is precisely the locality structure which elevates it from a bunch of prescriptions to a complete theory in which e.g. the particle- field dichotomy, the (time dependent) scattering theory and other pivotal aspects of the interpretation follow from the theory itself and need not to be imposed from the outside in form of recipes. The string idea is not a theory in this sense.

There is another more speculative issue of potential agreement. It is the issue of nonperturbative thermal and entropical behavior. LQP attributes a thermal aspect to all local algebras in particular if the boundary of the localization region defines a classical (Killing) horizon as in the case of black holes. In fact this phenomenon was discovered in a pure classical manner via the notion of a classical entropy and the use of thermodynamic relations. Attempts to interprete this classical entropy in the quantum sense led to the conjecture of holographic behavior of degrees of freedom for such situations. Although there exists by now a fundamental general understanding of the relation between horizon (with or without Killing vectors), the direct understanding of entropy in LQP is still shaky, despite our previous arguments concerning holographic behavior. The situation is not any better in M-theory with light cone quantization. The only thing which one can presently say is that if those speculations turn out to be correct, than also the entropical properties have nothing to do with CST classical Killing vectors but are general properties of LQP. Since in that case the definition of entropy required a good phase space behavior ("nuclearity") of QFT, it is interesting to ask what string people say on this subject. Their counting is that of a (nonrelativistic) spatial box quantization i.e. the standard finite number per unit of phase space volume. LQP would challenge the physical relevance of this concept since the correct localization in LQP is the modular space-time localization and not the box localization which in the relativistic context is the same as the unphysical Newton-Wigner localization. As we have emphasized, the nuclearity property (which even for free fields cannot be improved) assigns an infinite albeit "small" number (see mathematical appendix).

An even more basic discrepancy is the role of supersymmetry which in the quantization approach is used to tame perturbative short distance behavior and in this way to prevent getting too far into the quantum domain. It is an essential symmetry for many recent results. On the other hand LQP does not ask for such a symmetry and even questions its physical significance. One reason is, as we already mentioned on several occasions that it is a completely accidental symmetry which collapses under a contact with a heat bath. Here accidental means that it plays no role in the understanding of charge sectors, e.g. all known 2-dimensional soluble (tricritical Ising etc.) models can be solved without using SUSY although it is not wrong to use it. It also plays no role in the modular interpretation of space-time symmetries. It is true that in low dimensions the charge sector structure gets intermingled with the covering aspects of spacetime, but SUSY has no part in this nontrivial "marriage" which among other things is responsible for the existence of 3-mf invariants. The mentioned thermal collapse is what one expects of a accidental symmetry, but of course not of a bona fide inner or space-time symmetry, e.g. the Lorentz symmetry only suffers the usual spontaneous breakdown and not a collapse. If one is conservative in one's judgement one may say that the physics of SUSY (if there is any physics) has nothing to do with any known concept of symmetry and its possible meaning may be outside of LQP. But such a mystification for maintaining a property which entered physics in a completely formal way (and afterwards changed the motivation for its original introduction), is probably going a bit too far. Saying such things one is of course easily accused of ignorance about the marvelous achievements by supersymmetric n=4 gauge theories which are d=1+3 conformal invariant field theories. This if true, is indeed somewhat sensational and it would be nice to see a physical gauge invariant correlation function (of necessarily composite fields) even in lowest order perturbation theory, because then one could study the thermal stability problems and understand to what degree SUSY was necessary to find such a remarkable situation. Unfortunately despite the 15 year existence no such result is available (certainly not because of lack of physical interest).

The biggest discrepancy between LQP and string theory however lies in the physical philosophy of why one introduces a certain structure. In LQP this is done because one wants to extend the range of a principle. For example the introduction of extended operators which have a more general localization than that of pointlike fields or compactly localized operators would not be done for its own sake. The physical (Bohr-Heisenberg-Wigner) esthetics is in the introduction of those structures which are necessary to generalize the range of applications of (existing) principles, in this case the causality or locality principle. For theories with a mass gap in their positive energy spectrum, the causality principle allows for charge carriers with semiinfinite spacelike string localization and therefore they are introduced and not because one wants to investigate string-like structures in their own right. In fact in d=1+2 one finds that these objects must obey braid group statistics; if one limits the carriers to be compactly localized one falls back onto Fermions and Bosons and misses an important physical realization of the same principle. The esthetics underlying string theory and more generally any quantization approach is Dirac's (differential) geometric esthetics which in the modern mathematical setting favors geometrically natural generalizations. Gedankenexperiments which could explain notions like stringiness versus QFT-ness play no role. What if the infinite tower of particles in string theory converts in higher order in the genus as in Feynman's perturbation into resonances in the second Riemann sheet of the S-matrix; why is such a situation not compatible with LQP? Does stringiness meant in the above sense of LQP charge carriers with noncompact localization? What is the concept of localization, if any, in string theory? What mathematical terms (correlation functions, algebras, states) should one use for the characterization of string theory and what are the intrinsic properties and related principles which allow an intrinsic characterization independent of the prescriptions and recipes which went into its construction? More questions than answers.

Let us add some concluding remarks. Although a *direct* comparison of nonperturbative QFT with string theory is presently not possible as a result of the lack of an intrinsic physical definition, we tried an indirect evaluation based on what may be called "circumstantial evidence". As a geometrically based approach, string theory is strongest in the quasiclassical domain. It need classical Killing horizons in order to perceive thermal and entropical nonperturbative properties. Its similarity to classical Kaluza-Klein ideas and its selection of high dimensions (26, 10, 11 etc.) probably has the same explanation: what in low dimensions remains genuinely nonperturbative quantum, becomes more classical in higher dimensions and hence more easily detectable by the string formalism. It is only logical that it puts more confidence in differential geometric (and since recently also noncommutative geometric) Diracian structure than in the conceptual physical Bohr-Heisenberg mode of thinking.

Chapter 7

Introduction to Algebraic QFT

7.1 Some Useful Theorems

Algebraic QFT requires more mathematical rigor and physical depth than QM or even than standard perturbative QFT. In the latter case one has already achieved a reasonable intuitive understanding whereas in the d=1+3 nonperturbative QFT we only have untested scenarios. So one tries to safeguard the shaky intuition with mathematical rigor and conceptual clarity. Once a good nonperturbative intuition has been achieved, such an attitude may well appear as pedantic.

This means among other things, that the postulated physical requirements should be rigorous properties of the resulting theory. This, as was already mentioned before, is certainly not the case in the quantization approach. Even the strongest defenders of Lagrangians in QFT are perfectly aware that canonical commutation relations, functional integral representations etc. are only mental marks for the inspiration, their purpose is to suggest formal tricks and recipes which eventually lead to correct structures. They are almost never properties of the constructed theory, and therefore quantization is an art and not a mathematical theory. LQP on the other hand only uses intrinsic properties which are valid throughout the process of construction and maintains its validity for the resulting theory. The original postulates may bring about a lot of other surprising properties, but they never get lost in the results. Lagrangians and functional integral representations are in general not reconstructible from e.g. the physical correlation functions.

A representation (π, U) of a C^* -algebra and an automorphism group (\mathcal{A}, α_t) is a representation π of \mathcal{A} in a Hilbert space H together with a strongly continuous unitary representation U of \mathbf{R} in H:

$$U(t)\pi(A)U(t)^{-1} = \pi(\alpha_t(A)), \quad U(t) = e^{iHt}, \ H \ge 0$$
(7.1)

For such representations the following theorems hold.

Theorem 27 (Reeh-Schlieder) Let $\{\mathcal{A}(\mathcal{O})\}_{\mathcal{O}\in\mathcal{K}}$ be a local net with translation symmetry and \mathcal{O}_0 a space-time region such that there is $\mathcal{O}_0 \subset \mathcal{O}$ and a neighborhood of zero \mathcal{V} with: $\mathcal{O} \supset \mathcal{O}_0 + \mathcal{V}$ and additivity: $\bigvee_x \mathcal{A}(\mathcal{O}_0 + x) = \mathcal{A}''$. It follows that: $\overline{\mathcal{A}(\mathcal{O})\psi} = \overline{\mathcal{A}\psi}$ for all $\psi \in \mathcal{D}(e^{aP})$, $a \in V^{\uparrow}$

Remark 7 The physical content is that there exist no annihilation operators as long as the localization region allow a nontrivial causal complement. The cyclicity is equivalent to the dual situation of absence of annihilation operators in the causal disjoint region which is contained in the commutant algebra.

Proof. One shows that for any vector $\phi \perp \mathcal{A}(\mathcal{O})$ implies $\phi = 0$

For any such vector we have $(\phi, \alpha_{x_1}(A_1)..., \alpha_{x_n}(A_n)\psi) = 0$, or in terms of boundary values of analytic functions in the tube:

$$\lim_{x_i \to z_i} (\phi, e^{iz_1 P} A_1 e^{i(z_2 - z_1)P} A_2 \dots A_n e^{-(iz_n + a)P} e^{aP} \psi) = 0$$
(7.2)

where the limit is taken from the inside the tubes

$$\mathcal{T}^{(n,a)}: Im(z_{i+1} - z_i) \in V^{\uparrow}, \ z_0 = 0, \ z_{n+1} = ia$$
(7.3)

in which the matrix element is an analytic function $F_{\phi,\psi}(z_1,...,z_n)$. Different boundary orders correspond to different orders in the operator product. Since Fvanishes in an open set on the boundary it is (thanks to the generalized Schwarz reflection principle) identically zero. This proves the theorem, since ϕ is then orthogonal to a dense set of vectors and hence on all vectors and therefore the cyclicity of ψ follows.

Taking now locality into account we conclude that if $A\psi = 0$ for a vector ψ as above, and $A \in \mathcal{A}(\mathcal{O}')$; from this one concludes $BA\psi = AB\psi = 0$ for all $B \in \mathcal{A}(\mathcal{O})$ and hence A = 0 on a dense set and therefore $A \equiv 0$ or in words: ψ is cyclic and separating. The separating property of $\mathcal{A}(\mathcal{O})$ is equivalent to the cyclicity of $\mathcal{A}(\mathcal{O})'$ (with respect to the same vector Ω) and therefore guarantied by the cyclicity for the spacelike complement $\mathcal{A}(\mathcal{O}') \subset \mathcal{A}(\mathcal{O})'$.

This is a characteristic property in QFT of finite energy states (in particular of the vacuum Ω) with respect to local algebras $\mathcal{A}(\mathcal{O})$ such that O' is not empty i.e. it does not hold for state vectors in standard QM.

If not handled with great care, one can easily get into pitfalls with causality¹. In fact most of the more sophisticated apparent "violations of causality" in which quantum mechanical properties are attributed to local algebras (including those through tunneling) are due to some conceptual misunderstanding of QFT.

Literally speaking the R-S theorem says that by applying suitable operators which have some time duration in a spatially limited laboratory, one can approximate a state which describes the instantaneous creation of matter here and

 $^{^1 \}rm One$ of the more "spectacular" recently published claims about apparant causality violation has been critical reviewed in [74]

antimatter "behind the moon". A closer look reveals that such a sequence of approximations from the dense R-S set require more and more exotic (increasing energy-momentum costs) pieces of hardware. This suggests that the limited localization in phase space (i.e. a field theoretic analogue of the finite number of degrees of freedom of standard QM) and not just the localization in space-time becomes relevant for the cost balance. Indeed, the precise formulation of this idea in the form of the "nuclearity property" (of the degrees of freedom counting) has turned out to be extraordinarily useful. Theories which do not obey this requirement, as e.g. those with a e.g. "Hagedorn temperature", have physical pathologies. The appropriate definition² is the following:

Definition 6 [3] A positive energy representation of an observable net is said to fulfill the "nuclearity property" if the set of vectors of the form:

$$N_{\beta,r} := e^{-\beta H} \mathcal{A}^{(1)}(\mathcal{O}_r) \Omega$$

is nuclear (intuitively: close to a finite set) and its nuclear index (see mathematical appendix) is suitably bounded by β and the size r of the double cone \mathcal{O}_r .

The notation is as follows: $\mathcal{A}^{(1)}(\mathcal{O})$ is the unit ball of $\mathcal{A}(\mathcal{O})$, i.e. observables with operator norms ≤ 1 . Nuclear means that the set of vectors in H is contained in the image of $TH^{(1)}$ where T is a trace class operator, and $H^{(1)}$ the unit ball in H, i.e. that the infinite dimensional sphere in Hilbert space becomes an ellipsoid with suitably decreasing higher semi-axis. There exists a equivalent formulation of nuclearity, in which the global Hamiltonian is replaced by modular operators which have a direct affiliation to local regions, which is known under the name "modular nuclearity" [3].

All free theories and several interacting theories, for which it was feasible to check such difficult (remote from pointlike fields) requirements, fall into the range of validity of this property. On the other hand for models with infinite towers of stable particles, one should be prepared for violations. The nuclearity property not only secures the nonperturbative existence of temperature states of infinite systems (whose existence in the vacuum state has been assumed), but also give rise to a number of interesting and surprising physical concepts. The most prominent is the so called "split property":

Definition 7 For double cones \mathcal{O}_1 and \mathcal{O}_2 (or wedges in case of massive theories) with strict inclusion $O_1 \ll O_2$ (no touching of boundaries), there exists a canonically constructed type I factor \mathcal{N} with:

$$\mathcal{A}(\mathcal{O}_1) \subset \mathcal{N} \subset \mathcal{A}(\mathcal{O}_2) \tag{7.4}$$

Its existence can be used in order to split the apace and the algebras in terms of tensor products as follows:

 $^{^{2}}$ Some of the properties already made their first appearance in the chapter 3 on free fields.

$$\exists W \ H \ \to \ H \otimes H \ s.t. \ H = W^*(H \otimes H)W \ s.t.$$

$$\mathcal{A}(\mathcal{O}_1) = W^*(\mathcal{A}(\mathcal{O}) \otimes \mathbf{1})W \subset W^*(\mathcal{B}(H) \otimes \mathbf{1})W = \mathcal{N}$$

$$\mathcal{A}(\mathcal{O}_2)' = W^*(\mathbf{1} \otimes \mathcal{A}(\mathcal{O}_2)')W \subset W^*(\mathbf{1} \otimes \mathcal{B}(H))W = \mathcal{N}'$$

$$(7.5)$$

This split property follows from the nuclearity property (Haag). It is deeply related to one of the oldest concepts in QFT, the vacuum polarization (first studied by Heisenberg and later elaborated by Weisskopf). One of the observations in the old days was that a sharp spatial cutoff (e.g. by a characteristic step function of a spatial or even a space-time region) leads to infinite large vacuum fluctuations. This poses the question whether there is a smoother causal way of splitting into \mathcal{O} and the causal complement, such that the Hilbert space factorizes in a manner well-known from the QM box quantization (where such splits into a 3-dim. inside and outside region are frequently used). If one leaves a "collar" around \mathcal{O} , then the above theorem yields such a factorization. In ordinary QM, the carved out collar would prevent the tensor product space from being equal to the full space (or rather one would have to work with the tensor product of three factors including a factor for the collar). Thanks to the Reeh-Schlieder theorem, the QFT situation is better. The fact that one needs the collar in order to achieve the spatial split is very much related to the hyperfinite type III₁ property of the $\mathcal{A}(\mathcal{O})$ local algebras. For the proof of hyperfiniteness, the nuclearity enters in an essential way. Hyperfinite type III_1 factors are unique (always meant modulo isomorphisms), and therefore the nuclearity property leads to a universality of local algebras, thus convincingly confirming the ideas of the founding fathers of algebraic QFT: a single algebra is void of physical meaning (the "no hair" property of algebraic QFT) and all physical properties reside in the net relations, i.e. in the position of the algebras relative to each other.

Another technically important property is the following "property B" which is due to Borchers [3].

Theorem 28 Let E be a local projector $E \in \mathcal{A}(\mathcal{O})$. Then there exists an isometry V localizable in a possibly slightly bigger region $\widetilde{\mathcal{O}} \supset \mathcal{O}$ with $E = VV^*$.

Again this theorem points towards properly infinite (i.e. type III) algebras; if it would not be for the possible enlargement of \mathcal{O} , the statement $E = VV^*$, $V^*V = 1$ (isometry) yields $E \sim 1$, i.e. all the projectors are "infinite" and therefore $\mathcal{A}(\mathcal{O})$ is indeed of type III. All explicitly known local QFT algebras are actually hyperfinite factors of type III₁ in the refined classification theory of A. Connes. The subscript 1 refers to complete outerness of the action of modular groups on the algebra, whereas "hyperfinite" is somewhat loosely speaking a property of approximatability by finite degrees of freedoms (prerequisite for lattice approximations) which can be shown to arise from the QFT phase space structure which results from the nuclearity requirement. It holds for the local algebras, but does not necessarily apply to globalizations as GNS-representations of \mathcal{A}_{quasi} and \mathcal{A}_{univ} . Type III are the "biggest" von Neumann factors in the sense that they absorbs any other tensor factor. For the wedge region one can actually prove that $\mathcal{A}(W)$ is a hyperfinite type III₁ factor. Factor algebras are very natural also in physics since they generalize the notion of irreducibility in those cases where their intrinsic impurity prevents irreducibility. On the opposite end one finds type II_1 factors which are absorbed into any other tensor factor and hence smallest in this sense. The latter are big enough in order to incorporate all the generalizations of group symmetry which recently emerged in V. Jones inclusion theory of subfactors. In algebraic QFT only the so-called intertwiner subalgebras (associated with the composition and reduction of charge sectors of observable algebras) are of this kind. These subalgebras give rise to combinatorial or so called topological QFT. Such small algebras type II_1 algebras can only appear because the space-time symmetries remain outside the intertwiner algebras. External symmetries, in particular translations, require the presence of infinite projectors (typically type I) as defined above, and finally restrictions to local algebras (with only partial space-time symmetries) allow only infinite projectors (type III_1).

In connection with the limitation of energy-momentum and the nuclearity formalism (mentioned below) it is convenient to have a mathematical framework which makes precise the concept of energy-momentum transfer. This was elaborated [41] by the mathematical physicist Borchers and the mathematician Arveson. The idea is to first introduce a notion of spectrum of the automorphism. The automorphism α_t of the C^* -algebra may be extended via (7.1) to the enveloping von Neumann algebra $\pi(\mathcal{A})''$. It should not lead to any confusion if we stay sometimes with the same symbols for the extended objects. With the help of L. Schwartz test functions $f \in \mathcal{S}(\mathbf{R})$ we form $\alpha_f(A) = \int dt f(t) \alpha_t(A)$. It is easy to see that the extended automorphism, and therefore α_f , maps also the von Neumann extension into itself (since it commutes with elements from the commutant $\pi(\mathcal{A})'$ inside matrix elements). One now defines the (Arveson-) spectrum of $A \in \pi(\mathcal{A})''$ as:

$$spec_{\alpha}(A) = \left\{ \omega \in \mathbf{R} \mid \forall nbhs \ N \ of \ \omega, \exists f \in \mathcal{S}(\mathbf{R}) \ s.t. \ supp \tilde{f} \subset N, \ \alpha_f(A) \neq 0 \right\}$$
(7.6)

The size of the individual $\alpha_f(A)$ -contributions is evidently limited by $supp\tilde{f}$. We can manufacture operators A with $spec_{\alpha}(A) \in I, I$ given, by smoothening a given B with $f, supp\tilde{f} \subset I$:

$$A := \alpha_f(B)$$

The (algebraic) subspaces with energy transfer $\geq E$ are defined as:

$$\mathcal{A}_E = \{ A \in \mathcal{A} \mid sp_\alpha(A) \subset [E, \infty] \}$$
(7.7)

The usefulness of these concepts begins to show up if one relates this with projection operators in the Hilbert space H of the representation (π, U) :

$$P_E := proj \ on \ \bigcap_{E' < E} \mathcal{A}_{E'} H \tag{7.8}$$

These projectors define a spectral family since they fulfill $P_E = 1$ for $E \leq 0$ and $\lim_{E\to\infty} P_E = 0$, and are in addition upper continuous. Hence one may associate a "Hamiltonian" **H** with α_t :

$$\mathbf{H} = \int E dP_E, \ V(t) := e^{i\mathbf{H}t} = \int e^{iEt} dP_E$$
(7.9)
$$\sim \ \alpha_t(A) = V(t)AV(t)^{-1}$$

Since: $\pi(\mathcal{A})'\mathcal{A}_E H \subset \mathcal{A}_E H$, we find:

$$A' P_E = P_E A' P_E = P_E A',$$

$$\sim [P_E, A'] = 0, \forall A' \in \pi(\mathcal{A})'$$

or using more physical terminology: the infinitesimal generator \mathbf{H} of the symmetry α_t may always be chosen in such a way that \mathbf{H} is associated to the algebra (the Arveson-Borchers theorem). The algebraically determined \mathbf{H} is called the "minimal" generator. Although the general situation, unlike the well-known explicit Sugawara expression for the chiral translation generator in chiral conformal QFT, does not lead to a concrete functional expression, this should nevertheless be interpreted as the generalized analogue of the situation known in conformal QFT. The extension to abelian groups with several parameters should be obvious.

In the same vein, but taking in addition locality into account and using more powerful analyticity tools ("edge of the wedge techniques"), one proves the following four interesting theorems [41].

Theorem 29 (Locality and the shape of the spectrum) Let $\{\mathcal{A}(\mathcal{O}), \mathcal{A}, \mathbf{R}^d, \alpha\}$ be a local net with translation symmetry and positive energy. Let V(a) denote the above minimal positive energy representation. Then the lower bound of specV is Lorentz-invariant.

This is the counterpart of the classical fact that causal propagation can only be satisfied with L-covariant equations. As a result of this inexorable link in the classical theory, Einstein himself never separated the issue of L-invariance from causality. The content of this theorem is that these notions continue to stay inexorably linked in the local quantum theory setting.

Theorem 30 (General cluster property) Let $\{\mathcal{A}(\mathcal{O}), \mathcal{A}, \mathbb{R}^d, \alpha_a\}$ be a local net as before and ω a translation invariant state and $\{\pi, H, \Omega, U(a)\}$ the GNSrepresentation with $U(a)\Omega = \Omega$. Denote by P_0 the projector onto the subspace of pointwise invariant vectors i.e. $\Omega \in P_0H$. Assume furthermore that the center of $\pi(\mathcal{A})''$ is pointwise invariant under α_a . Then we have the following relation:

$$\lim_{\lambda \to \infty} (\Omega, \pi(A_1)\pi(\alpha_{\lambda b}B_1)\pi(A_2)...\pi(A_n)\pi(\alpha_{\lambda b}B_n)\Omega)$$
(7.10)
= $(\Omega, \pi(A_1A_2...A_n)P_0\pi(B_1B_2...B_n)\Omega)$
= $(\Omega, \pi(B_1B_2...B_n)P_0\pi(A_1A_2...A_n)\Omega), \quad b \ spacelike$

Naturally some of the A_i, B_j may be omittable identity operators, which allows to have $\#A \neq \#B$. In the case of the unique vacuum and a spectral mass gap one may prove the strong (faster than any inverse power in λ) approach of the right hand side which is the standard form of the cluster property. This is then the starting point of the derivation of scattering theory.

Theorem 31 (Additivity of spectrum) Let $\{\pi, H, U(a)\}$ be a factor representation (a von Neumann algebra with trivial center: $\mathcal{Z} = \pi(\mathcal{A})' \cap \pi(\mathcal{A})'')$ of a theory of local observables fulfilling the spectrum condition and assume that U(a) is the minimal representation. Then if p_1 and p_2 are in specP, so is p_1+p_2 . moreover if the mass spectrum consists of a discrete part $m_0 < m_1 < \dots$ and a continuum starting at $m_c > m_i$ then:

$$3m_0 \ge m_c$$

The expected (from scattering theory) relation $m_c = 2m_0$ remains still unproven.

Theorem 32 (Absence of classical fields) There exist no classical field theories (*i.e.* abelian algebras) which fulfill the spectrum condition.

Theorem 33 Let $\{\mathcal{A}(\mathcal{O}), \mathcal{A}, \mathbf{R}^d, \alpha\}$ be a theory of local observables and define:

$$\mathcal{A}(a) = \bigcap_{a \in \mathcal{O}} \mathcal{A}(\mathcal{O}) \tag{7.11}$$

Then $\mathcal{A}(a) \subset \mathcal{Z}(\mathcal{A})$.

Clearly this may be interpreted as an algebraic generalization of the famous Bohr-Rosenfeld argument on the nonexistence of finite electro-magnetic quantum field strength at a point i.e. the necessity for smearing (or avaraging) quantum field strength in order to avoid singularities. This theorem which is due to Wightman has generalizations to subsets of Minkowski space. For spacelike 3-d hypersurfaces and for time-like segments the analogously defined algebras are nontrivial and equal to the algebras of their causal completions i.e. $\mathcal{A}(\mathcal{O}'')$

7.2 Abstracting Superselection Principles

If we use fields in standard QFT in order to define local nets of field algebras $\{\mathcal{F}(\mathcal{O})\}\)$, we find the following properties³ (see also chapter 5):

• \tilde{P} -covariance, positive energy and uniqueness of the vacuum.

 \exists a strongly continuous representation U of the covering of the Poincaré group P_+^\uparrow :

$$U(L)\mathcal{F}(\mathcal{O})U(L)^{-1} = \mathcal{F}(L\mathcal{O})$$
(7.12)

 $^{^{3}}$ The reader interested in technical and conceptual details should follow the historical path and look at Haag's book as well as the original articles.

and the generators P_{μ} of the translations satisfy the spectrum condition $spec \in V^{\uparrow}$ with $P\Omega = 0$, Ω being the unique vacuum.

- \exists a compact (global gauge) group⁴ G and strongly continuous faithful representation U of G which commutes with the Poincaré group (factorization of internal and external symmetries) s. t.:: $U(g)\mathcal{F}(\mathcal{O})U(g)^{-1} = \mathcal{F}(\mathcal{O}), \quad U(g)\Omega = \Omega$
- $\exists \kappa \in G$ of order two i.e. $\kappa^2 = 1$ s. t.. with $\mathcal{F} = \mathcal{F}_+ + \mathcal{F}_-$, $\alpha_{\kappa}(\mathcal{F})_{\pm} = \pm \mathcal{F}$ and spacelike separated \mathcal{O}_1 and \mathcal{O}_2 ; the following graded (or "twisted") locality relation holds:

$$\{\mathcal{F}_{-}(\mathcal{O}_{1}), \mathcal{F}_{-}(\mathcal{O}_{2})\} = 0$$

$$[\mathcal{F}_{+}(\mathcal{O}_{1}), \mathcal{F}_{-}(\mathcal{O}_{2})] = 0 = [\mathcal{F}_{+}(\mathcal{O}_{1}), \mathcal{F}_{+}(\mathcal{O}_{2})]$$
(7.13)

We write this in the condensed form:

$$\mathcal{F}(\mathcal{O}') = \mathcal{F}(\mathcal{O})^{tw}, \quad \mathcal{F}(\mathcal{O})^{tw} := K\mathcal{F}(\mathcal{O})'K^{-1} \quad (7.14)$$

$$V \ s.t. \ V\mathcal{F}(\mathcal{O})V^{-1} = \kappa(\mathcal{F}(\mathcal{O})) \ and \ K = \frac{1+iV}{1+i}$$

- Additivity: $\mathcal{F}(\mathcal{O}) = \bigvee_i \mathcal{F}(\mathcal{O}_i), \quad \mathcal{O} = \bigcup \mathcal{O}_i$
- Haag (twisted) Duality:

$$\begin{aligned}
\mathcal{F}(\mathcal{O}) &= \mathcal{F}(\mathcal{O}')^{tw} & \curvearrowright \\
\mathcal{A}(\mathcal{O}')' & \mid_{H_0} = \mathcal{A}(\mathcal{O})'' \mid_{H_0}, \quad \mathcal{A}(\mathcal{O}) := \mathcal{F}(\mathcal{O}) \cap U(G)'
\end{aligned}$$
(7.15)

where the observable algebra is defined by this invariance principle and the von Neumann algebra of a noncompact region \mathcal{N} , as e.g. the causal complement of a double cone \mathcal{O}' , are defined in terms of an additive covering by double cones \mathcal{O}_i together with von Neumann closure: $\mathcal{A}(\mathcal{N}) = \bigcup_i \mathcal{A}(\mathcal{O}_i)$

Some comments are in order. Although some of these properties are evident or at least plausible, I recommend to look up the proofs. The conclusion in the \rightarrow direction does not hold in the case of d=1+1 where the order-disorder duality makes its appearance (see chapter 3, section 7). The quantum intuition acquired from standard QT (as well as from Lagrangian quantization) is treacherous in local QFT, an area for which a good intuition still needs to be developed. This applies in particular to the duality structure.

The notation $|_{H_0}$ denotes the restriction to the vacuum sector H_0 with: $U(G)H_0 = H_0$ pointwise in H_0 . What is referred to as the observable algebra in these notes is not \mathcal{A} in H, but rather the smaller (irreducible) algebra $\mathcal{A} |_{H_0}$. The gauge invariant part can also be obtained via the conditional expectation

⁴Llocal gauge groups are not directly related to symmetries in the same theory.

(averaging with compact group):

$$m(F) \equiv \int_{G} dg \alpha_{g}(F), \quad \int_{G} dg = 1$$
(7.16)
with : (1) $m(\mathcal{F}(\mathcal{O})) = \mathcal{A}(\mathcal{O})$
(2) m is normal
(3) m commutes with α_{g}

The continuity property (2), defined in terms of the predual (appendix) which is also equivalent to m being " σ -continuous", allows the continuation of m to all operators B(H) i.e. it is the natural kind of continuity for operations on von Neumann algebras. We obtain $\mathcal{F}'' = B(H), \mathcal{A}'' = m(\mathcal{F}'')$ and hence $\mathcal{A}'' = m(B(H))$ as well as $\mathcal{A}' = U(G)''$

This gives us the desired tensor decomposition of the Hilbert space:

$$H = \bigoplus_{\sigma} H_{\sigma} \otimes H'_{\sigma} \tag{7.17}$$

where the first factor H_{σ} is the irreducible representation space for the irreducible representation $U_{\sigma}(G)$ of the internal symmetry group, and H'_{σ} denotes its (infinite dimensional) multiplicity space, which is an irreducible representation space of \mathcal{A} corresponding to π_{σ} . With other words we have:

$$A |_{H_{\sigma} \otimes H'_{\sigma}} = \mathbf{1}_{H_{\sigma}} \otimes \pi_{\sigma}(A) \quad A \in \mathcal{A}$$

$$U(g) |_{H_{\sigma} \otimes H'_{\sigma}} = U_{\sigma}(g) \otimes \mathbf{1}_{H'_{\sigma}} \quad g \in G$$

$$(7.18)$$

 \mathcal{A} in H contains generally many other irreducible representations π_{σ} besides the vacuum representation, and the primer into the theory of superselection sectors consists in classifying these π_{σ} , in particular to understand what properties they share. For this purpose we introduce minimal projectors in the algebra U(G)'':

$$E = \int dg U(g) \left(\phi, U_{\sigma}(g^{-1})\phi\right) \quad \phi \in H_{\sigma} \text{ arbitrary, } \|\phi\| = 1$$
(7.19)

Since according to the Reeh-Schlieder theorem $\mathcal{F}(\mathcal{O})$ acts cyclically on Ω , we always find elements $F \in \mathcal{F}(\mathcal{O})$ with $EF\Omega \neq 0$. The definition:

$$T\psi = EF\psi \quad \psi \in H_0$$

determines a partial intertwiner $T: H_0 \to EH$ with the intertwining property:

$$T\pi_0(A) = \pi_E(A)T, \quad A \in \mathcal{A}(\mathcal{O}')$$

$$(7.20)$$

The reader easily checks that the vectors $T\Omega \in H_E$ and $|T|\Omega \in H_0$ (since $T^*T : H_0 \to H_0$) have the same expectation values on $\mathcal{A}(\mathcal{O}')$ i.e. induce the same partial states. Using the Reeh-Schlieder cyclicity one shows that there are sufficiently many partial intertwiners such that the set of states over $\mathcal{A}(\mathcal{O}')$ in all representation obtained from the decomposition of $\pi(\mathcal{A})$ on H agree i.e. the restriction of the net \mathcal{A} to $\mathcal{A}(\mathcal{O}')$ gives the same folium (see mathematical appendix) of states independent of the charge sector σ .

Theorem 34 All irreducible subrepresentations π_{σ} satisfy the (DHR) condition:

$$\pi_{\sigma} \mid_{\mathcal{A}(\mathcal{O}')} \simeq \pi_0 \mid_{\mathcal{A}(\mathcal{O}')} \quad \forall \mathcal{O} \in \mathcal{K}$$

$$(7.21)$$

i.e. the representations of the observable algebra (obtained from an invariance principle on the field algebra) are unitarily equivalent in the causal complement of double cones (and more generally any space time region which admits a non-trivial causal complement).

This is taken as a definition of (DHR) compactly localizable representations for an arbitrary observable net.

7.3 Starting the Reverse: the DHR Endomorphisms

The previous DHR localization condition may be now be taken as the starting point for the elaboration of the pivotal part of algebraic QFT: the DHR superselection theory. Let us start with the classification of simple (abelian) sectors because they are also simpler in the everyday use of the word. In order to appreciate the following definitions, one should think of one-dimensional representations of a group G which form a subcategory of representations closed under compositions. In terms of the projectors E_{σ} on $H_{\sigma} \otimes H'_{\sigma}$ from the previous section which are elements of the center $\mathcal{Z}(U(G'')) = \mathcal{Z}(\mathcal{A}'')$, we have $U(g)E_{\sigma} = E_{\sigma}U(g) = \chi(g)E_{\sigma}$. In order to understand this property in terms of observables \mathcal{A} only (without \mathcal{F}), we convince ourselves that the representation π_{σ} satisfies the Haag duality property, which up to now we only met in connection with the vacuum representation:

$$\pi(\mathcal{A}(\mathcal{O}')'' = \pi(\mathcal{A}(\mathcal{O}))' \cap \pi(\mathcal{A})''$$
(7.22)

Here the left hand side should be understood to as the von Neumann algebra generated by $\mathcal{A}(\mathcal{O}_1)$ for all $\mathcal{O}_1 \subset \mathcal{O}'$. For π irreducible the relation is often written as $\pi(\mathcal{A}(\mathcal{O}')) = \pi(\mathcal{A}(\mathcal{O}))'$. Replacing = by \subset , we have the Einstein causality relation, therefore (7.22) represents a strengthening of causality (maximal, as it turns out).

A π_{σ} , as obtained in the previous section by restriction from a field algebra \mathcal{F} , fulfills Haag duality since $\mathcal{A}(O)' = (\mathcal{F}(\mathcal{O}) \cap U(G)')' = \mathcal{F}(O)' \vee U(G)''$ and acting with the projection E_{σ} as well as with m on both sides (those actions commute) the U(G)'' is killed and we finally obtain: $\pi_{\sigma}(\mathcal{A}(\mathcal{O}))' = E_{\sigma}(\mathcal{F}(\mathcal{O})'E_{\sigma} = E_{\sigma}(\mathcal{F}(\mathcal{O})' \cap U(G)')E_{\sigma} = \pi_{\sigma}(\mathcal{A}(\mathcal{O}'))''$. In the last step we used the twisted duality of \mathcal{F} . We will later see that representations of the observable net \mathcal{A} fulfill Haag duality iff they correspond to simple sectors.

Let us now start to do the reverse, namely to construct a charge-carrying field algebra \mathcal{F} from the observable algebra \mathcal{A} and its DHR (7.21) representations. We first must find some good mathematical concepts to classify the DHR localized representations. The unitary equivalence of $\pi(\mathcal{A}(\mathcal{O}'))$ with $\pi_0(\mathcal{A}(\mathcal{O}'))$

in (7.21) guaranties the existence of partial intertwiners i.e. isometries V: $H_0 \rightarrow H_{\pi}$ with:

$$V\pi_0(A) = \pi(A)V, \quad A \in \mathcal{A}(\mathcal{O}') \tag{7.23}$$

We define a representation $\hat{\pi}(\mathcal{A})$ in H_0 equivalent to $\pi(\mathcal{A})$ in H_{π} as

$$\hat{\pi}(A) := V^{-1}\pi(A)V, \quad A \in \mathcal{A}$$
(7.24)

By construction this representation agrees with π_0 in \mathcal{O}' . For sufficiently large regions namely $\mathcal{O}_1 \supset \mathcal{O}$, the range of $\hat{\pi}$ is contained in that of π_0 i.e. $\hat{\pi}(\mathcal{A}(\mathcal{O}_1)) \subset \pi_0(\mathcal{A}(\mathcal{O}_1))$, and hence a fortiori $\hat{\pi}(\mathcal{A}) \subset \pi_0(\mathcal{A})$. This follows by using (vacuum) Haag duality, namely: $[\pi_0(\mathcal{A}'), \hat{\pi}(\mathcal{A})] = \hat{\pi}[\mathcal{A}', \mathcal{A}] = 0$ for $\mathcal{A}' \in \mathcal{A}(\mathcal{O}'_1), \mathcal{A} \in \mathcal{A}(\mathcal{O})$ and hence $\hat{\pi}(\mathcal{A}) \in \pi_0(\mathcal{A}(\mathcal{O}'_1)' \subset \pi_0(\mathcal{A})$ by Haag duality. Therefore ρ defined by:

$$\rho := \pi_0^{-1} \circ \hat{\pi}, \quad \rho : \mathcal{A} \to \mathcal{A} \tag{7.25}$$

is an endomorphism of the C^* algebra $\mathcal A$ with the following remarkable properties:

- ρ is localized in \mathcal{O} (notation: $loc \rho \subset \mathcal{O}$), i.e. $\rho(A) = A, A \in \mathcal{A}(\mathcal{O}')$
- transportable, i.e. $\forall \mathcal{O}_1, \mathcal{O}_2$ with $\mathcal{O}_2 \supset \mathcal{O}_1 \cup \mathcal{O} \ \exists U \in \mathcal{A}(\mathcal{O}_2) \ s.t. \ U\rho(A)U^* = A \text{ for } A \in \mathcal{A}(\mathcal{O}'_1)$
- $\rho(\mathcal{A}(\mathcal{O}_1)) \subset \mathcal{A}(\mathcal{O}_1), \quad \forall \mathcal{O}_1 \supset loc\rho$

The very simple proof of these properties is left to the reader. We follow Haag and use the notation Δ for the set of such $\rho's$, and denote by $\Delta(\mathcal{O})$ the subset of $\rho's$ with $loc\rho \in \mathcal{O}$.

In the constructive approach based on the observable net and its endomorphisms with the above properties, one defines the *sectors* as the equivalence classes of $\rho's$ modulo inner automorphisms. The following structural investigation of localized transportable endomorphisms is independent of the dimensionality of the QFT i.e. holds as well for low dimensional theories. Let us first look at abelian sectors which by definition are equivalence classes of automorphism i.e. $\rho's$ with $\rho(\mathcal{A}) = \mathcal{A}$.

Theorem 35 ρ is automorphism $\Leftrightarrow \pi_{\rho} = \pi_0 \circ \rho$ is Haag dual $\Leftrightarrow \rho^2$ is irreducible (no branched fusion) \Leftrightarrow Ind $[A : \rho(A)] = 1$ (trivial Jones index)

The reader should try to prove it for himself and consult Haag's book, if he needs more than 5 lines.

In algebraic QFT the Jones index enters through the statistics operators ε which we explain briefly in the sequel. They are special intertwiners ("Verketter" in the sense of Schur). An intertwining operator is a $V \in B(H)$ which links a representation $\pi_0 \rho$ with a subrepresentation of $\pi_0 \sigma$ i.e. $V \cdot \pi_0 \rho(A) = \pi_0 \sigma(A) \cdot V$ $\forall A \in \mathcal{A}$. In case that ρ is irreducible, the two representations are equivalent and the intertwining operator becomes a "charge transporting" operator. By Haag duality⁵ one obtains $V = \pi_0(T)$ with $T \in \mathcal{A}$ and the intertwining relations:

$$T\rho(A) = \sigma(A)T \quad \forall \ A \in \mathcal{A} \tag{7.26}$$

The space of self-intertwiners: (ρ, ρ) is the commutant $\rho(\mathcal{A})'$ of $\rho(\mathcal{A})$ in \mathcal{A} and by Schur's lemma, equal to the scalars **C1** iff ρ is irreducible. Therefore, when ρ is irreducible, the linear space of intertwiners: $\rho \to \sigma$ is a *Hilbert space within* the algebra of local observables with the inner product $(T_1, T_2) := T_1^*T_2$. The notation for the space of intertwiners T from σ to ρ is $T \in (\rho, \sigma)$.

For every pair of DHR endomorphisms there is a unitary local intertwiner $\varepsilon(\rho, \sigma) : \rho\sigma \to \sigma\rho$ i.e. $\varepsilon \in (\sigma\rho, \rho\sigma)$. This flip operator is called the *statistic operator*. The collection of statistics operators is uniquely determined by the coherence relations with local intertwiners and among themselves:

$$\begin{split} \varepsilon(\sigma_1, \sigma_2)\sigma_1(T_2)T_1 &= T_2\rho_2(T_1)\varepsilon(\rho_1, \rho_2) \quad \forall \ T_i : \rho_i \to \sigma_i \qquad (7.27) \\ \varepsilon(\rho_1\rho_2, \sigma) &= \varepsilon(\rho_1, \sigma)\rho_1(\varepsilon(\rho_2, \sigma)) \\ \varepsilon(\rho, \sigma_1\sigma_2) &= \sigma_1(\varepsilon(\rho, \sigma_2))\varepsilon(\rho, \sigma_1) \end{split}$$

together with the "initial conditions"

=

$$\varepsilon(\rho, id) = \varepsilon(id, \rho) = 1$$

$$\varepsilon(\rho, \sigma) = 1 \quad \text{whenever } \sigma < \rho$$

$$(7.28)$$

where $\sigma < \rho$ means $\log \sigma$ is in the left spacelike complement of $\log \rho$. The Artin braid relation is a special consequence of the above coherence relations

$$\rho_3(\varepsilon(\rho_1, \rho_2))\varepsilon(\rho_1, \rho_3)\rho_1(\varepsilon(\rho_2, \rho_3))$$

$$\varepsilon(\rho_2, \rho_3)\rho_2(\varepsilon(\rho_1, \rho_3))\varepsilon(\rho_1, \rho_2)$$
(7.29)

In particular, by assigning the local operators $\rho^{i-1}(\varepsilon(\rho,\rho))$ to the standard Artin generators σ_i of the braid group B_n (see chapter1) we obtain a unitary representation of the braid group B_{∞} in \mathcal{A} which we call the statistics of the endomorphism ρ (for reasons which soon will become evident).

We introduce a conjugate endomorphism $\bar{\rho}$ to ρ by demanding that $\bar{\rho}\rho$ contains the vacuum sector, i.e. that there exists an intertwiner $R \in (id, \bar{\rho}\rho)$ which induces a standard left inverse ϕ of ρ

$$\phi(A) = R^* \bar{\rho}(A) R \quad \forall \ A \in \mathcal{A} \tag{7.30}$$

with finite statistics. Here we recall that the left inverse of an endomorphism ρ of \mathcal{A} is a normalized positive linear map satisfying the relation $\phi(\rho(A)B\rho(C)) = A\phi(B)C$. It is called regular if it is of the above form, and standard, if in addition the statistics parameter $\lambda_{\rho} := \phi(\varepsilon(\rho, \rho)) \in \rho(\mathcal{A})'$ is a nonvanishing multiple of a

⁵In the reverse approach which starts from the observable algebra \mathcal{A} , the Haag duality is postulated for the vacuum representation. If it does not hold for the original net, one passes to the dual net \mathcal{A}^d which fulfills Haag duality by construction.

unitary (which then depends only on the sector $[\rho]$). A sufficient condition for the existence of a standard left-inverse and therefore of a conjugate is that there is *some* left-inverse with statistics parameter $\lambda_{\rho} \neq 0$ ("finite statistics") and that ρ is translation covariant with positive energy condition. The uniqueness of the standard left inverse is a consequence of its definition. Any theory with a mass gap possesses a standard left inverse [93]. The standard left inverse of ρ turns out to be a trace on $\rho(\mathcal{A})'$. The inverse modulus of λ_{ρ} is called the *statistical* dimension $d(\rho) \equiv d_{\rho} \geq 1$. One easily proves that $\lambda_{\rho} = \lambda_{\bar{\rho}}$. For irreducible $\rho's$ we have $\lambda_{\rho} = \frac{\kappa_{\rho}}{d_{\rho}}$ with κ_{ρ} being the *statistical phase*. If one computes this numbers using the field formalism presented below, one finds $d_{\rho} = dimH_{\rho}$ and $\kappa_{\rho} = \pm 1$ for Bosons/Fermions. In fact for d=3+1 the statistics operator is easily shown on general grounds to fulfill $\varepsilon^2 = 1$ (i.e. absence of monodromies) which leads to permutation group statistics. The concepts are much richer in the case of braid group statistics. Even in that case one succeeds to prove the identity of the spin phase with the above statistics phase i.e. the spin-statistics theorem.

We will not enter a presentation of V. Jones inclusion (subfactor) theory, but just mention that $\text{Ind}[A:\rho(A)] = d_{\rho}^2$, i.e. the inclusion index is the square of the statistical dimension.

In order to understand the reconstruction in the case of proper endomorphisms i.e. for $\rho's$ with $\operatorname{Ind}[A:\rho(A)] > 1$, we need some more conceptual preparation. This is obtained by briefly returning to the field algebra \mathcal{F} in the case where \mathcal{A} is the fixed point algebra under a nonabelian G. In that situation an irreducible endomorphism ρ with $\pi_0 \circ \rho \simeq \pi_\rho$ and $\operatorname{loc} \rho \subset \mathcal{O}$ gives via $\omega = \omega_0 \circ \rho$ a pure state localized in \mathcal{O} . The big Hilbert space H in which \mathcal{F} acts has many vectors which induce ω :

$$H_{\omega} = \left\{ \phi \in H \mid (\phi, A\phi) = \omega(A) \left\| \phi \right\|^2 \right\}$$
(7.31)

As the notation already anticipates, H_{ω} is a Hilbert space, a fact which is easily verified using the purity of ω . Its dimension is equal to the dimension d_{ρ} of the H_{ρ} - tensor factor. Physical intuition tells us that such vectors in H_{ω} can be created from the vacuum by applying charge-carrying fields in \mathcal{F} . In fact we have:

Theorem 36 Every $\phi \in H_{\omega}$ determines uniquely a field operator $\psi \in \mathcal{F}(\mathcal{O})$ with $\psi^*\Omega = \phi$ and the intertwining property $\psi A = \rho(A)\psi$, $A \in \mathcal{A}$.

Encouraged by the intertwining relation in the previous theorem, we define the following linear subspace of \mathcal{F} :

$$H_{\rho} = \{ \psi \in \mathcal{F} \mid \psi A = \rho(A)\psi, A \in \mathcal{A} \}$$
(7.32)

The notation suggests the structure of a Hilbert space. Indeed for two vectors ψ_i , i = 1, 2 we have the following scalar product:

$$\psi_1^* \psi_2 \in \mathbf{C} \cdot \mathbf{1} \tag{7.33}$$

The reason is that the inclusion of $\mathcal{A} \in \mathcal{F}$ is irreducible i.e. $\mathcal{A}' \cap \mathcal{F} = \mathbf{C} \cdot \mathbf{1}$. This follows from $\mathcal{A}' \cap \mathcal{F} = U(G)'' \cap \mathcal{F}$ and the statement that for any element F_0 with $F_0 \mid_{H_0} = c\mathbf{1} \mid_{H_0}$ from the latter algebra the conditional expectation of F^*F with $F := F_0 - c\mathbf{1}$ vanishes i.e. $\pi_0(m(F^*F)) = 0$ hence $m(F^*F) = 0$, since π_0 is faithful. But the expectation values of m in any vector state $\phi \in H$ may be written as an average with a positive integrand:

$$0 = (\phi, m(F^*F)\phi) = \int dg(\phi, \alpha_g(F^*F)\phi)$$

$$\alpha_g(F^*F) = 0 \curvearrowright F^*F = 0 \curvearrowright F_0 = c\mathbf{1}, \ qed.$$
(7.34)

Algebraic Hilbert spaces of isometries inside von Neumann algebras can only occur for von Neumann algebras of properly infinite type.

Theorem 37 For any set of field operators $(F_i)_{i=1...d_{\rho}} \in \mathcal{F}(\mathcal{O})$ transforming like an irreducible tensor representation U_{ρ} there exists a $\rho \in \Delta(\mathcal{O})$ and a $B \in A(\mathcal{O})$ s.t. $F_i = B\psi_i$ with $(\psi_i) \in H_{\rho}$ a orthonormal system of isometries spanning H_{ρ} . The endomorphism ρ is implemented by the ψ'_i s:

$$\rho(F) = \sum_{i=1}^{d_{\rho}} \psi_i F \psi_i^*$$

$$\phi(F) \equiv \frac{1}{d_{\rho}} \sum_{i=1}^{d_{\rho}} \psi_i^* F \psi_i, \quad \phi(\rho(F)) = F$$
(7.35)

The last property is the reason why ϕ is called the left-inverse of ρ . The most interesting and useful emerging structure is the so called Cuntz algebra O_d i.e. the unique C^* -algebra generated by a family of isometries $\{\psi_i\}$ with a full range i.e. with $\sum \psi_i \psi_i^* = \mathbf{1}$. A detailed investigation (not done here) reveals that this is a \mathbf{Z} -graded simple C^* -algebra i.e. without two-sided ideals. Doplicher and Roberts observed that this algebra O_d makes a perfect model for a characterization of the group dual which is appropriate for the encoding of internal symmetries in QFT. The reason is that since each compact group G is a subgroup of some U(d) for sufficiently large N, there is a natural action α on O_d (summation convention):

$$\alpha_{q}(\psi_{i}) = \psi_{i}g_{ii}, \quad unitary \text{ in } H_{d}$$

$$(7.36)$$

The tensor product structure is naturally contained in O_d since $H^k \simeq \otimes^k H$. The fixed point algebra:

$$O_G = \{ A \in O_d \mid \alpha_g(A) = A \; \forall g \in G \}$$

gives rise to an inclusion $O_G \subset O_d$ which turns out to encode the group structure, in analogy with Galois theory. It naturally contains all the intertwiners of tensor representations $T : H^{\otimes r} \to H^{\otimes s}$. In terms of endomorphisms these may be characterized purely algebraically i.e. without tensor products:

$$T\rho^{r}(A) = \rho^{s}(A)T, \quad A, T \in O_{G}$$

$$(7.37)$$

C

This should be compared with the "classical" Tanaka-Krein theory of group duals in terms of representation spaces and intertwiners. In QFT based on observables \mathcal{A} only, one only knows the ρ 's of \mathcal{A} and neither the $H'_{\rho}s$ nor \mathcal{F} . So the question how to construct from an algebra of intertwiners a bigger algebra with a group action is a "baby version" of the QFT symmetry problem: how to reconstruct the symmetry from its shadow it leaves on the observables (in analogy to the famous problem of Marc Kac: "how to hear the shape of a drum?"). For a successful treatment we must make sure that our representation category i.e. the endomorphisms and their intertwiners are big enough in order to contain the conjugates (antiparticle representations in QFT). This is easily achieved by securing the existence of a faithful selfconjugate representation because the tensor products of such a representation contain every irreducible representation. Let us briefly look at the special case SU(d). The first tensor power which contains the identity representation is H^d , explicitly the first invariant is:

$$S = \frac{1}{\sqrt{d!}} \sum_{P \in S_d} sign(P)\psi_{P(1)}....\psi_{P(d)}$$
(7.38)

Hence:

$$\hat{\psi}_{i} = \frac{1}{\sqrt{(d-1)!}} \sum_{\substack{P \in S_{d} \\ P(1)=i}} sign(P)\psi_{P(2)}....\psi_{P(d)}$$
(7.39)

fulfill $\hat{\psi}_i = \sqrt{d}\psi_i^*S$ and therefore is a basis in H_d^{conj} and $\bar{\rho}(A) = \sum_{i=1}^d \hat{\psi}_i A \hat{\psi}_i$. Thus $O_{SU(d)}$ contains all irreducible representations of SU(d) and every intertwiner. But how do we recognize that a C*-algebra is isomorphic to $O_{SU(d)}$? The answer is surprisingly simple: in addition to the operator S we must find a copy of the (infinite) permutation group S_∞ . The model theory $O_{SU(d)} \subset O_d$ as such a presentation:

$$\varepsilon(P) = \sum \psi_{(\alpha)} \psi^*_{(\alpha_P)}, \quad P \in S_n \subset S_\infty, \quad S_n \hookrightarrow S_{n+1}$$
(7.40)
(\alpha) = (\alpha_1, ..., \alpha_n), (\alpha_P) = (\alpha_{P(1)}....\alpha_{P(n)})

where we used a multiindex notation. In particular the formula for the basic transposition is:

$$\varepsilon((12)) = \sum_{i,j} \psi_i \psi_j \psi_i^* \psi_j^* = \pm \rho(U^*) U, \quad \pm : F/B$$
(7.41)

where in this Fermi/Bose alternative U is an auxiliary charge transporter: $\psi'_i = U\psi_i$ which shifts the charge into $loc\psi'_i \subset (loc\psi_i)'$.

One easily checks that the ε,S and ρ are related by:

$$SS^* = E_d := \frac{1}{d!} \sum_{P \in S_d} sign(P)\varepsilon(P)$$

$$S^*\rho(S) = (-1)^{d-1} d^{-1} \mathbf{1}$$
(7.42)

$$\begin{split} \rho(\varepsilon(P)) &= \varepsilon(P'), \quad P' \in S_{n+1}, \quad P'(1) = 1, \ P'(i+1) = P(i) \ i = 1....n \\ \phi(\varepsilon(P)) &= \begin{cases} \varepsilon(P) & P(1) = 1 \\ \frac{1}{d}\varepsilon(P') & P(1) \neq 1, \quad P'(i) = ((1P(1))P)(i+1) \end{cases} \end{split}$$

Here E_d is the antisymmetric representation projector in the S_d group algebra and ρ and its left inverse ϕ implement right and (partial) left shifts on S_{∞} . The algebra O_{SUd} is generated by these permutation group elements $\varepsilon(P)$ and the S-intertwiners (the so-called Brower elements). If $G \subset SU(d)$ then $O_G \supset$ $O_{SU(d)}$ and therefore there are more generators.

We will give the DR characterization of G without proof:

Theorem 38 [43] Let \hat{O} be a simple C^* -algebra with an endomorphism $\hat{\rho}$ and a unitary representation $\hat{\varepsilon}$ of S_{∞} with the following properties:

- (i) $\hat{\varepsilon}(P) \in (\hat{\rho}^n, \hat{\rho}^n), P \in S_n$
- $\begin{array}{l} (ii) \quad \hat{\varepsilon}((12...n+1))\hat{T} = \hat{\rho}(\hat{T})\hat{\varepsilon}((12...m+1)), \ \hat{T} \in (\hat{\rho}^n, \hat{\rho}^n) \\ (iii) \ \exists \hat{S} \in (id, \hat{\rho}^d) \ with \ \hat{S}^* \hat{S} = \mathbf{1}, \ \hat{S}^* \hat{\rho}(\hat{S}) = (-1)^{d-1} \frac{1}{d} \mathbf{1} \ and \ \hat{S} \hat{S}^* = \hat{E}_d \end{array}$

(iv) \hat{O} is generated by the intertwiners $\hat{T} \in (\hat{\rho}^n, \hat{\rho}^m), n, m \in \mathbf{N}$

Then there is a closed subgroup $G \in SU(d)$ (unique up to conjugation) and an embedding of \hat{O} in O_d with $\hat{O} \in O_G$, s. t. $\rho \mid_{O_G} = \hat{\rho}, \hat{\varepsilon} = \varepsilon$, and $\hat{S} = S$ hold.

The basic idea on which the proof relies is actually reasonably simple: one takes a kind of amalgamated product \mathcal{B} of O with O_d amalgamated over $O_{SU(d)} \subset$ \hat{O} , i.e. we look for an algebra with the following relations:

- $\psi_i A = \hat{\rho}(A)\psi_i, \quad A \in \hat{O}, \quad \psi_i \in O_d, \ i = 1....d$
- $\hat{\varepsilon}(P) = \varepsilon(P)$
- $\hat{S} = S$

The SU(d) action on \mathcal{B} is:

$$\alpha_g(\psi_i) = \sum_i \psi_i g_{ji}, \quad \alpha_g(A) = A \tag{7.43}$$

For the special case that \hat{O} is generated by $\hat{\varepsilon}(P)$ and \hat{S} , and hence $\hat{O} = O_{SU(d)} \subset$ O_d, \mathcal{B} is obviously O_d , as expected. If \hat{O} is genuinely bigger, there exist intertwiner $\hat{O} \ni T \notin O_{SU(d)}$. The operators $\hat{T}_{(\alpha)(\beta)} := \psi^*_{(\beta)} \hat{T} \psi_{(\alpha)}$ commute with $A \in \hat{O}$ and hence $T_{(\alpha)(\beta)} \in \hat{O} \cap \mathcal{B}$. Actually such operators are automatically in the center of \mathcal{B} , i.e. $\hat{O}' \cap \mathcal{B} = \mathcal{Z}(\mathcal{B})$. This follows from the invariance of $O' \cap \mathcal{B}$ under the action of SU(d) which means that this subalgebra consists of invariant SU(d) tensors F_i . With F_i , i = 1...n being a tensor multiplet, we determine an orthonormal basis ψ_i , i = 1....n and obtain the representation:

$$F_i = B\tilde{\psi}_i, \quad B = \sum_i F_i \tilde{\psi}_i^* \tag{7.44}$$

If we could show that the F_i commute with the generators of the Cuntz algebra, we would be done. But this is an easy computational result of :

$$B \in (\hat{\rho}^n, id), \quad \tilde{\psi}_i \in (id, \rho^n) \tag{7.45}$$

which according to the assumption (ii) of the previous theorem yields:

$$\hat{\rho}(B) = B\hat{\varepsilon}(n+1,...,1)$$

$$\rho(\tilde{\psi}_i) = \varepsilon(1,...,n+1)\tilde{\psi}_i$$
(7.46)

and hence $\psi_j F_i = \psi_j B \tilde{\psi}_i = \hat{\rho}(B) \rho(\tilde{\psi}_i) \psi_j = \dots = \phi_i \psi_j$, Q.E.D..

Let us now return to the problem of construction of the field algebra. A helpful and informative intermediate construction is the introduction of the so-called "reduced field bundle". This is a bimodule over \mathcal{A} which allows to use the $\rho's$ in a direct manner.

Definition 8 As our Hilbert space for the reduced field bundle we take the direct sum of vacuum Hilbert spaces $H_{\alpha} := (\alpha, H_0)$ and define operators F(e, A) with $A \in \mathcal{A}, e = (\rho_{\beta}, \rho, \rho_{\alpha})$ where All the irreducible endomorphisms are taken from a pre-selected set with one endomorphism ρ_{α} per sector $[\rho_{\alpha}]$. We define the action of F as:

$$F(e,A) \cdot (\alpha,\psi) = (\rho_{\beta},\pi_0(T_e^*\rho_a(A)) \cdot \psi)$$
(7.47)

where $T_e \in \mathcal{A}$ are intertwiners from the space $T_e \in (\rho_{\beta}, \rho \rho_{\alpha})$. Here e may be pictured as an edge on a fusion graph whose vertices are set of "charge edges" *i*. e. a triples of source charge $s(e) = \rho_{\alpha}$ range charge $r(e) = \rho_{\beta}$ and transferred charge $c(e) = \rho$.

The F's generate a C^* algebra \mathcal{F}_{red} in $H = \bigoplus_{\alpha} H_{\alpha}$. The commutation relations have the form of an exchange algebra:

$$F(e_{2}, A_{2}) \cdot F(e_{1}, A_{1}) = \sum_{f_{1} \circ f_{2}} R^{e_{2} \circ e_{1}}_{f_{1} \circ f_{2}}(\pm) \cdot F(f_{1}, A_{1}) \cdot F(f_{2}, A_{2})$$
(7.48)
$$R^{e_{2} \circ e_{1}}_{f_{1} \circ f_{2}}(+) = T^{*}_{e_{2}} T^{*}_{e_{1}} \rho_{\alpha}(\varepsilon(\rho_{2}, \rho_{1}) T_{f_{2}} T_{f_{1}}, \quad R(-): \rho_{1} \Leftrightarrow \rho_{2}$$

whenever F_2 is localized in the right /left spacelike complement of F_1 . In low dimensional theories where there is an invariant distinction the R's are related to the braid group whereas in d=3+1 one deals with the special case, the permutation group.

This reduced field bundle only agrees with the field algebra of the standard approach if G is abelian. Its Hilbert space lacks the group theoretic multiplicities incorporated in the formula (7.17) and the net inclusion $\mathcal{A} \subset \mathcal{F}_{red}$ is reducible and its index is the square of the index of the irreducible inclusion $\mathcal{A} \subset \mathcal{F}$ (assuming finite G). For d=3+1 this is the parastatistics⁶ description which deals with higher Young tableaux, but without an internal symmetry group.

 $^{^{6}\}mathrm{Beware}$ that Kadanoff et al. use this terminology in conformal field theory with a different meaning.

The parastatistics fields are more noncommutative and do not allow an interpretation in terms of "quantization" (e.g. they have no Lagrangians). In case of the nonabelian braid group statistics of chiral conformal field theory and d=2+1plektons this is the only available description. In that case the F_{red} -algebra was also called the "exchange algebra" [82].

In case of d=3+1 theories and for the subcategory of permutation group statistics sectors in d<3+1 there exists the famous canonical Doplicher-Roberts construction of a genuine field algebra in the sense of the beginning of this section.

Let us first mention the special case of only automorphisms and assume d=3+1, i.e. permutation group statistics. Since for automorphisms $d_{\rho} = 1$, the permutation group representation is abelian, we are dealing with Bosons/Fermions. In this case some very simple modifications of the reduced field bundle will give the field algebra. The interested reader is referred to [3] page 185, 186. This case may also be obtained by specialization of the construction of the reduced field bundle.

For the above mentioned nonabelian representations one applies the DR theorem [43] about the construction of the group from a subalgebra of the Cuntz algebra with a distinguished endomorphism. Without loss of generality, we may assume that there exists a ρ with statistical dimension d s. t.. $id \subset \rho^d$ which can always be achieved by adding conjugates. A C^* -algebra \hat{O} , as needed in the theorem, may be obtained via the inductive limit of intertwiner spaces:

$${}^{0}\hat{O} := \bigcup_{n,m \ge 0} (\rho^n, \rho^m) \tag{7.49}$$

where the induction uses the "right" embedding $(\rho^n, \rho^m) \to (\rho^{n+1}, \rho^{m+1})$: $T \to T \times 1_{\rho}$. This leads to a natural composition of two operators S and Tby embedding both in a space sufficiently shifted to the right. The algebra contains the statistics operators $\varepsilon(P)$ (still in bosonized form) as well as an isometry $S \in (id, \rho^d)$, $SS^* = E_d$. The endomorphism ρ acts in a natural way on ${}^0\hat{O}$ and $\phi(.) = S^*\rho(.)S$ defines a left inverse of ρ . The properties of $\phi, \rho, \varepsilon(P)$ and S are easily checked by computation. ${}^0\hat{O}$ has a unique C^* -norm and no ideal (i.e. \hat{O} is simple). Now the DR theorem leads to the identification with a subalgebra $O_G \subset O_d$ with a subgroup G of SU(d) which is determined up to conjugation. The field algebra is now simply the free product of the observable algebra \mathcal{A} with the Cuntz algebra O_d amalgamated over the subalgebra O_G with the following relations:

$$\psi_i A = \rho(A)\psi_i, \quad A \in \mathcal{A} \tag{7.50}$$

7.4 Remarks on Broken Symmetries

The idea of spontaneously broken symmetries originated during the 60^{ies} in Lagrangian QFT (Goldstone, Nambu). There were parallel developments in condensed matter physics, in which case the understanding of the phase transitions in the Heisenberg model was the main goal. Already by the end of this

decade there was a general model independent understanding [105] within the framework of QFT's possessing conserved quantum Noether currents independent of their (Lagrangian or non-Lagrangian) origin. The main theorem of this more general approach within the Wightman framework was the relation of the nonexistence of the global charges (as a result of large distance infrared divergencies in the spatial integrals over currents) with the long distance property of the matrix elements of the current operator between certain vector states as a result of the presence of "Goldstone Bosons" in the energy-momentum spectrum. The nice feature of these rigorous methods is that they apply to composite "Goldstones" (i.e. they go beyond the family of Goldstone Lagrangians for which a perturbative approach to the broken phase is possible) as well. In this way the statement became a structural theorem of general QFT.

Algebraic QFT offers an even more profound physical picture which we are going to explain in the sequel. The starting point is the DR reconstruction theory of the previous section. That theory deals with an unbroken symmetry G because only those transformations are in a one to one correspondence with the superselection sectors. Where to look for the bigger spontaneously broken group Γ ? The answer is contained in the breakdown of the vacuum Haag duality of \mathcal{A} [117]. The physical reason for this is that certain operators which, if one only looks at their local properties, carry charges and transform according to Γ -multipletts, globally "condense" into the vacuum sector.

We have met a special case of this phenomenon in connection with the d=1+1 order/disorder discussion in the last section of chapter 3. The main point there was that the original net violated the vacuum Haag Duality and the order/disorder fields were required precisely in order to restore it. By definition we called the field which did not belong to the original vacuum representation, but has a nonvanishing vacuum expectation "disorder". It is the adjunction of this field, which enlarges the observable net \mathcal{A} to the Haag dual net \mathcal{A}^d . Adjusting this to the situation at hand, we assume that our original observable net \mathcal{A} is smaller than its unique dual extension \mathcal{A}^d i.e.:

$$\mathcal{A} \subset \mathcal{A}^d \subseteq \mathcal{F} \tag{7.51}$$

where \mathcal{F} is the unique DR field algebra determined by the superselection theory of \mathcal{A}^d . The DR group G is the unbroken gauge group $\Gamma \supset G$, with Γ defined to be the group of automorphisms of \mathcal{F} which leave \mathcal{A} pointwise fixed. G is the unbroken part of Γ . The following theorem demonstrates the correctness of this interpretation.

Theorem 39 [78]

(i) Each $\gamma \in \Gamma$ leaves $\mathcal{F}(\mathcal{O})$ for each $\mathcal{O} \in \mathcal{K}$ globally stable and is locally normal.

(ii) G is the \mathcal{F} -vacuum stabilizer in Γ

(iii) The normalizer of G in Γ is the invariance subgroup which act automorphically on \mathcal{A}^d

The Goldstone theorem (or better the theorem on the Goldstone mechanism), i.e. the prediction of a special kind of zero mass particle as a result of spontaneous symmetry breaking, only *follows* under more stringent conditions and the *standard situation of a conserved Noether current* is certainly one such possibility. In order to understand better the physics involved, let us look at the vacuum expectation of the derivation defined by the generator δ of the automorphism, using one-parametric subgroup:

$$\delta(F) = \lim_{\lambda \to 0} \lambda^{-1} (\gamma_{\lambda}(F) - F) \tag{7.52}$$

The criteria for a spontaneous symmetry breaking in the general setting of algebraic QFT without or with Goldstone particles are then formulated in terms of behavior of the vacuum expectation of $\delta(F)$ for increasing localization regions.

In low-dimensional QFT the charge sectors associated to an observable net generally cannot be described in terms of group theoretical notions. Lacking a group theoretic characterization of the position of the observable algebra as the fixed points inside the field algebra under a compact group action, one takes the breakdown of Haag duality for double cones in the vacuum representation as the definition of "broken quantum symmetry". A discussion in terms of Noether currents a la Goldstone and Nambu, if possible at all in this case, is still missing.

From Lagrangian field theories one knows another mechanism of symmetry breaking which was first conjectured and exemplified by Schwinger and then brought into a perturbative setting by Higgs. Since it needs the formalism of gauge theory, its intrinsic content has never been spelled out; up to this date there is no rigorously known property of the observable part of the theory which tells us that a massive particle received its mass in such a way. Most of the folklore around this mechanism is not quite correct. For example the idea that the mechanism could be thought of as a "fattened" Goldstone boson is contradicted by the Schwinger model, because in d=1+1 there aren't any Goldstone bosons but there is the Schwinger-Higgs mechanism. In fact different from the previous Goldstone-Nambu mechanism of spontaneously broken symmetries, the terminology "spontaneously broken gauge invariance" is more a mnemotechnical calculational device than a physical meaningful intrinsic property of the theory. However this does not mean that there are no consistent nonperturbative conjectures which have some chance to be proven in algebraic QFT. One appealing idea is to think of a "would be" charged field with a Maxwellian (i.e. very nonlocal) charge ⁷ which is carried by a nonlocal (string-localized) matter field and becomes screened as a consequence of the emergence of a massive vector meson. To argue that such a "phase" of Maxwellian QFT exists was an important contribution of Schwinger. The rigorous theorem about the charge screening \leftrightarrow massive "photon" relation is again due to Swieca [105] In fact in order to show the existence of this mechanism, Schwinger invented the "Schwinger model". In chapter 4.5 we demonstrated, that the closely related opposite interpretation of "charge liberation" is actually the more natural one and that there is no intrinsic physical meaning to (Higgs) condensates. The understand-

⁷Formally a semiinfinite extended object formed from a Dirac field of QED modified by a Mandelstam like A_{μ} -flux to spatial ∞ serves as a candidate for a local gauge invariant, but global U(1) charge carrying field.

7.4. REMARKS ON BROKEN SYMMETRIES

ing of the massless limit of the selfinteracting massive vectormeson is actually the difficult part in perturbation theory, since it requires the transition from local to nonlocal string-like matter field coordinates (in order to get a matter field which allows a massless limit). Only in the Schwinger model this process has been carried out [79]. As a result the semiinfinite string-like charge carrier disappears from the physical stage and the resulting massive theory will be more local (good infrared properties, in the sense of the LSZ mass gap framework)) in the usual sense. There is still a lot of nonunderstood physics hidden behind the deceiving terminology "spontaneously broken gauge symmetries".

The concept of spontaneous symmetry breaking is also very important for thermal QFT. The exploration of the physical consequences is however more subtle than in the case of ground states as a result of the loss of Wigner's particle concept..

All internal and external symmetries with one exception, suffer at most a spontaneous symmetry breaking. The exception is supersymmetry which, as we already mentioned in the introduction, collapses completely. The reason for this radically different behavior is that supersymmetry is an "accidental symmetry". By this we mean a symmetry of a field algebra, which is not visible in the structure of charge sectors of the observable algebra and plays no important role in the understanding of the model. An illustrative example is the tricritical Ising model of chiral conformal field theory. Its observable algebra completely determines all its (finite number of) charge sectors. The fact that one can extend the energy-momentum tensor algebra by fermionic sectors and obtain in this way an action of supersymmetry is only an ornamental attribute and does not make the model any simpler than the other models in the same family of minimal models. The physical content of the model can be explored without taking notice of its supersymmetry. This is a general observation: whenever supersymmetry leaves the twighlight of its folklore and enters the clear air of controllable models (which presently only happens in low-dimensional QFT), it reveals its accidental structure. Too little is known about lowest order gauge invariant supersymmetric gauge theories in order to check whether the claimed properties (scale invariance, short distance compensations,...) are characteristic for supersymmetry.

The modular Tomita-Takesaki theory which, as we have seen, unravels internal (charge sectors) as well as external (Poincaré, conformal) symmetries of the observable algebras, also never points in the direction of supersymmetry. Whereas all such standard symmetries suffer at most a spontaneous breaking in thermal (KMS) states, supersymmetry is unstable under contact with a heat bath⁸; it suffers a spontaneous "collapse" [42], as one expects for an accidental symmetry. All symmetries with noncausal Noether currents (as the fermionic currents of supersymmetry) are potential candidates for accidental symmetries with unstable thermal behavior.

 $^{^8{\}rm The}$ Lorentz boosts applied to the rest frame of the heat bath system are the only spontaneously broken transformations of the Poincaré group.

7.5 Chiral Conformal Algebraic QFT

Chiral conformal QFT has turned out to be an ideal theoretical laboratory for algebraic (nonperturbative) QFT. Not only conformal QFT has profited from this close relation, but the confidence in the algebraic method has also significantly increased. To cite a recent example, within chiral conformal QFT, one was able to rigorously prove the equivalence of the standard approach using pointlike covariant fields with the net approach [80]. This is important because in formulating the net approach one did not intend to widen the physical content, but rather only to put the advanced theory of von Neumann algebras to the use for exploring the physical principles of local QFT.

Since the literature on the subject, even if restricted by the above guideline, is quite formidable, I will limit my attention to two points:

- What is charge structure and quantum symmetry after conformal compactification?
- How does one classify chiral conformal QFT?

The compactification of chiral conformal QFT is most efficiently done in terms of a universal C*-algebra $\mathcal{A}_{uni}(S^1)$ which is different from the noncompact DHR quasilocal algebra $\mathcal{A}(\mathbf{R})$. In order to understands its construction, we note that the net $\{\mathcal{A}(I)\}_{I \subset S^1}$ is not directed (as the nets of double cones in Minkowski space) towards infinity. Therefore we should think of a globalization which is different from the inductive limit. For this we use the following definition universal algebra \mathcal{A}_{univ} :

Definition 9 \mathcal{A}_{univ} is the C^* algebra which is uniquely determined by the system of local algebras $(A(I))_{I \in \mathcal{T}}$, $\mathcal{T} = family$ of proper intervals $I \subset S^1$ and the following universality condition:

(i) there are unital embeddings $i^{I} : \mathcal{A}(I) \to \mathcal{A}_{univ}$ s. t..

$$i^{J}|_{\mathcal{A}(I)} = i^{I} \quad if \quad I \subset J, \ I, J \in \mathcal{T}$$

$$(7.53)$$

and \mathcal{A}_{univ} is generated by the algebras $i^{I}(\mathcal{A}(I)), \ I \in \mathcal{T};$

(ii) for every coherent family of representations $\pi^{I} : \mathcal{A}(I) \to \mathcal{B}(H_{\pi})$ there is a unique representation π of \mathcal{A}_{univ} in H_{π} s. t..

$$\pi \circ i^I = \pi^I \tag{7.54}$$

The universal algebra inherits the action of the Möbius group as well as the notion of positive energy representation through the embedding.

The universal algebra has more global elements than the quasilocal algebra of the DHR theory: $\mathcal{A}_{quasi} \equiv \mathcal{A} \subset A_{univ}$ with the consequence that the vacuum representation π_0 ceases to be faithful and the global superselection charge operators which are outer for \mathcal{A} become inner for \mathcal{A}_{univ} . From this observation emerges the algebra of Verlinde which originally was obtained by geometric rather than local quantum physics arguments. The removal of a point ξ from S^1 (this removal recreates the infinity of \mathcal{A}_{quasi}) forces \mathcal{A}_{univ} to shrink to \mathcal{A} .

Most of this new features can be seen by studying global intertwiners in \mathcal{A}_{univ} . Let $I, J \in T$ and $\xi, \zeta \in I' \cap J'$ (i.e. two points removed from the complements) and choose ρ and σ s. t.. $\operatorname{loc}\rho$, $\operatorname{loc}\sigma \subset I$ and $\hat{\rho} \in [\rho]$ with $\operatorname{loc}\hat{\rho} \subset J$. Then the statistics operators $\varepsilon(\rho, \sigma)$ and $\varepsilon(\sigma, \rho) \in \mathcal{A}(I) \subset A_{\xi} \cap A_{\zeta}$ are the same (i.e. they don't need a label ξ or ζ) independently of whether we use the quasilocal algebra \mathcal{A}_{ξ} or \mathcal{A}_{ζ} for their definition. By Haag duality a charge transporter $V : \pi_0 \rho \to \pi_0 \hat{\rho}$ lies both in $\pi_0(\mathcal{A}_{\xi})$ and $\pi_0(\mathcal{A}_{\zeta})$. However its pre-images with respect to the embedding are different. In fact:

$$V_{\rho} \equiv V_{+}^{*}V_{-} \text{ with } V_{+} \in \mathcal{A}_{\xi}, \quad V_{-} \in \mathcal{A}_{\zeta}$$

$$V_{\rho} \in (\rho, \rho)_{alob}$$

$$(7.55)$$

is a global selfintertwiner, which is easily shown to be independent of the choice of V and $\hat{\rho}$. The representation of the statistics operators in terms of the charge transporters $\varepsilon(\rho, \sigma) = \sigma(V_+)^* V_+$, $\varepsilon(\sigma, \rho)^* = \sigma(V_-)^* V_-$ leads to:

$$\sigma(V_{\rho}) = \varepsilon(\rho, \sigma) V_{\rho} \varepsilon(\sigma, \rho) \quad \curvearrowright \pi_0 \sigma(V_{\rho}) = \pi_0 \left[\varepsilon(\rho, \sigma) \varepsilon(\sigma, \rho) \right]$$
(7.56)

The first identity is very different from the relation between $\varepsilon's$ due to local intertwiners. The global intertwiner V_{ρ} is trivial in the vacuum representation, thus showing its lack of faithfulness with respect to \mathcal{A}_{univ} . The global aspect of V_{ρ} is only activated in charged representations where it coalesces with monodromy operators. From its definition it is clear that it represents a charge transport once around the circle S^{19} . As a result of its existence, the monodromy which is defined as the above two-fold iteration of the braid generator, takes on some of its geometric meaning which it has e.g. in the theory of complex functions. The left hand side of the first equation in (7.56) expresses a transport "around" in the presence of another charge σ , i.e. a kind of charge polarization. Let us look at the invariant version of V_{ρ} namely the global "Casimir" operators $W_{\rho} = R_{\rho}^* V_{\rho} R_{\rho} : id \to id$. This operator lies in the center $\mathcal{A}_{univ} \cap \mathcal{A}'_{univ}$ and depend only on the class (=sector) $[\rho]$ of ρ . By explicit computation[89] one shows that after the numerical renormalization $C_{\rho} := d_{\rho} W_{\rho}$ one encounters the fusion algebra:

(i)
$$C_{\sigma\rho} = C_{\sigma} \cdot C_{\rho}$$
 (7.57)
(ii) $C_{\rho}^{*} = C_{\overline{\rho}}$
(iii) $C_{\rho} = \sum_{\alpha} N^{\alpha} C_{\alpha}$ if $\rho \simeq \oplus_{\alpha} N^{\alpha} \rho_{\alpha}$

Verlinde's modular algebra emerges upon forming matrices with row index equal to the label of the central charge and the column index to that of the sector in

⁹Note that in A_{univ} which corresponds to a compact quantum world it is not possible to "dump" unwanted charges to "infinity" (as in the case for A_{quasi}), but instead one encounters "polarization" effects upon charge transportation once around.

which it is measured:

$$S_{\rho\sigma} := \left| \sum_{\gamma} d_{\gamma}^2 \right|^{-\frac{1}{2}} d_{\rho} d_{\sigma} \cdot \pi_0 \sigma(W_{\rho}) \tag{7.58}$$

In case of nondegeneracy of sectors, which expressed in terms of statistical dimensions and phases means $\left|\sum_{\rho} \kappa_{\rho} d_{\rho}^{2}\right|^{2} = \sum_{\rho} d_{\rho}^{2}$, the above matrix S is equal to Verlinde's matrix S [118]which together with the diagonal matrix $T = \kappa^{-1} Diag(\kappa_{\rho})$, with $\kappa^{3} = (\sum_{\rho} \kappa_{\rho} d_{\rho}^{2}) / \left|\sum_{\rho} \kappa_{\rho} d_{\rho}^{2}\right|$ satisfies the modular equations of the genus 1 mapping class group

$$SS^{\dagger} = 1 = TT^{\dagger}, \quad TSTST = S$$

$$S^{2} = C, \quad C_{\rho\sigma} \equiv \delta_{\bar{\rho}\sigma}$$

$$TC = CT = T$$

$$(7.59)$$

It is remarkable that these properties are common to chiral conformal theories and to d=2+1 plektonic models even though the localization properties of the charge-carrying fields are quite different. In the former case one has the additional phase relation:

$$\frac{\kappa}{|\kappa|} = e^{-2\pi i c/8} \tag{7.60}$$

where c is the Virasoro constant which measures the strength of the two-point function of the energy-momentum tensor. This relation may be derived by studying the (modular) transformation properties of the Gibbs partition functions for the compact Hamiltonian L_0 of the conformal rotations under thermal duality transformations $\beta \rightarrow 1/\beta$. For d=2+1 plektons, no simple physical interpretation is known.

Lemma 40 The matrix S is similar to the character matrix in section 2 of the first chapter. However in distinction to nonabelian finite groups (which also yield a finite set of charge sectors of the fixed point observable algebra) the present nonabelian sectors produce a symmetric "character" matrix S which signals a perfect auto-duality between charge measurers $\{Q\}$ and charge creators $\{\rho\}$. Furthermore the algebra Q generated by the central charges and the action of the endomorphisms on those charges¹⁰ do not contain the old "group theoretical stuff" since the phenomenon of charge "polarization" only perceives endomorphisms with nontrivial monodromy.

This strongly suggests to try to understand the new "quantum symmetry" property in terms of the structural properties of Q. As a generalization of S one finds for the Q's in the presence of more than one polarization charges the entries of the higher genus mapping class group matrices [74]. The reason is that in addition to the the process:

$$vacuum \xrightarrow{split} \rho\bar{\rho} \xrightarrow[selfintertw.]{global} \rho \xrightarrow{\rho} \xrightarrow{fusion} vacuum$$
(7.61)

¹⁰This action leads out of the center and generates a global subalgebra of \mathcal{A}_{univ} .

which led to the global intertwiner $W_{\rho} = R_{\rho}^* V_{\rho} R_{\rho}$, there is the more involved global intertwiner associated with the process in which the global selfintertwining occurs after a split of nonvacuum charge σ and a later fusion to μ which appear in a $\rho\bar{\rho}$ reduction:

$$\sigma \xrightarrow{split} \alpha \beta \xrightarrow{global \alpha}_{intertw.} \alpha \beta \xrightarrow{fusion} \mu, \quad \sigma, \mu \subset \rho \bar{\rho}$$

$$(7.62)$$

with the global intertwiner $V_{\alpha} \in (\alpha, \alpha)_{glob}$ being used in: $T_{e(\sigma)}^* V_{\alpha} T_{e(\mu)}$ where $T_{e(\mu)}$ is the $\alpha\beta \to \mu$ fusion intertwiner and the hermitian adjoints represent the corresponding splitting intertwiner. As in the vacuum case, the selfintertwiners V become only activated after the application of another endomorphism say η , i.e. in the presence of another charge η (hence the name "polarization" mechanism). It can be shown that the following operators are the building blocks of the mapping class group matrices $T_{e(\sigma)}^* V_{\alpha} T_{e(\mu)}$ which have multichargemeasurer column and multicharge-creator row indices:

$$\phi_{\lambda}((T_{g(\eta)}\eta(T_{e(\sigma)}^*V_{\alpha}T_{e(\mu)})T_{f(\eta)}^*))) \tag{7.63}$$

. here $T_{f(\eta)}$ and $T_{g(\eta)}$ are the intertwiners corresponding to the charge edges $f(\eta) : \lambda \sigma \to \eta$ and $g(\eta) : \lambda \mu \to \eta$, whereas ϕ_{λ} is the left inverse of the endomorphism λ . Besides the global intertwiners V, we only used the local splitting intertwiners and their hermitian adjoints which represent the fusion intertwiners. The main question is: why do we organize the numerical data of the global charge-measurer and charge-creator algebra Q as entries in a multiindex matrix? What is the physical role of these matrices in a d=2+1 plektonic theory?

Closely related to these structures are the knot theoretical invariants of 3manifolds. These objects also appear by analyzing certain formal functional integrals with the hindsight of geometry and topology [77]. But in the context of algebraic QFT, the physical interpretation is quite different because the new properties have nothing to do with the "living space" (in the sense of quantum theoretical localization) of fields or algebras, but are rather manifestations of the inexorable link between external (space-time) and internal symmetries which one encounters in low dimensional real time (Minkowski space) QFT. It appears that they generalize in some sense the angular momentum decompositions and one would expect them to play a useful role in the understanding of e.g. the analysis of scattering of d=2+1 plektons. Although these ideas of linking "quantum symmetry" with a kind of universal mapping class group [74] (containing all genii) are highly seductive, I did not yet find an convincing argument for why one should read the numerical aspects of those polarization charges as entries of mapping class matrices acting on "something". This open problem is closely related to the previously asked questions.

It should be mentioned here that most attempts in the direction of quantum symmetry have been aimed towards modified ("weak"...) Hopf algebras, thus remaining near the spirit of the DR theory [81]. None of these attempts was yet successful. One expects from a useful quantum symmetry concept a clarification of the following two points:

- A better understanding why in low dimensions the link between external/internal symmetries is so strong, whereas in d=3+1 there was no possibility to bring them together in any nontrivial way. This aims in particular at a better physical understanding of the physical interpretation of knot- and 3-manifold- invariants.
- A simplification of the problem of computing correlations of "free" plektons, i.e. the freest objects (in d=2+1 preferably with vanishing cross sections) which fulfill the new braid group statistics. Since even the free plektonic charges have an analytically more complicated spacetime structure as a result of their semiinfinite localization, a symmetry concept which does not split internal/external aspects is expected to have a better chance to be useful for their understanding.

Concerning the classification of chiral conformal QFT's, it is reasonable to approach this problem in two steps:

- Classification of the physically admissable braid group representations which go with the category of finitely many localizable sectors ("rational" representations).
- Construction of representative 4-point functions for the different plektonic families.

The basic techniques for this two step approach is quite old [82] and have been elaborated for the unitary braid group representations affiliated with the special family of the Jones, Temperly-Lieb algebras. The more general representations are those affiliated with the Hecke algebra and with the Birman-Wenzl algebra explained in the next section. Even if one does not know anything about these mathematical construction, the concepts of algebraic field theory are so strong that they will lead us there.

Before we present the known (and presumably complete) set of families, it is helpful to notice the changes in the structure of the exchange algebra which result from the compactification through the \mathcal{A}_{uni} -globalization. The main difference to the standard exchange algebra associated with \mathcal{A}_{quasi} lies in the concept of localization. The criterion of: $locF(e, A) \subset \mathcal{O} \Leftrightarrow F$ commutes with all observables $B \in \mathcal{A}(\mathcal{O}')$, or in the notation of the reduced field bundle formalism:

$$\pi_{\beta}(B)F(e,A) = F(e,A)\pi_{\alpha}(B), \ c(e) = \rho, \ s(e) = \alpha, \ r(e) = \beta$$
(7.64)

where we used the previously introduced notation concerning edges e on a fusion graph consisting of the source charge α , the range charge β and the charge ρ transported by F. Written more explicitly, this commutation relation is equivalent to the existence of a local unitary charge transporter U which simultaneously transports ρ and A into \mathcal{O} : $locAd_U \circ \rho \subset \mathcal{O}$, $UA \in \mathcal{A}(\mathcal{O})$. Hence the localization in \mathcal{A}_{quasi} depends only on the pair (ρ, A) . For the \mathcal{A}_{uni} localization this characterization is too rough, because it ignores the possibility of carrying charges several times around the circle S^1 . We refine the definition as follows:
Definition 10 Let $J \subset \mathbf{R}$ be an intervall (of extension $< 2\pi$) of the universal covering \mathbf{R} of S^1 . For $\rho \in \Delta_{red}$ the pair (ρ, A) is said to be localized in J if there is a local operator $C \in \mathcal{A}(I)$ s. t. all F(e, A) with charge ρ are obtained from operators of the form F(e, C) through Moebius-transformations. Here I is an interval in the first sheet of R (i.e. inside the 2π interval which includes the zero).

The following theorem shows how the new localization concept adapted to the observable algebra \mathcal{A}_{uni} changes the structure of the exchange algebra in section 7.3.

Theorem 41 Let (ρ_1, A_1) and (ρ_2, A_2) be localized in intervals J_1 and J_2 on **R** which project onto disjoint intervals in S^1 . Define a relative winding number N s.t. $J_1 + 2\pi N < J_2 < J_1 + 2\pi (N+1)$. Then:

$$\begin{split} F(e_2, A_2) \cdot F(e_1, A_1) &= \sum_{f_1 \circ f_2} R_{f_1 \circ f_2}^{e_2 \circ e_1}(N) \cdot F(f_1, A_1) \cdot F(f_2, A_2) \\ R_{f_1 \circ f_2}^{e_2 \circ e_1}(N) &= T_{e_2}^* T_{e_1}^* \rho_\alpha(\varepsilon_N(\rho_2, \rho_1) T_{f_2} T_{f_1} \\ \varepsilon_N(\rho_2, \rho_2) &= \rho_1(Y_{\rho_2}^N) \varepsilon(\rho_2, \rho_1) Y_{\rho_2}^{-N}, \quad Y_\rho = e^{2\pi i h_\rho} V_\rho \end{split}$$

Here V_{ρ} is the selfintertwiner (7.55). The $\varepsilon'_N s$ are associated with the cylinder ribbon braid group[89].

Conformal QFT are in some way sophisticated free theories, where the sophistication refers to their charge structure. Physically there can be no genuine interaction on one light cone and mathematically one expects the braid group statistics together with some gross features of the charge structure to fix the observable algebras and their associated charge carrying operators ¹¹. The simplest family of plektonic chiral conformal theories are the so-called minimal theories; here minimal is used in the sense of the smallest statistical dimensions.

The knowledge about the admissible braid group representations and in particular about the statistical dimensions together with a bit of hindsight on short distance behavior allows in many cases the determination of plektonic 4-point functions [82]. There are other artistic methods to fix the 4-point functions through there monodromy properties by an ad hoc formal modification of the Coulomb-gas and contour representations. Since we are not interested in conformal theory per se, but only in its aspects of presenting a theoretical laboratory for LQP, we will not pursue this matter. A potentially successful method in the spirit of these notes has been sketched in chapter 6.8.

7.6 Classification of Admissable Particle-Statistics

The DHR method of classifying the admissable permutation group S_{∞} representations by defining sequences of projectors which are terminating after finitely

 $^{^{11}{\}rm This}$ is analogous to the determination of d=3+1 free fields with a given internal symmetry (a given charge structure).

many terms (the consequence of positivity) was extended by Ocneanu and Wenzl in such a way that it became a powerful tool for large classes of families of subfactor models. The formalism is applicable to the physically admissable braid-group representations in low-dimensional QFT and leads to the famous Markov-trace formalism on the ribbon braid group RB_{∞} . There are plausibility arguments (but no proof) that apart from exceptional cases, the Markov traces on the Hecke algebras and the Birman-Wenzl-Murakami algebras exhaust all possibilities. The physical origin of the Hecke algebra is "two-channel plektonic statistics" and its composites. This means that we are studying plektonic endomorphisms ρ with $\rho^2 \simeq \rho_1 \oplus \rho_2$ i.e. $d_{\rho}^2 = d_{\rho_1} + d_{\rho_2}$ with ρ, ρ_1, ρ_2 irreducible as well as all its higher composites ("fusion") [93]. The BWM-algebras result from plektonic 3-channel endomorphisms $\rho^2\simeq\rho_1\oplus\rho_2\oplus\alpha$ with α an automorphism i.e. $d_{\rho}^2 = d_{\rho_1} + d_{\rho_2} + 1$. Structural arguments based on the 4-point function suggest that each braid group representation family has two field theoretic realizations, one in which the associated observable algebra contains fields which transform under an internal symmetry group (the current or Kac-Moody algebras) and the other without such group theoretic multiplicities. In the following we present the arguments for this claims.

Our main tool is the application of the (iterated) left inverse ϕ on the intertwiner algebras. In this way one obtains tracial states on those algebras. Left inverses and some of its properties were already introduced in section 3. Their use in this analysis is synonymous with the more physical notion of "conjugate" or antiparticle. The latter is a particle with the same Poincaré-group representation but "opposite" internal symmetry behavior i.e. with possibility to annihilate into a state with the quantum numbers of the vacuum. A sufficient condition for the existence of antiparticles is the spectral gap (isolated one particle hyperboloids) which is also the standard assumption of scattering theory. Let us briefly remind ourselves that translated into the setting of endomorphisms of algebraic QFT this means $\bar{\rho}\rho \supset id$ i.e. the existence of an localized intertwiner R with:

$$R \cdot id(A) = \bar{\rho}\rho(A) \cdot R, \qquad A \in \mathcal{A} \tag{7.65}$$

where as a result of the localization of R the global algebra \mathcal{A} is either the quasilocal or the bigger universal algebra. Then:

$$\phi(A) := R^* \bar{\rho}(A) R \tag{7.66}$$

is a (unique for ρ irreducible) left inverse of ρ i.e. a positive linear map with $\phi(\rho(A)B\rho(C)) = A\phi(B)C$ for $A, B, C \in \mathcal{A}$. The complex number $\lambda_{\rho}\underline{1} = \phi(\varepsilon_{\rho})$ (for irreducible ρ) is called the statistics parameter and it is written as $\lambda_{\rho} = \frac{1}{d_{\rho}}\kappa_{\rho}$ with d_{ρ} the statistical dimension and $\kappa_{\rho} = e^{2\pi i h_{\rho}}$ the statistical phase. We note in passing that the spin-statistics theorem relates this to the spin-phase (in conformal QFT h_{ρ} is related to the scale-dimension). Under this assumption of irreducibility of ρ (always assumed in the rest of this section) ϕ maps the commutant of $\rho^2(\mathcal{A})$ in \mathcal{A} into the complex numbers:

$$\phi(A) = \varphi(A)\underline{1}, \quad A \in \rho^2(\mathcal{A})' \tag{7.67}$$

and by iteration a faithful tracial state φ on $\cup_n \rho^n(\mathcal{A})'$ with:

$$\phi^{n}(A) = \varphi(A)\underline{1}, \quad A \in \rho^{n+1}(\mathcal{A})'$$

$$\varphi(AB) = \varphi(BA), \quad \varphi(\underline{1}) = 1$$

Restricted to the $\mathbf{C}RB_n$ algebra generated by the ribbon braid-group which is a subalgebra of $\rho^n(\mathcal{A})'$ the φ becomes a tracial state, which can be naturally extended $(B_n \subset B_{n+1})$ to $\mathbf{C}RB_\infty$ in the above manner and fulfills the "Markovproperty":

$$\varphi(a\sigma_{n+1}) = \lambda_{\rho}\varphi(a), \quad a \in \mathbf{C}RB_n \tag{7.68}$$

The terminology is that of V. Jones and refers to the famous russian probabilist of the last century as well as to his son, who constructed knot invariants from suitable functionals on the braid group. The "ribbon" aspect refers to an additional generator τ_i which represents the vertical 2π rotation of the cylinder braid group (\simeq projective representation of B_n)[89].

It is interesting to note that the Markov-property is the combinatorial relict of the cluster property which relates the n-point correlation function in local QFT to the n-1 point correlation or in QM the physics of n particles to that of n-1 (rendering one particle a spectator by removing it to infinity. The infinite permutation- and braid groups are the only groups behaving like a russian "matrushka" i.e. the smaller ones are naturally contained in the bigger. This picture is similar to that of cluster properties which was already used in our attempts to understand statics in the nonrelativistic setting of the first chapter. The existence of a Markov trace on the ribbon braid group of (low dimensional) multi-particle statistics is the imprint of the cluster property on particle statistics. As such it is more basic than the notion of internal symmetry. It precedes the latter and according to the DR theory it may be viewed as the other side of the same coin on which one side is the old (compact group-) or new (quantum-) symmetry. With these remarks the notion of internal symmetry becomes significantly demystified.

Let us now return to the above 2-channel situation. Clearly ε_{ρ} has maximally two different eigenprojectors since otherwise there would be more than two irreducible components of ρ^2 . On the other hand ε_{ρ} cannot be a multiple of the identity because ρ^2 is not irreducible. Therefore ε_{ρ} has exactly two different eigenvalues λ_1, λ_2 i.e.

$$(\varepsilon_{\rho} - \lambda_1 \underline{1})(\varepsilon_{\rho} - \lambda_2 \underline{1}) = 0 \tag{7.69}$$

$$\leftrightarrow \varepsilon_{\rho} = \lambda_1 E_1 + \lambda_2 E_2 , \quad E_i = (\lambda_i - \lambda_j)^{-1} (\varepsilon_{\rho} - \lambda_j), \quad i \neq j$$
 (7.70)

which after the trivial re-normalization of the unitaries $g_k := -\lambda_2^{-1} \rho^{k-1}(\varepsilon_{\rho})$ yields the generators of the Hecke algebra:

$$g_{k}g_{k+1}g_{k} = g_{k+1}g_{k}g_{k+1}$$

$$g_{k}g_{l} = g_{l}g_{k} , \quad |j-k| \ge 2$$

$$g_{k}^{2} = (t-1)g_{k} + t , \quad t = -\frac{\lambda_{1}}{\lambda_{2}} \neq -1$$
(7.71)

The physical cluster property in the algebraic form of the existence of a tracial Markov state leads to a very interesting "quantization" ¹². Consider the sequence of projectors:

$$E_i^{(n)} := E_i \wedge \rho(E_i) \wedge \dots \wedge \rho^{n-2}(E_i) , \quad i = 1, 2$$
(7.72)

and the symbol \wedge denotes the projection on the intersection of the corresponding subspaces. The notation is reminiscent of the totally antisymmetric spaces in the case of Fermions. The above relation $g_1g_2g_1 = g_2g_1g_2$ and $g_1g_n = g_ng_1$, $n \geq 2$ in terms of the E_i reads:

$$E_{i}\rho(E_{i})E_{i} - \tau E_{i} = \rho(E_{i})E_{i}\rho(E_{i}) - \tau\rho(E_{i}) , \quad \tau = \frac{t}{(1+t)^{2}} \quad (7.73)$$
$$E_{i}\rho^{n}(E_{i}) = \rho^{n}(E_{i})E_{i} , \quad n \ge 2$$

The derivation of these equations from the Hecke algebra structure is straightforward. The following recursion relation is however tricky and will be given in the sequel.

Proposition 42 (Wenzl, DHR) The projectors $E_i^{(n)}$ fulfill the following recursion relation $(t = e^{2\pi i \alpha}, -\frac{\pi}{2} < \alpha < \frac{\pi}{2})$:

$$E_{i}^{(n+1)} = \rho(E_{i}^{(n)}) - \frac{2\cos\alpha\sin n\alpha}{\sin(n+1)\alpha}\rho(E_{i}^{(n)})E_{j}\rho(E_{i}^{(n)}) , \quad i \neq j, \quad n+1 < q(7.74)$$
$$E_{i}^{(q)} = \rho(E_{i}^{(q-1)}) \quad , \quad q = \inf\{n \in \mathbf{N}, n \mid \alpha \mid \ge \pi\} \quad if \; \alpha \neq 0, \; q = \infty \; if \; \alpha = 0$$

The DHR recursion for the permutation group S_{∞} is obtained for the special case t=0 i.e. $\alpha = 0$. In this case the numerical factor in front of product of three operators is $\frac{n}{n+1}$.

The proof is by induction. For n=1 the relation reduces to the completeness relation between the two spectral projetors of ε_{ρ} : $E_i = 1 - E_j$, $i \neq j$. For the induction we introduce the abbreviation $F = E_j \rho(E_i^{(n)}) = \rho(E_i^{(n)})E_j$ and compute F^2 . We replace the first factor $\rho(E_i^{(n)})$ according to the induction hypothesis by:

$$\rho(E_i^{(n)}) = \rho^2(E_i^{(n-1)}) - \frac{2\cos\alpha\sin(n-1)\alpha}{\sin n\alpha}\rho^2(E_i^{(n-1)})\rho(E_j)\rho^2(E_i^{(n-1)}) \quad (7.75)$$

We use that the projector $\rho^2(E_i^{(n-1)})$ commutes with the algebra $\rho^2(\mathcal{A})'$ (and therefore with $E_j \in \rho^{(2)}(\mathcal{A})'$), and that its range contains that of $\rho(E_i^{(n)})$ i.e. $\rho^2(E_i^{(n-1)})\rho(E_i^{(n)}) = \rho(E_i^{(n)})$. Hence we find:

$$\frac{F^2 = E_j \rho(E_i^{(n)}) - \frac{2\cos\alpha\sin(n-1)\alpha}{\sin n\alpha} \rho^2(E_i^{(n-1)}) E_j \rho(E_j) E_j \rho(E_i^{(n)}) \quad (7.76)$$

292

 $^{^{12}}$ In these notes we use this concept always in the original meaning of Planck as a discretization, and not in the modern form of a deformation.

Application of (7.73) with $\tau = \frac{1}{2 \cos \alpha}$ to the right-hand side yields:

$$F^{2} = E_{j}\rho(E_{i}^{(n)}) - \frac{\sin(n-1)\alpha}{2\cos\alpha\sin\alpha}\rho^{2}(E_{i}^{(n-1)})E_{j}\rho(E_{i}^{(n)}) = \frac{\sin(n+1)\alpha}{2\cos\alpha\sin\alpha\alpha}F$$
 (7.77)

where we used again the above range property and a trigonometric identity. For n = q-1 the positivity of the numerical factor fails and by $F^2E_j = (FF^*)^2$ and $FE_j = FF^*$ the operator F must vanish and hence E_j is orthogonal to $\rho(E_j^{(q-1)})$ which is the second relation in (7.74). For n < q-1 the right-hand side of the first relation in (7.74) with the help of (7.77) turns out to be a projector which vanishes after multiplication from the right with $\rho^k(E_j), k = 1, ..., n-2$ as well as with E_j . The remaining argument uses the fact that this projector is the largest with this orthogonality property and therefore equal to $E_i^{(n+1)}$ by definition of $E_i^{(n+1)}$ Q.E.D..

The recursion relation (7.74) leads to the desired quantization after application of the left inverse ϕ :

$$\phi(E_i^{(n+1)} = E_i^{(n)} \left(1 - \frac{2\cos\alpha\sin n\alpha}{\sin(n+1)\alpha} \eta_j \right), \quad i \neq j$$

$$\eta_j = \phi(E_j), \ 0 \le \eta_j \le 1, \ \eta_1 + \eta_2 = 1$$
(7.78)

From this formula one immediately recovers the permutation group DHR quantization for $\alpha = 0$. In that case positivity of the bracket restricts η_j to the values $\frac{1}{2}(1 \pm \frac{1}{d}), d \in \mathbf{N} \cup 0$. For $\alpha \neq 0$ one first notes that from the second equation (7.74) one obtains (application of ϕ):

$$\eta_j E_i^{(q-1)} = \phi(E_j \rho(E_i^{(q-1)})) = \phi(E_j E_i^{(q)}) = 0, \quad i \neq j$$
(7.79)

where the vanishing results from the orthogonality of the projectors. Since $\eta_1 + \eta_2 = 1$ we must have $E_i^{(q-1)} = 0$ for i=1,2, $q \ge 4$, because $E_i^{(q-1)} \ne 0$ would imply $\eta_j = 0$ and $E_j^{(q-1)} = 0$. This in turn leads to $E_j \equiv E_j^{(2)} = 0$ which contradicts the assumption that ε_{ρ} processes two different eigenvalues. This is obvious for q = 3 and follows for q > 3 from the positivity of ϕ (7.78) for n=2:

$$\phi(E_j^{(3)}) = -\frac{\sin\alpha}{\sin 3\alpha} E_j^{(2)} \quad \curvearrowright E_j^{(2)} = 0 \quad \curvearrowright E_i^{(q-1)} = 0, \ i = 1, 2, \ q \ge 4$$
(7.80)

Using (7.78) iteratively in order to descend in n starting from n = q-2, positivity demands that there exists an $k_i \in \mathbb{N}, 2 \leq k_i \leq q-2$, with:

$$\eta_i = \frac{\sin(k_i + 1)\alpha}{2\cos\alpha\sin k_i\alpha}, \ i = 1, 2 \quad \curvearrowright \sin(k_1 + k_2)\alpha = 0 \tag{7.81}$$

where the relation results from summation over *i*. Since the only solutions are $\alpha = \pm \frac{\pi}{q}$, $k_1 = d$, $k_2 = q - d$, $d \in N$, $2 \leq d \leq q - 2$, one finds for the statics parameters of the plektonic 2-channel family the value:

$$\lambda_{\rho} = \sum_{i=1}^{2} \lambda_{i} \eta_{i} = -\lambda_{2} \left[(t+1) \eta_{1} - 1 \right] = -\lambda_{2} e^{\pm \pi i (d+1)/q} \frac{\sin \pi/q}{\sin d\pi/q}$$
(7.82)

a formula which allows for a nice graphical representation. We have established the following theorem:

Theorem 43 . Let ρ be an irreducible localized endmorphism such that ρ^2 has exactly two irreducible subrepresentations. Then:

• ε_{ρ} has two different eigenvalues λ_1, λ_2 with ratio

$$\frac{\lambda_1}{\lambda_2} = -e^{\pm 2\pi i/q}, \quad q \in \mathbf{N} \cup \{\infty\}, \, q \ge 4 \tag{7.83}$$

• The modulus of the statistics parameter $\lambda_{\rho} = \phi(\varepsilon_{\rho})$ has the possible values

$$|\lambda_{\rho}| = \begin{cases} \frac{\sin \pi/q}{\sin d\pi/q}, & q < \infty \\ \frac{1}{d}, 0 & q = \infty \end{cases}, d \in N, \ 2 \le d \le q - 2 \tag{7.84}$$

• The representation $\varepsilon_{\rho}^{(n)}$ of the braid group B_n which is generated by $\rho^{(k-1)}(\varepsilon_{\rho}), k = 1, ..., n-1$ in the vacuum Hilbert space is an infinite multiple of the Ocneanu-Wenzl representation tensored with a one dimensional (abelian) representation. The projectors $E_2^{(m)}$ and $E_1^{(m)}$ are "cutoff" (vanish) for $d < m \le n$ and $q - d < m \le n$

respectively.

• The iterated left inverse $\varphi = \phi^n$ defines a Markov trace tr on B_n :

$$tr(b) = \varphi \circ \varepsilon_{\rho}(b) \tag{7.85}$$

The "elementary" representation which is characterized by two numbers d and q gives rise to a host of composite representation which appear if one fuses the ρ , ρ_1 , ρ_2 and reduces etc. We will not present the associated composite braid formalism.

The problem of 3-channel braid group statistics has also been solved with the projector method in case that one of the resulting channels is an automorphism τ :

$$\rho^2 = \rho_1 \oplus \rho_2 \oplus \tau \tag{7.86}$$

In that case ε_{ρ} has 3 eigenvalues λ_i which we assume to be different:

$$(\varepsilon_{\rho} - \lambda_1)(\varepsilon_{\rho} - \lambda_2)(\varepsilon_{\rho} - \lambda_3) = 0 \tag{7.87}$$

The relation to the statistics phases ω_{ρ} , ω_i is the following: $\mu_i^2 = \frac{\omega_i}{\omega^2}$. In addition to the previous operators $G_i = \rho^{i-1}(\varepsilon_{\rho}) = (G_i^{-1})^*$ we define projectors:

$$E_i = \rho^{i-1} (TT^*)$$

294

where $T \in (\rho^2 | \tau)$ is an isometry and hence E_i the projector onto the eigenvalue $\lambda_3 = \lambda_{\tau}$ of G_i . In fact one finds the following relations between the G_i and E_i :

$$E_{i} = \frac{\mu_{3}}{(\mu_{3} - \mu_{1})(\mu_{3} - \mu_{2})} (G_{i} - (\mu_{1} + \mu_{2}) + \mu_{1}\mu_{2}G_{i}^{-1})$$
(7.88)
$$E_{i}G_{i} = \mu_{3}E_{i}$$

This together with the trilinear relations between the $G'_i s$ and $E'_i s$ as well as the commutation of neighbors with distance ≥ 2 gives upon renormalization the operators g_i and e_i which fulfill the defining relation of the Birman-Wenzl algebra which again depends on two parameters. The Markov tracial state classification again leads to a quantization of these parameters except for a continuous one-parameter solution with statistical dimension d = 2 which is realized in conformal QFT as sectors on the fixed point algebra of the U(1) current algebra (which has a continuous one-parameter solution) under the action of the charge conjugation transformation (called "orbifolds" by people who prefer geometrical pictures to physical concepts).

There exists another method which allows to construct Markov-traces on the braid group B_{∞} which is a deformation theory of groups. It yields a kind of Hopf-algebra called q-deformed group or "Quantum Group" (beware: here the word "quantum" does not have the physical meaning given by Planck and Heisenberg!). This deformation theory carries the tracial states on the centralizer algebras of tensor representations of compact ie-groups into Markov-traces on the braid group. If the value of the deformation parameter approaches a unit root, these Markov-traces go smoothly over into the above Markov tracial states on the braid group algebra whereas the "quantum group" suffers a nonanalytic behavior.. This is a useful technique which allows to obtain a rather rich family of physically admissable B_{∞} representations by more familiar Lie-group methods which avoids the systematic DHR multi-channel classification. This is a very fine method as long as one does not confuse the q-deformation technique with the new quantum symmetry.

It is an interesting question, to what extend the combinatorial data of the hyperfinite type II_1 intertwiner algebra fixes the chiral conformal field theory. Whereas it is quite easy to see by the aforementioned methods that a given admissable braid group representation of the previous type always has a current algebra realization and a W-algebra (generalization of minimal models) realization, the problem of finding all theories corresponding to given fusion laws and braid group matrices is more complicated. Since the charge-induction algebra introduced in the previous section contains the c-value modulo eight of the energy-momentum algebra, one expects a discrete family of theories. Indeed Fuchs [90] identified such a class of "Ising-like" models which have the same fusion and braid-group structure as the Ising model. The following two statements are presently only conjectures:

Conjecture 44 Conformal QFT are fixed by their associated type II_1 intertwiner algebras (combinatorial, topological data) up to a possibly discrete ambiguity related to the above c mod 8 phenomenon. **Conjecture 45** "Free" d=2+1 plektons are in a one to one correspondence with chiral conformal QFT.

Here the restriction "free" is necessary, because braid group commutation relations and associated combinatorial data do not fix general d=2+1 theories, inasmuch as the internal symmetry structure of d=3+1 fermions and bosons does not determine interacting theories (which, depending on the framework of description, require also the specification of couplings and coupling strengths). "Free" plektons are not defined in terms of free equation of motions (which, as will be shown in the next section, would contradict the plektonic commutation relations) but rather in terms of absence of "on-shell" processes. This will be explained in the next section.

7.7 Constructive Aspects of Plektons

In view of the spin-statistics connection which, as we saw in the previous section, extends for low dimensional theories beyond bosons and fermions to plektons, it is reasonable to first investigate the properties of massive d=2+1 Wigner representations with abelian U(1)-spin in the sense of the little group. In passing we note that our discussion of covariant free fields from Wigner's canonical momentum space representation in chapter 3 holds with small modifications for U(1)-spin, s=semiinteger. Here the u and v intertwiners intertwine between these abelian representations and covariant representations of $\widetilde{SL(2, R)}$, where only two-fold coverings have to be considered. The formulas for free fields in Fock-space and the phenomenon of their nonuniqueness (leading again to Borchers classes) are completely analogous to chapter 3. In particular the more restrictive quantization method using Lagrangians and the canonical- or functional-formalism for constructing free Bosons or Fermions works the same way.

As in the case of d=3+1 theories there is however also the possibility to avoid the u, v intertwiners in favor of *introducing localization directly* via modular theory.

This modular localization approach is the only one for $s \neq \frac{n}{2}$ i.e. for any(spin)ons. The definition of the net of real Hilbert spaces which have their localization in wedges is the same as in chapter 3 (especially theorem..and will not be repeated here). But the descend to compact space time regions fails. In fact it turns out that the problem already starts with the noncompact spacelike cones, which may be obtained by intersecting just two wedges. In principle two things could happen: such intersections could be empty (or too small) or the spacelike cone based real subspaces could fail to fulfill isotony. According to a recent result of J.Mund [91], it is the latter possibility which actually happens. So the prerequisite for having a functor from subspaces of the Wigner representation space to von Neumann algebras fails. A different argument with the same result on the impossibility of having a Fock space structure can be obtained by generalizing the J-S theorem [91]. From this one learns that there are no on-shell x-space string-like (spacelike cone with core-direction n and apex x) localized anyon fields, for which the combination of $A(x, n_1)A(y, n_2)$ and $A(y, n_2)A(x, n_1)$ appropriately weighted by a relative phase vanishes for space-like string separation. Rather free anyons must have a virtual particle creation (annihilation) structure. In this property free anyon fields are more similar to the fields of d=1+1 dimensional fields of factorizable models, than to d=3+1 free fields. Their two-point function must have a continuous spectral mass contribution in addition to the on-shell Wigner component, a fact well-known from from factorizable models which have real particle conservation and (unlike free Bosons and Fermions) virtual particle nonconservation.

An extremely useful additional information comes from the structure of incoming and outgoing scattering states. In the following we will try to give a systematic derivation starting from the structural analysis of plektons with the algebraic method. We start with the charge sector theory on d=2+1 observable algebras. The following theorem and its extension to d=2+1 theories is crucial for plektonic sectors:

Theorem 46 [72] Covariant representations of an observable net \mathcal{A} which fulfill the spectral mass gap hypothesis admit localizations in spacelike cones S of arbitrary small width; in brief, the a-priori best possible localization is semiinfinite string-like (the core of these cones):

$$\pi |_{\mathcal{A}_0(S')} \cong \pi_0 |_{\mathcal{A}_0(S')} \quad S := a + \bigcup_{\lambda > 0} \lambda \mathcal{O}$$
(7.89)

Here are some notational explanations. $\mathcal{A}_0 \equiv \mathcal{A}_{quasi}$ is the quasilocal algebra i.e. the C^* -algebra generated by the norm closure of the union $\bigcup \mathcal{A}(\mathcal{O})$ of local algebras. The C^* -algebra belonging to noncompact regions as S and S' is defined analogously. The \cong means unitary equivalence on the indicated subalgebras where the apex a and the width of \mathcal{O} is arbitrary as long as the cones are proper i.e. their causal completions do not fill the whole space-like region. The construction of the π -associated vacuum representation π_0 is part of the proof [3]. The positive effect which spectral gaps have on localization properties is of course intuitivly plausible (and frequently used by condensed matter physicists), however a convincing proof requires the application of a sizable part of analytic techniques of algebraic QFT and will not be given here. It was also shown there that the charge-carrying particles are necessarily Bosons and Fermions with a internal compact symmetry structure i.e. the standard DR-situation . However in d=2+1 this conclusion cannot be drawn and braid-group statistics may occur . If the localization is DHR (instead of S a compact double cone \mathcal{O}) then the fields and the associated particles are necessarily Bosons or Fermions with a possible compact internal group symmetry.

Using now the vacuum representation on the vacuum GNS-space $H_{\pi_0} \equiv H_0$ as the defining representation \mathcal{A}_0 and making the standard assumption of Haag duality on the corresponding von Neumann algebras: $\mathcal{A}(S') = \mathcal{A}(S)'$, one may again trade π for a homomorphism $\rho : \mathcal{A}_0(M) \to B(H_0)$ which however as a result of the noncompact nature in this case is not necessarily "endo" i.e. $\rho(\mathcal{A}_0(M)) \subsetneq \mathcal{A}_0(M)$ and hence the composition of such sectors is ill-defined. An elegant way out is to follow the construction of a larger universal observable algebra \mathcal{A}_{uni} of the previous section. Since one only has to replace the proper intervals I on the circle by proper spacelike cones S, we will not repeat that construction. Now the net is generated by the $\mathcal{A}(S)$ and the universality condition is formulated in terms of coherent families of representations $\pi^S(S)$. The definition of the reduced field bundle as a $\rho(\mathcal{A}) - \mathcal{A}$ bimodule with a C^* -algebra structure is as before denoted by pairs F(e, A), with $e = (\rho_\alpha, \rho, \rho_\beta)$ which act on the reduced field bundle vectors $\{\rho_\gamma, \psi\}$ with $\rho'_{\alpha}s$ being representative endomorphisms (one chosen from each equivalence class):

$$F(e,A)\left\{\rho_{\gamma},\psi\right\} = \delta_{\alpha\gamma}\left\{\rho_{\beta},\pi_{0}(T_{e}^{*}\rho_{\alpha}(A)\psi\right\} \quad T_{e} \in (\rho_{\alpha}\rho,\rho_{\beta})$$
(7.90)

The exchange algebra relations follow in complete analogy to the circular case. Again one has to pay attention to the localization concept for operators in the exchange algebra. As in the circular case (which has the same global topology), the localization of pairs (ρ, A) has a topological subtlety in that it depends on path classes:

Definition 11 (ρ, A) with $A \in A$ is said to be localized in the path class of spacelike cones $[S_0, ..., S_n]$ with either $S_i \subset S_{i-1}$ or $S_i \supset S_{i-1}$ if there exists a sequence of unitary charge transporters $U_i \in \mathcal{A}(S_i \cup S_{i-1})$ such that $loc(AdU_i...U_1 \circ \rho) \in S_i$ and $U_n...U_1 A \in \mathcal{A}(S_n)$; shortly: $loc(\rho, A) \subset [S_0, ..., S_n]$.

Here we already anticipated the result that the localization depends only on the homotopy class of paths with the beginning S_0 and the end S_n fixed. In the vacuum representation π_0 , the *U*-transport once around is trivial (there are no nontrivial vacuum selfintertwiners), but in nonvacuum sectors one expects a nontrivial representation of the first homotopy-group.

All these consideration may look unnecessarily pedantic to readers familiar with manipulations in supersymmetry or with differential geometric methods. Why does one not write down the commutation relations of the exchange algebra right away? Well, if anybody succeeds to guess the right answer, he should go ahead. The point which I want to stress here is that one is entering a new terrain in QFT, where the intuition based on known algebraic structures or Lagrangians is of not much help. Therefore, as already emphasized before, a careful presentation of arguments in algebraic QFT is the result of a pragmatic and not a pedantic attitude.

Two operators $F_{1,2}$ with $S_0^{(1)} = S_0^{(2)}$ are called mutually spacelike localized if the "endpoints" of their paths also agree $S_{n_1}^{(1)} = S_{n_2}^{(2)}$. The path topology is best pictured by shifting the apex of the S's into the origin and considering the intersection of the shifted S with the unit spacelike hyperboloid (the shadow which S casts on the 2-dim. de Sitter space). The topology (in particular the first homotopy group) of the resulting path model is the same as that of the circle. Naturally the R-matrix structure of the exchange algebra i.e. the commutation relations between the two $F_{1,2}$ is the same as in the conformal case, namely the R-matrix structure constants represent the generators of the cylinder braid group:

$$F(e_2, A_2) \cdot F(e_1, A_1) = \sum_{f_1 \circ f_2} R_{f_1 \circ f_2}^{e_2 \circ e_1}(N) \cdot F(f_1, A_1) \cdot F(f_2, A_2)$$
(7.91)

where the R-matrices, the statistics operators and the T- intertwiners are defined as in chiral conformal QFT and the N is the relative path- winding on de Sitter hyperboloid. These d=2+1 exchange C^{*}-algebras are, unlike chiral conformal QFT's but similar to d=3+1 bosonic or fermionic fields with fixed internal symmetry, expected to have (for fixed R's) continuously many representations corresponding to the idea of charge structure preserving coupling constant deformations. We are interested in the "free" plektons and therefore we now look at the inner product of scattering states which is associated with the above exchange algebra. It is well-known (see section on scattering theory) that the Haag-Ruelle scattering theory determine the Fock space structure of the incoming multiparticle states, in particular their inner products. The resulting (±symmetrized) tensor product structure of the multi-particle states in terms of the one-particle Wigner space cannot be maintained in the presence of the above exchange algebra. For the computation of the asymptotic limits for plektons we choose a formulation of the Haag-Ruelle scattering theory which makes the independence of the limits on the Lorentz-frame manifest . We define:

$$f_t(x) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int d^3 p e^{-ipx+i(\frac{p^2-m^2}{2m})t} \tilde{f}(p), \quad f(x) \in S(\mathbf{R}^3) \quad (7.92)$$

$$F(t) = \int d^3 x f_t(x) \alpha_x(F(e,A))$$

and $locF(e, A) \subset \tilde{I}$, $\tilde{I} = [S_0....S_n]$, the path-class localization. The F is chosen in the standard fashion of H-R scattering theory namely the spectrum of $F\Omega$ contains an isolated mass shell and $supp\tilde{f} \cap specF\Omega \subset H_m$, the mass shell i.e. $F\Omega \equiv \psi$ is a one particle state. The quasilocal operators F(t) can be (as in the standard Haag-Ruelle theory) approximated by operators $F_{\varepsilon}(t)$ localized in $\tilde{I} + tV_{\varepsilon}(f)$ s. t.. $||F(t) - F_{\varepsilon}(t)|| < c |t|^{-N}$ where $V_{\varepsilon}(f)$ is the velocity support of f surrounded by an ε -safety collar. We chose the F'_is and f'_is in such a way that the $F_{\varepsilon,i}(t)$ are spacelike for large t. Then the following generalized Haag-Ruelle theorem holds.

Theorem 47 ([92]) The sequence of vectors $F_n(t)....F_1(t)\Omega$, with loc F_i relatively spacelike, converges to a L-covariant limit with braiding properties:

$$\lim F_n(t) \dots F_1(t)\Omega = (\psi_n, \tilde{I}_n) \times \dots \times (\psi_1, \tilde{I}_1)$$
$$U(L)(\psi_n, \tilde{I}_n) \times \dots \times (\psi_1, \tilde{I}_1) = (U(L)\psi_n, L\tilde{I}_n) \times \dots \times (U(L)\psi_1, (\tilde{I}_1))$$
$$(\psi_{\sigma(n)}, \tilde{I}_{\sigma(n)}) \times \dots \times (\psi_{\sigma(1)}, \tilde{I}_{\sigma(1)}) = \varepsilon(b)(\psi_n, \tilde{I}_n) \times \dots \times (\psi_1, \tilde{I}_1)$$
(7.94)

Some comments are in order. In the standard theory there is no dependence on the localization path, and we have a tensor product structure. In our present case we have a similar situation, but only if we keep the localizations I fixed, as indicated by the above notation. Applying the operators F_i in a different order, the result can be written as a unitary representer of a certain cylinder braid b_n applied to the original n-particle vector. The braid b_n is from the groupoid of colored (by the charge sectors) braids on the cylinder. It depends on the permutation σ and the spacelike paths $\tilde{I}_1, ..., \tilde{I}_n$ and can be obtained by a simple 3-dim. geometrical construction. Asymptotically only the directions of the I_i is relevant, i.e. the hyperbolic angle of the translated I_i with apex translated to zero intersects with the unit spacelike hyperboloid. With other words the path topology is that of the 2-dim. de Sitter space which is topologically equivalent to S^1 . This de Sitter space is also the set of points at spatial infinity by which one has to extend the Minkowski-space $M \to M$ in order to characterize the universal algebra in geometrical terms as $\mathcal{A}_{uni} = \mathcal{A}(\tilde{M})$ in analogy with the universal algebra of chiral conformal QFT $\mathcal{A}_{uni} = \mathcal{A}(S^1)$. These directions together with the non-coinciding velocity space as well as the inner product in terms of the Markov trace on the cylinder ribbon braid group may be elegantly encoded into an n-particle momentum space structure of vector valued wave functions on the covering space, which obey a covariance property [92]. With the structure of the in-space understood, the remaining problem is now the construction of the n-particle modular localization subspaces and the associated plektonic nets. We expect that similar ideas based on thermal methods for wedge localization as in chapter 6.8 will be useful. Here our voyage ends for the time being.

Literature to chapter 7:

R.Haag "Local Quantum Physics". Springer 1992

S.Doplicher, J.E.Roberts: Why is there a Field Algebra with Compact Gauge Group describing the Superselection Structure in Particle Physics? Commun.Math.Phys **131**, 51(1990).

K.H.Rehren and B.Schroer "Einstein Causality and Artin Braids" Nucl.Phys. **B312**, 715 (1989)

K.Fredenhagen, K.H.Rehren and B.Schroer, Rev.Math.Phys., Special Issue (1992) 113.

K.Fredenhagen, Proceedings of the 1991 Cargese Summer Institute.

Chapter 8

Tentative Resume and Outlook

The most fruitful times in theoretical physics were those of clash with principles. The resolution of a deep paradoxon can lead to an enormous amount of progress, as exemplified by the rapid emergence of QM as a result of Bohr's atomic model and the ensuing paradoxical situation with respect to classical physics.

The post electro-weak period of stagnation in particle physics and QFT did not lead yet to such a clear-cut clash. The present situation in QFT and high energy physics seems to be somewhat similar to the years preceding the progress of "renormalization" of QFT. Then there was also no clash with fundamental principles, even though many physicist thought that the correct handling of the ultraviolet problem requires radical new inventions.

In my opinion the present situation is the result of a significant disequilibrium between two modes of thinking whose delicate balance is the hall-mark of good progress in theoretical physics. I am thinking of Dirac's method build on mathematics and esthetical appeal, versus the Bohr-Einstein-Heisenberg-Wigner approach based on physical conceptual analysis. The modern "Diracian" approach [5] is that of inventions, based particularly on the formal *geometric extension of formalism*, leaving behind the physical principles which originally led to this formalism. Recent illustrations of such "Diracian inventions" are the introductions of supersymmetry, strings ore membranes.

In such a situation it seemed to be helpful to revisit those old ideas in a critical spirit, which were essential to the conceptual development of QFT. Gauge theories also belong to this category. Even though they were intensely investigated for almost three decades, there exists presently no intrinsic characterization, i.e. if one would receive the physical gauge invariant correlation on a divine silver plate, one would have no means to find out whether they have a gauge theory behind without asking the person who manufactured it. This is not to say that one lacks conjectures; one educated guess is that if one needs charge carriers which have a semiinfinite string-like extension, then the charges are Maxwellian charges of a gauge theory. According to our discussion in chapter 4.5 every theory involving spin=1 fields are gauge theories in which case the word (broken) "gauge" in LQP does not carry more information than renormalizable QFT of spin one. Semiinfinite massive charge carriers play an important role in the description of d=2+1 particles with braid group statistics (plektons). In that case the asymptotic string direction is not frozen by infrared photon clouds but rather fluctuating (topological charges in the sense of algebraic QFT). The only way to picture this within a Lagrangian "straight jacket" is through Mandelstam integrals over vectorpotentials which is a gauge theoretical picture (but not yet a computational scheme).

We emphasized in these notes that despite its conservative way of dealing with physical principles, algebraic QFT leads to a radical change of paradigm. Instead of the Newtonian view of a space-time filled with a material content one enters the reality of Leibnitz created by relation (in particular inclusions) between "monades" (~ the hyperfinite type III₁local von Neumann factors $\mathcal{A}(\mathcal{O})$ which as single algebras are nearly void of physical meaning). Related to this is a very new and surprising esthetics, namely the art of compressing relations between very big objects as type III₁von Neumann factors¹) into extremely simple structures (which is very reminiscent to the esthetics of the V. Jones subfactor theory). I expect that this new esthetics will be important in understanding the connection of localization and entropy, i.e. the generic incorporation of the Bekenstein-Hawking quasiclassical observations into local quantum physics.

Closely related is the different mathematical way of thinking about a local description of states. Whereas the standard approach uses the geometrical idea of sections in fibre-bundles, algebraic QFT deals with sheafs. This is because a partial state over $\mathcal{A}(\mathcal{O})$ is an equivalence class of states on \mathcal{A} which yield the same restriction on $\mathcal{A}(\mathcal{O})$ [3]. The algebraic net has the dual description in terms of a co-sheafs. Whereas the fibre bundle approach and Lagrangian QFT is limited to "quantization algebras" as CCR and CAR i.e. algebras over classical function (or section) spaces and their infinitesimal perturbative deformations, the algebraic approach is supposed to cover the unknown territory beyond (the generic nets are not indexed by classical function spaces).

Another important distinction between the standard approach and algebraic QFT is that the former deals already in its very formulation with global concepts as e.g. functional integrals (the restriction of integration to a region does not describe the physics of that region), whereas the latter starts with local concepts and makes contact with global aspects (as global topology) only in a later stage. Those aspects of the vacuum structure, which through the spontaneous breaking of localizable symmetries and superselection rules are related to local properties of the theory, are correctly accounted for. However vacuum degeneracies without any presently visible local origin (as e.g. the vacuum structure in the Seiberg-Witten duality construction), are presently out of reach by methods of algebraic

¹Here "big" is meant in the sense they absorbe any tensor factor with another von Neumann algebra. Although V.Jones formulated his subfactor theory in terms of the "smallest" infinite dimensional algebras (type II_1 which gets absorbed by any other tensor factor), his theory applies with only a few modifications to the present setting.

QFT.

It was our intention to apply concepts of algebraic QFT to those problems which in our view are not appropriately taken care of by the standard quantization formalism or which may even contain paradoxes and physically fruitful contradictions. Examples are the local gauge concept, QFT's in curved spacetime, the structure of nonperturbative low dimensional QFT's, and the role of various forms of "Duality" as well as "Quantum Symmetry". Even in cases where definite answers are still missing, algebraic QFT certainly casts a different and physically interesting light on those problems. We reviewed the two notion of temperature, the *standard* one being generated by a heath bath in a Lorentz frame and the second one by a *loss of information* through the creation of a horizon (the Hawking-Unruh temperature or mathematically: the Tomita KMS temperature originating from the vacuum representation of local observable algebras), the latter having a surprising relation to the crossing symmetry which represents the on shell aspect of causality and which played an important role at the cradle of string theory.

The deep crisis in QFT and elementary particle physics at the end of this century (with its ever increasing crave on sophisticated entertainment), is plainly evident if one just looks into the titles of hep-theory preprints of recent years. Never in its long and fascinating history has physics as a human activity of fundamental endeavor been threatened in its existence as presently. The crisis is to a large degree man made and its main cause is the adoption of market ideology in science. One of the consequences in particle physics is the emergence of (un)physical monocultures with the danger of long term loss of deep knowledge and the demoralization of those few young people who are willing to make profound intellectual investments without the promise of instant return. The monocultures which are cultivated within big electronically connected groups are easily identifyable as supersymmetry² with string theory on its back. The danger is not so much that many physicist adhere to fashions which lead into physically fruitless directions. This has happened before in this century and physics has survived such situations. The most dangerous aspect is the quasi-religious zeal and the signs of mass psychosis against which great intellectual brilliance is no antidote, a phenomenon which hitherto was restricted to the great political catastrophes of this century. The problem which threatens the future of particle theory is not the work on those subjects itself, but the monomaniacal zeal with which the protagonists make sure that the reign of their monoculture will be total. The followers do not investigate problems because they are led by intrinsic physical logic but rather because one of their gurus drops a word e.g. the present fashion to apply a doses of noncommutative geometry to strings. The picture the reader finds at the end of this notes (in an attached ps file) is more serious than he thinks at first glance: the theory divisions of almost all european and

 $^{^{2}}$ To avoid any misunderstanding, I am not talking about supersymmetry as a theoretical structure whose investigation is completely covered by intellectual curiosity and academic freedom, but I am pointing to the fact that after 25 years of marginal scientific merits but great employment capacity (since it can be easily manipulated in calculations) it become the cropping ground for a second much more dangerous monoculture: string theory.

national research institutions are headed by either protagonists or sponsors of those monocultures. The future of particle physics as we have known it, is extremely precarious indeed.

These notes are an (unfashionable and probably futile) attempt to go against this tide and to recapture some of the earlier conceptual spirit which views the laws of nature as the realization of physical principles. Whether such attempts can lead to rational discourse, as they did in better times, remains a matter of hope.

Acknowledgment: I thank José Helayel-Neto for the invitation and the hospitable atmosphere at the CNBPF, where some of the sections were written in the southern summer of 1997 and the rest in 1998. My thanks for reading parts of the manuscript and for interesting e-mail communications goes to Karl-Henning Rehren. I am also indebted for some exchange of letters with Raymond Stora concerning whose unflinching critical spirit has been a source of encouragement.

Bibliography

- [1] E.P.Wigner, Ann. Math.40, 149 (1939)
- [2] B. Schroer, Nucl. Phys. B (Physical Mathematics) 499, (1997), 519
- [3] R. Haag, "Local Quantum Physics" Springer Verlag 1992
- [4] J.-P. Eckmann and K. Osterwalder, J. Funct. Anal., 13, (1973), 1
 P. Leyland, J. Roberts and D. Testard, "Duality for Quantum Free Fields" CNRS 1978 preprint, unpublished.
- [5] B. Schroer, Annals of Physics Vol. 255, No.2, 270 (1997)
- [6] J.J. Bisognano and E.H. Wichmann, J. Math. Phys. 16, 985 (1975); 17, 303 (1976)
- [7] H.-J. Borchers, Commun. Math. Phys. 143, (1992), 315
- [8] See the contributions of H.-J. Borchers and H.-W. Wiesbrock in "Operator Algebras and Quantum Field Theory", eds. S. Doplicher, R. Longo, J. E. Roberts ans L. Zsido, academia Nazionale dei Lincei Roma, July 1-6, 1996
- [9] G.L. Sewell, Phys. Rev. Lett. **79A**, 23 (1980)
 G. L. Sewell, Ann. Phys. **141**, 201 (1982)
 G. L. Sewell, Phys. Lett. **122A**, 309 (1987)
- [10] S. J. Summers and R. Verch, Lett. Mat. Phys. 37, (1996) 145
- [11] H. Narnhofer, "Entropy Density for Relativistic Quantum Field Theories", in Advanced Series in Mathematical Physics Vol.20, World Scientific, eds. M. Aizenman and H. Araki
- B. Kay, "Entropy Defined, Entropy Increase and Decoherence Understood, and Some Black-hole Puzzles Solved", hep-th/98021172
- [13] F. Larsen and F. Wilszek, Phys. Rev. Lett. B375, (1995), 37
- [14] D. Bigatti and L. Susskind "Review of Matrix Theory", hep-th/9712072
 T. Banks, Lecture Notes, "Matrix Theory", hep-th/9710231

- [15] Unpublished notes of H.-W. Wiesbrock, FU-Berlin February 1998.
- [16] M. Duetsch and K. Fredenhagen, in preparation.
- [17] K. H. Rehren, hep-th/9711085
- [18] R.F. Streater and A.s. Wightman, "PCT, Spin and Statistics and All That" New York, Benjamin 1964
- [19] A. S. Wightman, in "High Energy Interactions and Field Theory", ed. M. Levy, Cargese Lecture in Theoretical Physics", 1964 Gordon and Breach, New York 1966.
- [20] B. Schroer, T. T. Truong and P. Weisz, Ann. Phys. 102, (1976) 156
- [21] A.B. Zamolodchikov, Int. J. Mod. Phys. A1 (1989) 4235
- [22] B. Schroer, T.T. Truong and P.Weisz, Phys. Lett. 63B (1976) 422
- [23] B. Berg, M. Karowski, H.J. Thun, T.T. Truong and P. Weisz, Phys. Lett. 67B (1977) 321
- [24] A.B. Zamolodchikov, Comm. Math. Phys.55 (1977) 183, ITSP (1977) 12.
 L. D. Faddeev, Sov. Sci., Rev. Math. Phys. C1, (1980) 107
- [25] V. Kurak and J.A. Swieca, Phys. Lett. 92 (1979) 289
- [26] V. Jones, Vaughn F.R. Jones, American Math Society publication No 80, 1991
- [27] F.A. Smirnov, Adv. Series in Math.Phys. 14, World Scientific1992
- [28] H. Babujian, A. Fring and M. Karowski, "Form Factors of the SU(N)-Chiral Gross Neveu Model", in preparation.
- [29] C.N. Yang, Phys.Rev.Lett. B86, (1977) 209
- [30] M. Karowski and P. Weisz, Nucl. Phys. **B139** (1978) 445
- [31] B. Schroer and H.-W. Wiesbrock, in preparation
- [32] B. Schroer and J.A. Swieca, Nucl. Phys. **B121** (1977) 505
- [33] A. Fring, I.J.of Mod. Phys. A 11, (1996) 1337
- [34] M. Yu Lashkevich, "Sectors of Mutually Local Fields in Integrable Models of QFT" Landau Institute Preprint, hep-th 9406118
- [35] R. Brunetti and K. Fredenhagen, "Interacting Quantum Field Theories in Curved Space: "Renormalizability of φ^{4} "

[36] M. Duetsch, T. Hurt, F.Krahe and G. Scharf, Nuovo Cimento A 107 (1994), 375

M. Duetsch, Nuovo Cim. 109 A, No.8, 1145 (1996)

A. Aste and G. Scharf, "Non-abelian gauge theories as a consequence of perturbative quantum gauge invariance", hep-th/9803011.

- [37] A.Klein and L.Landau, Pacific J. Math. 94, 341 (1981)
 A. Klein and L. J. Landau, J. Funct. Anal. 42, 368 (1981)
- [38] B. Schroer, Nucl. Phys. B ("Phys. Mathematics") 499, (1997), 547
- [39] B. Schroer, hep-th/9712124, to be published in AOP.
- [40] The relation between the KMS cyclicity relation and crossing symmetry was independently discovered by M. Niedermaier, "A Derivation of the Cyclic Form Factor Equation", MPI-97-31 Draft.
- [41] H:J: Borchers "Translation Group and Particle Representations in Quantum Field Theory" Springer Lecture Notes in Physics 1996.
- [42] D. Buchholz and I. Ojima, "Spontaneous Collapse of Supersymmetry" March 1997 University of Kyoto preprint.
- [43] S. Doplicher and J. E. Roberts, Commun. Math. Phys. 131, 51 (1990)
- [44] N. P. Landsman, Int. J. Mod. Phys. A30, 5349
- [45] W. H. Zurek, Phys. Today 44, No. 10, 36 (1991)
- [46] M. Brune et al., Phys. Rev. Lett. 77, No. 24, (1996)
- [47] A. Amann, Synthese 97: 125 (1993)
- [48] S. Summers and R. Werner, Ann. Inst. Henri Poincare 49, 215 (1988)
- [49] D. Buchholz, Commun. Math. Phys. 85, 49 (1982)
 Phys. Lett. B174, 331 (1986)
- [50] H. Epstein, Nuovo Cimento 27, 886 (1963)
 B. Schroer, 1962 unpublished, see ref 27 in [18]
- [51] J. Bros and D. Buchholz, Nucl. Phys. **B429**, 291 (1994)
- [52] R. Brout et al., Physics Reports **260** (1995), 329-446
- [53] M. Müger, Superselection Structure of QFT in 1+1 Dimensions, Desy preprint 97-073, ISSN 0418-9833
- [54] D. Buchholz, G. Mack and I. Todorov Nucl. Phys B (Proc. Suppl.) 5B, 20 (1988)

- [55] R. M. Wald, "Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics", University of Chicago Press 1994.
- [56] B. Schroer and J. A. Swieca, Phys. Rev. **D10**, 480 (1974)
- [57] B. Schroer, J. A. Swieca and A. H. Völkel, Phys. Rev. D11, 1509 (1975)
- [58] G. Mack, "Introduction to conformal invariant quantum field theory in two and more dimensions", Cargese 1987, and references therein.
- [59] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, Nucl. Phys. B 247, (1984) 83
- [60] B. Schroer, "Modular theory and symmetry in QFT", in Proc. Workshop on Mathematical Physics Towards the 21st Century, eds. R. N. Sen and A. Gersten, Ber-Sheva, Israel, 1993, Ben-Gurion University of the Negev Press, 1994
- [61] S.Weinberg, "The Quantum Theory of Fields", Vol. I, Cambridge University Press 1995
- [62] C. Itzykson and J.-B. Zuber, "Quantum Field Theory", McGraw-Hill 1980
- [63] W. Zimmermann, 1970 Brandeis lectures, eds. S. Deser, M. Grisaru and H. Pendleton, MIT Press, Cambridge, Mass., 1971

J. H. Lowenstein, in "Renormalization Theory", Proceedings of the NATO Advanced School of Mathematical Physics Erice 1975, eds. G. Velo and A. S. Wightman

O. Piguet and S. P. Sorella, "Algebraic Renormalization", Lecture Notes in Physics, Vol m26, Springer Verlag

- [64] Laurie M. Brown, "Renormalization" From Lorentz to Landau (and beyond), Springer Verlag 1993
- [65] C. Becchi, A. Rouet and R. Stora, Ann. Phys. 98, 287 (976)
- [66] I. T. Kugo and I. Ojima, Suppl. Prog. Theor. Phys. 66, (1979) 1
- [67] A. Aste, M, Duetsch and G. Scharf, hep-th/9602053A. Aste and G. Scharf, hep-th/9803011
- [68] J. Schwinger, IAE Report, Vienna, 1962, lecture presented at the "Seminar on Theoretical Physics, Trieste, 1962, and Phys Rev. 128, 2425 (1962)
- [69] D. Buchholz and R. Verch; Rev. Math. Phys. 7, (1995) 1195
- [70] See the contribution of Lewellyn-Smith to the Proc. of the 5th Hawai Topical Conf, Particle Physics, Ed. P. N. Dobson et al., University of Hawai Press Honolulu, Hawai 1993.

- [71] M. Radzikowski, PhD thesis, Princeton University, October 1992.
 M. Köhler, Class. Quant. Grav., 12, 1413 (1995)
- [72] D. Buchholz and K. Fredenhagen, Comm. Math. Phys. 84, 1 (1982)
- [73] R. Jost, "General Theory of Quantized Fields", American Math. Soc. Publications, 1963
- [74] B. Schroer, Rev. Math. Phys. 7, 645 (1995), and references quoted therein.
- [75] N. Yu. Reshetikhin and V. G. Tuarev, Inv. Math. 103, 547 (1991)
 V. G. Turaev and O. Y. Viro, Topology 31, 865 (1992)
- [76] M. Karowski and R. Schrader, Commun. Math. Phys. 151, 355 (1993)
- [77] E. Witten, Comm. Math. Phys. **121**, 351 (1989)
- [78] D. Buchholz S. Doplicher R. Longo and J. E. Roberts, Rev. Math. Phys. Special issua, 47 (1992)
- [79] D. Buchholz, Nucl. Phys. B469, 333 (1996)
- [80] M. Jörss, "The Construction of Pointlike Charged Fields from Conformal Haag-Kastler Nets" Desy preprint 1996
- [81] K.-H. Rehren, "Weak C* Hopf symmetry" Desy preprint 1996
- [82] K.-H. Rehren and B. Schroer, Nucl. Phys. B312, 715 (1989)
- [83] J. Glimm and A. Jaffe, "Quantum Physics. A functional integral point of view" Springer Verlag
- [84] B. Schroer, T.T. Truong, and P.Weisz, Ann. of Physics **102**, 156 (1976)
- [85] H.-W. Wiesbrock, Comm. Math Phys. 157, 83 (1993); Lett. Math. Phys. 29, 107 (1993)
- [86] H. Araki and L. Zsido, "Extension of the structure theorem of Borchers and its application to half-sided modular inclusions" to appear.
- [87] D. Schlingemann, Rev. Math. Phys., 8, 1187 (1996)
- [88] W. Kunhardt, "On Infravacua and Superselection Theory", hepth/9704212
- [89] K. Fredenhagen, K.-H. Rehren ans B. Schroer, Reviews in Math Phys., Special Issue (1992), 113
- [90] J. Fuchs, Nucl. Phys.B **328**, (1989), 585
- [91] J. Mund, "No-Go Theorem for "Free" Relativistic Anyons in d=2+1", FU-Berlin preprint, submitted to Lett. Math. Phys.; and private communication.

- [92] K. Fredenhagen, M. Gaberdiel and S.M. Rüger, Commun. Math. Phys. 175, (1996), 319
- [93] K. Fredenhagen, K.-H. Rehren and B. Schroer, Commun. Math. Phys. 125, (1989), 201
- [94] S. Takagi, Progr. of Theoretical Physics, **74**, (1985), 142
- [95] J.C. Barata and K. Fredenhagen, Commun. Math. Phys. 138, (1991) 175
- [96] G. Morchio and F. Strocchi, Jour. Math. Phys. 34, 899, (1993)
- [97] D. Buchholz, S. Doplicher, G. Morchio, J.E. Roberts and F. Strocchi, "A Model for Charges of Electromagnetic Type", Conference on Operator Algebras and Quantum Field Theory (Rome, 1996)
- [98] C. P. Staskiewicz, "Die lokale Struktur auf dem Kreis", FU thesis, Berlin 1995
- [99] G. C. Hegerfeldt, Phys. Rev. Lett. 72, (1994), 596
- [100] D. Buchholz and J. Yngvason, Phys. Rev. Lett. 73, (1994) 613
- [101] K. Fredenhagen and M. Jörss, Commun. Math. Phys., 175, (1996), 541
- [102] B. Berg, M Karowski and P. Weisz, Phys. Rev. D19, (1979) 2477
- [103] H. M. Babujian, M. Karowski and A. Zapletal, J. Phys. A: Gen. 30, (1997) 6425
- [104] D. R. Grigore, J. Math. Phys. 37, 460 (1996)
- [105] J. A. Swieca, "Goldstone's Theorem and Related Topics", Cargese Lectures in Physics, Vol.4, 215 (1970)
- [106] J. Glimm and A. Jaffe, "Quantum physics. A functional integral point of view". Springer 1987
- [107] J. H. Lowenstein, Commun. Math. Phys. 6, 49 (1967) and references therin.
- [108] J. H. Lowenstein and B. Schroer, Phys. Rev. D 3, Number 8, 1981 (1971)
- [109] Ferrara, Gatto and Grillo, Phys. Rev. D5, 3102 (1972)
- [110] M. A. Virasoro, Phys. Rev. D1, 2933 (1970)
- [111] B. Schroer, Nucl. Phys.B160, 330 (1979)
 A. M. Polyakov and P. B. Wiegmann, Phys. Lett. B 131, 121 (1963)
- [112] B. Schroer, "Modular Localization and the d=1+1 Formfactor program", hep-th/971214, To be published in the August 1998 issue of AOP

- [113] B. Schroer and H. W. Wiesbrock, "The Inverse Problem in Local Quantum Physics", in preparation
- [114] S. Doplicher and R. Longo, Invent. math. 75, 443 (1984)
- [115] E. Cremmer and J. Scherk, Nucl. Phys. B118, (1977), 61; for more historical details see also the superstring bible.
- [116] T. Banks, W. Fischler, N. Seiberg and L. Susskind, Phys. Rev. D55 (1997) 5117
- [117] J. Roberts, "Spontaneously broken gauge symmetries and superselection rules" in Proceedings of the International School of Mathematical Physics, Camerino, 1974, ed. by G. Galavotti,1976
- [118] E. Verlinde, Nucl. Phys. **300**, 360 (1988)

BIBLIOGRAPHY

312

Appendix A

Mathematical Appendix

A.1 Introduction

Even at the risk of exaggeration, it is tempting to compare the role of modular theory in von Neumann algebras with respect to local quantum physics (LQP) with that of calculus in relation to Newtonian mechanics. I hope that I succeed to convince the reader in this little mathematical appendix, that this comparision is more than marketing. Whenever it helps to understand the mathematical formalism and highlight the physical content, I will (as I already did for the representation theory of groups in the first chapter) use physical terminology in addition to the mathematical expressions. I try to keep the notation of the main text and use large calligraphic letters for nets of vNa, large Latin letters¹ as M, N... for vNa (and occasionally also calligraphic latin letters as $\mathcal{L}(H), \mathcal{A}, \mathcal{C}...$ for individual vNa's and C*-algebras), small Latin letters x, y... for individual operators and Greek letters for vectors and states on C^* -algebras. Nets of vNa will always be denoted by $\mathcal{A}, \mathcal{B}...$ With a few exceptions I omitted proofs, so the reader is asked to look up the literature given at the end and to use the appendix mainly as an orientation aid (with a mathematical physics bias) for the study of mathematical literature.

A.2 States on C*-algebras and representations

Since (concrete) von Neumann algebras (abbreviated: vNa) are special operator algebras in Hilbert² space H (always a complex Hilbert space unless stated otherwise), we first remind the reader of some definitions and properties of single operators. If nothing is said, an operator will always be a bounded linear map of H into itself (bounded \simeq continuous) and $\mathcal{L}(H)$ or B(H) is the notation for the vNa (by definition) of all such operators. Operators have an adjoint and $x \to x^*$

¹With all appologies for the intrusion into stringy and brany terminology.

 $^{^{2}\}mathrm{A}$ Hilbert space will always have a positive definite inner product and in the context of these notes it will be separable.

is an involution which fulfills the so called C^* identity: $||xx^*|| = ||x||^2$. The star is inherited from the sesquilinear nature of the inner product (for physicist antilinear in the left hand vector) by shuffling x from the "ket to the bra" vector. There are three types of operators which every physicist knows from QM: projections P with $P^2 = P$, selfadjoint operators $x = x^*$ and unitary operators $u^* \cdot u = u \cdot u^* = 1$. Projectors in operator algebras are always selfadjoint and project onto closed (Hilbert)subspaces. A positive operator $x \ge 0$ is a special kind of selfadjoint operator which quantum physicist meet in the tracially normalized form of density matrices (for the description of impure states in QM). In QFT more general normal operators $(x \cdot x^* = x^* \cdot x)$ as well as special nonnormal operators in the form of isometries $(v^* \cdot v = 1, v \cdot v^* = P)$ make their appearance. Another operator concept which is more important in QFT than in QM, is the polar decomposition of a normal operator into a unitary ("phase") and positive ("radial") part: $x = u \cdot h$ with $h = (x^* \cdot x)^{\frac{1}{2}} \equiv |x|$. This exists also for antilinear operators x in which case u will be antiunitary.

The spectrum of x : specx is the set of $\lambda' s$ such that $(x - \lambda)^{-1}$ is not a bounded operator. For selfadjoint operators If it comes to unbounded operators S, only those which have a dense domain of definition, and are closable feature in these notes (and in physics). Such operators are most conveniently handled by the method of their graph: $G(S) = \{(\xi, S\xi); \xi \in domS\}$ with $G(S)^-$ denoting the closure. For a closed operator the graph is a closed subspace of $H \oplus H$. Closable operators are defined as those whose graph presents no $(0, \xi \neq 0)$ obstruction in the closure (i.e. the closure is again the graph of an operator). The graph of the adjoint operator is $G(S^*) = \{(-S\xi,\xi); \xi \in domS\}^{\perp}$, where the orthogonal complement refers to $H \oplus H$. In this way one also sees that the denseness of the domain and the property of having adjoints are "dual" properties under hermitian conjugation. At this point it is also helpful to remind the reader on the relation between families of projectors $e(\lambda)$ which are indexed by subsets $(-\infty, \lambda)$ of the spectrum of a selfadjoint operator and the selfadjoint operator itself, which certainly everybody knows under the name of spectral resolution or spectral decomposition of x:

$$x = \int \lambda de(\lambda) \longleftrightarrow$$

$$\langle \eta, x\xi \rangle = \int \lambda d\mu_{\eta,\xi}(\lambda), \ \mu_{\eta,\xi}(\lambda) = \langle \eta, e(\lambda)\xi \rangle$$
(A.1)

This can be expanded to a functional calculus $f \in L^{\infty}(specx, \mu) \to f(x) \in \mathcal{L}(H)$.

The terminology selfadjoint is applied to unbounded operators only if they are densely defined (hence closable). In that case the measure μ decides when a vector ξ belongs to the domain of x by $\int |\lambda|^2 d\mu_{\xi,\xi}(\lambda) < \infty$. The extension of the polar decomposition of unbounded densely defined closed operators is also straightforward. The physically most important application is the decomposition of the Tomita (antilinear) involution S. Quite generally, unbounded operators in algebraic QFT are are mostly (in these notes exclusively) entering via modular theory i.e. they are not individual operators in the algebra but rather rather describe the modular characteristics of the representation of the algebra or of the pair (algebra, reference vector) where the most prominent example of a reference vector is the vacuum of QFT.

Most concrete calculations in algebraic QFT take place in von Neumann algebras. These are special operator algebras for which two prominent examples already have been given: the algebra of all bounded operators $\mathcal{L}(H)$, and the commutative algebra of all essentially bounded functions on a measure space X: $L^{\infty}(X,\mu)$. QM leads to the first kind of vNa, and the phase space observables of classical mechanics or the functional calculus of the spectral representation theory on $L^2(specA, \mu)$ of a selfadjoint operator A to the second. The reason why vNa are more important for QT than any other type of algebra (and in fact were introduced by von Neumann for this purpose) is that they implement the physical idea of compatibility in the theory of quantum measurements ("simultaneously measurable") through their notion of commutant. Von Neumann knew about other types of von Neumann algebras besides the two mentioned example. Together with Murray he got the first rough classification in terms of types I, II, and III in terms of projectors (see next section), but he thought that only the type I $\mathcal{L}(H)$ algebra is of physical relevance. The type II algebras received most of the mathematical attention of Murray and von Neumann and their picture of type III was even mathematically very vage. We know now that all three types occur in physics, type III becomes important through the localization concept in local quantum physics and type II enters in two ways: one is the intertwiner calculus (of localized charges) together with its natural tracial states (in the differential geometric Lagrangian approach called "topological field theory", but here the more appropriate name is combinatorical QFT) and in magnetic QM for constant magnetic external fields (the Hofsteadter theory). In form of the noncommutative torus it also has more recent applications in which physicist were involved. Type II₁ is too "small" to accomodate spacetime symmetries, but large enough (and excellently suited) to incorporate inner symmetries described by compact groups and "quantum symmetry" generalizations which are encountered in low-dimensional QFT.

Von Neumann algebras are special C*-algebras. A C*-algebra is a normed Banach algebra which fulfills the C* relation $||x^* \cdot x|| = ||x||^2$. The conceptual relation of C*-algebras to vNa (for a physicist) is similar to that of abstract groups³ to their concrete representations. In the case of topologically nontrivial QT with quantum ambiguities in the form of θ -angles chapter 1.2), one wants to consider the inequivalent theories for different $\theta's$ as different representations of one abstract object which is a C*-algebra C, a viepoint which has no counterpart in the geometric fibre-bundle approach where different θ -angles correspond to different fibre bundles. Therefore C*-algebras play an important conceptual role, since they allow a unifying point of view. In particular in the exploration

 $^{^{3}}$ The relation of vNa to group theory is also very close from another point of view: a vNa is generated by the group of its unitary elements. However in mathematics the analogy of vNa with measure theory and that of C*algebras with topology is more fruitful (noncommutative geometry).

of new symmetry concepts beyond group theory, the C*-algebras which can be related to a bimodule have attracted a lot of attention. But most of the calculations are done in the von Neumann closures of their representations. Although one knows considerably less about C*-algebras than von Neumann algebras, this did not hamper progress in algebraic QFT because the von Neumann extensions of their representations make the rich body of structural theorems and computational tools of vNa'a available to LQP.

The representation theory of C*-algebras is done via states. A state is a normalized linear positive functional on a C*-algebra⁴ and the space of states is simply a normalized convex cone inside the dual space \mathcal{C}^* . Therefore all the notions of state decomposition theory, as pure state, central decomposition etc. apply. The special feature of a vNa \mathcal{A} , which distinguishes it within the C^{*}setting is the fact that \mathcal{A} is the dual of a Banach space \mathcal{A}_* (the "predual") which contains the convex normalized cone $\mathcal{A}^{+(1)}_*$. In the case of the vNa $\mathcal{L}(H)$, this is the space of normalized positive trace operators i.e. density matrices in physical terminology. For the standard commutative vNa $L^{\infty}(\mathbf{R}, dx)$ one has $L^{\infty}(\mathbf{R}, dx)_* = L^1(\mathbf{R}, dx)$. Since (concrete) vNa also have the standard definition of being weakly closed *-subalgebras of of $\mathcal{L}(H)$, some states on \mathcal{A} may be simply obtained by restricting the density matrices to $\mathcal{A} \subset \mathcal{L}(H)$. At this place the reader should acquaint himself with the notion (the ideal) of compact operators $K(H) \subset \mathcal{L}(H)$ and the property $K(H)^* = \mathcal{L}(H)_*$. He should also develop an awareness for the existence of three different topologies: norm topology (operator norm), strong topology and weak topology (only defined for Hilbertspace operators). The convergence of sequences of operators on density matrices gives a slightly stronger topology than the weak and is called " σ -weak". It is the cause for the existence of the predual. The reader will find these definitions and related theorems in the first chapters of any book on operator algebras; some of them are quoted at the end of the appendix. More general than normal states on vNa's would behave in a very pathological manner, whereas on C*-algebras which are not vNa's, the notion of good and bad or rather regular and singular depends on the C*-algebra. For example the Heisenberg-Weyl algebra which underlies standard QM possesses singular states in the sense that there are no continuous translations e^{ipa} , because the lack of continuity does not allow to define an infinitesimal momentum operator p. In order to exclude such possibilities which violate the Stone-von Neumann unicity theorem for the Schrödinger representation, one has to insist in regular representations such that the translation U(a) has continuity properties at a = 0. In this way the so called topological (combinatorical) field theories (type II_1) are thought to originate from algebras which formally (e.g. in attempting to interpret them as the quantization of something) have spacetime aspects, but loose them through singular representations.

The crucial role of the GNS representation as well as the outline of a proof already appeared in the first chapter of the text. The difference in the infinite

 $^{^4 {\}rm States}$ can be more generally defined on *-algebras e.g. the Wightman theory and the closely related Borchers-Uhlmann tensor algebras of test functions.

dimensional case to the finite dimensional one is only in the topological aspects and the adaption may be done by the reader. The simplest situation is of course that of a faithful state for which the necessity to form equivalence classes in order to represent vectors is absent.

The GNS construction is the most valuable construction method for vNa, because the weak closure of the C^* -representation in the GNS Hilbert space is a natural vNa extension of C. The state is then automatically "normal" with respect to this generated vNa. (defined as continuity in terms of its predual), however other states on C do not have this property unless they lie in the same "folium" of states (i.e. states which correspond to density matrices on the vNa extension of the representation of C), in which case the vNa's are quasiequivalent. By this circumstantial definition it should already be clear to the reader that normality is some continuity property which uses a specific topology of vNa's (in fact it is the " σ -weak continuity"). The GNS construction permits to transfer properties from the vNa closure of the canonical GNS representation to the states and representations on the C*-algebra, e.g. one speaks of factorial representations and factor states of C. An important property in QT is the dominance of one state by another:

Theorem 48 $\omega_T(A) \leq \omega(A) \,\forall A \in A \iff \exists \text{ unique } T > 0 \in \pi_\omega(A)', ||T|| \leq 1$ with $\omega_T(A) = (T\Omega_\omega, \pi_\omega(A)\Omega) = (T^{\frac{1}{2}}\Omega_\omega, \pi_\omega(A)T^{\frac{1}{2}}\Omega_\omega)$

If a vNa is concretely given as an operator algebra in a Hilbert space, there are many ways to represent states by spatial vectors i.e the ambiguity is much larger than just a change of a representing vector by a phase. The modular theory furnishes additional concepts which delimit these possibilities. In particular the "natural cone representation" of states leads to a unique correspondence of states and vectors (presented in a later section).

As in elementary QM the notion of discrete eigenvectors was generalized by Dirac to the "improper" vectors in the continuous spectrum (δ -functions, or in more recent times also "rigged" Hilbert space), the notion of state finds its generalization in the theory of weights on C^* -algebras. In order to give meaning to the value ∞ being taken by "improper states", one should start with a functional which is only defined on the positive cone C_+ . There is a very good mathematical reason for introducing weights. There is also at least one physica reason. But in these notes we did not take up those issues.

The C*-algebra setting and the formalism of vNa is more fundamental for local quantum physics than the setting of functional integrals or other quantization methods as well as the fibre bundle methods of differential geometry. Besides to Schrödinger QM, the latter essentially apply only to those theories which allow a classical localization concept as the QFT version of Weylalgebras and CAR-algebras⁵. These are noncommutative C*-algebras over classical (test) function spaces. Apart from some extremely simple models (example

 $^{^{5}}$ This is why in QM and elementary QFT there is no need to leave the naive Hilbert space formulation which identifies (pure) states with unit rays created by vectors.

Schwinger's model), these methods just amount to a transcription of the perturbation theory into the language of euclidean field theory. Its extreme theoretical limitation (already the construction of the d=1+1 factorizing models explained in these notes is outside these quantization schemes) is only alleviated by the ease which with the greatest historical successes of QFT, namely the measured effects of QED and its electro-weak unification, were incorporated in that perturbative quantization framework. Again for historical and sociological reasons, the mathematical sophistication of the majority of quantum physicists during the last two decades developed more in the direction of differential geomety than towards operator algebras, as they were created with the inspiration from quantum physics by von Neumann. This is the main reason why presently a text on local quantum physics based on operator algebra methods needs a mathematical appendix as this.

A.3 Von Neumann Algebras and their Classification

Most of the following coarse grain classification of vNa's was already accomplished in the three papers of the founding fathers Murray and von Neumann . The strong connection with von Neumann's previous work on the foundation of QT and the measurement process is plainly visible in the key role of the projectors which are the simplest quantum observables.

We already mentioned the crucial notion of the commutant:

$$M' = \{ y \in \mathcal{L}(H) \mid [x, y] = 0, \forall x \in M \}$$
(A.2)

and its deep root in physics of quantum measurement as well as quantum localization⁶. The physical interpretation of the relation M'' = M is obvious. The fundamental "double commutant theorem" of von Neumann $Q'' = \overline{Q}^w$ (\overline{Q}^w is the weak closure of a *subalgebra Q of operators in $\mathcal{L}(H)$) converts aspects of topology into algebraic properties. A physically important algebraic method for manufacturing vNa's is to take commutants of representations of C*-algebras $M = \pi(C)'$, the most prominent case being a group algebra: $\mathcal{C} = \mathbb{C}G$ in which case one obtains the trivial case $M = \{\lambda \underline{1}\}$ only for irreducible representations. Most interest is focused on factors (vNa's with trivial center)

$$M \cap M' = \{\lambda \underline{1}\}\tag{A.3}$$

because general vNa's have a factorial decomposition. The von Neumann extension of an irreducible representation of a C*-algebra is a special case of a factor and the factorial property is transferred to the commutant. In physics the notion of "indecomposability" of a system of observables is both related to

⁶In LQP observables are called localized in a spacetime region \mathcal{O} if they commute with every spacelike disconnectedly localized observable, i.e. the localization in \mathcal{O} and its causal opposite (or dual) are inexorably linked. For same observable algebras (CCR, CAR), this can be implemented by the classical notion of support of functions.

irreducibility or factoriality, depending of whether one deals with global algebras which have pure states (e.g. vacuum, ground state of lowest energy) or local algebras which in LQP are typically factors with thermal properties. The only type of an vNa which admits pure states is $\mathcal{L}(H)$. They correspond to minimal one-dimensional projectors. The system of all projectors $\mathcal{P}(M)$ in a vNa form a lattice. It is clear that unitarily equivalent projectors should be considered as part of an equivalence class and the first aim would be to understand the class structure. In order to have coherence of this equivalence notion with additivity of orthogonal projectors, one need to follow Murray and von Neumann and enlarge the class of equivalent projectors in the following way:

Definition 12 Let $e, f \in \mathcal{P}(M)$, then

- 1. the two projectors are equivalent $e \sim f$ if there exists an partial isometry such that e and f are the source and range projectors: $u^*u = e$, $uu^* = f$
- 2. e is subequivalent to f, denoted as $e \leq f$ if $\exists g \in \mathcal{P}(M)$ such that g is dominated by f and equivalent to $e : e \sim g \leq f$

One easily checks that this definition indeed gives a bona fida equivalence relation in $\mathcal{P}(M)$. Via the relation between projectors and subspaces, these definitions and the theorems of the Murray von Neumann classification theory can be translated into relations between subspaces. The main advantage to restrict to factors is the recognition that any two projectors are then (sub)equivalent. One calls a vNa finite if a projector is never equivalent to a proper subprojector. Example: in $\mathcal{L}(H)$ infinite dimensional spaces allow a partially isometric mapping on infinite dimensional subspaces and therefore this factor is infinite. $Mat_n(C)$ is of course a finite factor. It was a great dicovery of Murray and von Neumann, that there are infinite dimensional finite factors. In fact they defined:

Definition 13 A factor M is said to be one of the following three types:

- 1. *I*, if it pocesses pure normal states (or minimal projectors).
- 2. II, if it not of type I and has nontrivial finite projectors.
- 3. III, if there are no nontrivial finite projectors.

Murray and von Neumann were able to refine their classification with the help of the trace. In more recent terminology a trace without an additional specification is a weight Tr with $Trxx^* = Trx^*x \ \forall x \in M$. A tracial state is a special case of a tracial weight.

The use of tracial weights gives the following refinement:

Definition 14 Using normal tracial weight one defines the following refinement for factors:

1. type I_n if $ranTr\mathcal{P}(M) = \{0, 1, ..., n\}$, the only infinite type I factor is type I_{∞} . Here the tracial weight has been normalized in the minimal projectors (for finite n this weight is in fact a tracial state).

- 2. type II_1 if the Tr is a tracial state with $ranTr\mathcal{P}(M) = [0, 1]$; type II_{∞} if $ranTr\mathcal{P}(M) = [0, \infty]$.
- 3. type *III* if no tracial weight exists i.e. if $ranTr\mathcal{P}(M) = \{0, \infty\}$.

In particular all nontrivial projectors (including 1) are Murray-von Neumann equivalent.

The classification matter rested there, up to the pathbreaking work of Connes which in particular led to an important gain in understanding and partially resolving type *III* factors. Since the modular theory is heavily used, we will mention some results later on.

1. Although nowhere explicitly stated, all indications are that von Neumann believed on the basis of his analysis of physical observables, that only the type I with its pure states⁷ was of physical interest and that in particular type II and the associated idea of a "continuous geometry" (because of the continuous range of the notion of "dimension") was of exclusive mathematical importance. We know nowadays that the most important algebra of LQP is the hyperfinite type III_1 factor. It is the carrier of the physical localization properties (including the thermal aspects) and the spacetime symmetries and is (from the modular point of view) the "most noncommutative" factor in the sense that its folium contains only "very noncommutative states". It also appears for heat bath thermal representations in the thermodynamical limit. Type I reappears through the representation theory of globalizations of nets. One of the special very comforting properties of typeIII is that any isomorphism or endomorphism (in particular automorphism) is unitarily implementable in a standard representation. Type *II* do not enter physics directly because they are too small in order to incorporate spacetime symmetries and localization. However they enter via the algebras generated from intertwiners in the decomposition theory of localized charges by "freezing" the localization regions of the latter, together with "Markov traces" on these intertwining algebras, which results from the physical cluster decomposition property after freezing. These combinatorical algebras are better known under the name of topological field theories because this is the way they appear if one tries to construct them by functional quantization of classical Chern-Simons Lagrangians. Since localization is a condition sine qua non for physical interpretation, it is always important (for physics) to know their position inside a typeI or III factor from which they originate. TypeII vNa's also appear in Hofsteadters work on QM in a constant magnetic background (noncomutative tori, well-known from the work of Connes and Rieffel). It is believed that singular states on extended Weyl like C^* -algebras, associated with Chern-Simons Lagrangians via quantization, lead directly to $TypeII_1$ vNa's.and

⁷In the general mathematical theory the notion of purity of states is not an important one. For the decomposition theory of states the concept of factor (central) or in physical terms into superselected charge sectors is more important.

constitute the operator algebra version behind Witten's differential geometric receipes on functional integrals involving Chern-Simons actions which lead to topological field theories outside the standard setting of euclidean Feynman-Kac representations of real time QFT. The subalgebras on which singular states become regular should then agree with the alias "gauge invariant" subalgebras.

A.4 Modular Theory

Returning to the analogy with the commutative case $L^{\infty}(X,\mu) \sim B(H) = \mathcal{L}(H), L^2(X,\mu) \sim H$, and adding the illustrative example of the type II₁ tracable algebra, we realize that there are different grades of noncommutativity. Whereas the first case is entirely commutative and the second case noncommutative, the type II₁-algebra's with a trace is somewhere in the middle, because although it is noncommutative, the operators of course commute inside the tracial state. For these noncommutative algebras the Tomita-Takesaki modular theory allows to measure the degree of noncommutativity with respect to a GNS state. Since there are no tracial states or even weights on type *III*, they are in a way the most noncommutative algebras. Among these the physically relevant (as a carrier of spacetime localization) type *III*₁ sticks out since all states are equally noncommutative i.e. there are no inner relations in the modular group for different states.

Take as an example the I_{∞} factor (the modular concepts for the I_n factor were already explained in chapter1 of the main text). In order to describe it as the result of the GNS construction with a faithful state φ on $\mathcal{L}(H)$, let ρ be a trace class operator such that in its spectral representation in terms of a orthonormal basis $|i\rangle$ it has strictly positive (nonvanishing) components λ_i for all $i = 1...\infty$.

$$\rho = \sum_{i} \lambda_{i} |i\rangle \langle i|, \sum_{i} \lambda_{i} = 1$$

$$\varphi(x) = tr\rho x$$
(A.4)

This garanties the faithfulness of the state and in complete analogy to the matrix case in chapter1, the GNS construction leads to a Hilbert space of Hilbert-Schmidt operators \mathcal{H} which is generated by the algebra $\mathcal{L}(H)$ together with the inner product $(x; y) = tr\rho x^* y$. The reader immediatly realizes that in addition to the GNS left action of $\mathcal{L}(H)$ on $\mathcal{H} \equiv L^2(\mathcal{L}(H), \varphi)$ one also has an opposite right action such that $\mathcal{L}(H)' = \mathcal{L}(H)_r^{opp}$. In fact at this point it is helpful to take some hint from the standard Gibbs heat bath description by imagining a $\rho = \frac{e^{-\beta K}}{tre^{-\beta K}}$ with an hamiltonian K with discrete positive spectrum such that the trace exists. Then the same calculation as done in chapter1 with matrices will lead to:

$$Sx\rho = x^*\rho$$

$$(A.5)$$

$$\sim S = J\Delta^{\frac{1}{2}}$$

where in complete analogy to the finite dimensional case the J represents a antiunitary flip from the left action to its opposite and $\Delta^{\frac{1}{2}} = \pi_l(\rho) \otimes \pi_r(\rho^{-1})$. The last formula for $\Delta^{\frac{1}{2}}$ corresponds to the physical fact that $\Delta^{it} = e^{-i\beta K_{th}}$ with the thermal hamiltonian (the generator of time translations in the thermal state) K_{th} is the sum of the left hand minus the right hand action of the original hamiltonian K on:

$$H \otimes H \simeq L^2(\mathcal{L}(H), \varphi)$$
 (A.6)

as in the finite dimensional case. Whereas in the adjoint action on the observable algebra $\mathcal{L}(H)$ the difference cancels, in the applications to vectors in \mathcal{H} , only H_{th} gives the correct time propagation without (in the thermodynamic limit $V \to \infty$) thermal fluctuations in the reference state ρ . This is the setting which led Haag, Hugenholz and Winnik to there formulation of the KMS condition and their emphasis of the fact that this condition survives the thermodynamical limit (which is generally outside type I) and is responsible for the stability of these thermal states. Another easily accessible illustration is that of a $typeII_1$ factor for which one can directly work with the (unique) tracial state. The situation leads to similar formulae i.e. the algebra acts left and right (the opposite) on itself and the antiunitary J act in the same way but now $\Delta =$ 1, i.e. the KMS formula is trivial. In both cases the GNS construction led to a GNS tripel $(M; H, \Omega)$ with Ω a cyclic and separating vector in H of M. The separating property results from the faithfulness of the state and means that $x\Omega = 0 \curvearrowright x = 0$. A vNa with these extra trimmigs is called "in standard form" (strictly speaking this terminology is only used if in addition one uses the natural cone described below). Now we are well-prepared to state the result of the Tomita-Takesaki theory in the most general setting:

Theorem 49 Let M be a vNa in standard form (all vNa's with separable preduals permit this standard form). Then the Tomita involution:

$$Sx\Omega \equiv x^*\Omega$$
 (A.7)

is closable and hence permits a unique polar decomposition:

$$S = J\Delta^{\frac{1}{2}} \tag{A.8}$$

The adjoint action of the antiunitary Tomita involution J and the bounded one parametric group generated by Δ^{it} on the algebra is the following:

$$adJM = M'$$

$$ad\Delta^{it}M = \sigma_t(M) = M$$
(A.9)

where σ_t is called the modular automorphism group. The modular construction can be generalized to the case of any separable vNa with a faithful normal weight.

The proof of this theorem is simple for type I and II_1 but somewhat demanding in the type III cases, with the type III_1 being the most demanding. In the most general case with unbounded S one must use analyticity techniques (which the older generation QFT-physicists is familiar with because similar analytic methods have been used in Wightman QFT) which entered through the relation with the KMS theory. We refer the reader to the literature at the end of this appendix. The form of modular operators for type I factors is known whereas the general form for type *III* factors is unknown. More profound than the above relation with heat bath physics is the relation with local algebras $\mathcal{A}(\mathcal{O})$ of the nets of LQP. These algebras are hyperfinite type III_1 and therefore belong to the nontrivial application of the T-T modular theory (the subscript refers to Connes modular refinement of the type classification. The unboundedness of Δ and the fact that the modular automorphism is σ_t outer (for all values of t) with respect to M, makes this algebra more noncommutative than the other types and it turns out, that it is precisely this situation which is necessary in order to have modular groups with relations to spacetime symmetry. The KMS condition allows a direct check if an automorphism σ_t is a modular automorphism of a given state φ on M by checking that it satisfies the KMS boundary condition.

Definition 15 A state φ is said to fulfil the KMS boundary condition (at inverse temperature $\beta = 1$) with respect to a an automorphism σ_t of a C^{*}-algebra C if $\varphi(\sigma_t(x)y)$ is a continuous (in t) boundary value of a strip analytic function F(z), -1 < Imz < 0 with:

$$F(t) = \varphi(\sigma_t(x)y)$$

$$F(t+i) = \varphi(y\sigma_t(x))$$
(A.10)

In extending the state to the vNa $M \equiv \pi_{\varphi}(\mathcal{C})''$ there arises the question whether the automorphism extends to this vNa in a natural way or in technical terms whether it has has a σ -weak continuous extension. This turns out to have an affirmative answer if the C*-algebra is unital and φ is a σ_t invariant state. It furthermore turns out that this property is characteristic i.e. such a σ_t which fulfils the KMS condition for a faithful state is necessarily the modular automorphism group a T-T theory of any associated GNS representation. The importance of the KMS condition in quantum statistical mechanics of open systems (i.e. in the thermodynamic limit) is related to their stability against local perturbations.

Whereas the Δ^{it} depends on the realization of the GNS representation in the class of the unitarily equivalent realizations, the modular automorphism only depends on the state and not on its representing vector. In order to have the converse, one need to generalize from states to weights. In that case Connes showed that each cocycle related modular group $u_t^*\sigma_t(\cdot)u_t$ is the modular group of a unique normal weight

$$\omega_u(\cdot) \equiv \lim_{t \to \frac{i}{2}} \omega(u_t \cdot u_t^*) \tag{A.11}$$

A useful generalization is the relative modular theory which refers to two cyclic and separating vectors say Ω, Ω' simultaneously. One defines the conjugate linear operator:

$$S_{\Omega',\Omega} x \Omega = x^* \Omega, \ x \in M \tag{A.12}$$

Again the closability is easily established and the polar decomposition leads to a positive operator $\Delta_{\Omega',\Omega}^{\frac{1}{2}}$ which is called the relative modular operator of the pair Ω', Ω . If Ω' is in the natural cone of Ω (see below), $J_{\Omega',\Omega} = J_{\Omega,\Omega} = J$. Defining the unitaries $U_{\Omega',\Omega}(t) \equiv (\Delta_{\Omega',\Omega})^{it}$ one finds the following symmetric (between M and M') relations:

$$U_{\Omega',\Omega}(t)xU_{\Omega',\Omega}^{*}(t) = U_{\Omega'}(t)xU_{\Omega'}^{*}(t), \ x \in M$$

$$U_{\Omega',\Omega}(t)x'U_{\Omega',\Omega}^{*}(t) = U_{\Omega}(t)xU_{\Omega}^{*}(t), \ x' \in M'$$
(A.13)

with $U_{\Omega'}(t) \equiv \Delta_{\omega'}^{it}, U_{\Omega}(t) \equiv \Delta_{\omega}^{it}$. Using another "spectator" cyclic separating vector Ω'' the definition

$$u_{\Omega',\Omega}(t) \equiv U_{\Omega',\Omega''}(t)U^*_{\Omega,\Omega''} \tag{A.14}$$

gives a unitary which commutes with M' and therefore lies in M. The spectator Ω'' was chosen because this unitary turns out to be independent of Ω'' . This operator is the famous Connes cocycle which relates the modular operators of the two states $U_{\Omega'}(t) = u_{\Omega',\Omega}(t)U_{\Omega}(t)$:

$$u_{\Omega',\Omega}(t_1 + t_2) = u_{\Omega',\Omega}(t)\sigma_{t_1}^{\omega}(u_{\Omega',\Omega}(t_2))$$

$$\sigma_t^{\omega'}(\cdot) = u_{\Omega',\Omega}(t)\sigma_t^{\omega}(\cdot)u_{\Omega',\Omega}^{*}(t)$$

$$(D\omega' : D\omega)(t) = u_{\Omega',\Omega}(t)$$
(A.15)

The notation of the last equation suggest to view the Connes cocycle as the noncommutative generalization of the classical Radon-Nikodym concepts. It was already mentioned before that given the modular group σ_t^{ω} and a Connes cocycle, one can construct a state ω' such that its modular group $\sigma_t^{\omega'}$ is cocycle related to σ_t^{ω} with one caveat: the state may be improper i.e. a weight.

The modular theory can be used for strengthening the relation between states ω and representing vectors $\xi(\omega)$. To this end one starts from the vNa in standard form (M, H, Ω) and defines in H the "natural cone" $\mathcal{P}^{\#}$ as the weak closure of a convex set:

$$\mathcal{P}^{\#} = \frac{\overline{\Delta^{\frac{1}{4}} M_{+} \Omega}}{\{J x J x \Omega, x \in M\}}$$
(A.16)

The equality of the two expressions is derived from the relation $\Delta^{\frac{1}{4}}xx^*\Omega = \sigma_{-\frac{i}{4}}(x)J\sigma_{-\frac{i}{4}}(x)\Omega$. For entire analytic operators $x \in M_0$ i.e. such that $\sigma_t(x)$ has an analytic continuation to an entire function $\sigma_z(x)$ this relation is easily derived and with $y \equiv \sigma_{-\frac{i}{4}}(x) \in M$ we obtain the desired identity which, as a result of the denseness of M_0 in M can be continued to all M. One then has the following unicity theorem:

Theorem 50 $\xi(\omega)$ is the only vector in $P^{\#}$ such that

$$(\xi(\omega), x\xi(\omega)) = \omega(x), \ x \in M$$

$$(A.17)$$

$$(\xi(\omega), x'\xi(\omega)) = \omega(AdJ(x'^*)), \ x' \in M'$$
the map $\xi \in \mathcal{P}^{\#} \to \omega_{\xi} \in M_{*+}$ is a homeomorphism $\mathcal{P}^{\#} \to M_{*+}$ in the norm topology with $\|\xi - \eta\|^2 \leq \|\omega_{\xi} - \omega_{\eta}\| \leq \|\xi - \eta\| \|\xi + \eta\|$ and the inverse map $\omega \to \xi(\omega)$ is monotonously increasing and concave with respect to the natural ordering of the two cones.

In the case of dominance $\omega \leq \omega_{\Omega}$, one obtains an explicit formula for $\xi(\omega)$:

$$\xi(\omega) = \left| A' \Delta^{-\frac{1}{2}} \right| \Omega \tag{A.18}$$

Here $A' \in M'$ is the unique operator which corresponds according to the dominance theorem to the state ω namely $\omega(\cdot) = (A'\Omega, \cdot A'\Omega)$, whereas |B| denotes the positive part in the polar decomposition of B. With a little extra efford one can proof that $A' = u_{\frac{1}{2}}$

Another physically useful aspect of the natural cone formalism is the fact that automorphism (and endomorphisms) $\alpha \in Aut(\mathcal{A})$ can be naturally unitarily implemented, i.e. $\exists U(\alpha)$ with:

$$U(\alpha)AU^{*}(\alpha) = \alpha(A)$$

$$U(\alpha)\mathcal{P}^{\#} \subseteq \mathcal{P}^{\#}$$

$$[U(\alpha), J] = 0$$
(A.19)

In fact $U(\alpha)\xi(\omega) = \xi(\alpha^{-1*}(\omega))$ where $(\alpha^*\omega)(A) \equiv \omega(\alpha(A))$. The natural cone theory is also the starting point for Connes reconstruction of algebras from the spatial data of the position of the natural cone inside the Hilbert space $\mathcal{P}^{\#}$ $\subset H_{\Omega}$. Reconstructions of (nets of) algebras from (nets of) positions of real subspaces $H_R = \mathcal{P}^{\#} + -\mathcal{P}^{\#}$ is an important issue in chapter 6. In the main text it was not necessary to use this construction (because of the existence of a reference algebra of the "in net" supplied by scattering theory permitted the introduction of a modular Møller operator).

The application of modular theory to the problem of classification of von vNa's led Connes to a complete understanding of all hyperfinite vNa's. Hyperfinite means that they can be approximated by finite dimensional vNa's or equivalently that the algebra is in the range of a conditional expectation in $\mathcal{L}(H)$. The physical intuitive content is that the associated physical systems allow an interpretation as a thermodynamic limit of a sequence of finite systems or equivalently, in case of properly infinite LQP, an approximation in terms of QM (type I) systems. All local vNa's $\mathcal{A}(\mathcal{O})$ are hyperfinite, but there globalizations sometimes involve "free product" constructions which lead out of hyperfiniteness.

A.5 C*-Algebras related with Bimodules

In chapter 7 of the main text we have seen that the DHR theory leads to the (reduced) field bundle which is really an algebraic bimodule on which the observable algebra acts from the right through the identity representation and to the left through its endomorphic image. This theory led to the theory of which define a structure. The DR analysis in the case of d=1+3 QFT finally led to a complete understanding of an associated field algebra with a compact group action (symmetry). For this result to emerge algebraic Hilbert spaces and associated Cuntz C*-algebras played an important role. For d $\leq 1 + 2$ QFT one neads a yet unknown generalization. It is a natural idea to look for generalizations within the setting of bimodules (containing Hilbert spaces as special cases) and their naturally associated algebras. Assume that $\mathcal{A} \subset \mathcal{B}$ and denote by \mathcal{C} the relative commutant $\mathcal{C} = \mathcal{A}' \cap \mathcal{B}$. Define a subset of \mathcal{B} :

$$X_{\rho} = \{ \psi \in \mathcal{B} \mid \psi A = \rho(A)\psi \}$$
(A.20)

where for the endomorphism of \mathcal{A} we used the standard notation ρ . Define now the \mathcal{C} -valued inner product:

$$\left\langle \psi, \psi' \right\rangle_{\mathcal{C}} = \psi^* \psi' \tag{A.21}$$

such that $\|\langle \psi, \psi \rangle_{\mathcal{C}}\| = \|\psi\|_{\mathcal{B}}^2$. If X_{ρ} is finite projective, we call ρ inner in \mathcal{B} . One immediatly realizes that by setting $\mathcal{C} = C\mathbf{1}$ but allowing $A \neq B, A' \cap B = C\mathbf{1}$ i.e. irr. inclusion, X_{ρ} is a Hilbert space in \mathcal{B} and ρ the restriction to \mathcal{A} of an inner endomorphism in \mathcal{B} i.e. $\rho(B) = \sum_{i=1}^{n} \psi_i B \psi_i^*$, with $\psi_i, i = 1, ..., n$ an orthonormal basis in X_{ρ} . In that case the DH-theory shows that \mathcal{B} is a crossed product of \mathcal{A} by the action of a compact group. However the universal algebra of the algebraic compactification of $d \le 1 + 2$ QFT's with braid group statistics in chapter 7 is a C*-algebra with nontrivial center and there are indications that a would be field algebra \mathcal{B} may be such that the relative commutant is nontrivial $A' \cap B \neq C\mathbf{1}$. In that case it may be interesting to look for a symmetry concept which is related to a more general crossed product \mathcal{B} associated to the pair $\{\mathcal{A}, \rho\}$ and to find reasonable conditions for \mathcal{C} which insure uniqueness of \mathcal{B} . A special situation of this type, which is mathematically interesting in its own right is the situation where X is given as a Hilbert C*-bimodule with coefficients in \mathcal{C} , i.e. a right Hilbert \mathcal{C} -module with a monomorphism of C into $\mathcal{L}(X)$ defining the left action. In that case the bimodule tensor powers X^r may be considered as the objects of a C*-category (with those adjointable right module maps which commute with the left action of \mathcal{C} being the arrows of the category). There is a functorial construction (used in the DR theory) which relates a C^{*}-algebra \mathcal{O}_X with such a situation. Algebras as this and their subalgebras (example: Cuntz-Krieger algebras) from such a bimodule viewpoint have been first studied by Exel and Pimsner. For a presentation which is most close to the spirit of LQP I refer to the paper of Doplicher et. al. below (where also furthergoing ivestigations closer to the symmetry problem of LQP have been announced).

A.6 Conditional Expectations, Canonical Endomorphisms

The knowledge about conditional expectations on vNa's of QFT physicist is in the majority of cases limited to the abelian case which is of course probabilistic cradle of this concept. The renormalization group manipulations in eucledian field theory where one often decimates degrees of freedom as well as Nelson's Markov property uses either tacitly or explicitely such concepts. As in the abelian case of measure spaces and the associated $L^{\infty}(X,\mu)$ algebra, one defines a conditional expectation $E: M \to N$ from M to a subalgebra $N \subset M$ as a projection of norm 1, i.e. a completely positive normalized (unit preserving) map⁸ E with $E(y^*xy) = y^*E(x)y, y \in N$. The prototype noncommutative example is obtained by having a group G act on M (example free field of chapter 3 of a multicomponent field with a SU(N) action) and using for N the fixpoint algebra $N = M^G$. A conditional expectation for this situation is given by the Haar average as in chapter 7. If the group is not compact, one obtains an improper version of E which is called a operator-valued weight and has the analog relation to a bona fide E as a weight has to a state.

There is one crucial difference to the commutative situation: a conditional expectation (or its improper version) does not exist for each pair $N \subset M$. Instead one finds a fascinating relation between conditional expectations and modular theory as described in the following theorem of Takesaki:

Theorem 51 A modular group σ_t of (M, Ω) leaves a subfactor N invariant iff the inclusion $N \subset M$ possesses a normal (σ -weak continuous) Ω -invariant conditional expectation E iff the modular group σ_t of M restrict to the corresponding modular objects of N on $\overline{N\Omega}$ and the equality $\overline{N\Omega} = \overline{M\Omega}$ holds only iff M = N. In that case also the modular conjugation of M restricts to that of N.

Let us indicate the proof. Assuming that the modular theory restricts, we define $E: M \to N$ as PxP = E(x)P where P projects on $\overline{N\Omega}$. This definition would be reasonable if we can show that $PxP = yP, y \in N$, because then by using the existence of a U with $U^*U = P, UU^* = P^{\perp}$, one defines $E(x) = PxP + P^{\perp}UxU^*P^{\perp}$ which verifies the above relation and the properties of a conditional expectation. In order to show the required property we start from the evident inequality $NP \subseteq PMP$ and show the opposite inequality \supseteq : $PMP \subseteq PJ_MN'J_MP = J_MP'N'PJ_M = J_NPN'PJ_N = NP$ where in the first step we used $M \subseteq J_M N' J_M$. This shows the \rightarrow part. Reversely, let us assume the existence of an $E: M \to N$ preserving the vector state Ω . Define $Pa\Omega = E(a)\Omega \in H_N$ and obtain $(a - E(a))\Omega \in H_N^{\perp}$. From $a \in M, b, c \in N$ and $(\Omega, c^* PaPb\Omega) = (\Omega, c^* E(ab)\Omega) = (\Omega, c^* E(a)b\Omega)$ obtain PaP = E(a)P. Now study the Tomita operator on $H_N: S_M P a \Omega = S_M E(a) \Omega = E(a^*) \Omega = P S_M a \Omega$. From the denseness of states $M\Omega$ obtain $S_M P = PS_M$ which means that S_M leaves H_N invariant and restricts to S_N . The rest follows by polar decomposition.

As weights constitute a generalization of states, the natural generalization of conditional expectations are the operator valued weights. In both cases the generalizations are necessary in order to be able to invert or dualize certain

 $^{^{8}}$ All of our endomorphisms, operator valued weights and endomorphism are always assumed to be injective so that they can be inverted on their image.

relations. The conversion back of a cocycle related modular group to a state of which it is the modular group will have an obstruction which is removed by allowing weights. Similarly another theorem of Connes and Hagerup states that the dual inclusion $M' \subset N'$ for any conditional expectation of the original inclusion $E \subset C(M, N)$ possesses a dual $E^{-1} : N' \to M'$ which generally turns out to be an operator valued weight $E^{-1} \subset P(N', M')$. The latter is a singular (not defined on 1) generalization of a conditional expectation in a similar sense as weights are singular generalizations of states.

We will see in the next section that the group average example for a conditional expectation is in a certain sense typical; conditional expectations are always related to what a physicist calls an internal symmetry which structure the relation between observable nets and field nets. A different kind of inclusion which in the next section is shown to be relates to space-time symmetries is given in terms of the so-called canonical endomorphism which was introduced by Longo. The construction is as follows. Start from an inclusion of properly infinite vNa's $N \subset M$ on a separable Hilbert space H. Then there exist vectors $\Omega \in H$ which are cyclic and separating for both algebras (for finite algebras such vectors do definitely not exist and therefore there is no canonical endomorphism in the sense below). With the help of the corresponding $J_{M,N}$ and their adjoint actions $j_{M,N}$ we define the canonical endomorphism γ :

$$\gamma = j_N j_M |_M \in End(M) \tag{A.22}$$

It maps M into a subalgebra N_1 of N and the j's can also be used in order to define a canonical extension:

restr. :
$$N_1 \equiv j_N j_M(M) \subset N$$
 (A.23)
ext. : $M \subset M_1 \equiv j_M j_N(N) = \gamma^{-1}(N)$

In fact without changing the Hilbert space one can keep on going into both directions by defining $\rho \equiv j_{N_1} j_N \in End(N), \gamma_1 \equiv j_M j_{M_1} \in End(M_1)$ with the standardness of the new algebras with respect to Ω always being maintained. The equality between modular conjugations of a vNa and its commutant leads to:

$$J_{N_1}J_N = J_N J_N J_{N_1} J_N = J_N j_N (J_{N_1}) = J_N J_M$$
(A.24)
= = J_M J_{M_1}

and hence $\rho = \gamma|_N$ and $\gamma = \gamma_1|_M$. Finally we notice that the inclusions (A.23) are isomorphic thanks to $N = \gamma_1(M_1)$ and $N_1 = \gamma(M) = \gamma_1(M)$. For plausible reasons they are called "dual" relative to $N \subset M$. They are anti-isomorphic to $M' \subset N'$ by application of j_N and in the same vein $M'_1 \subset M'$ is the anti-isomorphic image of $N \subset M$ under j_M . The "tower" and "tunnel" created by positive and negative powers γ^n applied to M resp. N become the even resp. odd parts of the Jones tower/tunnel associated with subfactors if the inclusion $N \subset M$ permits a conditional expectation.

A.7 Deep Inclusions and (spacetime) Geometry

Geometric inclusions, as e.g. a double cone localized algebra in LQP containing a localized algebra of a genuinly smaller double cone, do not permit vacuum invariant conditional expectations because the modular groups are very different. In the sense of the geometrical interpretation coming from LQP they are "deep"i.e they have a large relative commutant $N' \cap M$. There is a particular deep family of inclusions which is susceptible to profound investigations by modular methods. They are the so-called "half-sided modular" inclusions as well as the "modular intersections". Since many of their properties were already mentioned in the main part of the notes, we will be content with referring to some additional remarks in the small print at the end of chapter 3.5.

A.8 Shallow Inclusions and (internal) Symmetry

The existence of conditional expectations is the prerequisite for the Jones theory of subfactors, as well as for its older LQP analogon, the DHR theory of (localized) endomorphisms. In more recent times, there has been a merger of the two into the sector theory of Longo, which is with increasing frequency also used by mathematicians:. It is very closely related with Connes theory of correspondences (bimodules). Bimodules also have been used in the 1970 work of DHR on field bundles. In order to have unimpeded unitary implementation of endomorphisms we will assume type *III* factors. Let $L^2(M)$ be the standard Hilbert space with the modular conjugation J_M . Let us denote $L^2(M)$ as H_{ρ} if we use it as a representation space of $M - M^{opp}$ acting as $m_1 \cdot \xi \cdot m_2 = \rho(m_1) J_M m_2^* J_M$. In this way an endomorphism and $L^2(M)$ give rise to a bimodule H_{ρ} . Since according to our assumption every isomorphism is spatial, one easily sees that each M - M bimodule, whether originating in the indicated way or not, can always be written in the form of an H_{ρ} and unitary equivalent bimodules have inner conjugate $\rho's$. We define:

$$Sect(M) = End(M)/Int(M)$$
 (A.25)

i.e. sectors are equivalence classes of *End* or *Bim* according to taste. Physicists prefer the *End* because the composition rule follows more standard procedures (ordinary composition of endomorphisms):

$$H_{\rho_1} \otimes_M H_{\rho_2} = H_{\rho_2 \rho_1} \tag{A.26}$$

The natural setting for this bimodule calculus is the Connes spatial modular theory. The sector notation is the standard equivalence class notation: $[\rho] \in Sect$, if $\rho \in End$.

Intertwiners are isometric operators V in M which relate endomorphisms $V: \rho_1 \to \rho_2$. They form linear spaces:

$$(\rho_1, \rho_2) = \{ V \in M; V \rho_1(x) = \rho_2(x) V \}$$
(A.27)

$$= Hom(H_{\rho_1}, H_{\rho_2})$$

In case ρ_1 is irreducible, i.e. $M \cap \rho_1 = \mathbf{C} \cdot \mathbf{1}_M$ we can define an inner product:

$$\langle V|W\rangle \mathbf{1}_M \equiv V^*W, \ V, W \in (\rho_1, \rho_2)$$
 (A.28)

In particular the selfintertwiners correspond to relative commutants:

$$(\rho, \rho) = M \cap \rho(M)' \tag{A.29}$$

The index of an inclusion of factors is defined if $N \subset M$ possesses a conditional expectation E. In terms of this conditional expectation, the definition is:

$$Ind(E) := E^{-1}(1)$$
 (A.30)

where E^{-1} denotes the operator valued weight which corresponds to E by applying the previously mentioned Connes bijection between operator valued and their duals. The right hand side lies in the center of M and therefore is a multiple of unity: $Ind(E) \in [1, \infty]$ with ∞ if the identity is not in the range of the operator valued weight E^{-1} . If it is finite, then in fact the operator valued weight E^{-1} can be normalized by $E' = Ind(E)^{-1}E^{-1}$ and defines the dual conditional expectation which belongs to $M' \in N'$ and fulfils $E'^{-1} = Ind(E)E$ and hence Ind(E') = Ind(E) with a central valued Ind(E). The set of normal conditional expectations consists of more than one element if the inclusion is not irreducible. In that case one uses the so called minimal index which corresponds to the previously defined minimal conditional expectation E_0 :

$$[M:N] := \inf_{E} Ind(E) = Ind(E_0) \tag{A.31}$$

This index has a series of remarkable properties which are best formulated and understood by introducing the concept of statistical dimension:

Definition 16 The statistical dimension of an inclusion is the square root of its minimal index $d \equiv [M:N]^{\frac{1}{2}}$

Using this concept in the above fusion and decoposition formalism of finite index endomorphisms we find that d_{ρ} follows precisely the rules of a dimensional function in that it is additive and multiplicative:

$$d_{\rho} = \sum_{i} d_{\rho_{i}}, \ \rho = \sum_{i} \rho_{i}$$
(A.32)
$$d_{\rho_{1}\rho_{2}} = d_{\rho_{1}}d_{\rho_{2}}$$

It is one of the advantages of type *III*, that one can also freely add endomorphisms. For adding *n* of them one selects *n* isometries $v_i, i = 1...n$ with orthogonal ranges $v_i^* v_j = \delta_{ij}$ and total range one $\sum_i v_i v_i^* = \mathbf{1}$ and defines the direct sum endomorphism ρ :

$$\rho = v_1 \rho_1 v_1^* \oplus \dots \oplus v_n \rho_n v_n^* \tag{A.33}$$

330

The presentation of sectors would be incomplete without introducing the conjugate $\bar{\rho}$. For a LQP physicist, the conjugate is represented by the "anticharge" which is carried by the antiparticle. The latter if brought together with the particle can annihilate into the vacuum sector $p + \bar{p} \rightarrow vac + \dots$ (In the case of abelian charges the resulting charge is the vacuum charge without additional contributions).

Theorem 52 Let $\theta \in Sect(M)$ be an irreducible sector with $\theta \overline{\theta} = \overline{\theta} \theta$. Then θ has finite index iff the sector $\theta \overline{\theta}$ contains the identity. In that case $\theta \overline{\theta}$ and $\overline{\theta} \theta$ contain the identity with multiplicity one and if θ' is any other sector such that $\theta' \theta \supset id$, then $\theta' = \overline{\theta}$

In the case of finite statistical dimensions there exists a canonical way to implement conditional expectations by isometries. An important special case is the following:

Theorem 53 Let $N \subset M$ be an irreducible inclusion with finite index. Then there exists two isometries $v \in M, w \in N$ with $vm = \gamma(m)v \ \forall m \in M$ and $wn = \gamma(n)w \ \forall n \in N$ such that

$$w^* v = [M:N]^{-\frac{1}{2}} \mathbf{1} = w^* \gamma(v)$$
 (A.34)

$$E(m) = w^* \gamma(m) w : M \to N \tag{A.35}$$

Furthermore M can be represented in terms of N by adjunction of a single basis element $v: m = [M:N] \cdot E(mv^*)v$, i.e. M = Nv.

Remark 8 The last representation is the typeIII adaption of the well known representation in terms of a Popa-Pimsner basis which for finite factors consists if a basis whose (minimal) dimension is related to the index of the inclusion.

Having explained the meaning of conjugate for finite index endomorphisms, one may formulate the Frobenius reciprocity relations for intertwiner spaces. They are completely analogous to their classic group theoretical counterpart:

$$(\rho_{\gamma}, \rho_{\alpha}\rho_{\beta}) \xrightarrow{\eta} (\rho_{\gamma}\bar{\rho}_{\beta}, \rho_{\alpha}) \tag{A.36}$$

Here the Frobenius isomorphism η is a linear map which can be explicitly in terms of an intertwiner basis and allows (as all other intertwiner relations) a nice graphical representation. In fact the intertwiners and their properties form a so called C^{*}-category.

From the point of view of subfactor theory the endomorphisms are are special realizations of a given inclusion $\rho(M) = N \subset M$. Any type *III* inclusion allows for a endomorphic representation: just compose any isomorphism of M with N with the injection map $N \hookrightarrow M$. Since type *III* factors have myriads of isomorphisms, the endomorphic representations are highly arbitrary. In terms

of an ρ the canonical endomorphism can be written as $\gamma = \rho \bar{\rho}$ and the two "charge-anticharge intertwiners $R \in (id, \rho \bar{\rho})$ and \bar{R} may be shown to fulfil:

$$\rho(R^*)\bar{R} = d_{\rho}^{-1}\mathbf{1} = \bar{\rho}(\bar{R}^*)R \tag{A.37}$$

This situation invites to define a left inverse ϕ of ρ . This is a completely positive normal linear map defined by:

$$\phi(x) = R^* \bar{\rho}(x) R, \ x \in M \tag{A.38}$$

It gets its name from the relation $\phi \rho = id$ which is a consequence of the easily established relation:

$$\phi(\rho(x_1)x\rho(x_2)) = x_1\rho(x)x_2, \ x_i, x \in M$$
(A.39)

Whereas automorphisms have an inverse, injective endomorphisms only possess a left inverse in the above sense unique under the assumed finite index and irreducibility conditions). The left inverse is an important tool which the physicist introduced in order to obtain natural trcial states on the $(\rho^n, \rho^n), n = 1, 2, ...\infty$ intertwiner algebras which contain the multiparticle statistic operators. The inverse product $E = \rho \phi$ is easily shown to define a conditional expectation $M \to \rho(M)$. for the reducible case there are several left inverses and conditional expectations and one defines the minimal left inverse which is also called the standard left inverse) such that it corresponds to the minimal conditional expectation which was related to the minimal index [M:N] (Its square root is the statistical dimension d_{ρ}).

A.9 Split Property, Localizing Map and Nuclearity

As a generalization of the standardness of a representation of a single vNa (always with separable predual), one calls a pair (N, M) of vNa M and sub vNa $N \subset M$ standard, if there exists a faithful normal state ω such that the GNS representation $(\pi_{\omega}, \Omega_{\omega}, H_{\omega})$ gives rise to an inclusion $\pi_{\omega}(N) \subset \pi_{\omega}(M)$ which is standard to the vector Ω_{ω} . This last terminology means that in addition to $\pi_{\omega}(M)$, this vector is also standard with respect to and the "collar" $\pi_{\omega}(N)' \cap \pi_{\omega}(M)$. We will identify the abstract vNa and their GNS representations π_{ω} .

An inclusion (N, M) is called split if there exists an intermediate type I factor B:

$$N \subset B \subset M \tag{A.40}$$

A type *I* factor defines a tensor product split of the Hilbert space $H = H_1 \bar{\otimes} H_2$ such that $B = B(H_1)$. In this case there exists an isomorphism Φ of $N \vee M'$ (the commutant of the collar) to the tensor product $N \bar{\otimes} M'$

$$\Phi(n \cdot m') = n\bar{\otimes}m', \ n \in N, \ m' \in M' \tag{A.41}$$

332

The additional property resulting from standard split is that the middle factor B can be adjusted so that this isomorphism can be canonically implemented. For this we use the standardness of $N \cap M'$ with respect to Ω and that of $N \otimes M'$ with respect to $\Omega \otimes \Omega$. This results in the unitary implementability of Φ in terms of the following unitary:

$$U_{can}n \cdot m'\Omega = n\Omega \bar{\otimes} m'\Omega \qquad (A.42)$$
$$U_{can}: H \rightarrow H_1 \bar{\otimes} H_2$$
$$H_1 = \overline{N\Omega} \quad , H_2 = \overline{M'\Omega}$$

Using the modular involution J of the collar $N' \cap M$, the canonical type I factor is defined as:

$$B_{can} = N \lor JNJ \tag{A.43}$$

$$= U_{can}^* B(H_1) \bar{\otimes} 1_{H_2} U_{can} \tag{A.44}$$

One of the most useful consequences of the split property is the existence of a universal localizing map U_{can} . It gets its name from the remarkable property of "localizing" global automorphisms of B(H) inside M such that the action on the subalgebra N is identical to the restriction of the global action. In the context of the geometric action of the Poincaré group on the net, where M is the algebra localized in e.g. a bigger double cone, the definition:

$$U_{loc}(g) \equiv U_{can}^* U(g) \bar{\otimes} 1 U_{can} \tag{A.45}$$

has the correct property and may be viewed as a kind of algebraic pre-version of the Quantum Noether theorem.

In QFT applications this property is garantied for inclusions of localized algebras with a causal collar between them if the local degrees of freedom have a reasonable phase space behavior called "Nuclearity Property". Phase space degree of freedom counting in LQP is based on the concept of ε -content $N(\varepsilon)$ of a map Θ between Banach spaces. $N(\varepsilon)$ is defined as the maximal number of elements E_i in the unit ball of the source space such that $\|\Theta(E_i - E_j)\| > \varepsilon$ for $i \neq j$. The order q of the map is then defined to be:

$$q = \lim_{\varepsilon \searrow 0} \sup \frac{\ln \ln \varepsilon}{\ln \frac{1}{\varepsilon}} \tag{A.46}$$

In heat bath LQP the map would be $\Theta : \mathcal{A}(\mathcal{O}) \longrightarrow H$, given by $\Theta_{\Omega,\beta}(\mathcal{A}) = e^{-\beta \mathbf{H}}\Omega, A \in \mathcal{A}(\mathcal{O})$ with H the Hamiltonian, whereas in case of thermality through localization one takes $\Theta_{\Omega,\Delta}(A)\Omega = \Delta_{\hat{O}}^{\frac{1}{4}}A\Omega, A \in \mathcal{A}(\mathcal{O})$ with Δ the modular operator of the localization region. For the derivation od the split property from the nuclear properties of these maps with a certain value of q (which measures the strength with which the number of relativistic degrees of freedom per unit phase space volume approaches infinity) we refer to [3].

A.10 Inverse Problem of Modular Theory

Mathematically the inverse problem of modular theory is the construction of a vNa in standard position i.e. pairs (M, Ω) such that a given would be pair of modular operators (Δ_0, J_0) of (M_0, Ω_0) , with M isomorphic to M_0 , are also the modular operators of (M, Ω_0) . The appearantly weaker (more general) version in which one only demands that the new algebra (M, Ω_0) shares the same Δ_0 turns out to be reducible to the previous problem and is the kind of inverse problem close to the physical one described below. There are other inverse problems which have been solved, the most prominent being Connes inverse problem of reconstructing vNa's from the position of their natural cones in Hilbert space which (as the below physical problem) has a unique solution.

The above isomorphy classes $IC(M_0, \Omega_0)$ have many elements. For example $M = VM_0V^*$ with a unitary V commuting with Δ_0, J_0 and obeying $V\Omega_0 = \Omega_0$ gives a family of vNa's. This construction does not exhaust the possibilities. Already for matrices i.e. I_n factors one can construct yet another family of inverse problem classes by choosing V as $V = V_1U$ with V_1 having the same commutation properties as above but $V_1\Omega_1 = \Omega_0$. Here Ω_1 is such that (Δ_0^{-1}, J_0) are the modular objects of (M_0, Ω_1) and U commutes with J_0 and obeys $\Delta_0^{-1} = U^*\Delta_0U, U\Omega = \Omega, U\Omega_1 = \Omega_1$. This construction only works for I_n factors because only there Δ_0^{-1} can be the modular operator of a pair (M_0, Ω_1) . For type I factors one has a rather complete solution of the inverse problem. For type III on the other hand extremely little is known.

The physical aspects of the inverse problem which deal with a special setting within hyperfinity type III_1 factors or in LQP terminology wedge or double cone localized algebras. There one has always a given reference algebra which in the case of massive particles would be the wedge algebra associated to the incoming free fields say Δ_0 . The first step in the construction of interacting algebras is the recognition that the would be interacting factor algebra M for the wedge shares the modular Δ_0^{it} which is the Lorentz boost. This is a corrolar of the fact that the interaction does not manifest itself in the representation of the (connected) Poincaré group but only in the reflections as explained in chapter5. On the other hand the J of the interacting M is related to J_0 by the physical S-matrix $J = S_{scat} J_0$. In this case there are sufficiently many additional requirements from LQP so that the interacting wedge algebra is uniquely determined by the physical S-matrix. The unitary equivalence transformation between M_0 and M is analogous to the Møller operator from scattering theory and called the modular Møller operator. It is very interesting that in this physical case the solution of the inverse problem in the sense of modular theory coalesces with that of the inverse problem of LQP: the unique relation of the (physically admissable) S-matrix with the associated QFT.

The inverse problem of modular theory and its nonuniqueness, the field theoretic setting and relation to the inverse problem of QFT (admissable scattering matrix $S \rightarrow local \ QFT$), the modular Møller operator and the nonperturbative constructive approach.

Suggested Literature to this Appendix

O. Bratteli and D. W. Robinson, "Operator Algebras and Quantum Statistical Mechanics" I, Springer, New York 1979

V. S. Sunder, "An Invitation to von Neumann Algebras", Springer, New York 1987

R. V. Kadison and J. R. Ringrose, "Fundamentals of the Theory of Operator Algebras" I ans II, Academic Press, New York 1986

M. Takesaki, "Conditional expectations in von Neumann algebras", J. Funct. Anal. **9**, 306-321 (1972)

R. Longo and K.-H. Rehren, "Nets of subfactors", Rev. Math. Phys., 7, (1995), 567

S. Doplicher, C. Pinzari and R. Zuccante, "*The C*-algebra of a Hilbert Bi-module*", funct-an/9707006 and previous work of Rieffel, Exel and Pimsner cited therein.

S. Doplicher and R. Longo, "Standard and split inclusions of von Neumann algebras", Invent. math. 75, 493 (1984)

To my Brazilian collegues: olha para a home page do Ruy Exel: http://www.ime.usp.br/~exel/ A. Connes, "Noncommutative Geometry" Academic Press 1994



