Generalized Calogero-Sutherland systems from many-matrix models

Alexios P. Polychronakos¹

Institutionen för teoretisk fysik, Box 803 S-751 08 Uppsala, Sweden

and

Physics Department, University of Ioannina 45110 Ioannina, Greece

Abstract

We construct generalizations of the Calogero-Sutherland-Moser system by appropriately reducing a model involving many unitary matrices. The resulting systems consist of particles on the circle with internal degrees of freedom, coupled through modifications of the inverse-square potential. The coupling involves SU(M) non-invariant (anti)ferromagnetic interactions of the internal degrees of freedom. The systems are shown to be integrable and the spectrum and wavefunctions of the quantum version are derived.

PACS: 03.65.Fd, 71.10.Pm, 11.10.Lm, 03.20.+i

¹E-mail: poly@teorfys.uu.se

The inverse-square interacting particle system [1, 2, 3] and its spin generalizations [4, 5, 6, 7, 8, 9] are important models of many-body systems, due to their exact solvability and intimate connection to spin chain systems [10, 11, 12, 13], 2-dimensional Yang-Mills theories [14, 15, 16] etc.

A useful route for studying these systems is through reductions of appropriate matrix models [17, 18]. Although this does not give rise to all known models in the quantum case, it is still a very direct way to show integrability and solvability. In this paper, we will explore this approach further and show how, starting with many matrices effectively coupled through constraints, we can derive further generalizations of these systems with internal degrees of freedom.

The starting point will be the many-matrix lagrangian

$$L = \operatorname{tr} \sum_{i=1}^{M} -\frac{1}{2a_i} (U_i^{-1} \dot{U}_i)^2$$
(1)

The U_i are $N \times N$ unitary matrices depending on time t and overdot stands for time derivative. The a_i are real positive parameters. For convenience, we shall normalize the a_i such that

$$\sum_{i=1}^{M} a_i = 1 \tag{2}$$

The above system is clearly integrable and solvable, being nothing more than a collection of independent unitary matrix models. In the absence of further constraints it would give rise to a collection of independent spin-generalizations of the Sutherland model. It is, however, possible to reduce it in a different way: the above model is invariant under (independent) left- and right-multiplications of the U_i by constant matrices. This manifests in the existence of conserved quantities, namely

$$R_{i} = -\frac{i}{a_{i}} U_{i}^{-1} \dot{U}_{i} , \qquad L_{i} = \frac{i}{a_{i}} \dot{U}_{i} U_{i}^{-1}$$
(3)

The above conserved hermitian matrices Poisson-commute with themselves to the U(N) algebra and mutually to zero. We will choose to fix the values of the following time-independent quantities:

$$-\frac{i}{a_i}U_i^{-1}\dot{U}_i + \frac{i}{a_{i+1}}\dot{U}_{i+1}U_{i+1}^{-1} = P_i$$
(4)

where we have adopted periodic conditions in i, that is, $M + 1 \equiv 1, 0 \equiv M$. Clearly the chosen expressions for the P_i , being the sum of independent U(N) generators, also constitute mutually commuting Poisson-U(N) matrices.

The key to a successful reduction of this system to particles lies in an appropriate parametrization of the U_i . We will choose

$$U_i = W_{i-1}^{-1} \Lambda^{a_i} W_i \tag{5}$$

where W_i are unitary matrices and $\Lambda = diag(e^{i\theta_n})$ is a diagonal unitary matrix. The reason for the choice of exponents in (5) will be apparent in the sequel. From the relation

$$U \equiv U_1 \cdots U_N = W_N^{-1} \Lambda W_N \tag{6}$$

we see that W_N and $e^{i\theta_n}$ are the diagonalizer and the eigenvalues of the matrix U, and the remaining W_i are determined recursively from (5). The above parametrization has the redundancy generated by left-multiplication of all W_i by the same diagonal matrix. This will lead to a 'gauge constraint' later on.

With the above parametrization the expressions for the lagrangian and the P_i become, after some algebra,

$$L = \operatorname{tr}\sum_{i=1}^{M} \frac{1}{2} \left\{ -(\Lambda^{-1}\dot{\Lambda})^{2} + \left(\frac{1}{a_{i}} + \frac{1}{a_{i+1}}\right) \mathcal{L}_{i}^{2} - \frac{2}{a_{i}} \mathcal{L}_{i-1} \Lambda^{a_{i}} \mathcal{L}_{i} \Lambda^{-a_{i}} \right\}$$
(7)

$$W_i P_i W_i^{-1} \equiv K_i = \left(\frac{1}{a_i} + \frac{1}{a_{i+1}}\right) \mathcal{L}_i - \frac{1}{a_i} \Lambda^{-a_i} \mathcal{L}_{i-1} \Lambda^{a_i} - \frac{1}{a_{i+1}} \Lambda^{a_{i+1}} \mathcal{L}_{i+1} \Lambda^{-a_{i+1}}$$
(8)

where we defined

$$\dot{W}_i W_i^{-1} = i \mathcal{L}_i \tag{9}$$

From the structure of the above lagrangian, we see that the generator of left-rotations for W_i (found by differentiating L with respect to $i\dot{W}_iW_i^{-1}$) is exactly K_i as expressed in (8). Therefore, the K_i are also mutually commuting Poisson-U(N) generators. This justifies the choice of exponents in (5), since any other choice would spoil the decoupling and U(N) nature of the K_i . We stress, however, that the K_i are no more conserved and are, thus, dynamical quantities. Further, from (8) we see that the diagonal elements of K_i must satisfy

$$\sum_{i=1}^{M} K_{inn} = 0 \quad \text{for each } n \tag{10}$$

which is the 'Gauss law' originating from the 'gauge invariance' of the parametrization (5) as stated previously. The off-diagonal elements of \mathcal{L}_i satisfy

$$\sum_{j=1}^{M} \mathbf{M}_{ij}(i\theta_{mn}) \mathcal{L}_{jmn} = K_{imn}$$
(11)

where $\theta_{mn} \equiv \theta_m - \theta_n$ and the matrix $\mathbf{M}(x)$ is defined as

$$\mathbf{M}_{ij}(x) = \left(\frac{1}{a_i} + \frac{1}{a_{i+1}}\right)\delta_{ij} - \frac{1}{a_i}e^{-a_ix}\delta_{i,j+1} - \frac{1}{a_{i+1}}e^{a_{i+1}x}\delta_{i+1,j}$$
(12)

(again, taking $\delta_{i,M+1} = \delta_{i,1}$). In terms of the matrices K_i and the angles θ_n the hamiltonian (which is identical to the lagrangian since it only contains kinetic terms)

becomes

$$H = \frac{1}{2} \sum_{n=1}^{N} p_n^2 + \frac{1}{2} \sum_{m \neq n=1}^{N} \sum_{i,j=1}^{M} \mathbf{M}_{ij}^{-1}(i\theta_{mn}) K_{imn} K_{jnm} + \frac{1}{2} \sum_{n=1}^{N} \sum_{i,j=1}^{M} \tilde{\mathbf{M}}_{ij}^{-1}(0) K_{inn} K_{jnn}$$
(13)

where $p_n = \dot{\theta}_n$ is the canonical momentum of θ_n and $\tilde{\mathbf{M}}(0)$ is the matrix $\mathbf{M}(0)$ projected to the subspace orthogonal to its zero-eigenvalue eigenvector.

In order to invert the matrix \mathbf{M} , we note that it is essentially a discrete second derivative on a set of points at distances a_i , conjugated with an exponential factor. Generalizing from the case of equally-spaced points (that is, the Cartan matrix of SU(M)), where the inverse is proportional to the distance |i - j|, we define

$$b_i = \sum_{j=1}^i a_j , \quad b_{ij} = b_i - b_j$$
 (14)

and try a form

$$\mathbf{M}_{ij}^{-1}(x) = e^{-b_i x} \left\{ A(x) + B(x)b_{ij} + C(x)|b_{ij}| \right\} e^{b_j x}$$
(15)

The elements with i, j either 1 or M need special attention due to the 'wrap around' properties of \mathbf{M} . (Note that b_i is not periodic in its index, but rather $b_{i+M} = b_i + 1$.) This fixes the coefficients A, B, C and the inverse of \mathbf{M} becomes

$$\mathbf{M}_{ij}^{-1}(x) = e^{-b_{ij}x} \left\{ \frac{1}{(1-e^x)(1-e^{-x})} + \frac{b_{ij}(1+e^x)}{2(1-e^x)} - \frac{|b_{ij}|}{2} \right\}$$
(16)

The matrix $\mathbf{M}(0)$ has a zero eigenvalue corresponding to the eigenvector $(1, \ldots 1)$. In order to compute the inverse of $\tilde{\mathbf{M}}$ it suffices to take the limit $x \to 0$ in (16) and discard pieces that are constant in i, j since they clearly project on the zero-eigenvalue eigenspace. We find

$$(\mathbf{M}(x)^{-1})_{ij} = -\frac{1}{x^2} + \frac{1}{12} + \frac{b_{ij}^2 - |b_{ij}|}{2} + O(x)A_{ij}$$
(17)

Discarding the constant piece and taking $x \to 0$ we obtain \mathbf{M}^{-1} . Putting everything together, we finally obtain

$$H = \frac{1}{2} \sum_{n} p_n^2 + \frac{1}{2} \sum_{i,j;m \neq n} V_{ij}(\theta_{mn}) K_{imn} K_{jnm} + \frac{1}{2} \sum_{i,j;n} \tilde{V}_{ij} K_{inn} K_{jnn}$$
(18)

with the potentials V_{ij} and \tilde{V}_{ij} defined as

$$V_{ij}(x) = e^{-ib_{ij}x} \left(\frac{1}{4\sin^2 \frac{x}{2}} + i\frac{b_{ij}}{2}\cot \frac{x}{2} - \frac{|b_{ij}|}{2} \right) \quad , \qquad \tilde{V}_{ij} = \frac{b_{ij}^2 - |b_{ij}|}{2} \tag{19}$$

This is a generalization of the Sutherland model of particles on the circle with spin degrees of freedom, encoded by the K_i , where the different K_i couple through a

parametric modification of the inverse-sine-squared potential. Interestingly, this is the same classical model as the one obtained by Blom and Langmann starting from the 2-dimensional Yang-Mills point of view [19]. The standard spin-Sutherland model is recovered either as the limit where all K_i but one are zero, or as the limit where all a_i but one are zero, in which case only the sum of all K_i appears in the hamiltonian.

The quantum version of this model proceeds along similar lines. The constant matrices P_i become, now, U(N) generators transforming under some fixed representations of SU(N) and carrying some U(1) charge (since they are not necessarily traceless). We shall choose them to be irreducible, the reducible case being simply the direct sum of models with one irreducible component for each P_i . Therefore, the model is labeled by a set of irreps r_i and charges q_i carried by the P_i . The matrices K_i become time-varying SU(N) generators in the same irrep r_i and with the same charge q_i as P_i . In a more standard notation, putting

$$K_i = \sum_{a=1}^{N^2 - 1} T^a K_i^a + \frac{q_i}{N}$$
(20)

with T^a the fundamental SU(N) generators, the K_i^a obey the SU(N) algebra for each *i* and commute for different *i*, while the q_i are central. The hamiltonian (18) remains valid for the quantum operators as well; the only term requiring ordering, namely $V_{ii}K_{imn}K_{inm}$, is automatically symmetrized by the summation over m, n.

To turn the K_i into internal degrees of freedom we follow the standard construction [15] (see also [20]). The K_i can be realized à la Jordan-Wigner in terms of bosonic oscillators

$$K_{imn} = \sum_{k=1}^{d_i} \left(A_{ikm}^{\dagger} A_{ikn} - \frac{1}{N} \delta_{mn} \sum_{l=1}^{N} A_{ikl}^{\dagger} A_{ikl} \right) + \frac{q_i}{N} \delta_{mn}$$
(21)

where d_i is the number of rows in the Young tableau of r_i . The A_{ikm} are a collection of commuting bosonic ladder operators

$$[A_{ikm}, A_{jln}^{\dagger}] = \delta_{ij} \delta_{kl} \delta_{mn} \tag{22}$$

The above imbeds r_i , as well as all other irreps with up to d_i rows, in the Fock space of A_{ikm} . To simplify the notation and interpretation of the model we will choose $d_i = 1$ for all K_i . We can always achieve the case $d_i \ge 1$ as a model with $M' = \sum_{i=1}^{M} d_i$ matrices with all $d_i = 1$. Choosing

$$a_j = 0$$
 for $\sum_{k=1}^{i-1} d_k < j \le \sum_{k=1}^i d_k$ (23)

makes all the K_j in this range, overall d_i in number, appear only through their sum in the hamiltonian, reproducing the original $d_i > 1$ matrix. We thus drop the summation and index k in (21) and (22).

The gauge constraint (10) in terms of (21) implies

$$\sum_{i=1}^{M} A_{in}^{\dagger} A_{in} = \text{constant} \equiv \text{m} \text{ for all } n, \qquad \sum_{i=i}^{M} q_i = 0$$
(24)

From the above we see that the Fock states generated by the M oscillators A_{in} for each n transform in the m-fold symmetric irrep of SU(M). We therefore define the N mutually commuting SU(M) generators

$$S_{nij} = A_{in}^{\dagger} A_{jn} - \frac{\mathrm{m}}{M} \delta_{ij} \tag{25}$$

carrying the m-fold symmetric irrep of SU(M). In terms of these, substituting (21) in (18), we obtain after some algebra

$$H = \frac{1}{2} \sum_{n} p_{n}^{2} + \frac{1}{2} \sum_{m \neq n} \left(\sum_{ij} V_{ij}(\theta_{mn}) S_{mij} S_{nji} + V_{o}(\theta_{mn}) \frac{\mathrm{m}(\mathrm{m} + M)}{M} \right) + \frac{1}{2} \sum_{n} \sum_{ij} \tilde{V}_{ij} S_{nii} S_{njj} + \frac{1}{2N} \sum_{ij} \tilde{V}_{ij} \left(q_{i}q_{j} - S_{ii}S_{jj} \right)$$
(26)

where we defined the total spin $S = \sum_{n=1}^{N} S_n$ and $V_o(x) \equiv V_{ii}(x) = 1/(2 \sin \frac{x}{2})^2$. The above hamiltonian refers to a system of particles with m-fold symmetric SU(M)spins S_n . Due, however, to the existence of the matrix V(x) in the coupling of spins, the interactions above are not SU(M)-invariant (the total spin S is not conserved). Only the diagonal elements (that is, the Cartan part) of the total spin are conserved. The appearance of the total spin in (26) can be eliminated by choosing the charges

$$q_i = S_{ii} = n_i - \frac{mN}{M} \tag{27}$$

where n_i is the number of boxes in r_i . (In this construction the r_i are symmetric, so n_i are their lengths.) This is a natural choice, identifying the U(1) charge to the Z_N charge of r_i and subtracting the average Z_N charge of all irreps.

In terms of the original matrix problem, the hamiltonian is the sum of independent laplacians over the space U(N). The U(1) part will lead to charges Q_i (the momentum of the 'center-of-mass' of each U_i) which, by (4) and (6), have to satisfy

$$Q_i - Q_{i+1} = q_i \quad , \qquad \sum_i a_i Q_i = \sum_n p_n \equiv P \tag{28}$$

This fixes the Q_i in terms of the q_i and the total momentum P. Factoring out the U(1) part for each U_i , we are left with laplacians on the SU(N) manifold. It is known that the matrix elements of each irrep R of SU(N) form a degenerate eigenstate multiplet of the laplacian with eigenvalue given by the quadratic Casimir $C_2(R)$ of the irrep. Therefore, the eigenstates of the matrix hamiltonian are

$$\Psi(\{U_i\}) = \prod_{i=1}^{M} (\det U_i)^{\frac{Q_i}{N}} R_i(U_i)_{\alpha_i \beta_i}$$
(29)

where R_i are irreps of SU(N) (α_i, β_i label their matrix elements). The energy eigenvalue E and degeneracy D corresponding to this state are

$$E = \sum_{i=1}^{M} \frac{a_i}{2} \left\{ C_2(R_i) + \frac{Q_i^2}{N} \right\} \quad , \qquad D = \prod_{i=1}^{M} (dimR_i)^2 \tag{30}$$

Each $R_i(U_i)_{\alpha_i\beta_i}$ transforms under left-rotations of U_i in the R_i irrep (acting on the left index) and under right-rotations of U_i in the conjugate irrep \bar{R}_i (acting on the right index). The condition (4), however, tells us that the sum of the right-generator for U_i and the left-generator for U_{i+1} are constrained to be in the r_i irrep of SU(N) carried by P_i . Therefore, we must project the states β_i and α_{i+1} , transforming in the \bar{R}_i and R_{i+1} respectively, to the subspace of states transforming in the r_i . Call $G(\bar{R}_i, \beta_i; R_{i+1}, \alpha_{i+1} | r_i, \gamma_i)$ the Clebsch-Gordan coefficient that projects these states to the γ_i state of r_i . Then the final energy eigenstate wavefunction for this model becomes

$$\Psi(\{U_i\}) = \sum_{\{\alpha_j,\beta_j\}} \prod_{i=1}^M (\det U_i)^{\frac{Q_i}{N}} R_i(U_i)_{\alpha_i\beta_i} G(\bar{R}_i,\beta_i;R_{i+1},\alpha_{i+1}|r_i,\gamma_i)$$
(31)

The indices γ_i do not imply a degeneracy of states corresponding to the dimensionality of r_i , since these are states fixed by the constraints (4) and are not summed over. The degeneracy of each state labeled by the R_i is given by the number of times that the irreps r_i are contained in the direct products $\bar{R}_i \times R_{i+1}$ or, equivalently, the number of times that R_{i+1} is contained in $R_i \times r_i$. Calling this integer $D(R_i, r_i; R_{i+1})$, we have for the total degeneracy

$$D = \prod_{i=1}^{M} D(R_i, r_i; R_{i+1})$$
(32)

Clearly, an irrep R_i can appear in Ψ only if it is contained in the product $R_i \times r_1 \cdots \times r_N$. This requires that the total Z_N charge of the r's be zero, that is, the total number of boxes in the Young tableaux of r_i should be a multiple of N. This is indeed the case, the total number being n = mN. The above, upon proper reinterpretation, gives the spectrum and wavefunctions in each corresponding sector of the particle-spin model as classified by the Cartan elements of the total spin.

We conclude by mentioning that a construction similar to (21) in terms of fermionic oscillators would give rise to a model as in (26) but with negative sign (ferromagnetic) spin interactions.

I would like to thank E. Langmann for comments and for communicating his results [19] prior to publication.

References

- F. Calogero, J. Math. Phys. 10 (1969) 2191 and 2197; 12 (1971) 419; Lett. Nuovo Cimento 13 (1975) 411; F. Calogero and C. Marchioro, Lett. Nuovo Cimento 13 (1975) 383.
- [2] B. Sutherland, Phys. Rev. A4 (1971) 2019; 5 (1972) 1372; Phys. Rev. Lett. 34 (1975) 1083.
- [3] J. Moser, Adv. Math. 16 (1975) 1.
- [4] J. Gibbons and T. Hermsen, *Physica* **D11** (1984) 337.
- [5] S. Wojciechowski, *Phys. Lett.* A111 (1985) 101.
- [6] Z.N.C. Ha and F.D.M. Haldane, *Phys. Rev.* B46 (1992) 9359.
- [7] N. Kawakami, *Phys. Rev.* **B46** (1992) 1005 and 3191.
- [8] J.A. Minahan and A.P. Polychronakos, *Phys. Lett.* **B302** (1993) 265.
- [9] K. Hikami and M. Wadati, *Phys. Lett.* A173 (1993) 263.
- [10] F.D.M. Haldane, *Phys. Rev. Lett.* **60** (1988) 635; **66** (1991) 1529.
- [11] B.S. Shastry, *Phys. Rev. Lett.* **60** (1988) 639; **69** (1992) 164.
- [12] M. Fowler and J.A. Minahan, *Phys. Rev. Lett.* **70** (1993) 2325.
- [13] A.P. Polychronakos, Phys. Rev. Lett. 70 (1993) 2329; Nucl. Phys. B419 (1994) 553.
- [14] A. Gorsky and N. Nekrasov, Nucl. Phys. B414 (1994) 213.
- [15] J.A. Minahan and A.P. Polychronakos, *Phys. Lett.* **B326** (1994) 288.
- [16] E. Langmann, M. Salmhofer and A. Kovner, Mod. Phys. Lett. A A9 (1994) 2913.
- [17] D. Kazhdan, B. Kostant and S. Sternberg, Comm. Pure Appl. Math. 31 (1978) 481.
- [18] M.A. Olshanetsky and A.M. Perelomov, *Phys. Rep.* **71** (1981) 314; **94** (1983)
 6.
- [19] J. Blom and E. Langmann, solv-int/9804007.
- [20] J. Avan and A. Jevicki, *Nucl. Phys.* **B469** (1996) 287.