## SU-ITP-98-48

## INITIAL CONDITIONS IN PRE-BIG BANG<sup>a</sup>

N. KALOPER

*Department of Physics, Stanford University, Stanford, CA 94305-4060, USA E-mail: kaloper@leland.stanford.edu*

This note is a summary of the work reported in [hep-th/9801073](http://arxiv.org/abs/hep-th/9801073). We give a brief discussion of the fine tuning problem in pre-big bang cosmology. We use the flatness problem as our test case, and in addition to the exact numerical limits on initial conditions, we highlight the differences between pre-big bang and standard inflation. The main difference is that in pre-big bang the universe must be smooth and flat in an exponentially large domain already at the beginning of the dilatondriven inflation.

It has been known at least for a decade that it is extremely hard to construct a working mechanism for inflation in string theory. The moduli fields, and in particular, the dilaton, tend to roll rapidly in the presence of vacuum energy, and this rolling slows down the cosmological expansion of the particle horizon. Hence the standard inflationary scenarios cannot work, since the horizon and flatness problems cannot be solved.

One hopes that this problem is only temporary and that it will be eventually solved with the increasing understanding of string theory. But there is also a possibility that a different implementation of inflation might be possible. One of the most interesting suggestions in this direction has been the pre-big bang (PBB) scenario<sup>[1](#page-4-0)</sup>. This scenario attempts to use the rolling dilaton to run inflation. Due to this, there appear solutions which inflate superexponentially for some time in the string frame. In the course of this expansion, the curvature and the coupling would become large, and the evolution proceeds towards the curvature singularity (and not away from it). If one would assume the existence of a mechanism which could saturate the growth of the curvature and coupling, and overturn superexponential expansion into a subluminal power law one, one could match the solution onto a late time Friedmann-Robertson-Walker (FRW) one. The phase of strong curvature has been argued to resemble the Big Bang and the one preceding it has been dubbed the pre-big bang<sup>[1](#page-4-0)</sup>.

Nevertheless, it has not been clear just how successful PBB is in explaining  $\alpha$ cosmological problems. In fact already in<sup>[1](#page-4-0)</sup> it has been noted that at the onset of

<sup>a</sup>Talk given at PASCOS-98 and at CAPP-98.

<span id="page-1-0"></span>the dilaton-dominated inflating phase, the size of the homogeneous domain had to be large in string units,  $L_H >> l_s$ , and the string coupling had to be small,  $g^2$  < 1, but this has not been considered a real problem. More recently, this has been revisited in<sup>[2](#page-4-0)</sup>, where it has been argued that in a universe which was not always dilaton-dominated, the requirements for sufficient inflation would require fine tuning of initial conditions. The main point of  $2$  was that the presence of matter other than the dilaton could delay the onset of inflation and hence infringe on the efficiency of the scenario. Counterarguments have been given in <sup>[3](#page-4-0),[4](#page-4-0),[5](#page-4-0)</sup>, where it has been claimed that there exist initial conditions for which this does not happen.

We have undertaken a comprehensive analysis of these issues in  $6$ , and have found a precise quantitative formulation of the fine tuning problem in PBB. Our results show that if PBB is to solve the horizon and flatness problems, at the onset of the dilaton domination, the horizon size of the inflating domain must be  $L_H \geq 10^{19}l_p$ . Simultaneously, the string coupling must satisfy 0 <  $g^2 \leq 10^{-53}$ . Perhaps the clearest way to see the severity of these constraints is to consider the kinetic energy of the dilaton at the onset of inflation. In terms of the integral of motion  $B = -\dot{\Phi}a^3$ , to be defined precisely below, this requires  $B \ge 10^{38} g^{-2} \ge 10^{91}$ . These numbers are just a reformulation, and not a solution, of the cosmological problems. The task of PBB must be to explain the presence of these numbers.

Let us now give a brief derivation of these constraints. The classical dynamics of PBB is defined by the action

$$
S = \int d^4x \sqrt{g} \frac{e^{-\sigma}}{l_s^2} \left(\frac{1}{2}R + \frac{1}{2}(\nabla\sigma)^2 + \mathcal{L}_m(\mathcal{Y}, g_{\mu\nu}, \sigma)\right) \tag{1}
$$

where  $g_{\mu\nu}$  and  $\sigma$  are the string-frame metric and the dilaton, and Y stands for all other matter degrees of freedom. This action gives a valid description of the dynamics only up to the regions where  $\exp(-\sigma) \sim 1$  and/or  $R \sim 1/l_s^2$ , where higher order corrections in coupling and curvature are expected to be important, invalidating the truncation of the theory that led to (1). In an FRW background

$$
ds^{2} = -dt^{2} + a^{2}(t)\left(\frac{dr^{2}}{1 - kr^{2}} + r^{2}d\Omega\right)
$$
 (2)

one can write the equations of motion coming from (1) as follows:

$$
\ddot{\sigma} + 3H\dot{\sigma} - \dot{\sigma}^2 = l_s^2 e^{\sigma} (3p - \rho) \qquad \dot{H} = H\dot{\sigma} - 3H^2 - 2\frac{k}{a^2} + l_s^2 p e^{\sigma}
$$

$$
\dot{\rho} + 3H(\rho + p) = 0 \qquad \dot{\sigma}^2 + 6H^2 - 6H\dot{\sigma} + 6\frac{k}{a^2} - 2l_s^2 e^{\sigma} \rho = 0 \qquad (3)
$$

If one ignores the influence of the curvature and the matter stress-energy, one can find two classes of expanding solutions, which are accelerating for  $t < 0$ and decelerating for  $t > 0$ , and are separated by a curvature singularity at  $t = 0$ . Strictly speaking, the solutions are valid for only for times  $|t| > l_s$ . For  $|t| \leq l_s$  $|t| \leq l_s$  $|t| \leq l_s$ , the corrections are large and so the action ([1\)](#page-1-0) does not apply. In PBB, one assumes the existence of some corrections-driven graceful exit mechanism which avoids the singularity, and further assumes that for  $t \geq l_s \sim l_p$  the string coupling is roughly a constant of  $O(1)$ . The period of inflation can take place between the moment of the onset of dilaton domination, defined roughly by  $\dot{\sigma}^2 \sim 6k/a^2$ , and at most the end of the string phase. During this time, the evolution mostly obeys the superexponential law. If we define  $\Phi = \exp(-\sigma)/l_s^2$ , we can write

$$
a_{+} = |t_{0}| \left| \frac{t}{t_{0}} \right|^{1/\sqrt{3}} \qquad \Phi = \frac{1}{l_{p}(0)^{2}} \left| \frac{t}{t_{0}} \right|^{1/\sqrt{3}+1} \tag{4}
$$

Here  $l_p(0) = 1/\sqrt{\Phi(t_0)}$  is the Planck length at the onset of inflation, as opposed to the Planck length  $l_p \sim l_s$  at the end.

We can now consider the flatness problem as our test case. Requiring that inflation solves the flatness problem means that the the time of the exit,  $|\Omega - 1| \leq 10^{-60}$ . Since in PBB the exit is roughly at the Planck time, we find that at this time  $a_f H_f \sim 10^{30}$ . On the other hand, if we compute the particle horizon during PBB, we find that  $L_f / a = \text{const} \sim O(1)$ . Also, by continuity  $H_f \sim 1/l_s$ , and hence to solve the flatness problem we must have  $L_f \geq 10^{30} l_s \sim e^{70} l_s$ . At the onset of inflation, by the equations of motion,  $L_i \sim a_i \sim H_i^{-1} \sim |t_i|$ . Using  $L_f/a_f = L_i/a_i$ , we obtain the constraint on  $L_i$ at the onset of inflation:  $L_i = a_i/a_f L_f \geq 10^{30} a_i/a_f$ , or equivalently

$$
\left(\frac{\Phi_i}{\Phi_f}\right)^{1/\sqrt{3}} = \left(\frac{|t_i|}{l_s}\right)^{1+1/\sqrt{3}} \ge 10^{30} \tag{5}
$$

This constraint can be translated into the following requirements for the horizon, mass, entropy of the universe, and the coupling constant  $g$  at the onset of inflation:

$$
L_i \ge 2 \times 10^{19} l_s \qquad M_i \ge 10^{72} M_s \qquad S \ge 2 \times 10^{38} \qquad g_0^2 \le 10^{-53} \qquad (6)
$$

If the universe at the onset of PBB inflation were to appear accidentally, it would be extremely unlikely to satisfy (6). The probability for such an event, by standard inflationary arguments, would be very low,  $P \leq \exp(-10^{38})$  at best. Further, it would already have to be homogeneous over length scales of order of at least  $10^{19}l_s$ , in contrast to the case of standard inflation, which requires homogeneity over length scales of only few  $l_s$ .

From inspecting the equations of motion [\(3](#page-1-0)), it is easy to see that the quantity  $B = -\dot{\Phi}a^3$  is an integral of motion during th PBB phase. Translated in terms of this independent parameter, the inflationary constraint reads

$$
B \ge 4 \times 10^{38} g_0^{-2} \ge 4 \times 10^{91} \tag{7}
$$

Thus we see that if PBB is to successfully solve cosmological problems, it must explain the origin of these numbers. Merely saying that these numbers are required for inflation only amounts to restating the cosmological problems in a different fashion, and no more.

We note that since this talk has been given two more works have appeared, attempting to explain some of the large numbers we have discussed here  $^{7,8}$  $^{7,8}$  $^{7,8}$  $^{7,8}$  $^{7,8}$ . This newest version of the PBB scenario considers the collapse of classical dilaton waves, which form black holes. However the black holes so formed must be extremely massive,  $M \sim 10^{72} M_S$ , to fit our universe inside of them. We should also note that one might be tempted to argue that the application of the flatness problem to PBB might be misleading, since after all at the onset of dilaton domination, the dilaton kinetic energy and the curvature energy are of the same order of magnitude,  $\dot{\sigma}^2 \sim 6k/a^2$ . This might resemble the standard inflation, where at the onset of inflation  $E(\text{vacuum}) \sim k/a^2$ . However, in open PBB this condition is put in by hand, by requiring that the spacetime before inflation was precisely flat. In closed PBB, this condition comes about only after requiring that the universe was at least  $\sim 10^{8}l_s$  across at the original singularity, again an input. Thus the fact that the dilaton kinetic energy and the curvature energy were approximately the same at the onset of PBB does not explain the flatness problem.

One may wonder if these conditions could be produced by cosmological evolution preceding inflation. The picture of the evolution before the dilaton domination depends very strongly on the dominant source at that time. If the universe before PBB inflation had been dominated by a positive curvature term (i.e. had been closed) it would have had to start from another space time singularity before the stringy phase of PBB. At this, pre-stringy stringy time, it would have had to be homogeneous over length scales  $L \geq 10^8 l_s$ . This may seem better than Big Bang without inflation, but is still exponentially unlikely, with the probability  $P \sim \exp(-10^{24})$ . If the universe had been dominated by negative spatial curvature (i.e. had been open), it would have corresponded to the collapse of a homogeneous distribution of scalar kinetic energy for an infinite time, and of infinite size. This universe would have to be dominated by the homogeneous scalar fluctuation for nearly ever, in order to solve the cosmological problems. In particular, if one considers inhomogeneous quantum fluctuations of the scalar field, with wavelengths  $\lambda \sim |t|$  and amplitudes

<span id="page-4-0"></span> $\delta \Phi \sim \sqrt{\Phi} |t|^{-1}/(2\pi)$ , one could see that they would dominate over the classical homogeneous perturbations  $\Delta \Phi \sim B/t^2$  for all times  $|t| \geq Bl_p(0)$ . Hence in the very far past such quantum fluctuations would backreact with the background more strongly than the homogeneous classical perturbations, and could completely destroy the large-scale homogeneity for the success of PBB. We underline here that we are not addressing the problem of growth of perturbations in this context. We do not consider the small perturbations  $\delta \rho \leq \rho$ , or the issues of Jeans instability, which may or may not exist in PBB. Rather we argue that there may be initial inhomogeneous quantum fluctuations which can completely destroy the classical Minkowski background, making it look like a collection of open and closed universes. Since we do not have the necessary space to give an exhaustive discussion of these issues here, we refer the reader interested in a more detailed account of this problem to  $6,8$ .

We want to emphasize that the constraints  $g^{-2} \geq 10^{53}$  and  $B \geq 10^{38} g^{-2} \geq$  $10^{91}$  on the parameters of the PBB theory follow from the requirement of flatness of the universe, so they will remain intact even if it is possible to solve the homogeneity problem in the PBB cosmology. We believe that the need to introduce such extraordinarily large parameters represents a major challenge to the PBB cosmology.

## Acknowledgments

The author wishes to thank R. Bousso and A. Linde for collaboration on <sup>6</sup> and for useful conversations, A. Linde for very helpful comments on the manuscript, and T. Damour and G. Veneziano for useful conversations and clarification of their ideas. This work has been supported in part by NSF Grant PHY-9219345.

## References

- 1. G. Veneziano, Phys. Lett. B 265, 287 (1991); M. Gasperini and G. Veneziano, Astropart. Phys. 1, 317 (1993).
- 2. M.S. Turner and E. Weinberg, Phys. Rev. D 56, 4604 (1997).
- 3. G. Veneziano, Phys. Lett. B 406, 297 (1997).
- 4. A. Buonanno, K.A. Meissner, C. Ungarelli and G. Veneziano, Phys. Rev. D **57**, 2543 (1998).
- 5. M. Maggiore and R. Sturani, Phys. Lett. B 415, 335 (1997).
- 6. N. Kaloper, A. Linde and R. Bousso, eprint [hep-th/9801073](http://arxiv.org/abs/hep-th/9801073).
- 7. A. Buonanno, T. Damour and G. Veneziano, eprint [hep-th/9806230](http://arxiv.org/abs/hep-th/9806230).
- 8. A. Ghosh, G. Pollifrone and G. Veneziano, eprint [hep-th/9806233.](http://arxiv.org/abs/hep-th/9806233)