One-loop effective potential of $N=1$ supersymmetric theory and decoupling effects

I.L. Buchbinder^{*}, M. Cvetič⁺ and A.Yu. Petrov^{*}

[∗]Department of Theoretical Physics, Tomsk State Pedagogical University 634041 Tomsk, Russia

⁺Department of Physics and Astronomy, University of Pennsylvania Philadelphia, PA 19104–6396, USA

Abstract

We study the decoupling effects in $N = 1$ (global) supersymmetric theories with chiral superfields at the one-loop level. Examples of gauge neutral chiral superfields with minimal (renormalizable) as well as non-minimal (non-renormalizable) couplings are considered, and decoupling in gauge theories with $U(1)$ gauge superfields that couple to heavy chiral matter is studied. We calculate the one-loop corrected effective Lagrangians that involve light fields and heavy fields with mass of order M . Elimination of heavy fields by equations of motion leads to decoupling effects with terms that grow logarithmically with M . These corrections renormalize light fields and couplings in the theory (in accordance with the "decoupling theorem"). When the field theory is an effective theory of the underlying fundamental theory, like superstring theory, where the couplings are calculable, such decoupling effects modify the low energy predictions for the effective couplings of light fields. In particular, for the class of string vacua with an "anomalous" $U(1)$, the vacuum restabilization triggers decoupling effects, which can significantly modify the low energy predictions for couplings of the surviving light fields. We also demonstrate that quantum corrections to the chiral potential depending on massive background superfields and corresponding to supergraphs with internal massless lines and external massive lines can also arise at the two-loop level.

Contents

1 Introduction

This paper is devoted to the calculation of the one-loop effective action for several models of the global $N = 1$ supersymmetric theory with chiral superfields and a subsequent study of some of their phenomenologically interesting aspects. In particular, we investigate in detail the decoupling effects due to the couplings of heavy and light chiral superfields in the theory and subsequent implications for the low energy effective action of light superfields.

In principle the decoupling effects of heavy fields in field theory are well understood. According to the decoupling theorem [\[1](#page-54-0), [2\]](#page-55-0) (for additional references see, e.g.,[[3\]](#page-55-0)) in the field theory of interacting light (with masses m) and heavy fields (with masses M) the heavy fields decouple; the effective Lagrangian of the light fields can be written in terms of the original classical Lagrangian of light fields with loop effects of heavy fields absorbed into redefinitions of new light fields, masses and couplings, and the only new terms in this effective Lagrangian are non- renormalizable, proportional to inverse powers of M (both at tree- and loop-levels).

In a field theory as an effective description of phenomena at certain energies, the rescaling of the fields and couplings due to heavy fields does not affect the structure of the couplings, since those are free parameters whose values are determined by experiments. On the other hand if the field theory is describing an effective theory of an underlying fundamental theory, like superstring theory, where the couplings at the string scale are calculable, the decoupling effects of the heavy field can be important and can significantly affect the low energy predictions for the couplings of light fields at low energies. Therefore the quantitative study of decoupling effects at the loop-level in effective supersymmetric theories is important; it should improve our understanding of such effects for the effective Lagrangians from superstring theory and provide us with calculable corrections for the low energy predictions of the theory. We also note that as the decoupling theorem is based on finite renormalization of fields and parameters as all parameters in the effective theory (fields, masses, couplings) are determined from the corresponding string theory and hence cannot be renormalized. Therefore we will use consistence with the decoupling theorem only to check the results.

Effective theories of $N = 1$ supersymmetric four-dimensional perturbative string vacua can be obtained by employing techniques of two-dimensional conformal field theory [\[4\]](#page-55-0). In particular the kählerian and the chiral (super-) potential can be calculated explicitly at the tree level. While the chiral potential terms calculated at the string tree-level are protected from higher genus corrections (for a representative work on the subject see, e.g., [\[5](#page-55-0), [6](#page-55-0)], and references therein), the kählerian potential is not. Such higher genus corrections to the kählerian potential could be significant; however, their structure has not been studied very much. In this paper we shall not address these issues and assume that the string theory calculation provides us with a (reliable, calculable) form of the effective theory at M_{string} , which would in turn serve as a starting point of our study.

One of the compelling motivations for a detailed study of decoupling effects is the phenomenon of vacuum restabilization[[7](#page-55-0)] for a class of four-dimensional (quasi-realistic) heterotic superstring vacua with an "anomalous" $U(1)$. (On the open Type I string side these effects are closely related to the blowing-up procedure of Type I orientifolds and were recently studied in[[8\]](#page-55-0).) For such string vacua of perturbative heterotic string theory, the Fayet-Iliopoulos (FI) D- term is generated at genus-one[[9\]](#page-55-0), thus triggering certain fields to acquire vacuum expectation values (VEV's) of order $M_{String} \sim g_{gauge} M_{Planck} \sim 5 \times 10^{17}$ GeV along D- and F- flat directions of the effective $N = 1$ supersymmetric theory. (Here g_{gauge} is the gauge coupling and M_{Planck} the Planck scale.) Due to these large string-scale VEV's a number of additional fields obtain large string-scale masses. Some of them in turn couple through (renormalizable) interactions to the remaining light fields, and thus through decoupling effects affect the effective theory of light fields at low energies. (For the study of the effective Lagrangians and their phenomenological implications for a class of such four-dimensional stringvacua see, e.g., $[10]$ $[10]$ $[10]$ – $[12]$ $[12]$ $[12]$ and references therein.)

The tree level decoupling effects within $N = 1$ supersymmetric theories, were studied withinan effective string theory in [[13\]](#page-55-0). In a related work [[14](#page-55-0)] it was shown that the leading order corrections of order $\frac{1}{M}$ are to the effective chiral potential, however there are also

important next order effects in the effective chiral potential[[15\]](#page-55-0). In addition, in[[15\]](#page-55-0) the nonrenormalizablemodifications of the kählerian potential (as was also pointed out in $[16]$ $[16]$ $[16]$) were systematically studied. These tree level decoupling effects (as triggered by, e.g., vacuum restabilization for a class of string vacua) lead to new nonrenormalizable interactions which are competitive with the nonrenormalizable terms that are calculated directly in the superstring theory.

In this paper we consider one-loop decoupling effects in $N = 1$ supersymmetric theory. We study both the effects on chiral (gauge neutral) superfields and on the effects of gauge superfields. (In another context see [\[17\]](#page-56-0).) It turns out that an essential modification of low energy predictions takes place not only for chiral superfields[[18\]](#page-56-0) but also for gauge superfields. As stated earlier such effective Lagrangians arise naturally due to the vacuum restabilization for a class of supersymmetric string vacua and trigger couplings between heavy fields with massscale $M \sim 10^{17}$ GeV and the light (massless) fields [[19\]](#page-56-0). (Note however, that we do not include supergravity effects which could also be significant.)

As a result we find that the one-loop effective action after a redefinition of fields, masses and couplings coincides with the one-loop effective action of the corresponding theory where heavy superfields are completely absent, in accordance with the decoupling theorem. However, since the masses and the couplings of the fields are calculable in string theory (at the mass scale M_{string} , the decoupling effects add additional corrections to the effective action of the light superfields. These corrections grow logarithmically with M (mass of the heavy superfields) and modify the effective couplings in an essential way, which for a class of string vacua under consideration can be significant.

Another interesting result presented in this paper pertains to the chiral effective potential. When the chiral potential depends on massive superfields, quantum corrections due to these fields appear earlier than in the case when one considers light fields only.

This paper is organized as follows. In Section 2 the general structure of the effective action studied is given and the general approach to addressing the decoupling effects is presented. Section 3 is devoted to the study of the effective action for the "minimal" model with one heavy and one light (gauge neutral) chiral superfield. In Section 4 the leading order decoupling corrections to the effective action for non-minimal models (with more general couplings) are considered. Section 5 is devoted to the investigation of the one-loop decoupling effects in $N = 1$ supersymmetric theory with $U(1)$ gauge vector superfields and chiral superfields charged under $U(1)$. A summary and discussion of the obtained results are given in Section 6. In Appendix A details of the calculation of the one-loop kählerian effective potential for the minimal model are presented, in Appendix B the calculation of the one-loop kählerian effective action via diagram technique for the minimal model is described, and in Appendix C details of the calculation for the effective action of non-minimal models are given.

2 Effective action in the model of interacting light and heavy superfields

2.1 General structure of the effective action

 $N = 1$ supersymmetric actions with chiral supermultiplets arise as a subsector of an effective theory of $N = 1$ supersymmetric string vacua. Such calculations are carried out for perturbative string vacua primarily by employing conformal field theory techniques. (Though less powerful techniques, e.g., sigma-model approach, in which the integration over massive string modes is carried out in the the background of the ten-dimensional manifold with the structure $M^4 \times K$ where M^4 is a four-dimensional Minkowski space and K is a suitable sixdimensional compact (Calabi-Yau) manifold, can also be employed.) The resulting effective theories contain as an ingredient $N = 1$ chiral superfields Φ^i with action

$$
S[\Phi, \bar{\Phi}] = \int d^8 z K(\bar{\Phi}^i, \Phi^i) + (\int d^6 z W(\Phi^i) + h.c.) \tag{2.1}
$$

Here $\Phi^i = \Phi^i(z)$, $z^A \equiv (x^a, \theta_\alpha, \bar{\theta}_\alpha); a = 0, 1, 2, 3; \alpha = 1, 2, \dot{\alpha} = 1, 2, d^8 z = d^4 x d^2 \theta d^2 \bar{\theta}.$ Real function $K(\bar{\Phi}^i, \Phi^i)$ is called the kählerian potential, the holomorphic function $W(\Phi^i)$ is called the chiral potential[[20](#page-56-0)]. Expression (2.1) represents the most general action of gauge neutral chiral superfields which does not contain higher derivatives at a component level[[20\]](#page-56-0). We refer to this action as the chiral superfield model of a general form. In a special case $K(\bar{\Phi}^i, \Phi^i) = \Phi \bar{\Phi}$, $W(\Phi^i) \sim \Phi^3$ we obtain the well-known Wess-Zumino model. For $W(\Phi^i) = 0$ the present theory represents itself as a $N = 1$ supersymmetric four-dimensional sigma-model (see, e.g., [[6\]](#page-55-0)). The action (2.1) , which originates from superstring theory, can be treated as a classical effective action of the fundamental theory, suitable for description of phenomena at energies much less than the Planck scale. Such models of chiral superfields are widely used for the study of possible phenomenological implications of superstring theories (see, e.g., recent papers [\[8](#page-55-0), [10](#page-55-0), [11](#page-55-0), [12,](#page-55-0) [18\]](#page-56-0) and references therein). One of the most important aspects of the study of these models pertains to the investigation of the decoupling effects, which is the main subject of the present paper.

The starting point in the study of the decoupling effects is the model with the classical action (2.1) and, for the sake of simplicity, two chiral superfields: a light one, ϕ , and a heavy one, Φ , i.e. $\Phi^i = {\Phi, \phi}$. The aim is to to calculate the low-energy effective action in the oneloop approximation and to compute the one-loop corrected effective action of light superfield, only.

We refer to the model in which the kählerian potential is of the canonical (minimal) form:

$$
K(\Phi, \bar{\Phi}, \phi, \bar{\phi}) = \Phi \bar{\Phi} + \phi \bar{\phi}
$$
\n(2.2)

as the minimal model, and the model in which

$$
K(\Phi, \bar{\Phi}, \phi, \bar{\phi}) = \Phi \bar{\Phi} + \phi \bar{\phi} + \tilde{K}(\Phi, \bar{\Phi}, \phi, \bar{\phi})
$$
\n(2.3)

with $\tilde{K} \neq 0$ – as the non-minimal one (in analogy with [[10\]](#page-55-0)). We assume that the function $\tilde{K}(\Phi, \bar{\Phi}, \phi, \bar{\phi})$ can be expanded into power series in superfields $\Phi, \bar{\Phi}, \phi, \bar{\phi}$ where the leading order term is at least of the third order in the chiral superfields (and thus proportional to at least one inverse power of M)

$$
\tilde{K}(\Phi, \bar{\Phi}, \phi, \bar{\phi}) = \frac{\phi \bar{\Phi}^2}{M} + \frac{\Phi \bar{\phi}^2}{M} \dots \tag{2.4}
$$

The chiral potential M is taken to be of the form:

$$
W = \frac{M}{2}\Phi^2 + \tilde{W}(\Phi, \phi)
$$
\n(2.5)

where \tilde{W} is also at least of the third order in $\{\Phi, \phi\}$ superfields. The functions \tilde{W} could have the following structure

$$
\tilde{W}(\Phi,\phi) = \phi\Phi^2 + \Phi\phi^2 + \phi^3 + \Phi^3 + \frac{\phi\Phi^3}{M} + \dots \tag{2.6}
$$

with M as a massive parameter. Hence the possible vertices of interaction of superfields have the form

$$
\phi\Phi^2,\Phi\phi^2,\frac{\Phi\bar\phi^2}{M},\phi^3,\Phi^3,\frac{\phi\Phi^3}{M}\ldots
$$

The effective action $\Gamma[\Phi, \bar{\Phi}, \phi, \bar{\phi}]$ is defined as the Legendre transform from the generating functionalof connected Green functions [[21](#page-56-0)] $W[J, \bar{J}]$:

$$
\exp(\frac{i}{\hbar}W[J,\bar{J},j,\bar{j}]) = \int \mathcal{D}\varphi \mathcal{D}\bar{\varphi} \mathcal{D}\bar{\Psi} \mathcal{D}\Psi \exp(\frac{i}{\hbar}(S[\Psi,\bar{\Psi},\varphi,\bar{\varphi}] + \langle \int d^6z(J\Psi + j\varphi) + h.c.)))
$$
\n
$$
\Gamma[\Phi,\bar{\Phi},\phi,\bar{\phi}] = W[J,\bar{J}] - (\int d^6z(J\Phi + j\phi) + h.c.)
$$
\n(2.7)

Γ[Φ, $\bar{\Phi}$, ϕ , $\bar{\phi}$] can be calculated using the loop-expansion method. This method employs the splitting of all the chiral superfields into a sum of the background superfields Φ , ϕ and the quantum ones Φ_q , ϕ_q , using the rule

$$
\Phi \rightarrow \Phi + \sqrt{\hbar} \Phi_q
$$
\n
$$
\phi \rightarrow \phi + \sqrt{\hbar} \phi_q
$$
\n(2.8)

As a result the action([2.1](#page-4-0)) after such changes can be written as

$$
S_q = \int d^8 z K(\Phi + \sqrt{\hbar} \Phi_q, \bar{\Phi} + \sqrt{\hbar} \bar{\Phi}_q, \phi + \sqrt{\hbar} \phi_q, \bar{\phi} + \sqrt{\hbar} \bar{\phi}_q) +
$$

+
$$
[\int d^6 z W(\Phi + \sqrt{\hbar} \Phi_q, \phi + \sqrt{\hbar} \phi_q) + h.c.]
$$
 (2.9)

and the effective action takes the form:

$$
\exp(\frac{i}{\hbar}\Gamma[\Phi,\bar{\Phi},\phi,\bar{\phi}]) = \int \mathcal{D}\phi_q \mathcal{D}\bar{\phi}_q \mathcal{D}\bar{\Phi}_q \mathcal{D}\Phi_q \exp(\frac{i}{\hbar}S[\Phi+\sqrt{\hbar}\Phi_q,\bar{\Phi}+\sqrt{\hbar}\bar{\Phi}_q, \phi+\sqrt{\hbar}\phi_q, \bar{\phi}+\sqrt{\hbar}\phi_q] - \sqrt{\hbar}\left(\int d^6z(\frac{\delta\Gamma}{\delta\Phi(z)}\Phi_q(z)+\frac{\delta\Gamma}{\delta\phi(z)}\phi_q(z)) + h.c.)\right)
$$
(2.10)

(fordetails see [[20](#page-56-0), [21\]](#page-56-0)). The effective action [\(2.10](#page-5-0)) can be cast in the form $\Gamma[\Phi, \bar{\Phi}, \phi, \bar{\phi}] =$ $S[\Phi, \bar{\Phi}, \phi, \bar{\phi}] + \tilde{\Gamma}[\Phi, \bar{\Phi}, \phi, \bar{\phi}]$. Here $\tilde{\Gamma}[\Phi, \bar{\Phi}, \phi, \bar{\phi}]$ is a quantum correction in effective action which can be expanded into power series in \hbar as

$$
\tilde{\Gamma}[\Phi, \bar{\Phi}, \phi, \bar{\phi}] = \sum_{n=1}^{\infty} \hbar^n \Gamma^{(n)}[\Phi, \bar{\Phi}, \phi, \bar{\phi}]
$$
\n(2.11)

The one-loop quantum correction $\Gamma^{(1)}$ to the effective action is defined through the following expression[[21\]](#page-56-0):

$$
e^{i\Gamma^{(1)}} = \int \mathcal{D}\Phi_q \mathcal{D}\bar{\Phi}_q \mathcal{D}\phi_q \mathcal{D}\bar{\phi}_q \exp(iS_q^{(2)}) \tag{2.12}
$$

Here $S_q^{(2)}$ corresponds to the part of S_q [\(2.9](#page-5-0)) which is quadratic in quantum superfields. It is of the form

$$
S_q^{(2)} = \int d^8 z (K_{\Phi \bar{\Phi}} \Phi_q \bar{\Phi}_q + K_{\phi \bar{\Phi}} \phi_q \bar{\Phi}_q + K_{\Phi \bar{\phi}} \Phi_q \bar{\phi}_q + K_{\phi \bar{\phi}} \phi_q \bar{\phi}_q) +
$$

+
$$
[\int d^6 z W_{\Phi \Phi} \Phi_q^2 + W_{\phi \Phi} \Phi_q \phi_q + W_{\phi \phi} \phi_q^2 + h.c.]
$$
 (2.13)

As a result we arrive at the one-loop effective action of the form

$$
\Gamma[\Phi, \bar{\Phi}, \phi, \bar{\phi}] = S + \hbar \Gamma^{(1)} = \int d^8 z K(\Phi, \bar{\Phi}, \phi, \bar{\phi}) + [\int d^6 z W(\Phi, \phi) + h.c.] +
$$

+ $\hbar (\int d^8 z K^{(1)}(\Phi, \bar{\Phi}, \phi, \bar{\phi}) + (\int d^6 z W^{(1)}(\Phi, \phi) + h.c.))$ (2.14)

Here we suppose that the one-loop correction in the effective action $\Gamma^{(1)}$ can be represented in the form

$$
\Gamma^{(1)} = \int d^8 z K^{(1)}(\Phi, \bar{\Phi}, \phi, \bar{\phi}) + (\int d^6 z W^{(1)}(\Phi, \phi) + h.c.) + \dots \tag{2.15}
$$

Dots denote terms that depend on the supercovariant derivatives of the chiral superfields.

The loop corrected effective action has the following structure

$$
\Gamma[\Phi, \bar{\Phi}, \phi, \bar{\phi}] = \int d^8 z L_{eff}(\Phi, D_A \Phi, D_A D_B \Phi, \bar{\Phi}, D_A \bar{\Phi}, D_A D_B \bar{\Phi},
$$

$$
\phi, D_A \phi, D_A D_B \phi, \bar{\phi}, D_A \bar{\phi}, D_A D_B \bar{\phi}) + (\int d^6 z L_{eff}^{(c)}(\Phi, \phi) + h.c.) + ...
$$
 (2.16)

Here D_A are supercovariant derivatives, $D_A = (\partial_a, D_\alpha, \bar{D}_{\dot{\alpha}})$. L_{eff} is the effective super-Lagrangian that we write in the form

$$
L_{eff} = K_{eff}(\Phi, \bar{\Phi}, \phi, \bar{\phi}) + \dots
$$

$$
K = K(\Phi, \bar{\Phi}, \phi, \bar{\phi}) + \sum_{n=1}^{\infty} \hbar^n K^{(n)}
$$
(2.17)

and $L^{(c)}$ is the effective chiral Lagrangian

$$
L^{(c)} = W_{eff}(\Phi, \phi) + \dots \tag{2.18}
$$

 K_{eff} is the kählerian effective potential that depends only on the chiral superfields Φ , $\bar{\Phi}$, ϕ , $\bar{\phi}$ but not on their (covariant) derivatives. W_{eff} is the chiral effective potential that depends on on (holomorphic) chiral superfields $\{\Phi, \phi\}$, only. Dots denote the terms that depend on the the space-time derivatives of chiral superfields only. Furthermore, one can prove that the one-loop correction to the chiral potential is zero (for the $N = 1$ supersymmetric theory which does not include gauge superfields). However, higher corrections can exist (cf. [\[22](#page-56-0)]–[[24](#page-56-0)]), i.e.

$$
W_{eff}(\Phi,\phi) = W(\Phi,\bar{\phi}) + \sum_{n=2}^{\infty} \hbar^n W^{(n)}(\Phi,\phi)
$$
 (2.19)

Here $K^{(n)}$ and $W^{(n)}$ are loop corrections to the kählerian and chiral potential, respectively.

Since in this paper we concentrate on the one-loop corrected effective action only, we are mainly interested in the correction to the kählerian potential which is the leading term in the one-loop corrected low-energy effective action. (At low energies $(E \ll M)$ higher derivative terms are suppressed.)

Our ultimate goal is to obtain the effective action for light superfields, only. For that purpose one must eliminate heavy superfields from the one-loop effective action $\Gamma[\Phi, \Phi, \phi, \phi]$ [\(2.14\)](#page-6-0) by means of the effective equations of motion. These equations can be solved by an iterative method up to a certain order in the inverse mass M of heavy superfield. Substituting a solution of these equations into the effective action [\(2.14](#page-6-0)) we then obtain the one-loop corrected effective action of light superfields only. In the following subsection we shall describe the procedure in detail.

2.2 The effective equations of motion

The effective equations of motion for heavy superfields in the model with the effective action $(2.14, 2.15)$ are of the form

$$
\frac{\delta \Gamma}{\delta \Phi} = 0 \quad : \quad -\frac{1}{4} \bar{D}^2 \left(\frac{\partial K}{\partial \Phi} + \frac{\partial K^{(1)}}{\partial \Phi} \right) + \frac{\partial W}{\partial \Phi} = 0
$$
\n
$$
\frac{\delta \Gamma}{\delta \bar{\Phi}} = 0 \quad : \quad -\frac{1}{4} \bar{D}^2 \left(\frac{\partial K}{\partial \bar{\Phi}} + \frac{\partial K^{(1)}}{\partial \bar{\Phi}} \right) + \frac{\partial \bar{W}}{\partial \bar{\Phi}} = 0
$$
\n
$$
(2.20)
$$

The effective equations of motion for light superfields have an analogous form. We consider the case when the interactions with the gauge superfields are absent (see however Section 5) and $W^{(1)} = 0$ (which is absent at one-loop level (cf., discussion above)).

The equations (2.20) can be solved via an iterative method, described below. We can represent the heavy superfield Φ in the form

$$
\Phi = \Phi_0 + \Phi_1 + \ldots + \Phi_n + \ldots \tag{2.21}
$$

where Φ_0 is zeroth-order approximation, Φ_1 is first-order one, etc.. We assume that $|D^2\Phi| \ll$ $M\Phi$ since the superfield Φ is heavy, and thus the assumption is valid. The zeroth-order approximation Φ_0 can be found from the condition

$$
\frac{\partial W}{\partial \Phi}|_{\Phi = \Phi_0} = 0
$$

After a substitution of the expansion([2.21\)](#page-7-0) into equations [\(2.20](#page-7-0)) we arrive at the following equation for the $(n+1)$ -th-order solution for $\bar{\Phi}_{n+1}$

$$
\left(\frac{\partial \bar{W}}{\partial \bar{\Phi}}\right)|_{\Phi = \Phi_0 + \dots + \Phi_{n+1}} - \left(\frac{\partial \bar{W}}{\partial \bar{\Phi}}\right)|_{\Phi = \Phi_0 + \dots + \Phi_n} =
$$
\n
$$
\bar{D}^2 \Big|_{\phi = \Phi_0 + \dots + \Phi_{n+1}} - \left(\frac{\partial \bar{W}}{\partial \bar{\Phi}}\right)|_{\Phi = \Phi_0 + \dots + \Phi_n} =
$$
\n
$$
\frac{\partial K^{(1)}}{\partial K^{(1)}} \Big|_{\phi = \Phi_0 + \dots + \Phi_{n+1}} - \left(\frac{\partial \bar{W}}{\partial \bar{\Phi}}\right)|_{\Phi = \Phi_0 + \dots + \Phi_n} =
$$
\n
$$
\frac{\partial K^{(1)}}{\partial K^{(1)}} \Big|_{\phi = \Phi_0 + \dots + \Phi_{n+1}} - \left(\frac{\partial \bar{W}}{\partial \bar{\Phi}}\right)|_{\Phi = \Phi_0 + \dots + \Phi_n} =
$$
\n
$$
\frac{\partial K^{(1)}}{\partial K^{(2)}} \Big|_{\phi = \Phi_0 + \dots + \Phi_{n+1}} - \left(\frac{\partial \bar{W}}{\partial \bar{\Phi}}\right)|_{\Phi = \Phi_0 + \dots + \Phi_n} =
$$
\n
$$
\frac{\partial K^{(1)}}{\partial K^{(2)}} \Big|_{\phi = \Phi_0 + \dots + \Phi_{n+1}} - \left(\frac{\partial \bar{W}}{\partial \bar{\Phi}}\right)|_{\Phi = \Phi_0 + \dots + \Phi_n} =
$$
\n
$$
\frac{\partial K^{(1)}}{\partial K^{(2)}} \Big|_{\phi = \Phi_0 + \dots + \Phi_{n+1}} - \left(\frac{\partial \bar{W}}{\partial \bar{\Phi}}\right)|_{\Phi = \Phi_0 + \dots + \Phi_n} =
$$
\n
$$
\frac{\partial K^{(1)}}{\partial K^{(2)}} \Big|_{\phi = \Phi_0 + \dots + \Phi_n} =
$$
\n
$$
\frac{\partial K^{(1)}}{\partial K^{(1)}} \Big|_{\phi = \Phi_0 + \dots + \Phi_n} =
$$
\n
$$
\frac{\partial K^{(2)}}{\partial K^{(2)}} \Big|_{\phi = \Phi_0 + \dots + \Phi_n} =
$$

$$
= \frac{2}{4} \left(\frac{\partial \mathbf{R}}{\partial \Phi} \middle| \Phi = \Phi_0 + ... + \Phi_n - \frac{\partial \mathbf{R}}{\partial \Phi} \middle| \Phi = \Phi_0 + ... + \Phi_{n-1} + \frac{\partial \mathbf{R}}{\partial \Phi} \middle| \Phi = \Phi_0 + ... + \Phi_n - \frac{\partial \mathbf{R}}{\partial \Phi} \middle| \Phi = \Phi_0 + ... + \Phi_{n-1} \right)
$$

and an analogous equation for Φ_{n+1} . These sets of equations can be used to calculate Φ_n for any $n \ge 1$. Since the chiral potential is of the form $W = \frac{M}{2}\Phi^2 + \tilde{W}$ (see eqs. [\(2.5-2.6](#page-5-0)), eq. (2.23)) can be rewritten in the form

$$
M\bar{\Phi}_{n+1} + (\frac{\partial \tilde{\tilde{W}}}{\partial \bar{\Phi}})|_{\Phi = \Phi_0 + \dots + \Phi_{n+1}} - (\frac{\partial \tilde{\tilde{W}}}{\partial \bar{\Phi}})|_{\Phi = \Phi_0 + \dots + \Phi_n} =
$$
\n
$$
\bar{\Phi}_{n+1} + (\frac{\partial \tilde{\tilde{W}}}{\partial \bar{\Phi}})|_{\Phi = \Phi_0 + \dots + \Phi_n} =
$$
\n
$$
(2.23)
$$

$$
= \frac{\bar{D}^2}{4} \left(\frac{\partial K}{\partial \Phi}\big|_{\Phi = \Phi_0 + \ldots + \Phi_n} - \frac{\partial K}{\partial \Phi}\big|_{\Phi = \Phi_0 + \ldots + \Phi_{n-1}} + \frac{\partial K^{(1)}}{\partial \Phi}\big|_{\Phi = \Phi_0 + \ldots + \Phi_n} - \frac{\partial K^{(1)}}{\partial \Phi}\big|_{\Phi = \Phi_0 + \ldots + \Phi_{n-1}}\right)
$$

with an analogous equation for Φ_{n+1} . As a result we can find Φ , $\bar{\Phi}$ in principle to any order.

Thus, a substitution of the heavy superfields Φ , $\bar{\Phi}$ expressed in terms of light superfields by means of equations (2.22-2.23) allows one to obtain the one-loop corrected effective action of light superfields only.

2.3 Calculation of the one-loop kählerian effective potential

Let us now consider the method to calculate the one-loop effective action. The most effective method consists of expressing the quantities in terms of matrices (matrix approach). First we write the heavy quantum superfield Φ_q and light one ϕ_q as a column vector: $\psi_q =$ $\int \Phi_q$ ϕ_q \setminus . As a result, the part of the classical action that is quadratic in quantum superfields([2.13\)](#page-6-0) takes the form

$$
S^{(2)}[\psi_q, \bar{\psi}_q] = \int d^8 z K_{\psi \bar{\psi}} \psi_q \bar{\psi}_q + (\int d^6 z W'' \psi_q^2 + h.c.) \tag{2.24}
$$

Here $K_{\psi\bar{\psi}}$ and \overline{W}'' are matrices of the form

$$
K_{\psi\bar{\psi}} = \begin{pmatrix} K_{\phi\bar{\phi}} & K_{\phi\bar{\Phi}} \\ K_{\Phi\bar{\phi}} & K_{\Phi\bar{\Phi}} \end{pmatrix} W'' = \begin{pmatrix} W_{\phi\phi} & W_{\phi\Phi} \\ W_{\phi\Phi} & W_{\Phi\Phi} \end{pmatrix}
$$
(2.25)

The one-loop effective action [\[21](#page-56-0)]([4.4\)](#page-24-0) is of the form

$$
exp(i\Gamma^{(1)}) = \int D\psi_q D\bar{\psi}_q S^{(2)}[\psi_q, \bar{\psi}_q]
$$
\n(2.26)

where $S^{(2)}[\psi_q, \bar{\psi}_q]$ is given by (2.24), $D\psi_q$ denotes the integration over both ϕ_q and Φ_q .

To perform the integration we employ the technique analogous to that used for the study of the Wess-Zumino model $[22, 23]$ $[22, 23]$ $[22, 23]$. First we consider a theory of a real scalar superfield q which represents itself as a column $q =$ $\int u$ \overline{v} \setminus . The action for q reads as

$$
S_q^0 = -\frac{1}{16} \int d^8 z q^T D^\alpha \bar{D}^2 D_\alpha q \tag{2.27}
$$

where q^T is a string (row) of the form $q^T = (u \, v)$. The effective action U_q for the superfield q can be obtained by the Faddeev-Popov procedure. We choose the gauge fixing functions $\chi[q] = \frac{1}{4}\bar{D}^2q - \psi$ and $\bar{\chi}[q] = \frac{1}{4}D^2q - \bar{\psi}$, hence the effective action \bar{U}_q is defined with the following integral:

$$
e^{iU_q} = \int Dq \delta(\frac{1}{4}D^2q - \bar{\psi}) \delta(\frac{1}{4}\bar{D}^2q - \psi) \exp(iS_v) det M_0 \qquad (2.28)
$$

Here M_0 is a Faddeev-Popov matrix

$$
M_0 = \begin{pmatrix} 0 & -1_2 \frac{1}{4} \bar{D}^2 \\ -1_2 \frac{1}{4} D^2 & 0 \end{pmatrix}
$$
 (2.29)

where 1_2 is an unit 2×2 matrix. By definition, U_q is constant: since the theory of q is gauge invariant the corresponding effective action does not depend on ψ (cf. [\[22, 23\]](#page-56-0)).

By multiplying the corresponding left-hand sides and right-hand sides of eqs.([2.26\)](#page-8-0) and (2.28) we obtain

$$
\exp(i\Gamma^{(1)}[\Phi,\bar{\Phi}] + iU_q) = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} Du Dv \delta(\frac{1}{4}D^2q - \bar{\psi}) \delta(\frac{1}{4}\bar{D}^2q - \psi) \exp(iS_q) \times \times det M_0 \exp(iS^{(2)}[\psi,\bar{\psi}])
$$
\n(2.30)

Then we integrate over $\psi, \bar{\psi}$ by means of the delta functions (cf. [\[22, 23, 20\]](#page-56-0)) and since $e^{iU_{\psi}}$ and $det M_0$ are constants we arrive at the following expression for the one-loop effective action for the superfields ψ, ψ :

$$
\exp(i\Gamma^{(1)}) = \int Dq \exp(iS[q]) \tag{2.31}
$$

which leads to

$$
\Gamma^{(1)} = \frac{i}{2} Tr \log \Delta \tag{2.32}
$$

Here Tr is a functional supertrace, and

$$
S[q] = \frac{1}{2}q\Delta q \tag{2.33}
$$

$$
\Delta = \Box 1_2 - \frac{1}{16} (K_{\psi\bar{\psi}} - 1) \{D^2, \bar{D}^2\} - \frac{1}{4} W'' \bar{D}^2 - \frac{1}{4} \bar{W}'' D^2 \tag{2.34}
$$

The terms proportional to the supercovariant derivatives of $K_{\psi\bar{\psi}}, W^{''}$ and $\bar{W}^{''}$ are omitted since the one-loop kählerian effective potential by definition does not depend on the derivatives of superfields.

In order to determine $Tr \log \Delta$ we use the Schwinger representation

$$
Tr \log \Delta = tr \int \frac{ds}{s} \exp(is\tilde{\Delta}) \exp(is\tilde{\Delta})
$$

Here tr denotes a trace of the matrix. Since Δ is a matrix operator we turn to the problem of calculating the exponent of the matrix. Let $\Omega = \exp(is\Delta))$. Then the matrix equation for Ω reads

$$
-i\frac{\partial}{\partial s}\Omega = \Omega\tilde{\Delta}
$$
 (2.35)

Here $\tilde{\Delta}$ is a matrix operator of the form

$$
\tilde{\Delta} = -\frac{1}{16}(K_{\psi\bar{\psi}} - 1)\{D^2, \bar{D}^2\} - \frac{1}{4}W''\bar{D}^2 - \frac{1}{4}\bar{W}''D^2
$$
\n(2.36)

In order to solve the equation (2.35) we represent Ω in the form of an expansion in terms of a spinor supercovariant derivatives

$$
\Omega(s) = 1_2 + \frac{1}{16}A(s)D^2\bar{D}^2 + \frac{1}{16}\tilde{A}(s)\bar{D}^2D^2 + \frac{1}{8}B^{\alpha}(s)D_{\alpha}\bar{D}^2 + \frac{1}{8}\tilde{B}_{\dot{\alpha}}(s)D^{\dot{\alpha}}D^2 + \frac{1}{4}C(s)D^2 + \frac{1}{4}\tilde{C}(s)\bar{D}^2
$$
\n(2.37)

The equation (2.35) leads to the following system of equations for the coefficients A, B, C ,

$$
\frac{1}{i}\dot{A} = F + AF\Box - CW''
$$
\n
$$
\frac{1}{i}\dot{B}^{\alpha} = \tilde{B}_{\dot{\alpha}}W''\partial^{\alpha\dot{\alpha}} + B^{\alpha}F\Box
$$
\n
$$
\frac{1}{i}\dot{C} = -\bar{W}'' - A\bar{W}''\Box + CF\Box
$$
\n(2.38)

and an analogous system of equations for $\tilde{A}, \tilde{B}, \tilde{C}$, which can be obtained from this one by changing W["] into \bar{W} " and vice versa. Here $F = 1 - K_{\psi\bar{\psi}}$. Since the initial condition for Ω is $\Omega|_{s=0} = 1$ the initial conditions for $A, \tilde{A}, B, \tilde{B}, C, \tilde{C}$ are $A|_{s=0} = \tilde{A}|_{s=0} = C|_{s=0} = \tilde{C}|_{s=0} = 0.$ The solution for B^{α} , $\tilde{B}^{\dot{\alpha}}$ evidently has the form $B^{\alpha} = \tilde{B}^{\dot{\alpha}} = 0$. The manifest form of the matrices $A, \tilde{A} C, \tilde{C}$, necessary for exact calculations, is of the form

$$
A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}; C = \begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix}
$$
 (2.39)

Here index 1 denotes the sector of heavy superfield Φ and 2 the sector of light superfield ϕ . Now let us solve the system for matrices A, C . The solution for $\overline{A}, \overline{C}$ can be easily obtained in an analogous way since the system with $B^{\alpha} = \tilde{B}^{\dot{\alpha}} = 0$ is invariant under the change $A \to \tilde{A}$, $C \to \tilde{C}$.

Let us study the solution for A, C which should be chosen in the form

$$
A = A_i + A_0 \tag{2.40}
$$

$$
C = C_i + C_0
$$

Here A_i , C_i is a partial solution of the inhomogeneous system, and A_0 , C_0 is a general solution of the homogeneous system. It is straightforward to see that $A_i = -1\Box^{-1}$, $C_i = 0$. And A_0 , C_0 should satisfy the system of equations

$$
\frac{1}{i}\dot{A}_0 = A_0 F \Box + C_0 W''
$$
\n
$$
\frac{1}{i}\dot{C}_0 = A_0 \bar{G} \Box + C_0 F \Box
$$
\n(2.41)

 A_0, C_0 should be chosen to be of the form $A_0 = a_0 \exp(i\omega s), C_0 = c_0 \exp(i\omega s)$ where a_0, c_0, ω are some functions of the background superfields and the d'Alembertian operator, but are independent of s. As a result we arrive at the equations for a_0, c_0 :

$$
a_0(\omega 1 - F\Box) + c_0 W'' = 0
$$

\n
$$
c_0(\omega 1 - F\Box) + a_0 \bar{W}'' \Box = 0
$$
\n(2.42)

This system of equations has a non-trivial solution at

$$
\det\left(\begin{array}{cc}\omega\mathbf{1}_2 - F\Box & W''\\ \bar{W}''\Box & \omega\mathbf{1}_2 - F\Box\end{array}\right) = 0\tag{2.43}
$$

In principle, parameters ω can be found from this equation. Their exact form is determined by the structure of the matrix W'' and F. It turns out that for the specific cases studied in detail the subsequent sections (minimal (Section 3) and non-minimal (Section 4) cases) these parameters are different. As a result the final solution can be cast in the form:

$$
A = \sum_{k} a_{0k} \exp(i\omega_k s) - \frac{1}{\Box} \tag{2.44}
$$

$$
C = \sum_{k} c_{0k} \exp(i\omega_k s) \tag{2.45}
$$

where ω_k are all the roots of the equation (2.43) and a_{0k} , c_{0k} are values of a_0 , c_0 which can be found from (2.42) for the corresponding root ω_k . a_{0k} , c_{0k} are fixed by means of initial conditions.

Sinceonly the coefficients A and A in ([2.35](#page-10-0)) contribute to the trace $Tr \log \Delta$ the one-loop kähleriancontribution to the effective action (2.32) (2.32) (2.32) can be written in the form

$$
K^{(1)} = -\frac{i}{2} \int_0^\infty \frac{ds}{s} tr(A + \tilde{A}) U(x, x'; s)|_{x = x'}
$$
 (2.46)

where $U(x, x'; s) = \exp(is\Box)\delta^4(x - x')$ (see [[22, 23](#page-56-0)]) and tr denotes a trace of the matrix. The operator $U(x, x'; s)$ satisfies the equation

$$
\Box U = \frac{1}{i} \frac{\partial}{\partial s} U \tag{2.47}
$$

and $U(x, x'; s)|_{x=x'} = \frac{1}{16\pi^2(1-x')}$ $\frac{1}{16\pi^2(i s)^2}$. A, and \tilde{A} are functions of \Box . Hence in order to calculate the one-loop kählerian effective potential it is necessary to find A and \tilde{A} and to expand them into a power series in \Box .

In this section we addressed in detail the techniques needed to calculate the one-loop corrected effective action, to eliminate the heavy fields and to obtain the effective action of light fields only. In the subsequent sections these techniques will be applied to obtain the explicit form of the one-loop corrected actions for specific models. In Section 5 we shall also include interactions with the $U(1)$ vector superfields and modify the procedure accordingly.

3 One-loop effective action for minimal models

In this section we study decoupling effects for the model with minimal kählerian potential [\(2.2](#page-4-0)) $K = \Phi \Phi + \phi \phi$. The first part consists of calculating the one-loop correction to the kählerian effective potential. In the second part we solve the effective equations of motion for the heavy superfields. As a result we arrive at the effective action of the light superfields.

3.1 Calculation of the effective action

3.1.1 The one-loop kählerian effective potential

Here we are going to calculate the one-loop contribution to kählerian effective potential by means of the effective equations of motion. We study the minimal model with the chiral potential in the form

$$
W = \frac{1}{2}(M\Phi^2 + \lambda\Phi\phi^2) + \frac{1}{3!}g\phi^3
$$
\n(3.1)

with the corresponding functions in W'' (see eq.[\(2.25](#page-8-0)))

$$
W_{\phi\phi} = \lambda \Phi + g\phi
$$

\n
$$
W_{\phi\Phi} = \lambda \phi
$$

\n
$$
W_{\Phi\Phi} = M
$$
\n(3.2)

The total classical action with the chiral potential (3.1) is of the form

$$
S = \int d^8 z (\phi \bar{\phi} + \Phi \bar{\Phi}) + [\int d^6 z (\frac{1}{2} (M \Phi^2 + \lambda \Phi \phi^2) + \frac{1}{3!} g \phi^3) + h.c.] \tag{3.3}
$$

Note that the chiral potential used is of the "minimal" form: it involves the renormalizable terms only and the renormalizable coupling between the light and heavy superfields is *linear* in the heavy fields, which yields a dominant contribution in the study of the decoupling effects. These types of couplings are typical for a class of effective string models after the vacuum restabilization was taken into account, and thus this minimal model provides a prototype example for the study of decoupling effects in $N = 1$ supersymmetric theories. (The results for this model and the physics consequences were presented in [\[18\]](#page-56-0). For the sake of completeness we present here the intermediate steps in the derivation.)

In order to find the one-loop kählerian effective potential we should determine the operator $\Omega(s)$ that satisfies the equation [\(2.35](#page-10-0)). For the case of this minimal model this equation leads to the following system of equations for matrices A, C :

$$
\frac{1}{i}\dot{A} = -CW''
$$
\n
$$
\frac{1}{i}\dot{C} = -\overline{W}'' - A\overline{W}'' \square
$$
\n(3.4)

with the analogous equations for \tilde{A}, \tilde{C} . Initial conditions are $A|_{s=0} = \tilde{A}|_{s=0} = C|_{s=0}$ $\tilde{C}|_{s=0} = 0$. Calculations described in Appendix A show that the one-loop contribution to kählerian effective potential is of the form:

$$
K^{(1)} = -\frac{1}{32\pi^2} \Big((|\lambda \Phi + g\phi|^2 + 2\lambda^2 \phi \bar{\phi} + M^2 +
$$

+ $\sqrt{(|\lambda \Phi + g\phi|^2 - M^2)^2 + 4|\lambda^2 \Phi \bar{\phi} + \lambda M\phi + \lambda g|\phi|^2)|^2} \Big) \times$
 $\times \log \frac{(|\lambda \Phi + g\phi|^2 + 2\lambda^2 \phi \bar{\phi} + M^2 + \sqrt{(|\lambda \Phi + g\phi|^2 - M^2)^2 + 4|\lambda^2 \Phi \bar{\phi} + \lambda M\phi + \lambda g|\phi|^2|^2})}{\mu^2} +$
+ $(|\lambda \Phi + g\phi|^2 + 2\lambda^2 \phi \bar{\phi} + M^2 - \sqrt{(|\lambda \Phi + g\phi|^2 - M^2)^2 + 4|\lambda^2 \Phi \bar{\phi} + \lambda M\phi + \lambda g|\phi|^2|^2}) \times$
 $\times \log \frac{(|\lambda \Phi + g\phi|^2 + 2\lambda^2 \phi \bar{\phi} + M^2 - \sqrt{(|\lambda \Phi + g\phi|^2 - M^2)^2 + 4|\lambda^2 \Phi \bar{\phi} + \lambda M\phi + \lambda g|\phi|^2|^2})}{\mu^2} \Big)$ (3.5)

and the effective action in the one-loop approximation is of the form

$$
\Gamma^{(1)} = \int d^8 z (\phi \bar{\phi} + \Phi \bar{\Phi}) + (\int d^6 z (\frac{1}{2} M \Phi^2 + \Phi \phi^2) + h.c.) - \frac{1}{32\pi^2} ((|\lambda \Phi + g\phi|^2 + 2\lambda^2 \phi \bar{\phi} + M^2 + \sqrt{(|\lambda \Phi + g\phi|^2 - M^2)^2 + 4|\lambda^2 \Phi \bar{\phi} + \lambda M \phi + \lambda g |\phi|^2)|^2}) \times
$$

$$
\times \quad \log \frac{(|\lambda \Phi + g\phi|^2 + 2\lambda^2 \phi \bar{\phi} + M^2 + \sqrt{(|\lambda \Phi + g\phi|^2 - M^2)^2 + 4|\lambda^2 \Phi \bar{\phi} + \lambda M \phi + \lambda g |\phi|^2 |^2})}{\mu^2} + (|\lambda \Phi + g\phi|^2 + 2\lambda^2 \phi \bar{\phi} + M^2 - \sqrt{(|\lambda \Phi + g\phi|^2 - M^2)^2 + 4|\lambda^2 \Phi \bar{\phi} + \lambda M \phi + \lambda g |\phi|^2 |^2}) \times
$$

$$
\times \quad \log \frac{(|\lambda \Phi + g\phi|^2 + 2\lambda^2 \phi \bar{\phi} + M^2 - \sqrt{(|\lambda \Phi + g\phi|^2 - M^2)^2 + 4|\lambda^2 \Phi \bar{\phi} + \lambda M \phi + \lambda g |\phi|^2 |^2})}{\mu^2})
$$

Here parameter μ is is the renormalization scale parameter (see Appendix A). This is the focal result of this section and is further used for the study of decoupling effects. Note also that the same result can be obtained via a diagrammatic approach, which is described in Appendix B.

3.1.2 Corrections to the chiral potential

Let us now address loop corrections to the chiral effective potential. By definition the chiral effective potential depends only on chiral background superfields, i.e. in order to consider corrections to the chiral effective potential one must set $\bar{\Phi} = \bar{\phi} = 0$. Possible vertices contributing to one-loop effective potential should be quadratic in quantum superfields[[21\]](#page-56-0). They have the form

$$
K_{\bar{\phi}\bar{\phi}}\bar{\phi}^2, K_{\phi\phi}\phi^2, (K_{\phi\bar{\phi}} - 1)\phi\bar{\phi}, \frac{1}{2}W_{\phi\phi}\phi^2,
$$

$$
K_{\bar{\Phi}\bar{\phi}}\bar{\phi}\bar{\Phi}, K_{\Phi\phi}\Phi\phi, (K_{\Phi\bar{\phi}})\Phi\bar{\phi}, \frac{1}{2}W_{\Phi\phi}\Phi\phi,
$$

$$
K_{\bar{\Phi}\bar{\Phi}}\bar{\Phi}^2, K_{\Phi\Phi}\Phi^2, (K_{\Phi\bar{\Phi}} - 1)\Phi\bar{\Phi}, \frac{1}{2}W_{\Phi\Phi}\Phi^2
$$

Here all derivatives of K and W are taken at $\bar{\Phi} = \bar{\phi} = 0$. The standard method to address these corrections ([\[24\]](#page-56-0)–[\[26](#page-56-0)]) is the following. Corrections contributing to the chiral potential should be of the form:

$$
\int d^8 z f(\Phi) \left(-\frac{D^2}{4\Box} \right) g(\Phi) \tag{3.6}
$$

Namely, after a transformation to an integral over the chiral superspace by the rule $\int d^8 z \tilde{F}(\Phi, \bar{\Phi}) = \int d^6 z \left(-\frac{\bar{D}^2}{4} \right)$ $\frac{\bar{D}^2}{4}$) $F(\Phi, \bar{\Phi})$ these corrections take the form

$$
\int d^6z f(\Phi)g(\Phi) \tag{3.7}
$$

Here $f(\Phi)$, $q(\Phi)$ are functions of the chiral superfield Φ , only.

However, factor \square^{-1} in [\(3.6](#page-13-0)) can arise only from propagators of massless superfields as propagators of massive superfields are proportional to $(\square - m^2)^{-1}$. Namely, in the case of the massive chiral superfields, instead of [\(3.6](#page-13-0)) we must consider the expression

$$
\int d^8 z f(\Phi) \left(- \frac{D^2}{4(\Box - m^2)} \right) g(\Phi) \tag{3.8}
$$

A transformation to the form of an integral over the chiral superspace leads to

$$
\int d^6 z f(\Phi) \left(\frac{\Box}{\Box - m^2}\right) g(\Phi) \tag{3.9}
$$

which after Fourier transformation is of the form

$$
\int \frac{d^4 p}{(2\pi)^4} f(\Phi) \left(\frac{p^2}{p^2 + m^2}\right) g(\Phi) \tag{3.10}
$$

This expression evidently vanishes for superfields slowly varying in space-time, $(p^2 \to 0)$ and $m \neq 0$. Hence supergraphs contributing to the chiral effective potential cannot contain massive propagators.

Let us now carry out a dimensional analysis of possible one-loop diagrams contributing to the chiral effective potential. Each loop corresponds to a contribution with a scaling dimension 2 since each loop includes an integration over momenta which contribute a scaling dimension 4, however each contraction of the loop to the point by the rule $\delta_{12}D_1^2\bar{D}_1^2\delta_{12}=16\delta_{12}$

decreases a number of D , \overline{D} -factors by 4 and the corresponding scaling dimension by 2. Each propagator of a massless superfield gives no contribution (scaling dimension 0) since it has the form (cf. [\[20](#page-56-0)])

$$
G(z_1, z_2) = -\frac{D_1^2 \bar{D}_2^2}{16\Box} \delta^8(z_1 - z_2)
$$
\n(3.11)

(It is more convenient for dimensional analysis to associate D-factors with propagators instead of vertices.) Each vertex proportional to an external chiral superfield corresponds to one factor \bar{D}^2 instead of two since one of them is used to transform a vertex to a form of an integral over $d^4\theta$. As a result such a vertex decreases a scaling dimension by 1. One of the factors $D²$ does not contribute to the scaling dimension since it is used to transform the expression([3.6\)](#page-13-0) to the form [\(3.7](#page-14-0)). Then, during such a transformation one more square of external momenta arises. Therefore the scaling dimension due to contributions from a one-loop supergraph has the form $2L + 1 - n_{W''}$, where L is the number of loops (in our case $L = 1$, and $n_{W''}$ is the number of external chiral lines W''. By definition the effective potential is an effective Lagrangian in the low energy (infrared) limit $p^2 \to 0$. Hence a non-trivial one-loop chiral effective Lagrangian can arise only for

$$
3 - n_{W''} = 0 \tag{3.12}
$$

Otherwise the contribution would either vanish or be singular in the infrared limit. We also note that the vertices proportional to derivatives of $K(\Phi, \bar{\Phi})$ do not contribute to the scaling dimension.

Furthermore, chiral potential contributions should contain one factor D^2 more than \bar{D}^2 after a transformation of all vertices to the form of integrals over $d^4\theta$. The reason for this is the following. After each pair $\bar{D}^2 D^2$ is transformed to a d'Alembertian operator, one factor D^2 should be employed to transform the integral over $d^4\theta$ into an integral over $d^2\theta$ (see $(3.6)-(3.7)$ $(3.6)-(3.7)$ $(3.6)-(3.7)$ $(3.6)-(3.7)$.

It is easy to see that one of two factors \bar{D}^2 associated with the vertex proportional to $K_{\Phi\Phi}$ can be transported only to an external line $K_{\Phi\Phi}$ during the "D-algebra" transformations. However, all the derivatives of K and W are considered at $\bar{\Phi} = 0$, and therefore $\bar{D}^2 K_{\Phi \Phi} = 0$. As a result we can put $K_{\Phi\Phi} = 0$.

Now let us consider the possible one-loop diagrams contributing to the chiral effective potential. At the one-loop level $n_{W''} = 3$ (see above) and each external line W'' corresponds to \bar{D}^2 . Hence the supergraph should contain four \bar{D}^2 -factors, i.e. two vertices proportional to $K_{\bar{\Phi}\bar{\Phi}}$. However, a straightforward construction shows that such a diagram must contain the line proportional to $\frac{\bar{D}_1^2 \bar{D}_2^2}{16} \delta_{12} = 0$. Hence a contribution of such a diagram is equal to zero, and a one-loop contribution to the chiral effective potential is absent: $W^{(1)}(\Phi) = 0$. We note that this situation is analogous to the general model of one chiral superfield studied in[[27\]](#page-56-0).

However, higher order (loop) corrections to the chiral effective potential can arise not only for diagrams with external massless lines but also for those with heavy external lines, in spite of the fact that it was commonly believed that quantum corrections to the chiral effective potential for massive superfields are absent. For example, consider the supergraph

Here a double line denotes the external superfield Φ , and a single line corresponds to the propagator $\langle \phi \phi \rangle$ of the massless superfield ϕ . A contribution of such a supergraph is of the form

$$
I = \int \frac{d^4 p_1 d^4 p_2}{(2\pi)^8} \frac{d^4 k d^4 l}{(2\pi)^8} \int d^4 \theta_1 d^4 \theta_2 d^4 \theta_3 d^4 \theta_4 d^4 \theta_5 \left(\frac{g}{3!}\right)^2 \lambda^3 \Phi(-p_1, \theta_3) \Phi(-p_2, \theta_4) \Phi(p_1 + p_2, \theta_5) \times
$$

\n
$$
\times \frac{1}{k^2 l^2 (k+p_1)^2 (l+p_2)^2 (l+k)^2 (l+k+p_1+p_2)^2} \times
$$

\n
$$
\times \frac{\bar{D}_3^2}{4} \delta_{32} \frac{D_1^2 \bar{D}_4^2}{16} \delta_{14} \delta_{42} \frac{D_1^2 \bar{D}_5^2}{16} \delta_{15} \delta_{52}
$$
\n(3.13)

After a tedious calculation analogous to that carried out in[[27\]](#page-56-0) we arrive at the expression:

$$
I = \left(\frac{g}{3!}\right)^2 \lambda^3 \int \frac{d^4 p_1 d^4 p_2}{(2\pi)^8} \frac{d^4 k d^4 l}{(2\pi)^8} \int d^2 \theta \Phi(-p_1, \theta) \Phi(-p_2, \theta) \Phi(p_1 + p_2, \theta) \times \times \frac{k^2 p_1^2 + l^2 p_2^2 + 2(kl)(p_1 p_2)}{k^2 l^2 (k + p_1)^2 (l + p_2)^2 (l + k)^2 (l + k + p_1 + p_2)^2}
$$
\n(3.14)

which can be cast in the form

$$
I = \left(\frac{g}{3!}\right)^2 \lambda^3 \int d^2\theta \int \frac{d^4p_1 d^4p_2}{(2\pi)^8} \bar{W}'''^2 \Phi(-p_1, \theta) \Phi(-p_2, \theta) \Phi(p_1 + p_2, \theta) S(p_1, p_2) \quad (3.15)
$$

Here p_1, p_2 are external momenta. The expression for $S(p_1, p_2)$ is of the form

$$
\int \frac{d^4k d^4l}{\left(2\pi\right)^8} \frac{k^2 p_1^2 + l^2 p_2^2 + 2(kl)(p_1 p_2)}{k^2 l^2 (k+p_1)^2 (l+p_2)^2 (l+k)^2 (l+k+p_1+p_2)^2}
$$

After Fourier transformation, eq. (3.15) has the form

$$
I = \left(\frac{g}{3!}\right)^2 \lambda^3 \int d^2\theta \int d^4x_1 d^4x_2 d^4x_3 \int \frac{d^4p_1 d^4p_2}{(2\pi)^8} \Phi(x_1, \theta) \Phi(x_2, \theta) \Phi(x_3, \theta) \times
$$

$$
\times \exp[i(-p_1x_1 - p_2x_2 + (p_1 + p_2)x_3)] S(p_1, p_2)
$$
(3.16)

The effective potential is obtained from the above expression by taking superfields that slowly vary in space-time, i.e. we take $\Phi(x_1, \theta) \Phi(x_2, \theta) \Phi(x_3, \theta) \simeq \Phi^3(x_1, \theta)$. As a result one obtains

$$
I = \left(\frac{g}{3!}\right)^2 \lambda^3 \int d^2\theta \int d^4x_1 d^4x_2 d^4x_3 \int \frac{d^4p_1 d^4p_2}{(2\pi)^8} \Phi^3(x_1, \theta) \times
$$

$$
\times \exp[i(-p_1x_1 - p_2x_2 + (p_1 + p_2)x_3)]S(p_1, p_2)
$$
(3.17)

Theintegration over $d^4x_2d^4x_3$ leads to delta-functions $\delta(p_2)\delta(p_1+p_2)$ and hence the eq. ([3.16\)](#page-16-0) takes the form

$$
I = \left(\frac{g}{3!}\right)^2 \lambda^3 \int d^2\theta \int d^4x_1 \Phi^3(x_1, \theta) S(p_1, p_2)|_{p_1, p_2 = 0}
$$
\n(3.18)

Therefore the result for the two-loop contribution to the chiral effective potential due to this diagram is of the form

$$
\frac{6}{(16\pi^2)^2}\zeta(3)(\frac{g}{3!})^2\lambda^3\Phi^3\tag{3.19}
$$

As the last step we took into account that

$$
\int \frac{d^4k d^4l}{(2\pi)^8} \frac{k^2 p_1^2 + l^2 p_2^2 + 2(k_1 k_2)(p_1 p_2)}{k^2 l^2 (k+p_1)^2 (l+p_2)^2 (l+k)^2 (l+k+p_1+p_2)^2} |_{p_1=p_2=0} = \frac{6}{(4\pi)^4} \zeta(3)
$$

We have therefore demonstrated that chiral corrections to the effective action that depend on massive superfields can also arise. However, such effects can appear at the two-loop level at most, and after solving the effective equations of motion for heavy fields such corrections are proportional to $\frac{1}{M^3}$. Therefore, they are highly suppressed compared to the decoupling effects due to the one-loop corrected kählerian effective potential. In the following we shall concentrate on the latter effects only.

By analogous reasoning, corrections to the chiral effective potential that involve a coupling of the heavy superfield to the light ones, are of the form $\frac{6}{(16\pi^2)^2}\zeta(3)(\frac{g}{3!})^2\lambda^3\Phi\phi^2$ which is of the same form as the corresponding classical chiral effective potential term. They are described by supergraphs analogous to those given above, but with one heavy external line and two massless ones. At the two-loop level $K_{eff} = K + K^{(1)} + K^{(2)}$, and $W_{eff} = W + W^{(2)}$ where $K^{(2)}$ and $W^{(2)}$ are two-loop corrections to the kählerian potential and chiral potential, respectively. $W^{(2)}$ depends non-trivially on Φ . Hence the presence of such a correction to W should modify the effective equations of motion([2.22,2.23\)](#page-8-0). After substitution of the solution of effective equations of motion into the effective action, $W^{(2)}$ turns out to be suppressed by 1 $\frac{1}{M}$, i.e. it can have the form ~ $\frac{\phi^4}{M}$ $\frac{\phi^*}{M}$. Hence the effective dynamics of the two-loop corrected theory allow us to conclude that a two-loop contribution to the effective superpotential can yield corrections of order $\frac{1}{M}$, i.e. it can be competitive with the leading tree-level (classical) decoupling effects.

3.2 The effective action for light superfields

Now we solve the effective equations of motion for this minimal model that allows us to eliminate heavy superfields from effective action up to some order in the inverse powers of the mass parameters, and as a consequence obtain the one-loop effective action of light superfields only. The one-loop corrected effective action is given by ([3.6\)](#page-13-0). The effective equations of motion for the heavy superfield with the chiral potential [\(3.1](#page-12-0)) have the form

$$
\frac{\delta \Gamma^{(1)}}{\delta \Phi} = 0: \quad -\frac{1}{4} \bar{D}^2 (\bar{\Phi} + \frac{\partial K^{(1)}}{\partial \Phi}) + M\Phi + \frac{\lambda}{2} \phi^2 = 0
$$

$$
\frac{\delta \Gamma^{(1)}}{\delta \bar{\Phi}} = 0: \quad -\frac{1}{4} D^2 (\Phi + \frac{\partial K^{(1)}}{\partial \bar{\Phi}}) + M\bar{\Phi} + \frac{\lambda}{2} \bar{\phi}^2 = 0
$$
(3.20)

As described in the previous Section these equations can be solved for the heavy superfield via an iterative method and a subsequent substitution of this solution into the one-loop corrected effective action. (Analogous equations for ϕ are not essential since our aim is to minimize the effective action with respect to heavy superfields.)

Now let us solve the equations (3.20) via the iterative method. Using $|D^2\Phi| \ll M\Phi$, since we consider the case with large $M \gg p$, one finds that the lowest (zeroth) order solution of these equations is

$$
\Phi_0 = -\lambda \frac{\phi^2}{2M} \tag{3.21}
$$

and with the corresponding value for $\bar{\Phi}_0$.

Thefirst order correction (cf. $(2.22)-(2.23)$ $(2.22)-(2.23)$ $(2.22)-(2.23)$ $(2.22)-(2.23)$) is obtained from the equation

$$
\Phi_1 = \frac{1}{4M}\bar{D}^2(\bar{\Phi}_0 + \frac{\partial K^{(1)}}{\partial \Phi}|_{\Phi = \Phi_0})\tag{3.22}
$$

where Φ_0 is given by (3.21). As for higher order corrections, they can be found by continuing the iterative procedure.

It turns out to be that for the case with $g \neq 0$, Φ_1 is of order $\frac{1}{M}$ (as Φ_0), and it is of the form

$$
\Phi_1 = -\frac{\bar{D}^2}{64\pi^2 M} \left(\lambda g \bar{\phi} (1 + \log \frac{2g^2 |\phi|^2}{\mu^2}) \right) + O(\frac{1}{M^2})\tag{3.23}
$$

On the other hand for the case with $g = 0$ (i.e. the self-interactions of the light superfield in the classical chiral potential are absent), Φ_1 is of the order $\frac{1}{M^2}$ and takes the form

$$
\Phi_1 = \frac{\lambda \bar{D}^2(\bar{\phi}^2)}{8M^2} + \frac{1}{32\pi^2} \frac{\lambda^3}{2M^2} \bar{D}^2 \left\{ |\phi|^2 (1 + \log \frac{2M^2}{\mu^2}) + \right. \\
\left. + \left. (\frac{\bar{\phi}^2}{2} + |\phi|^2)(1 + \log[5\lambda^4 \frac{|\phi|^4}{M^4} - \lambda^4 \frac{|\phi|^2 (\phi^2 + \bar{\phi}^2)}{2M^4})] \right) \right\} + O(\frac{1}{M^3})
$$

As for the second order correction Φ_2 , it is proportional to $\frac{1}{M^2}$ for $g \neq 0$ and to $\frac{1}{M^3}$ for $g = 0$. Note, the cases with $g = 0$ and $g \neq 0$ lead to essentially different expressions for the heavy superfield Φ in terms of the light one ϕ .

For the calculation of the effective action for the light superfield it is convenient to expand $K^{(1)}$ [\(3.5](#page-13-0)) into a power series in $\frac{1}{M}$ (up to the order $\frac{1}{M^3}$):

$$
K^{(1)} \;\; = \;\; -\frac{1}{32\pi^2} \Big\{ (4\lambda^2|\phi|^2 + 2\frac{\lambda^2}{M}(\lambda\Phi + g\phi + h.c.)|\phi|^2 + \frac{|\phi|^4}{M^2}(g^4 + 2\lambda^2g^2)) (1 + \log\frac{2M^2}{\mu^2}) + \right.
$$

+
$$
2\lambda^4 \frac{|\phi|^4}{M^2} (1 - \log \frac{2M^2}{\mu^2})
$$
 +
\n+ $[2|\lambda \Phi + g\phi|^2 - 2\frac{\lambda^2}{M} (\lambda \Phi + g\phi)|\phi|^2 + \frac{|\phi|^4}{M^2} (2\lambda^4 - g^4 - 2\lambda^2 g^2)] \times$
\n $\times \left(\log \left\{ \frac{2|\lambda \Phi + g\phi|^2}{\mu^2} - \frac{2\lambda^2}{M\mu^2} (\lambda \Phi + g\phi + h.c.)|\phi|^2 + \frac{|\phi|^4}{M^2\mu^2} (2\lambda^4 - g^4 - 2\lambda^2 g^2) \right\} \right\} + O(\frac{1}{M^3})$ (3.24)

Here we took into account that Φ is suppressed by at least one inverse power of M. In the following we shall substitute the solutions of effective equations of motion into the expression [\(3.24\)](#page-18-0).

3.2.1 Contribution of the self-interaction of the light superfield

First we consider the case $g \neq 0$. In this case we can write $\Phi = \Phi_0 + \Phi_1$ in the form (cf. [\(3.23\)](#page-18-0))

$$
\Phi = -\lambda \frac{\phi^2}{2M} - \frac{\bar{D}^2}{64\pi^2 M} \left(\lambda g \bar{\phi} (1 + \log \frac{2g^2 |\phi|^2}{\mu^2}) \right) + O(\frac{1}{M^2})
$$
\n(3.25)

Its substitution into [\(3.24](#page-18-0)) leads to the expression

$$
K^{(1)} = -\frac{1}{32\pi^2} 2g^2 |\phi|^2 \left\{ \log \frac{g^2 |\phi|^2}{\mu^2} + \frac{\lambda}{g |\phi|^2} (\bar{\phi}(-\frac{\lambda \phi^2}{2M} + \lambda g \frac{\bar{D}^2}{64\pi^2 M} [\bar{\phi}(1 + \mu + \log \frac{g^2 |\phi|^2}{\mu^2})] + h.c.) \right\} - \frac{\lambda^2 \phi + \bar{\phi}}{M} \frac{\phi}{g} - \frac{2\lambda^2 g(\phi + \bar{\phi}) |\phi|^2 \log \frac{g^2 |\phi|^2}{\mu^2} + O(\frac{1}{M^2})} \tag{3.26}
$$

Here a factor $e = \exp(1)$ is introduced to cancel superfluous factor -1 .

As for the chiral potential which has the form([3.1\)](#page-12-0) it can be written as (cf. (3.25)):

$$
V = \frac{g}{3!} \phi^3 - \frac{1}{8} \frac{\lambda^2 \phi^4}{M} + \frac{\lambda^2 g^2}{2M} \left\{ \frac{\bar{D}^2}{64\pi^2} (\bar{\phi} (1 + \log \frac{2g^2 |\phi|^2}{\mu^2})) \right\}^2 + O(\frac{1}{M^2}) \tag{3.27}
$$

The effective action for light superfields can now be written up to the order $\frac{1}{M^2}$ in the form

$$
S_{eff}[\phi,\bar{\phi}] = S_{eff}^{cl}[\phi,\bar{\phi}] + S_{eff}^{q}[\phi,\bar{\phi}]
$$
\n(3.28)

where $S_{eff}^{cl}[\phi, \bar{\phi}]$ is the classical part:

$$
S_{eff}^{cl}[\phi,\bar{\phi}] = \int d^8 z \phi \bar{\phi} + \left\{ \int d^6 z \left(\frac{g}{3!} \phi^3 - \frac{1}{8} \frac{\lambda^2 \phi^4}{M} + \frac{\lambda^2 g^2}{2M} \left\{ \frac{\bar{D}^2}{64\pi^2} (\bar{\phi} (1 + \log \frac{2g^2 |\phi|^2}{\mu^2})) \right\}^2 \right) +
$$

+ h.c. + O(\frac{1}{M^2}) \tag{3.29}

and $S^q_{eff}[\phi, \bar{\phi}]$ is the quantum part:

$$
S_{eff}^{q}[\phi,\bar{\phi}] = -\frac{1}{32\pi^{2}} \int d^{8}z \left[2g^{2}|\phi|^{2}\left\{\log\frac{g^{2}|\phi|^{2}}{\mu^{2}} + \frac{\lambda}{g|\phi|^{2}}(\bar{\phi}(-\frac{\lambda\phi^{2}}{2M} + \lambda g\frac{\bar{D}^{2}}{64\pi^{2}M}[\bar{\phi}(1+\log\frac{g^{2}|\phi|^{2}}{\mu^{2}})] + h.c.)\right) - \frac{\lambda^{2}}{M}\frac{\phi+\bar{\phi}}{g}\right\} - \frac{2\frac{\lambda^{2}}{M}g(\phi+\bar{\phi})|\phi|^{2}\log\frac{g^{2}|\phi|^{2}}{\mu^{2}} + O(\frac{1}{M^{2}})
$$
\n(3.30)

The main feature of the expressions $(3.28)-(3.30)$ is the fact that with increasing M the leading contribution of the decoupling effects grow logarithmically with M. Let us fix μ by imposing the condition $\log \frac{2M^2}{\mu^2} = -1$. This renormalization condition eliminates from the one-loop effective action the terms that are proportional to λ , i.e., the coupling of the heavy and light superfields in the classical action. As a consequence, S_d^q in eq. (3.30) is of the form:

$$
-\frac{1}{32\pi^2}2g^2|\phi|^2\log\frac{g^2|\phi|^2}{eM^2}
$$

This term *increases* as $M \to \infty$ and becomes leading in comparison with the classical con-tribution [\(3.29](#page-19-0)) which in the limit $M \to \infty$ is of the form:

$$
S_{eff}^{cl}[\phi,\bar{\phi}] = \int d^8z \phi \bar{\phi} + \left\{ \int d^6z \frac{g}{3!} \phi^3 + h.c. \right\} + O(\frac{1}{M})
$$

Alternatively, we do not fix μ and address the effective action [\(3.28](#page-19-0))-(3.30) at energy scales specified by the values of μ . In the leading (zeroth) order in $\frac{1}{M}$ expansion the quantum effective action (3.30) in the form

$$
S_{eff}^{q}[\phi,\bar{\phi}] = -\frac{1}{32\pi^{2}} \Big\{ 4\lambda^{2}|\phi|^{2} (1+\log\frac{2M^{2}}{\mu^{2}}) + 2|g\phi|^{2} \Big(\log\frac{2|g\phi|^{2}}{\mu^{2}} \Big) \Big\} + O(\frac{1}{M}) \quad (3.31)
$$

As μ is taken as a free parameter, describing the effective theory at energy scales μ , we see that the second term of this expression (proportional to $\log \frac{2M^2}{\mu^2}$) can become competitive with those of the classical action as $\mu \ll M \to \infty$. Therefore from both perspectives, i.e. when we fix μ and when we treat μ as a free parameter, the decoupling effects grow logarithmically with M and as $M \to \infty$ they become dominant. Hence the appearance of such terms is not an effect caused by a choice of normalization parameter μ but it is a genuine property of the model itself.

The total one-loop effective action can be finally written as

$$
S_{eff}[\phi,\bar{\phi}] = \int d^8 z (|\phi|^2 - \frac{\hbar}{32\pi^2} \{ 4\lambda^2 |\phi|^2 (1 + \log \frac{2M^2}{\mu^2}) + 2|g\phi|^2 \Big(\log \frac{2|g\phi|^2}{\mu^2} \Big) \}) +
$$

+
$$
[\frac{1}{3!} \int d^6 z g \phi^3 + h.c.] + O(\frac{1}{M})
$$
(3.32)

Here we took into account that all the terms that depend on the heavy superfield Φ , after the substitution of the solution of the effective equations of motion, are at least of the first order in the inverse mass M.

According to the decoupling theorem when we eliminate heavy superfields via effective equations of motion we can split any quantum correction to the effective action into a sum of two terms. The first term is a quantum correction in the corresponding theory describing dynamics of light superfields only, with heavy superfields put to zero, i.e. the standard Coleman-Weinberg potential associated with the interaction of the light fields only. The second one corresponds to a sum of terms that are proportional to at least one power of $\frac{1}{M}$.

To convince ourselves that our results are in accordance with the decoupling theorem, we introduce the following field and coupling redefinitions. First, we see that the kinetic energy term (proportional to $|\phi|^2$) of the one-loop effective action is of the form

$$
|\phi|^2 [1 - \frac{\hbar}{32\pi^2} \Big\{ 4\lambda^2 (1 + \log \frac{2M^2}{\mu^2}) \Big\}]
$$

We can redefine the kinetic energy term by the rule

$$
\tilde{\phi} = Z\phi \tag{3.33}
$$

where $\tilde{\phi}$ is a new chiral superfield, and Z is a finite renormalization constant equal to

$$
Z = \left[1 - \frac{\hbar}{32\pi^2} 4\lambda^2 (1 + \log \frac{2M^2}{\mu^2})\right] \tag{3.34}
$$

Then, if we also redefine coupling g by the rule $\tilde{g} = Z^{-3}g$ we see that the chiral potential $g\phi^3$ remains invariant. We note that the coupling λ corresponds to the interaction between light and heavy superfields, and after a redefinition of the light superfield (3.33) it will present itself only in terms that are at least first order in the inverse mass M. (Note also that analogous redefinitions of the superfield and couplings could be carried out at higher orders of the effective action.) As a result we arrive at the one-loop effective action of the form

$$
S_{eff}[\phi,\bar{\phi}] = \int d^8 z \{ |\tilde{\phi}|^2 - \frac{\hbar}{32\pi^2} 2Z^4 \tilde{g}^2 |\tilde{\phi}|^2 (\log \frac{2Z^4 \tilde{g}^2 |\tilde{\phi}|^2}{\mu^2}) \} + \left[\int d^6 z \frac{1}{3!} \tilde{g} \tilde{\phi}^3 + h.c. \right] + O(\frac{1}{M})
$$
\n(3.35)

However, we can expand Z (3.34) into a power series in \hbar as

$$
Z = 1 - \frac{\hbar}{64\pi^2} 4\lambda^2 (1 + \log \frac{2M^2}{\mu^2}) + O(\hbar^2)
$$

Substituting this expansion into (3.35) and taking into account only terms of zeroth and first order in \hbar we arrive at

$$
S_{eff}[\phi,\bar{\phi}] = \int d^8 z \{ |\tilde{\phi}|^2 - \frac{\hbar}{32\pi^2} 2\tilde{g}^2 |\tilde{\phi}|^2 (\log \frac{2\tilde{g}^2 |\tilde{\phi}|^2}{\mu^2}) \} + + [\int d^6 z \frac{1}{3!} \tilde{g} \tilde{\phi}^3 + h.c.] + O(\frac{1}{M})
$$
(3.36)

This effective action coincides at the zeroth order with the form of the one-loop effective action for the "pure light" theory where quantum corrections originate from the couplings of the light fields only. Therefore the result for the effective action of light superfields given by expressions([3.28–](#page-19-0)[3.30\)](#page-20-0) is in formal agreement with the decoupling theorem.

Note, however, that the parameters of the theory, i.e. fields, masses and couplings, are determined from the string theory; they are fixed-calculable at M_{string} and therefore they cannot be adjusted via redefinitions performed above. Thus, the corrections due to decoupling effects correct in a quantitative manner the predictions for the low-energy values of the couplings. For a class of perturbative string vacua with an anomalous $U(1)$ after the vacuum restabilization, the effective theory has couplings of the heavy and light fields of the type discussed in this section. Typical values of the couplings g and λ (at $M_{string} \sim 10^{17}$ GeV) are of the order of the gauge coupling ~ 0.8 and the mass parameter of the heavy fields $M \sim M_{string}$. The decoupling effects then modify the low-energy ($\mu \sim 1 \text{ TeV}$) predictions for the corresponding tri-linear couplings of the light fields by an amount of order $\frac{g^2}{16\pi^2} \log(\frac{M^2}{\mu^2}) \sim$ 0.25. corrections, and Thus for such classes of string vacua decoupling effects could correct the predictions for the couplings by $10\% - 50\%$.

3.2.2 Absence of the self-interaction of the light superfield

Nowlet us put $g = 0$ in ([3.1\)](#page-12-0), i.e. a self-interaction of light superfields is absent. In this case Φ_1 is of the order $\frac{1}{M^2}$ and $\Phi = \Phi_0 + \Phi_1$ takes the form

$$
\begin{array}{lll} \Phi&=&-\lambda\displaystyle\frac{\phi^2}{2M}+\displaystyle\frac{\lambda \bar{D}^2(\bar{\phi}^2)}{8M^2}+\\&+&\displaystyle\frac{1}{32\pi^2}\displaystyle\frac{\lambda^3}{2M^2}\bar{D}^2\Big\{(\frac{\bar{\phi}^2}{2}+|\phi|^2)(2+\log[5\lambda^4\displaystyle\frac{|\phi|^4}{M^2\mu^2}-\lambda^4\displaystyle\frac{|\phi|^2(\phi^2+\bar{\phi}^2)}{M^2\mu^2}])\Big\}+O(\displaystyle\frac{1}{M^3}) \end{array}
$$

The one-loop kählerian effective potential is of the form

$$
K^{(1)} = -\frac{1}{32\pi^2} \int d^8 z \{ 4\lambda^2 |\phi|^2 (1 + \log \frac{2M^2}{\mu^2}) + 4\lambda^4 \frac{|\phi|^4}{M^2} + \left[\frac{5}{2} \lambda^4 \frac{|\phi|^4}{M^2} + 2\lambda^4 \frac{|\phi|^2}{M^2} (\phi^2 + \bar{\phi}^2) \right] \times \times \left(\log \left[\frac{5}{2} \lambda^4 \frac{|\phi|^4}{M^2 \mu^2} - \lambda^4 \frac{|\phi|^2}{M^2 \mu^2} (\phi^2 + \bar{\phi}^2) \right] \right) + O(\frac{1}{M^3})
$$
\n(3.37)

As a result, in this expression the leading order correction is proportional to $\frac{1}{M^2}$. The classical Lagrangian takes the form

$$
S_0 = \phi \bar{\phi} + \Phi \bar{\Phi}
$$

which for this case looks like

$$
S_0 = \phi \bar{\phi} + \frac{\lambda^2 |\phi|^4}{4M^2}
$$
 (3.38)

and the chiral potential, which is of the special form:

$$
W = \frac{1}{2}(\lambda \Phi \phi^2 + M\Phi^2)
$$

can be written as

$$
W = -\frac{1}{8} \frac{\lambda^2 \phi^4}{M} + \frac{1}{4} \lambda \phi^2 \left\{ \frac{\lambda \bar{D}^2 (\bar{\phi}^2)}{8M^2} + \frac{\lambda^3}{64\pi^2 M^2} \bar{D}^2 \left(\left(\frac{\bar{\phi}^2}{2} + |\phi|^2 \right) (2 + \right. \\ + \left. \log \left[\frac{5}{2} \lambda^4 \frac{|\phi|^4}{M^2 \mu^2} - \lambda^4 \frac{|\phi|^2}{M^2 \mu^2} (\phi^2 + \bar{\phi}^2) \right] \right) \right\} + O(\frac{1}{M^3}) \tag{3.39}
$$

As a result the total effective action of light superfields for this case can be cast in the form

$$
S_{eff}[\phi,\bar{\phi}] = S_{eff}^{cl}[\phi,\bar{\phi}] + S_{eff}^{q}[\phi,\bar{\phi}]
$$
\n(3.40)

where $S_{eff}^{cl}[\phi,\bar{\phi}]$ is a classical part of the effective action which is of the form

$$
S_{eff}^{cl}[\phi,\bar{\phi}] = \int d^{8}z \{ \phi\bar{\phi} + \frac{\lambda^{2}|\phi|^{4}}{4M^{2}} \} -
$$

+
$$
\left(\int d^{6}z \left(-\frac{1}{8}\frac{\lambda^{2}\phi^{4}}{M} + \frac{1}{4}\lambda\phi^{2} \left\{ \frac{\lambda\bar{D}^{2}(\bar{\phi}^{2})}{8M^{2}} + \frac{\lambda^{3}}{64\pi^{2}M^{2}}\bar{D}^{2} \left(\left(\frac{\bar{\phi}^{2}}{2} + |\phi|^{2} \right) (2 + \frac{\lambda^{2}}{2}) \right) \right) \right\} + \log\left[\frac{5}{2}\lambda^{4} \frac{|\phi|^{4}}{M^{2}\mu^{2}} - \lambda^{4} \frac{|\phi|^{2}}{M^{2}\mu^{2}} (\phi^{2} + \bar{\phi}^{2})]) \right) \} \right) + h.c. + O(\frac{1}{M^{3}})
$$
(3.41)

$$
S_{eff}^{q} = -\frac{1}{32\pi^{2}} \int d^{8}z \{ 4\lambda^{2} |\phi|^{2} (1 + \log \frac{2M^{2}}{\mu^{2}}) + 4\lambda^{4} \frac{|\phi|^{4}}{M^{2}} + \left[\frac{5}{2} \lambda^{4} \frac{|\phi|^{4}}{M^{2}} + 2\lambda^{4} \frac{|\phi|^{2}}{M^{2}} (\phi^{2} + \bar{\phi}^{2}) \right] \times
$$

$$
\times \left(\log \left[\frac{5}{2} \lambda^{4} \frac{|\phi|^{4}}{M^{2} \mu^{2}} - \lambda^{4} \frac{|\phi|^{2}}{M^{2} \mu^{2}} (\phi^{2} + \bar{\phi}^{2}) \right] \right) + O(\frac{1}{M^{3}})
$$
(3.42)

To convince ourselves that the results are consistent with the decoupling theorem, we can again carry out the field redefinition([3.33\)](#page-21-0) with the finite renormalization constant [\(3.34](#page-21-0)). As a result we arrive at the one-loop effective action of light superfields which is a sum of (3.41) and (3.42) and has the form

$$
S_{eff}[\phi,\bar{\phi}] = \int d^8 z \phi \bar{\phi} + O(\frac{1}{M})
$$
\n(3.43)

Therefore we find that in the absence of self-interaction of the light superfield the effective action for ϕ , ϕ , after a suitable choice of μ and redefinitions of fields and couplings becomes a sum of the classical action of the light superfield and terms at the one loop level that are at least second order in the inverse mass parameter. Hence the result (3.41)-(3.42) for the effective action of the light superfield is again consistent with the decoupling theorem. However, note again, since the parameters of the theory (fields, masses, couplings) are determined from string theory, the one-loop decoupling effects again modify in a quantitative manner the low energy predictions for the couplings.

4 One-loop effective action for non-minimal models

In this section we generalize the discussion to the case of non-minimal models. We consider exampleswith the kählerian potential of the form (2.3) (2.3) . However, as it turns out an exact computation of the one-loop correction in the effective action for a general form of \tilde{K} leads to essential difficulties. Therefore we consider two specific examples.

4.1 The model with heavy quantum superfields and external light superfields

4.1.1 Calculation of the effective action

We consider for example the theory with the action

$$
S = \int d^8 z \left(\phi \overline{\phi} + (1 + \frac{\alpha}{M} (\phi + \overline{\phi})) \Phi \overline{\Phi} \right) +
$$

+
$$
\left(\int d^6 z \left[\frac{1}{2} (M + c\phi) \Phi^2 + \frac{\lambda}{2} \Phi \phi^2 \right] + h.c. \right)
$$
(4.1)

which corresponds to the functions K and W of the form

$$
W = \frac{1}{2}(M + c\phi)\Phi^2 + \frac{\lambda}{2}\Phi\phi^2
$$

$$
K = \Phi\bar{\Phi} + \frac{\alpha}{M}(\phi + \bar{\phi})\Phi\bar{\Phi}
$$
 (4.2)

In principle it is possible to consider the effective action for the general case when both light and heavy superfields are split into a sum of background and quantum parts. However, calculation in this case is very complicated; hence in order to simplify the computing we put light superfields ϕ , $\bar{\phi}$ to be pure background ones. Heavy superfields Φ , $\bar{\Phi}$ are split into a sum of background superfields Φ_0 , $\bar{\Phi}_0$ and quantum ones $\bar{\Phi}$, $\bar{\Phi}$ by a standard way

$$
\Phi \to \Phi_0 + \Phi, \ \bar{\Phi} \to \bar{\Phi}_0 + \bar{\Phi}
$$

As a result the quadratic action of quantum superfields looks like

$$
S = \int d^8 z \Big([1 + \frac{\alpha}{M} (\phi + \bar{\phi})] \Phi \bar{\Phi} \Big) + \Big(\int d^6 z \frac{1}{2} (M + c\phi) \Phi^2 + h.c. \Big) \tag{4.3}
$$

and does not depend on heavy background superfields. Therefore the one-loop quantum correction in the effective action is also Φ_0 , $\bar{\Phi}_0$ independent due to the special (quadratic) structure of action (4.1) in heavy superfields. In this case the column vector ψ (see Section 2.3) consists of the heavy superfield Φ only, and all the derivatives with respect to ψ are equivalent to those with respect to Φ only.

Φ,

The one-loop correction in the effective action $\Gamma^{(1)}$ can be defined in this case by the equation

$$
e^{i\Gamma^{(1)}} = \int D\Phi D\bar{\Phi} \exp(iS^{(2)}) \tag{4.4}
$$

where S_2 is the part of the action quadratic in quantum superfields Φ , $\bar{\Phi}$ (see Section 2.3)

$$
S_2 = \int d^8 z (K_{\Phi \bar{\Phi}} \Phi \bar{\Phi} + \frac{1}{2} K_{\bar{\Phi} \bar{\Phi}} \bar{\Phi} \bar{\Phi} + \frac{1}{2} K_{\Phi \Phi} \Phi \Phi) +
$$

+
$$
[\int d^6 z \frac{1}{2} W_{\Phi \Phi} \Phi \Phi + h.c.]
$$
 (4.5)

and $K_{\Phi\bar{\Phi}}$, $K_{\bar{\Phi}\Phi}$, $K_{\Phi\Phi}$ are functions of light background superfields ϕ , $\bar{\phi}$, only.

Using([4.4\)](#page-24-0) one can write $\Gamma^{(1)}$ as

$$
\Gamma^{(1)} = \frac{i}{2} Tr \log G^{(\psi)} \tag{4.6}
$$

The straightforward calculation of the effective potential based on (4.6) is very complicated because elements of the matrix $G^{(\psi)}$ are defined in different chiral superspaces hence their chiralities are mixed. However, in this model it is possible to employ a technique analogous to that one used in [\[22, 23\]](#page-56-0). Namely, let us consider a theory of a real scalar superfield with the standard action

$$
S_v = -\frac{1}{16} \int d^8 z v D^\alpha \bar{D}^2 D_\alpha v \tag{4.7}
$$

This theory can be quantized using the Faddeev-Popov procedure (cf. [\[22, 23](#page-56-0)]), and the effective action W_v corresponding to this theory can be determined by the following integral:

$$
e^{iW_v} = \int Dv \delta(\frac{1}{4}D^2v - \bar{\Phi}) \delta(\frac{1}{4}\bar{D}^2v - \Phi) \exp(iS_v) det M_0 \tag{4.8}
$$

Here M_0 is a Faddeev-Popov matrix,

$$
M_0 = \begin{pmatrix} 0 & -\frac{1}{4}\bar{D}^2 \\ -\frac{1}{4}D^2 & 0 \end{pmatrix}
$$
 (4.9)

It is evident that W_v is a constant: since the model (4.7) is gauge invariant its effective action doesnot depend on Φ (cf. [[22, 23\]](#page-56-0)).

If we multiply the corresponding left-hand sides and right-hand sides of([4.4\)](#page-24-0) and (4.8) we obtain

$$
\exp(i\Gamma^{(1)}[\phi,\bar{\phi}] + iW_v) = \int \mathcal{D}\Phi \mathcal{D}\bar{\Phi}Dv\delta(\frac{1}{4}D^2v - \bar{\Phi})\delta(\frac{1}{4}\bar{D}^2v - \Phi)\exp(iS_v)detM_0 \times
$$

$$
\times \exp(i\{\int d^8z(K_{\Phi\bar{\Phi}}\Phi\bar{\Phi} + \frac{1}{2}K_{\bar{\Phi}\bar{\Phi}}\bar{\Phi}\bar{\Phi} + \frac{1}{2}K_{\Phi\Phi}\Phi\Phi) +
$$

$$
+ \left[\int d^6z \frac{1}{2}W_{\Phi\Phi}\Phi\Phi + h.c.\right]\}) \tag{4.10}
$$

Here S_v is given by (4.7). Then we integrate over Φ , $\bar{\Phi}$ by means of delta functions (cf. [[22](#page-56-0), [23, 20\]](#page-56-0)) and since e^{iW_v} and $det M_0$ are constants we arrive at the following expression for the one-loop effective action for the superfields ϕ , ϕ :

$$
\exp(i\Gamma^{(1)}) = \int Dv \exp(\frac{i}{2}v\Delta v); \ \Gamma^{(1)} = \frac{i}{2}Tr \log \Delta \tag{4.11}
$$

where

$$
\Delta = \Box + \frac{1}{16} (K_{\Phi\bar{\Phi}} - 1) \{D^2, \bar{D}^2\} + \frac{1}{4} W_{\Phi\Phi} \bar{D}^2 + \frac{1}{4} \bar{W}_{\bar{\Phi}\bar{\Phi}} D^2 + + \frac{1}{4} (\bar{D}^2 K_{\Phi\Phi}) \bar{D}^2 + \frac{1}{4} (D^2 K_{\bar{\Phi}\bar{\Phi}}) D^2
$$
\n(4.12)

This expression can be used to calculate the explicit form of the one-loop effective action. We note that the last two terms in this expression depend on derivatives of the background superfields and therefore do not contribute to the kählerian effective potential.

To find the one-loop effective action from([4.11](#page-25-0)) we use the Schwinger representation

$$
Tr \log \Delta = Tr \int \frac{ds}{s} e^{is\Delta} \tag{4.13}
$$

Since the kählerian effective action (by definition) does not depend on supercovariant derivatives of superfields, one calculates the kählerian effective potential via proper-time method whereone can omit the last two terms in (4.12) (4.12) and obtain (cf. $[22, 23]$)

$$
K^{(1)} = \frac{i}{2} Tr \int \frac{ds}{s} e^{is\Delta t}
$$

= $\frac{i}{2} \int d^8 z_1 d^8 z_2 \delta^8 (z_1 - z_2) \int \frac{ds}{s} e^{is\Delta t} e^{is\Box} \delta^8 (z_1 - z_2)$ (4.14)

where

$$
\Delta' = \Box + \frac{1}{16} (K_{\Phi\bar{\Phi}} - 1) \{D^2, \bar{D}^2\} + \frac{1}{4} W_{\Phi\Phi} \bar{D}^2 + \frac{1}{4} \bar{W}_{\bar{\Phi}\bar{\Phi}} D^2 \equiv
$$

$$
\equiv \Box + H \{D^2, \bar{D}^2\} + \frac{1}{4} W'' \bar{D}^2 + \frac{1}{4} \bar{W}'' D^2
$$
(4.15)

Here $W'' = W_{\Phi\Phi}$, $H = \frac{1}{16}(K_{\Phi\bar{\Phi}} - 1)$. After calculations described in Appendix C we arrive at the renormalized one-loop correction in the kählerian effective potential in the form

$$
K^{(1)} = -\frac{1}{32\pi^2} \int d^8 z \frac{W'' \bar{W}''}{K_{\Phi\bar{\Phi}}^2} \Big[\log \Big\{ \frac{W'' \bar{W}''}{\mu^2 K_{\Phi\bar{\Phi}}^2} \Big\} \Big] \tag{4.16}
$$

Inthe case when K and W are given by (4.2) (4.2) , and we can write this correction as

$$
K^{(1)} = -\frac{1}{32\pi^2} \int d^8 z \frac{|(M+c\phi)|^2}{\left(1 + \frac{\alpha}{M}(\phi + \bar{\phi})\right)^2} \left[\log\left\{\frac{|(M+c\phi)|^2}{\mu^2(1 + \frac{\alpha}{M}(\phi + \bar{\phi}))^2}\right\}\right]
$$
(4.17)

We note that the result for the one-loop contribution to kählerian effective potential does not depend on λ . This result, however, is natural since the vertex $\lambda \Phi \phi^2$ is linear in the quantum superfield Φ and hence it can lead to one-particle-reducible diagrams, only. The same follows from (4.15) with K and W given by (4.2) .

To study the effective action of light superfields in the model one can expand the $K^{(1)}$ into a power series in the inverse mass.

4.1.2 Solving the effective equations of motion

Let us consider decoupling effects for the non-minimal model with the one-loop kählerian effective potential given by (4.17) . The result (4.17) can be represented as a power series in

1 $\frac{1}{M}$. The leading order terms in this expansion have the form

$$
K^{(1)} = -\frac{1}{32\pi^2} \Big\{ \phi\bar{\phi} [2(c^2 - 3\alpha c + 2\alpha^2) + (c^2 - 4\alpha c + 6\alpha^2) \log \frac{M^2}{\mu^2} + \frac{\phi\bar{\phi}(\phi + \bar{\phi})}{M} [\log \frac{M^2}{\mu^2}(-12\alpha^3 + 9c\alpha^2 - 2\alpha c^2) + \frac{(3\alpha^2 - 2\alpha c)(c - \alpha) + (6\alpha^2 - 4\alpha c + c^2)(c - \alpha) + \frac{1}{2}(c - 2\alpha)(\alpha^2 - c^2) + \alpha^2(c - \alpha)] + O(\frac{1}{M^2}) \Big\}
$$
(4.18)

For K , W of the form (4.2) the effective equations of motion $(2.22, 2.23)$ take the form:

$$
-\frac{1}{4}\bar{D}^{2}(\bar{\Phi}(1+\alpha\frac{\phi+\bar{\phi}}{M})) + (M+c\phi)\Phi + \frac{\lambda}{2}\phi^{2} = 0
$$

$$
-\frac{1}{4}D^{2}(\Phi(1+\alpha\frac{\phi+\bar{\phi}}{M})) + (M+c\bar{\phi})\bar{\Phi} + \frac{\lambda}{2}\bar{\phi}^{2} = 0
$$
(4.19)

The zeroth order solution is

$$
\Phi_0 = -\frac{\lambda}{2(M + c\phi)}\phi^2 = -\frac{\lambda}{2M}\phi^2 + O(\frac{1}{M^2})
$$

Dueto (3.20) (3.20) the *n*th order equation has the form

$$
-\frac{1}{4}\bar{D}^2(\Phi_{n-1}(1+\alpha\frac{\phi+\bar{\phi}}{M})) + (M+c\phi)\Phi_n = 0
$$
\n(4.20)

and the analogous equation for $\bar{\Phi}_n$. Hence the first order approximation can be written as

$$
\Phi_1 = \frac{1}{4(M + c\phi)} \bar{D}^2 \left(-\frac{\lambda}{2(M + c\bar{\phi})} \bar{\phi}^2 (1 + \alpha \frac{\phi + \bar{\phi}}{M}) \right) = -\frac{\lambda}{16M^2} \bar{D}^2 (\bar{\phi}^2) + O(\frac{1}{M^3})
$$

As a result we can substitute $\Phi = \Phi_0 + \Phi_1$ into the one-loop effective action and write the effective action for light superfields in the form

$$
S_{eff}[\phi,\bar{\phi}] = S_{eff}^{cl}[\phi,\bar{\phi}] + S_{eff}^{q}[\phi,\bar{\phi}] \tag{4.21}
$$

where $S_{eff}^{cl}[\phi,\bar{\phi}]$ is a classical part of the effective action which is of the form

$$
S_{eff}^{cl}[\phi,\bar{\phi}] = \int d^8 z \phi \bar{\phi} - \left\{ \int d^6 z \frac{3\lambda^2 \phi^4}{8M} + h.c. \right\} + O(\frac{1}{M^2}) \tag{4.22}
$$

and $S_{eff}^{q}[\phi,\bar{\phi}]$ is a quantum part of the effective action looking like

$$
S_{eff}^{q}[\phi,\bar{\phi}] = -\frac{\hbar}{32\pi^{2}} \int d^{8}z \{ \phi\bar{\phi}[2(c^{2} - 3\alpha c + 2\alpha^{2}) + (c^{2} - 4\alpha c + 6\alpha^{2}) \log \frac{M^{2}}{\mu^{2}}] +
$$

+
$$
\{ \frac{\phi\bar{\phi}(\phi + \bar{\phi})}{M}[-\frac{c^{2} - 3\alpha c + 2\alpha^{2}}{c^{2} - 4\alpha c + 6\alpha^{2}}(-12\alpha^{3} + 9c\alpha^{2} - 2\alpha c^{2}) +
$$

+
$$
(3\alpha^{2} - 2\alpha c)(c - \alpha) + (6\alpha^{2} - 4\alpha c + c^{2})(c - \alpha) +
$$

+
$$
\frac{1}{2}(c - 2\alpha)(\alpha^{2} - c^{2}) + \alpha^{2}(c - \alpha)] +
$$

+
$$
O(\frac{1}{M^{2}})\}
$$

(4.23)

To check that the result corresponds to the decoupling theorem we can renormalize the superfield ϕ according to ([3.33\)](#page-21-0) with the finite renormalization constant Z of the form

$$
Z = \left(1 - \frac{h}{32\pi^2} [2(c^2 - 3\alpha c + 2\alpha^2) + (c^2 - 4\alpha c + 6\alpha^2) \log \frac{M^2}{\mu^2}] \right)^{1/2}
$$
(4.24)

As a result the effective action being the sum of the classical part([4.22\)](#page-27-0) and the quantum one([4.23](#page-27-0)) takes the form

$$
S_{eff}[\phi,\bar{\phi}] = \int d^8 z \tilde{\phi}\tilde{\bar{\phi}} + O(\frac{1}{M})
$$
\n(4.25)

which is thus formally consistent with the decoupling theorem. (However, note again that fields and couplings α , c are determined from string theory, and thus the classical action gets essential one-loop corrections.)

4.2 The model with light and heavy quantum superfields and light and heavy external superfields

4.2.1 Calculation of the effective action

It turns out that for non-minimal models it is possible obtain explicit expressions for the the one-loop corrected action in the leading orders in $\frac{1}{M}$ when both light and heavy superfields have background and quantum parts, by employing the diagram approach.

Let us consider a specific non-minimal model with action

$$
S = \int d^{8}z [\phi \bar{\phi} + \Phi \bar{\Phi} + \frac{\lambda}{M} (W_{1}(\phi) \bar{\Phi} + \bar{W}_{1}(\bar{\phi}) \Phi)] +
$$

+
$$
[\int d^{6}z (\frac{g}{3!} \phi^{3} + \frac{M}{2} \Phi^{2}) + h.c.]
$$
 (4.26)

Thepropagators of the model have the standard form ([B.1](#page-49-0)). The zeroth order in $\frac{1}{M}$ the expansion of the one-loop kählerian effective potential corresponds to the case where vertices, proportional to $\frac{1}{M}$, are absent. The corresponding supergraphs consist of $\langle \phi \phi \rangle$ -propagators with alternating background fields ϕ and $\overline{\phi}$ (cf. [\[28](#page-56-0), [29](#page-56-0)]):

Here the external legs correspond to alternating ϕ and ϕ .

The result is given by a sum of all such supergraphs. After calculations that are completely parallel with those carried out above we arrive at the leading order correction $K_0^{(1)}$ displayed ineq. $(B.8)$ $(B.8)$ $(B.8)$. $(Cf. [29, 28]$ $(Cf. [29, 28]$ $(Cf. [29, 28]$. Other methods of calculating the kählerian effective potential weregiven in $|30|$.)

To study corrections that are higher orders in the $\frac{1}{M}$ expansion, it is more convenient to split superfields ϕ , Φ into a sum of background ϕ , Φ and quantum ϕ_q , Φ_q superfields. As a result we arrive at the following quadratic action of the quantum superfields ϕ_q , Φ_q in external superfields ϕ , Φ

$$
S = \int d^{8}z [\phi_{q}\bar{\phi}_{q} + \Phi_{q}\bar{\Phi}_{q} + \frac{\lambda}{M}(W'_{1}(\phi)\phi_{q}\bar{\Phi}_{q} + W''_{1}(\phi)\Phi\phi_{q}^{2} + h.c.)] +
$$

+
$$
[\int d^{6}z (\frac{g}{2}\phi\phi_{q}^{2} + \frac{M}{2}\Phi_{q}^{2}) + h.c.]
$$
(4.27)

Let us study corrections at different orders in $\frac{1}{M}$ expansion. It is evident that the presence of corrections at higher orders is due to the presence of vertices proportional to λ in the supergraphs. One can assume that corrections at higher orders in the $\frac{1}{M}$ expansion can arise in two cases. In the first case the diagram contains the background heavy superfield Φ and the vertex is of the form $\int d^8 z \overline{W}'' \Phi \overline{\phi}_q^2$. Note also that in the diagrams each factor D^2 corresponds to a quantum superfield ϕ_q . Therefore a contribution of such vertices is of the form

Here the double external line with sign \times corresponds to $\bar{W}''\Phi$ (an analogous fragment of a diagram arises for the contribution proportional to $W''\bar{\Phi}$). Then a transportation of any of the two D^2 -factors during D-algebra transformations leads to the factor $D^2\Phi$, i.e. diagrams containing such vertices do not contribute to the kählerian effective potential. Therefore corrections at the first and higher orders in the inverse mass expansion can appear only in supergraphs with light external field lines only.

Let us study the possible diagrams of that type. We must pay attention to the fact that the propagator of the light superfield ϕ depends on the background superfield. We denote it with a bold line which has the following diagrammatic representation

Here an external line denotes the background superfields ϕ , $\bar{\phi}$. Summation of this chain of diagrams allows one to show that the total propagator of the light superfield is of the form

$$
\langle \phi_q(z_1)\overline{\phi}_q(z_2)\rangle = -\frac{\overline{D}_1^2 D_2^2}{16(\Box - g^2|\phi|^2)} \delta_{12} \tag{4.28}
$$

Thepropagator of the heavy superfield Φ is given by ([B.1](#page-49-0)). One can easily see that in addition to the diagrams discussed above, which do not contain vertices proportional to $\frac{\lambda}{M}$, there are two other types of supergraphs. The first one consists of an arbitrary number of repeating links of the form $\langle \Phi \bar{\Phi} \rangle \langle \phi \bar{\phi} \rangle$, i.e.

where bold external lines with sign \times correspond to external $\frac{\lambda}{M}W'_{1}(\phi)$ term, and dashed internal ones to $\langle \Phi \bar{\Phi} \rangle$ propagators. The second type of diagrams consist of an arbitrary number of repeating links of the form

 $<\phi\bar{\phi}>_t<\Phi\bar{\Phi}><\phi\bar{\phi}>_t<\Phi\Phi><\phi\bar{\phi}>_t<\bar{\Phi}\Phi><\phi\phi>\sqrt{\phi}\bar{\Phi}>_t<\Phi\bar{\Phi}>$ which can be diagrammatically represented as

Here the dashed-and-dotted line denotes $\langle \Phi \Phi \rangle$ and the jagged line denotes $\langle \bar{\Phi} \bar{\Phi} \rangle$.

The contribution of a diagram of the first type, which consists of n such links, is of the form

$$
I_n = \frac{1}{n} \left\{ \frac{\lambda^2}{M^2} |W_1'|^2 < \phi \bar{\phi} > \phi \bar{\Phi} > \right\}^n \tag{4.29}
$$

and the contribution of a diagram of the second type that consists of n chains is of the form

$$
J_n = \frac{1}{n} \Big\{ \frac{\lambda^8}{M^8} |W_1'|^8 < \phi\bar{\phi} > \phi\bar
$$

Usingthe exact form of the propagator given by $(B.1)$ $(B.1)$ $(B.1)$ and carrying out D-algebra transformations we find that (4.29) and (4.30) can be cast in the form

$$
I_n = \frac{1}{\Box} \frac{1}{n} \left(\frac{\Box^2}{(\Box - M^2)(\Box - g^2|\phi^2|)} \right)^n \frac{\lambda^{2n}}{M^{2n}} |W_1'|^{2n}
$$
(4.31)

and

$$
J_n = \frac{1}{\Box} \frac{1}{n} \left(\frac{\Box^7}{\left(\Box - M^2 \right)^4 \left(\Box - g^2 | \phi^2 | \right)^4} \right)^n \frac{\lambda^{2n}}{M^{6n}} |W_1'|^{8n} \tag{4.32}
$$

respectively.

We can now find the leading terms in the inverse mass expansion for the above contributions. It is easy to see that as n grows the order in the inverse mass contribution increases. J_n is proportional to $M^{-6n}\Box^{-n-1}$. After Fourier transformation and the integration over momenta the leading term will be of the order M^{2-8n} , i.e. it will involve at least six powers of the inverse mass parameter. As for I_n it is proportional to $M^{-2n}\Box^{-1}$; after the integration the leading term is of the order M^{2-2n} . Therefore the zeroth and the second order are given by I_1 and I_2 . A straightforward calculation carried out in the framework of dimensional regularization yields

$$
I_1 = \lambda^2 \frac{|W_1'|^2}{32\pi^2} \left[\frac{2}{\epsilon} + \log \frac{M^2}{\mu^2} + 1\right] + \frac{\lambda^2}{32\pi^2 M^2} |W_1'|^2 g^2 |\phi|^2 \left[\frac{2}{\epsilon} + \log \frac{M^2}{\mu^2}\right]
$$

\n
$$
I_2 = -\frac{\lambda^4 |W_1'|^4}{32\pi^2 M^2} \left(\frac{2}{\epsilon} + \log \frac{M^2}{\mu^2} + 1\right)
$$
\n(4.33)

Therefore the leading term for the one-loop correction to the effective action of the nonminimal model with the classical action (4.26) is the sum of I_1 and I_2 (eq.(4.33)) as well as the correction([B.8](#page-50-0)) and it takes the form:

$$
K^{(1)} = -\frac{1}{32\pi^2} g^2 |\phi|^2 (\frac{2}{\epsilon} + \log \frac{g^2 |\phi|^2}{\mu^2}) + \lambda^2 \frac{|W_1'|^2}{32\pi^2} [\frac{2}{\epsilon} + \log \frac{M^2}{\mu^2} + 1] -
$$
(4.34)

$$
- \frac{1}{32\pi^2 M^2} [\lambda^4 |W_1'|^4 (\frac{2}{\epsilon} + \log \frac{M^2}{\mu^2} + 1) - \lambda^2 |W_1'|^2 g^2 |\phi|^2 {\frac{2}{\epsilon} + \log \frac{M^2}{\mu^2}}] + O(\frac{1}{M^3})
$$

To cancel the one-loop divergences we must add a counterterm of the form

$$
S_{ct}^{(1)} = -\frac{1}{16\pi^2 \epsilon} \Big(-g^2 |\phi|^2 + \lambda^2 |W_1'|^2 - \frac{1}{M^2} [\lambda^2 g^2 |W_1'|^2 |\phi|^2 - \lambda^4 |W_1'|^4] \Big) \tag{4.35}
$$

As a result we arrive at the final form of the renormalized one-loop correction in the kählerian effective potential:

$$
K^{(1)} = -\frac{1}{32\pi^2} g^2 |\phi|^2 \log \frac{g^2 |\phi|^2}{\mu^2} + \lambda^2 \frac{|W_1'|^2}{32\pi^2} [\log \frac{M^2}{\mu^2} + 1] -
$$

$$
- \frac{1}{32\pi^2 M^2} \Big[\lambda^4 |W_1'|^4 (\log \frac{M^2}{\mu^2} + 1) - \lambda^2 |W_1'|^2 g^2 |\phi|^2 \log \frac{M^2}{\mu^2} \Big] + O(\frac{1}{M^3})
$$
(4.36)

The essential feature of this correction is that terms that would depend on the background heavy superfield Φ are absent. This fact is due to the form of the vertex $\int d^8 z W(\phi) \bar{\Phi}$ whose presence necessarily leads to terms which depend on supercovariant derivatives of Φ.

4.2.2 Solution of the effective equations of motion

Now let us consider the effective action (4.36) corresponding to the model with classical action [\(4.26](#page-28-0)). The effective equation of motion corresponding to this model have the form

$$
M\Phi - \frac{1}{4}\bar{D}^2(\bar{\Phi} + \frac{\lambda}{M}\bar{W}_1(\bar{\phi})) = 0
$$
\n
$$
M\bar{\Phi} - \frac{1}{4}D^2(\Phi + \frac{\lambda}{M}W_1(\phi)) = 0
$$
\n(4.37)

We note that quantum corrections in this model do not depend on Φ. The zeroth order approximationfor the superfield Φ is $\Phi_0 = 0$ (since $\frac{\partial W}{\partial \Phi} = 0$ only at $\Phi = 0$) (see ([2.20](#page-7-0))). The first order approximation([3.20\)](#page-18-0) looks like

$$
\Phi_1 = \frac{\lambda}{4} \frac{\bar{D}^2}{M^2} \bar{W}_1(\bar{\phi}) \tag{4.38}
$$

It follows from [\(3.20](#page-18-0)) that for this model the *n*-th order approximation Φ_n is of the order $\frac{1}{M^{n+1}}$. Therefore the final result for the effective action of light superfields reads as

$$
\Gamma^{(1)} = \int d^8 z \Big(\phi \bar{\phi} - \frac{\hbar}{32\pi^2} \{ g^2 |\phi|^2 \log \frac{g^2 |\phi|^2}{\mu^2} + \lambda^2 |W_1'|^2 [1 + \log \frac{M^2}{\mu^2}] - \frac{1}{M^2} [\lambda^4 |W_1'|^4 (1 + \log \frac{M^2}{\mu^2}) - \lambda^2 |W_1'|^2 g^2 |\phi|^2 \log \frac{M^2}{\mu^2}] \} \Big) + \frac{\left(\int d^6 z \frac{g}{3!} \phi^3 + h.c. \right) + O\left(\frac{1}{M^3} \right)}{\Big(4.39 \Big)}
$$
\n
$$
(4.39)
$$

This is a final result for the one-loop corrected effective action of light superfields and applies to a general choice of W_1 .

We note that the result (4.39) is consistent with the decoupling theorem. If we choose for example $W_1 = \phi^2$, $W_1' = 2\phi$ (the minimal model), we can carry out the redefinition of the superfield ϕ by the rule ([3.33\)](#page-21-0) with

$$
Z = \left(1 - \frac{\hbar}{32\pi^2} [4\lambda^2 (1 + \log \frac{M^2}{\mu^2})]\right)^{1/2}
$$
\n(4.40)

Coupling g is redefined by the rule $\tilde{g} = Z^{-3}g$. As a result the effective action of light superfields takes the form

$$
\Gamma^{(1)} = \int d^8 z (\tilde{\phi}\tilde{\bar{\phi}} - \frac{\hbar}{32\pi^2} g^2 |\tilde{\phi}|^2 \log \frac{g^2 |\tilde{\phi}|^2}{\mu^2}) + (\int d^6 z \frac{\tilde{g}}{3!} \tilde{\phi}^3 + h.c.) + O(\frac{1}{M^2})
$$
(4.41)

The result is consistent with the decoupling theorem and coincides in form with the case that corresponds to the minimal model discussed in section 3. However, we note again that since parameters of the theory (fields, masses, couplings) are fixed from string theory, they cannot be redefined. Therefore the classical action is modified by one-loop corrections. The final result for the effective action of light superfields is given by([4.39](#page-31-0)).

5 Quantum corrections to the effective action in gauge theories

5.1 Gauge invariant model of massive chiral superfields

For the sake of completeness we would like to address the decoupling effects in the $N = 1$ supersymmetric gauge theory with chiral superfields that interact with the corresponding gauge superfields (i.e. chiral superfields that are charged under the gauge symmetry)¹. To simplify the consideration we choose the Abelian $U(1)$ gauge theory with the chiral superfields charged under $U(1)$. The simplest (minimal) model describing gauge invariant interactions between massive and light chiral superfields, and the $U(1)$ gauge superfield is of the form:

$$
S = \int d^{8}z (\bar{\Phi}_{1}e^{2gV}\Phi_{1} + \bar{\Phi}_{2}e^{-2gV}\Phi_{2} + \bar{\phi}e^{-gV}\phi) + + \frac{1}{64} \int d^{6}z W^{2} + (\int d^{6}z(M\Phi_{1}\Phi_{2} + \lambda \Phi_{1}\phi^{2}) + h.c.)
$$
 (5.1)

Here V is a real scalar superfield, Φ_1, Φ_2, ϕ are heavy chiral superfields, $W_\alpha = \bar{D}^2 D_\alpha V$ is a superfield strength. The model is invariant under the following gauge transformations with Λ is a chiral superfield, Λ is an antichiral one (c.f. [\[20](#page-56-0)]): denotes

$$
V \rightarrow V + i(\bar{\Lambda} - \Lambda)
$$

 $\overline{^{1}\mathrm{M.C.}}$ would like to thank L. Everett for a discussion on this topic.

$$
\begin{array}{rcl}\n\Phi_1 & \to & e^{2ig\Lambda}\Phi_1 \\
\Phi_2 & \to & e^{-2ig\Lambda}\Phi_2 \\
\phi & \to & e^{-ig\Lambda}\phi\n\end{array} \tag{5.2}
$$

For the sake of concreteness we chose the specific $U(1)$ charges for the chiral superfields such that there is an allowed tri-linear interaction term between Φ_i and ϕ . For the sake of simplicity we did not include self-interactions of the light fields in the chiral potential. In addition, we did not include explicitly the terms of possible other light chiral superfields which we chose not to interact with Φ_i and ϕ superfields; these additional chiral superfields however should be present, in order to cancel chiral anomalies.

The gauge-fixing term is of the form

$$
S_{gf} = -\frac{1}{16\xi} \int d^8 z D^2 V \bar{D}^2 V \tag{5.3}
$$

where ξ is a gauge parameter. The propagators for the fields of the model have the form (see, e.g., [\[20\]](#page-56-0))

$$
\langle \Phi_1 \bar{\Phi}_1 \rangle = -\frac{\bar{D}_1^2 D_2^2}{16(\Box - M^2)} \delta^8(z_1 - z_2)
$$

\n
$$
\langle \Phi_2 \bar{\Phi}_2 \rangle = -\frac{\bar{D}_1^2 D_2^2}{16(\Box - M^2)} \delta^8(z_1 - z_2)
$$

\n
$$
\langle \Phi_1 \Phi_2 \rangle = -\frac{M \bar{D}_1^2}{4(\Box - M^2)} \delta^8(z_1 - z_2)
$$

\n
$$
\langle \phi \bar{\phi} \rangle = -\frac{\bar{D}_1^2 D_2^2}{16 \Box} \delta^8(z_1 - z_2)
$$

\n
$$
\langle v v \rangle = (-\frac{D^\alpha \bar{D}^2 D_\alpha}{8 \Box^2} + \xi \frac{\{D^2, \bar{D}^2\}}{16 \Box^2}) \delta^8(z_1 - z_2)
$$
(5.4)

Possible vertices associated with the action [\(5.1](#page-32-0)) are

$$
g^n \bar{\phi} v^n \phi, \ g^l \bar{\Phi}_1 v^l \Phi_1, \ g^m \bar{\Phi}_2 v^m \Phi_2, \ \lambda \Phi_1 \phi^2, \ \lambda \bar{\Phi}_1 \bar{\phi}^2 \tag{5.5}
$$

The effective action can again be studied in the framework of the loop expansion. To carry out the calculations we split all the superfields into a sum of background Φ_1 , Φ_2 , ϕ , V and quantum $\Phi_{1q}, \Phi_{2q}, \phi_q, v$ superfields by the rule

$$
\begin{array}{rcl}\n\Phi_1 & \to & \Phi_1 + \Phi_{1q}, \ \Phi_2 \to \Phi_2 + \Phi_{2q} \\
\phi & \to & \phi + \phi_q, \ v \to V + v\n\end{array} \tag{5.6}
$$

As a result we arrive at the quadratic action of quantum superfields

$$
S_q = \int d^8 z \Big\{ \frac{1}{2} v \big[\Box + \big(\frac{1}{\xi} - 1 \big) \frac{1}{16} \{ D^2, \bar{D}^2 \} + g^2 (4 \bar{\Phi}_1 e^{2gV} \Phi_1 + 4 \bar{\Phi}_2 e^{-2gV} \Phi_2 + \bar{\phi} e^{-gV} \phi \big) \big] v +
$$

+
$$
\bar{\Phi}_{1q}e^{2gV}\Phi_{1q} + \bar{\Phi}_{2q}e^{-2gV}\Phi_{2q} + \phi_{q}e^{-gV}\bar{\phi}_{q} +
$$

\n+ $2g(\bar{\Phi}_{1}e^{2gV}v\Phi_{1q} + \bar{\Phi}_{1q}e^{2gV}v\Phi_{1}) - 2g(\bar{\Phi}_{2}e^{-2gV}v\Phi_{2q} + \bar{\Phi}_{2q}e^{-2gV}v\Phi_{2}) -$
\n- $g(\bar{\phi}e^{-gV}v\phi_{q} + \bar{\phi}_{q}e^{-gV}v\phi)\} +$
\n+ $\left(\int d^{6}z(M\Phi_{1q}\Phi_{2q} + \lambda\Phi_{1}\phi_{q}^{2} + 2\lambda\phi\Phi_{1q}\phi_{q}) + h.c.\right)$ (5.7)

In the following we employ this action to determine the one-loop corrections.

The one-loop contribution to the effective action $\Gamma^{(1)}$ is defined in the usual way (see [\[21\]](#page-56-0)):

$$
\exp(i\Gamma^{(1)}[\Phi_1, \Phi_2, \phi, V]) = \int D\Phi_{1q} D\Phi_{2q} D\phi_q Dv \exp(iS_q[\Phi_{1q}, \Phi_{2q}, \phi_q, v; \Phi_1, \Phi_2, \phi, V]) \quad (5.8)
$$

The low-energy leading terms in the effective action are the kählerian effective potential that depends only on $\Phi_1, \Phi_2, \phi, \bar{\Phi}_1, \bar{\Phi}_2, \bar{\phi}, V$ but not on their supercovariant derivatives, the chiral effective potential that is a holomorphic function of chiral superfields Φ_1 , Φ_2 , ϕ , and the field depending effective gauge coupling $f(\Phi_1, \Phi_2, \phi)$ holomorphic function of superfields that multiplies the term proportional to W^2 . They completely specify the low-energy effective action of $N = 1$ supersymmetric gauge theory (see, e.g., [\[29\]](#page-56-0)).

5.2 One-loop kählerian potential in supersymmetric gauge theory

Tostudy decoupling effects in the model (5.1) (5.1) we determine the one-loop kählerian effective potential and consider the effective equations of motion for the heavy superfields. We carry out the calculation of one-loop quantum contribution to kählerian effective potential on the base of the expression (5.8) with quadratic action of quantum superfields [\(5.7](#page-33-0)). The perturbative approach for the action [\(5.7](#page-33-0)) can be developed in the following way. We consider

$$
S_q = \int d^8 z \left\{ \frac{1}{2} v \left[\Box + \left(\frac{1}{\xi} - 1 \right) \frac{1}{16} \{ D^2, \bar{D}^2 \} \right] v + \bar{\Phi}_{1q} \Phi_{1q} + \bar{\Phi}_{2q} \Phi_{2q} + \phi_q \bar{\phi}_q + \right. \\ \left. + \left(\int d^6 z (M \Phi_{1q} \Phi_{2q} + h.c.) \right) \right\} \tag{5.9}
$$

as a quadratic part which leads just to the propagators([5.4\)](#page-33-0) and all other terms in [\(5.7](#page-33-0)) are treated as vertices of interaction.

Let us consider possible one-loop supergraphs. They can contain both $\langle vv \rangle$ and $<\phi\bar{\phi}>, \langle\Phi_1\bar{\Phi}_1>, \langle\Phi_2\bar{\Phi}_2>, \langle\Phi_1\Phi_2\rangle\rangle$ propagators. We will use the Landau gauge $\xi = 0$. The propagator $\langle v v \rangle$ is proportional to the projection operator $\Pi_1 = -\frac{D^{\alpha} \bar{D}^2 D_{\alpha}}{8\Box}$ in this gauge, and propagators $\langle \phi \phi \rangle$, $\langle \Phi_1 \bar{\Phi}_1 \rangle$, $\langle \Phi_2 \bar{\Phi}_2 \rangle$ are proportional to the projection operator $\Pi_0 = \frac{\bar{D}^2 D^2}{16 \Box}$ $\frac{16D^2}{16D}$ (see e.g.[\(5.4](#page-33-0))). The projection operators Π_0 and Π_1 are orthogonal to each other[[20\]](#page-56-0). As a result each supergraph containing propagators of both chiral superfields and gauge superfields has zero contribution (it is easy to see that the analogous situation takes place for supergraphs containing the propagator $\langle \Phi_1 \Phi_2 \rangle$ due to the identity $\bar{D}^2\Pi_0 = 0$. Therefore there are no supergraphs containing both propagators of chiral and gauge superfields. As a result, in the Landau gauge there can exist only two types of supergraphs: the first of them contains propagators of chiral superfields only, and the second one contains propagators of gauge superfields only. Hence in order to calculate the one-loop kählerian effective potential we can omit in action (5.7) the cross terms, i.e. those that include both the chiral and vector quantum superfields because such terms will lead to supergraphs containing both propagators of chiral and gauge superfields. As a result one can use the expression for the quadratic action

$$
S_q = \int d^8 z \left\{ \frac{1}{2} v \left[\Box + \left(\frac{1}{\xi} - 1 \right) \frac{1}{16} \{ D^2, \bar{D}^2 \} + g^2 (4 \bar{\Phi}_1 e^{2gV} \Phi_1 + 4 \bar{\Phi}_2 e^{-2gV} \Phi_2 + \bar{\phi} e^{-gV} \phi) \right] v + \bar{\Phi}_{1q} e^{2gV} \Phi_{1q} + 4 \bar{\Phi}_{2q} e^{-2gV} \Phi_{2q} + \bar{\phi}_q e^{-gV} \phi_q \right\} + \left(\int d^6 z (M \Phi_{1q} \Phi_{2q} + \lambda \Phi_1 \phi_q^2 + 2 \lambda \phi \Phi_{1q} \phi_q) + h.c. \right)
$$
\n(5.10)

instead of([5.7\)](#page-33-0). It leads to the same supergraphs as([5.7\)](#page-33-0). At the end of the calculations the parameter ξ must be put to zero.

To study the one-loop contribution to kählerian effective potential it is convenient to introduce background superfields $\tilde{\phi}$, $\tilde{\Phi}_1$, $\tilde{\Phi}_2$ and conjugated ones in the following way

$$
\tilde{\phi} = e^{-gV/2}\phi, \quad \tilde{\phi} = \bar{\phi}e^{-gV/2}
$$
\n
$$
\tilde{\Phi}_1 = e^{gV}\Phi_1, \quad \tilde{\bar{\Phi}}_1 = \bar{\Phi}_1 e^{gV}
$$
\n
$$
\tilde{\Phi}_2 = e^{-gV}\Phi_2, \quad \tilde{\bar{\Phi}}_2 = \bar{\Phi}_2 e^{-gV}
$$
\n(5.11)

The quantum superfields are redefined by the same way. Such redefinitions represent themselves as a sort of introduction of background covariantly chiral superfields [\[20](#page-56-0)]. We note that background superfields $\tilde{\phi}$, $\tilde{\Phi}_1$, $\tilde{\Phi}_2$ can be treated as constants. The quadratic action of quantum superfields takes the form

$$
S_q = \int d^8 z \left\{ \frac{1}{2} v \left[\Box + (\frac{1}{\xi} - 1) \frac{1}{16} \{ D^2, \bar{D}^2 \} + g^2 (4 \tilde{\bar{\Phi}}_1 \tilde{\Phi}_1 + 4 \tilde{\bar{\Phi}}_2 \Phi_2 + \tilde{\bar{\phi}} \tilde{\phi}) \right] v +
$$

+ $\tilde{\bar{\Phi}}_{1q} \tilde{\Phi}_{1q} + \tilde{\bar{\Phi}}_{2q} \tilde{\Phi}_{2q} + \tilde{\phi}_q \tilde{\bar{\phi}}_q \right\} +$
+ $\left(\int d^6 z (M \tilde{\Phi}_{1q} \tilde{\Phi}_{2q} + \lambda \tilde{\Phi}_1 \tilde{\phi}_q^2 + 2 \lambda \tilde{\phi} \tilde{\Phi}_{1q} \tilde{\phi}_q) + h.c. \right)$ (5.12)

Thepropagators $\langle \tilde{\phi} \tilde{\phi} \rangle$, $\langle \tilde{\Phi}_1 \tilde{\Phi}_1 \rangle$ and other ones have the standard form ([5.4\)](#page-33-0).

As a result we find the one-loop correction to the effective action to be of the form

$$
\Gamma^{(1)} = \Gamma_1^{(1)} + \Gamma_2^{(1)} \tag{5.13}
$$

where

$$
\exp(i\Gamma_1^{(1)}) = \int D\tilde{\Phi}_{1q} D\tilde{\Phi}_{2q} D\tilde{\phi}_q \exp\left[\int d^8 z \{\tilde{\Phi}_{1q} \tilde{\Phi}_{1q} + \tilde{\Phi}_{2q} \tilde{\Phi}_{2q} + \tilde{\phi}_q \tilde{\phi}_q\right] + + \left(\int d^6 z (M \tilde{\Phi}_{1q} \tilde{\Phi}_{2q} + \lambda \tilde{\Phi}_1 \tilde{\phi}_q^2 + 2\lambda \tilde{\phi} \tilde{\Phi}_{1q} \tilde{\phi}_q) + h.c.\right)] \exp(i\Gamma_2^{(1)}) = \int Dv \exp\left(i \int d^8 z \frac{1}{2} v [\Box + (\frac{1}{\xi} - 1) \frac{1}{16} \{D^2, \bar{D}^2\} + g^2 (4 \tilde{\Phi}_1 \tilde{\Phi}_1 + + 4 \tilde{\Phi}_2 \tilde{\Phi}_2 + \tilde{\phi} \tilde{\phi})]v\right)
$$
\n(5.14)

We note that since transformations of the quantum superfields [\(5.11](#page-35-0)) are local their Jacobian is equal to 1. The contributions $\Gamma_1^{(1)}$, $\Gamma_2^{(1)}$ can be calculated in a straightforward way. Correction $\Gamma_1^{(1)}$ is given in terms of supergraphs of two types. Diagrams of the first type consist only of propagators $\langle \tilde{\phi} \tilde{\phi} \rangle$, i.e. they can be represented as rings consisting of different numbers of repeating links

$$
\left\|\quad\right\|
$$

Each link has the form

$$
K = \frac{\lambda^2 |\tilde{\Phi}_1|^2 D^2 \bar{D}^2}{16k^4} \delta_{12}
$$
\n(5.15)

The contribution of a diagram consisting of n such links has the form

$$
J_n = \int d^4\theta \int \frac{d^4k}{\left(2\pi\right)^4} K^n \tag{5.16}
$$

The total contribution of all such diagrams is then of the form $J = \sum_{n=1}^{\infty} J_n$. After calculations analogous to those carried out in Section 4 we arrive at the explicit expression

$$
J = -\frac{1}{32\pi^2} \int d^4\theta \lambda^2 |\tilde{\Phi}_1|^2 \frac{\lambda^2 |\tilde{\Phi}_1|^2}{\mu^2}
$$
 (5.17)

The diagrams of the second type consist of repeating links of the form

where the bold line is a total propagator of a light superfield which depends on the background superfield $\tilde{\Phi}_1$. It can be obtained by a method analogous to the one used in the previous section (but with background, $\tilde{\Phi}_1$, $\tilde{\bar{\Phi}}_1$ instead of ϕ , $\bar{\phi}$) and has the form

$$
\langle \tilde{\phi}\tilde{\bar{\phi}} \rangle = \frac{\lambda^2 |\tilde{\Phi}_1|^2}{k^2 + \lambda^2 |\tilde{\Phi}_1|^2} \frac{D^2 \bar{D}^2}{16k^2} \tag{5.18}
$$

The dashed line corresponds to $\langle \tilde{\Phi}_1 \tilde{\Phi}_1 \rangle$, and the external line with sign \times denotes the background superfield ϕ . The link has the form $\lambda^2|\tilde{\phi}|^2 < \tilde{\phi}\tilde{\phi} > < \tilde{\Phi}_1 \tilde{\Phi}_1 >$ As a result the contribution of a diagram consisting of n such links is of the form

$$
I_n = \int d^4\theta \int \frac{d^4k}{\left(2\pi\right)^4} L^n \tag{5.19}
$$

where

$$
L = \frac{\lambda^2 |\tilde{\phi}|^2 D^2 \bar{D}^2}{16(k^2 + M^2)(k^2 + \lambda^2 |\tilde{\Phi}_1|^2)}
$$
(5.20)

The total contribution of all these diagrams, after calculations analogous to those carried out in the previous section, has the form

$$
I = \sum_{n=1}^{\infty} I_n = -\frac{1}{32\pi^2} \left\{ \frac{2}{\epsilon} (-\lambda^2 |\tilde{\phi}|^2) + [M^2 + \lambda^2 (|\tilde{\Phi}_1|^2 - |\tilde{\phi}|^2)] \log \frac{[M^2 + \lambda^2 (|\tilde{\Phi}_1|^2 - |\tilde{\phi}|^2)]}{\mu^2} - \lambda^2 |\tilde{\Phi}_1|^2 \log \frac{\lambda^2 |\tilde{\Phi}_1|^2}{\mu^2} \right\}
$$
(5.21)

After cancellation of a divergence proportional to $\frac{1}{\epsilon}$, via a suitable counterterm, we arrive at the following one-loop contribution to the kählerian effective potential:

$$
K_{1r}^{(1)} = I + J = -\frac{1}{32\pi^2} \Big\{ [M^2 + \lambda^2 (|\tilde{\Phi}_1|^2 - |\tilde{\phi}|^2)] \log \frac{[M^2 + \lambda^2 (|\tilde{\Phi}_1|^2 - |\tilde{\phi}|^2)]}{\mu^2} \Big\} \tag{5.22}
$$

This expression can be expanded into a power series in $1/M$ as

$$
K_1^{(1)} = -\frac{1}{32\pi^2} \Big\{ \lambda^2 (|\tilde{\Phi}|^2 - |\tilde{\phi}|^2)(1 + \log \frac{M^2}{\mu^2}) + \frac{1}{2M^2} \lambda^4 (|\tilde{\Phi}|^2 - |\tilde{\phi}|^2)^2 \Big\} + O(\frac{1}{M^4}) (5.23)
$$

Now we turn to the calculation of $\Gamma_2^{(1)}$. Using [\(5.14](#page-35-0)) it is easy to see that

$$
K_2^{(1)} = \frac{i}{2}tr \log \Delta \tag{5.24}
$$

where

$$
\Delta = \Box + (\frac{1}{\xi} - 1)\frac{1}{16}\{D^2, \bar{D}^2\} + g^2(4\tilde{\Phi}_1\tilde{\Phi}_1 + 4\tilde{\Phi}_2\tilde{\Phi}_2 + \tilde{\phi}\tilde{\bar{\phi}})
$$
(5.25)

Using a proper-time method (see Section 4) one derives

$$
\Gamma_2^{(1)} = \frac{i}{2} \text{tr} \int_0^\infty \frac{ds}{s} \exp(is\Delta)
$$
\n(5.26)

and a straightforward computation yields

$$
\exp(is\Delta) = \exp(is\Delta) \exp(isA)(1 + \sum_{n=1}^{\infty} \frac{is\Box(\frac{1}{\xi} - 1)^n}{n!} \frac{1}{16\Box} \{D^2, \bar{D}^2\})
$$

where $A = g^2(4\tilde{\Phi}_1\tilde{\bar{\Phi}}_1 + 4\tilde{\Phi}_2\tilde{\bar{\Phi}}_2 + \tilde{\phi}\tilde{\bar{\phi}})$. A subsequent calculation can be carried in a completely parallel fashion as that given in Appendix A. As a result we arrive at

$$
K_2^{(1)} = \frac{1}{32\pi^2} \int_0^\infty \frac{ds}{s^{2-\epsilon}} \exp(isA) = \frac{1-\xi}{16\pi^2} A^{1-\epsilon} (\frac{2}{\epsilon} - \gamma + 1)
$$
(5.27)

Here we introduced the dimensional regularization parameter ϵ since the integral is divergent. We also put $\xi = 0$, i.e. we work in the Landau gauge. Expanding this expression into a power series in ϵ we arrive at

$$
K_2^{(1)} = \frac{1}{32\pi^2} \left(\frac{2}{\epsilon} \left(g^2 (4\tilde{\Phi}_1 \tilde{\Phi}_1 + 4\tilde{\Phi}_2 \tilde{\Phi}_2 + \tilde{\phi} \tilde{\phi}) \right) - \right. \\
\left. - \left\{ g^2 (4\tilde{\Phi}_1 \tilde{\Phi}_1 + 4\tilde{\Phi}_2 \tilde{\Phi}_2 + \tilde{\phi} \tilde{\phi}) \log \frac{g^2 (4\tilde{\Phi}_1 \tilde{\Phi}_1 + 4\tilde{\Phi}_2 \tilde{\Phi}_2 + \tilde{\phi} \tilde{\phi})}{\mu^2} - \gamma + 1 \right\} \right. \tag{5.28}
$$

The divergent part can be cancelled via a suitable counterterm which leads to the renormalized correction of the form

$$
K_{2r}^{(1)} = -\frac{1}{32\pi^2} [g^2 (4\tilde{\bar{\Phi}}_1 \tilde{\Phi}_1 + 4\tilde{\bar{\Phi}}_2 \tilde{\Phi}_2 + \tilde{\bar{\phi}}_2 \tilde{\phi})] \log \frac{g^2 (4\tilde{\bar{\Phi}}_1 \tilde{\Phi}_1 + 4\tilde{\bar{\Phi}}_2 \tilde{\Phi}_2 + \tilde{\bar{\phi}}_2 \tilde{\phi})}{\mu^2} - \gamma + 1) \tag{5.29}
$$

Hence the final result for the one-loop kählerian effective potential rewritten in terms of standard background chiral superfields Φ_1, Φ_2, ϕ and background gauge superfield V is

$$
K^{(1)} = K_{1r}^{(1)} + K_{2r}^{(1)} =
$$

= $-\frac{1}{32\pi^2} \left\{ \lambda^2 (\bar{\Phi}_1 e^{2gV} \Phi_1 - \bar{\phi} e^{-gV} \phi)(1 + \log \frac{M^2}{\mu^2}) + \frac{1}{2M^2} \lambda^4 (\bar{\Phi}_1 e^{2gV} \Phi_1 - \bar{\phi} e^{-gV} \phi)^2 +$
+ $[g^2 (4\bar{\Phi}_1 e^{2gV} \Phi_1 + 4\bar{\Phi}_2 e^{-2gV} \Phi_2 + \bar{\phi} e^{-gV} \phi) \times$
 $\times (\log \frac{g^2 (4\bar{\Phi}_1 e^{2gV} \Phi_1 + 4\bar{\Phi}_2 e^{-2gV} \Phi_2 + \bar{\phi} e^{-gV} \phi)}{\mu^2} - \gamma + 1)] \right\}$ (5.30)

The effective equations of motion for massive superfields (with the action([5.1](#page-32-0)) and the quantum correction $K^{(1)}$ (5.30)) are of the form

$$
- \frac{\bar{D}^2}{4} (e^{2gV} \bar{\Phi}_1 + \frac{\partial K^{(1)}}{\partial \Phi_1}) + M \Phi_2 + \beta \phi^2 = 0
$$

$$
- \frac{\bar{D}^2}{4} (e^{-2gV} \bar{\Phi}_2 + \frac{\partial K^{(1)}}{\partial \Phi_2}) + M \Phi_1 = 0
$$
(5.31)

Here $\beta = \lambda(1 + \frac{\hbar g^2 \alpha}{16\pi^2})$ where

$$
\alpha = \int_0^1 dx \frac{\log[x(1-x)]}{1-x(1-x)}
$$

(see the next subsection) is a one-loop corrected coupling for the chiral interaction. (Quantum corrections to the chiral potential are considered in details in the next subsection.) Solutions of these equations have the form

$$
\Phi_1 = \frac{\bar{D}^2}{4M^2} \Big(\beta \bar{\phi}^2 e^{-2gV} \big[1 - 4 \frac{g^2 \hbar}{32\pi^2} \big(1 + \log \frac{g^2 \bar{\phi} e^{-gV} \phi}{\mu^2} \big) \big] \Big) + O(\frac{1}{M^3})
$$
\n
$$
\Phi_2 = -\frac{\beta \phi^2}{M} + O(\frac{1}{M^3})
$$
\n(5.32)

As a result it turns to be that the one-loop contribution to the kählerian effective potential can be expressed as

$$
K^{(1)} = -\frac{1}{32\pi^2} \left\{ -\lambda^2 \bar{\phi} e^{-gV} \phi (1 + \log \frac{M^2}{\mu^2}) + \right. \\
\left. + g^2 \bar{\phi} e^{-gV} \phi \log \frac{g^2 \bar{\phi} e^{-gV} \phi}{\mu^2} \right\} + O(\frac{1}{M^2})
$$
\n(5.33)

Here $\gamma + 1$ is absorbed into a redefinition of μ . It is easy to see that this correction is gauge invariant. We note that the quantum correction arises in the effective action of light superfields that depend on the gauge coupling. The expression [\(5.33](#page-38-0)) is quite analogous to the minimal model of gauge neutral chiral superfields where the self-interaction of the light chiral superfield is absent. Namely, the only contribution due to the gauge interactions (i.e. terms proportional to the gauge coupling g) corresponds to the standard Coleman-Weinberg potential (the second term, in the square brackets), which is due to the gauge interactions of the light fields only.

The redefinition $\phi \to \tilde{\phi} = Z^{1/2} \phi$ where

$$
Z = 1 + \lambda^2 \frac{\hbar}{32\pi^2} (1 + \log \frac{M^2}{\mu^2})
$$
\n(5.34)

in turn yields the action corresponding to the sector of chiral superfields only (i.e. with the background gauge superfields put to zero) and can be written as

$$
\Gamma = \int d^8 z |\tilde{\phi}|^2 + O(\frac{1}{M^2})\tag{5.35}
$$

This result is again formally consistent with the decoupling theorem. However, since all parameters (fields, masses, couplings) are determined from string theory it turns out to be that the classical action gets an essential quantum correction given by([5.33\)](#page-38-0).

5.3 Chiral potential corrections

We now turn to the study of possible chiral corrections to the effective action. It turns out that unlike in the model with interacting gauge neutral chiral superfields, where the first set of quantum chiral correction arises at the two-loop level (see Section 3), in $N = 1$ supersymmetric gauge theories the one-loop corrections to the chiral effective potential are possible. This situation is quite analogous to standard $N = 1$ super - Yang - Mills theory with chiral matter[[24](#page-56-0)]. However, we note that in the present theory chiral corrections to effective action depend on the massive superfield Φ_1 .

In order to study chiral corrections to the effective chiral potential action it is more suitable to use a Feynman gauge $\xi = 1$. In this gauge the propagator for the gauge superfield reads as

$$
\langle vv \rangle = \frac{1}{\Box} \delta^8 (z_1 - z_2) \tag{5.36}
$$

First, the chiral effective potential depends only on the chiral background superfields ϕ , Φ . Second, we note that diagrams containing massive propagators cannot contribute to the chiral effective potential (see Section 3). Therefore possible one-loop diagrams can contain only vertices of the type $\bar{\phi}v\phi$ where ϕ is a background superfield, and those of the type $\lambda\Phi_1\phi^2$ where Φ_1 is a background superfield. Let the number of vertices $\bar{\phi}\nu\phi$ be N_1 , and the number of $\lambda \Phi_1 \phi^2$ be N_2 . The number of quantum superfields ϕ and $\bar{\phi}$ is equal to $2N_2$, and N_1 , respectively. Since the only Green function for massless superfields is $\langle \phi \phi \rangle$, these two numbers should be equal, $2N_2 = N_1$. Each vertex $\lambda \Phi_1 \phi^2$ corresponds to one factor \bar{D}^2 since

it is purely chiral, and each vertex $\bar{\phi}v\phi$ with ϕ as the external field corresponds to one factor D^2 (see e.g., [\[20\]](#page-56-0)). Then, a diagram that can contribute to the chiral effective potential has to have a number of factors D^2 that is one larger than the number of factors \bar{D}^2 [[27\]](#page-56-0), i.e. $N_1 - N_2 = 1$. Therefore we conclude that $N_1 = 2$, $N_2 = 1$. The only supergraph with such a structure is given below:

This supergraph yields the following contribution

$$
J = \lambda g^2 \int d^4 \theta_1 d^4 \theta_2 d^4 \theta_3 \int \frac{d^4 k}{(2\pi)^4} \frac{1}{k^2 (k - p_1)^2 (k - p_2)^2} \Phi(p_1 + p_2, \theta_1) \phi(-p_1, \theta_2) \phi(-p_2, \theta_3) \times \times \delta_{12} \frac{\bar{D}_1^2 D_3^2}{12} \delta_{13} \frac{D_2^2}{4} \delta_{23}
$$
\n(5.37)

After D-algebra transformation, integration over momenta and Fourier transformation this contribution has in the infrared limit $p_1, p_2 \to 0$ the form (cf. [\[24](#page-56-0)])

$$
J = \frac{1}{16\pi^2} \lambda g^2 \beta \int d^6 z \Phi_1(z) \phi^2(z) \tag{5.38}
$$

where

$$
\alpha = \int_0^1 dx \frac{\log[x(1-x)]}{1-x(1-x)}
$$

is a constant encountered in[[24](#page-56-0)]. (Note the same constant is encountered in the previous subsection.) This contribution coincides in form with the classical chiral potential. The one-loop corrected chiral effective potential is then of the form:

$$
W^{(1)} = (\lambda + \frac{\hbar}{16\pi^2} \lambda g^2 \alpha) \Phi_1(z) \phi^2(z)
$$
\n(5.39)

However, since the heavy superfield Φ is at least of first order in the inverse mass (as a solution of the effective equations of motion) this term is also of first order in the inverse mass. Nevertheless, this result demonstrates that quantum corrections to the chiral effective potential can depend not only on massless superfields but also on massive ones. (It was commonly believed that quantum corrections to the chiral effective potential can depend only on massless superfields.)

5.4 Strength depending contributions in the effective action

In this subsection, we are going to consider quantum corrections to the gauge function, i.e. a holomorphic function that multiplies W^2 in the leading low-energy approximation to the effective action. To simplify the consideration we again use the Langau gauge $\xi = 0$. The loops can contain only either propagators for chiral superfields or propagators for vector superfields. However, since the external vector superfield can connect only to the vertex of the form $v\Phi_i\bar{\Phi}_i$ (or $v\phi\bar{\phi}$) the possible supergraphs must necessarily contain propagators for the chiral superfields and therefore at the one-loop level, only propagators for chiral superfields contribute. (We also note that in principle it is possible to study corrections proportional to W^2 in a general gauge.)

Now, we turn to the study of possible supergraphs with two external gauge lines. First, we note that the vertex $g\bar{\Phi}_2 v \Phi_2$ can be associated either with the propagator $\langle v v \rangle$ or with the external vector superfield line. However, as noted above, in the Landau gauge the loops of the form under consideration cannot contain propagators $\langle v v \rangle$. Therefore the only contributing supergraph containing the vertices $v\Phi_2\bar{\Phi}_2$ is of the form

− − − − D² D¯ ² D¯ ² D²

Recall that the dashed line denotes the propagator $\langle \Phi_2 \bar{\Phi}_2 \rangle$ (the same on as for $\langle \Phi_1 \bar{\Phi}_1 \rangle$), and the wavy line denotes the propagator $\langle v v \rangle$. The contribution of the diagram has the form

$$
I = 4g^2 \int \frac{d^4k}{\left(2\pi\right)^4} \int d^4\theta_1 d^4\theta_2 V(\theta_1) V(\theta_2) \frac{1}{\left(k^2 + M^2\right)^2} \frac{D_1^2 \bar{D}_2^2}{16} \delta_{12} \frac{\bar{D}_1^2 D_2^2}{16} \delta_{12} \tag{5.40}
$$

After D-algebra transformations and dimensional regularization this expression is equal to

$$
I = \frac{4g^2}{16\pi^2} \int d^2\theta \frac{1}{64} W^2 \int \frac{d^n k}{(2\pi)^n} \frac{1}{(k^2 + M^2)^2} =
$$

=
$$
\frac{4g^2}{16\pi^2} \int d^2\theta \frac{1}{64} W^2 \left(\frac{2}{\epsilon} - \gamma + \log \frac{M^2}{\mu^2}\right)
$$
(5.41)

where ϵ is the usual dimensional regularization parameter. The divergent part can be cancelled via a suitable counterterm, and the resulting correction is

$$
I = \frac{4g^2}{16\pi^2} \frac{1}{64} \int d^2\theta W^2 \log \frac{M^2}{\mu^2}
$$
 (5.42)

The Euler constant γ is here absorbed into a redefinition of the normalization parameter μ . As a result we see that this correction leads to a term proportional to $\log \frac{M^2}{\mu^2}$ which evidently increases logarithmically with M as $M \to \infty$.

Now we turn to the study of corrections proportional to $W²$ which do not contain propagators of the field Φ_2 . First of all, let us consider corrections at the zeroth order in the inverse mass expansion. Since the heavy superfields Φ_1 , Φ_2 (after solving the effective equations of motion) correspond to expressions that are of the first and/or higher orders in the inverse mass expansion (see the discussions above), the corrections that are of the zeroth order in 1 $\frac{1}{M}$ expansion therefore correspond to diagrams with massless external chiral superfields only. Such corrections are described by the supergraphs

Here the bold dashed line (in the following we will denote it as G_1) is a propagator of the form

and the bold wavy line (in the following we will denote it as G_2) is a propagator of the form

External lines denote superfields ϕ .

Consequently, the contributions are of the form

$$
G_1 = \langle \Phi_1 \bar{\Phi}_1 \rangle + \lambda^2 |\phi|^2 \langle \Phi_1 \bar{\Phi}_1 \rangle \langle \phi \bar{\phi} \rangle \langle \Phi_1 \bar{\Phi}_1 \rangle + \dots + \langle \Phi_1 \bar{\Phi}_1 \rangle (\lambda^2 |\phi|^2)^n (\langle \phi \bar{\phi} \rangle \langle \Phi_1 \bar{\Phi}_1 \rangle)^n + \dots
$$

\n
$$
G_2 = \langle \phi \bar{\phi} \rangle + \lambda^2 |\phi|^2 \langle \phi \bar{\phi} \rangle \langle \Phi_1 \bar{\Phi}_1 \rangle \langle \phi \bar{\phi} \rangle + \dots + \langle \phi \bar{\phi} \rangle (\lambda^2 |\phi|^2)^n (\langle \Phi_1 \bar{\Phi}_1 \rangle \langle \phi \bar{\phi} \rangle)^n + \dots
$$
\n(5.43)

The superpropagators $\langle \phi \phi \rangle$, and $\langle \Phi_1 \bar{\Phi}_1 \rangle$ are given by [\(5.4](#page-33-0)) and [\(5.18\)](#page-36-0), respectively.

A straightforward calculation yields

$$
G_1 = -\frac{D^2 \bar{D}^2}{16k^2} \frac{k^2 + M^2}{k^2 + M^2 + |\phi|^2}
$$

\n
$$
G_2 = -\frac{D^2 \bar{D}^2}{16(k^2 + M^2 + |\phi|^2)}
$$
\n(5.44)

As a result, contributions from the diagrams shown in Fig.2 and Fig.3 correspond to

$$
I_2 = g^2 \int \frac{d^4k}{(2\pi)^4} \int d^4\theta_1 d^4\theta_2 V(\theta_1) V(\theta_2) \frac{(k^2 + M^2)^2}{(k^2(k^2 + M^2 + |\phi|^2))^2} \frac{D_1^2 \bar{D}_2^2}{16} \delta_{12} \frac{\bar{D}_1^2 D_2^2}{16} \delta_{12}
$$

\n
$$
I_3 = 4g^2 \int \frac{d^4k}{(2\pi)^4} \int d^4\theta_1 d^4\theta_2 V(\theta_1) V(\theta_2) \frac{1}{(k^2 + M^2 + |\phi|^2)^2} \frac{D_1^2 \bar{D}_2^2}{16} \delta_{12} \frac{\bar{D}_1^2 D_2^2}{16} \delta_{12} \quad (5.45)
$$

respectively. After D-algebra transformations, integration over momenta, and a subtraction of divergences we find

$$
I_2 = \frac{g^2}{16\pi^2} \frac{1}{64} \int d^2\theta W^2 (\log \frac{M^2 + |\phi|^2}{\mu^2} + A)
$$

\n
$$
I_3 = \frac{4g^2}{16\pi^2} \frac{1}{64} \int d^2\theta W^2 (\log \frac{M^2 + |\phi|^2}{\mu^2})
$$
\n(5.46)

Here $A = 1 + \int_{\alpha}^{1} dx \frac{1-x}{x}$ is a constant that depends on the infrared regularization parameter α . We note that this parameter can also be absorbed into a redefinition of the renormalization parameter μ .

The total correction in the gauge sector is a sum of corrections I_1 ([5.42\)](#page-41-0), I_2 and I_3 [\(5.46](#page-42-0)). It has the form

$$
I = I_1 + I_2 + I_3 = \frac{g^2}{64} \frac{1}{16\pi^2} \int d^2\theta W^2 (4 \log \frac{M^2}{\mu^2} + (\log \frac{M^2 + |\phi|^2}{\mu^2} + A) +
$$

+
$$
4 \log \frac{M^2 + |\phi|^2}{\mu^2})
$$
 (5.47)

The leading term of this correction is

$$
I = \frac{9g^2}{64} \frac{1}{16\pi^2} \int d^2\theta W^2 \log \frac{M^2}{\mu^2} + O\left(\frac{1}{M^2}\right) \tag{5.48}
$$

Here the constant A is absorbed into a redefinition of the normalization parameter μ .

Let us now consider contributions that depend on external heavy chiral superfields. We assume that the supergraph can contain n_1 vertices of the form $\Phi_1 \phi_q^2$, \bar{n}_1 vertices of the form $\bar{\Phi}_1 \bar{\phi}_q^2$, n_2 vertices of the form $\phi \Phi_{1q} \phi_q$, and \bar{n}_2 vertices of the form $\bar{\phi} \vec{\Phi}_{1q} \bar{\phi}_q$. Thus, the number of external Φ_1 is equal to n_1 , and the number of external Φ is equal to \bar{n}_1 . We note that there are no propagators of heavy superfields Φ_2 in the supergraphs under consideration (in the Landau gauge), therefore the only possible supergraphs are $\langle \phi \phi \rangle$ and $\langle \Phi_1 \bar{\Phi}_1 \rangle$. Hence the number of quantum superfields ϕ and $\bar{\phi}$, and correspondingly, the number of Φ_{1q} and $\bar{\Phi}_{1q}$ must be equal, and we arrive at the relations $n_2 = \bar{n}_2$, $2n_1 + n_2 = 2\bar{n}_1 + \bar{n}_2$. These relations leads us to conclude that $n_1 = \bar{n}_1$ and therefore contributions of supergraphs must be proportional to $(\Phi_1 \bar{\Phi}_1)^{n_1}$. Since the solution of the effective equations of motion yields the result that the heavy superfield Φ_1 is at least of the second order in $\frac{1}{M}$ (see above) we conclude that terms with a non-trivial dependence on the background field Φ_1 must be of the fourth order in $\frac{1}{M}$.

As a result, the one-loop corrected effective action is the gauge sector if of the form:

$$
\Gamma^{(1)}[W] = \frac{1}{64} \int d^6 z W^2 (1 + 9g^2 \frac{\hbar}{16\pi^2} \log \frac{M^2}{\mu^2}) + O\left(\frac{1}{M^4}\right)
$$
(5.49)

This expression contains terms logarithmically increasing with M and thus the gauge function f can get significant corrections due to the contributions of the heavy fields. Note, however, that these corrections can be reinterpreted as the standard one-loop threshold corrections due to the contributions of the heavy fields. Namely, these leading effects are due to the coupling of the vector gauge fields to the heavy chiral superfields, and are therefore proportional to the gauge coupling g. On the other hand a contribution due to the coupling λ between light and heavy chiral superfields is absent at this order. (It appears only at higher orders in the 1 $\frac{1}{M}$ expansion.)

Again it is straightforward to show that the obtained result is in compliance with the decoupling theorem. We redefine the vector superfield V (and consequently the superfield strength W^{α}) by the rule $\tilde{V} = Z_1^{1/2}V$ where Z_1 is a finite renormalization constant

$$
Z_1 = 1 + 9g^2 \frac{\hbar}{16\pi^2} \log \frac{M^2}{\mu^2}
$$
 (5.50)

Then the expression [\(5.49](#page-43-0)) takes the form

$$
\Gamma^{(1)}[W] = \frac{1}{64} \int d^6 z W^2 + O\left(\frac{1}{M^2}\right) \tag{5.51}
$$

which is again of the same form as the corresponding term in the classical action. On the other hand the terms that include the interaction between the vector superfield and the chiral ones should stay invariant under such a redefinition of fields. Therefore we must redefine the gauge coupling g by the rule $\tilde{g} = Z_1^{-1/2}g$. After these redefinitions the one-loop effective action takes the form

$$
\Gamma^{(1)} = \frac{1}{64} \int d^6 z \tilde{W}^2 + \int d^8 z \tilde{\bar{\phi}} e^{-\tilde{g}\tilde{V}} \tilde{\phi} + O(\frac{1}{M})
$$
\n(5.52)

As a result the one-loop corrected effective action is of the same form as the classical action of light (massless) superfields ϕ, v , only, as expected by the decoupling theorem. However, since the values of fields and couplings are determined from string theory, they cannot be redefined, and the final result for the one-loop corrected effective action in the sector of gauge superfields is thus given by [\(5.49\)](#page-43-0) which allows us to conclude that the classical action in this sector gets essential corrections. couplings

6 Summary

In this paper we presented a systematic analysis of the decoupling effects in $N = 1$ supersymmetric theories with chiral superfields. We developed techniques to evaluate explicitly the one-loop corrected actions that involve both the heavy and light chiral superfields for different types of classical kählerian and chiral potentials. By a subsequent elimination of the heavy chiral superfields by their equations of motion, via an iterative procedure, the resulting effective actions of light chiral superfields include the quantitative decoupling corrections and the one-loop level. We considered different examples of theories describing dynamics of interacting light (massless) and heavy superfields. For these theories the one-loop kählerian effective potential was calculated and subsequently the heavy superfields were expressed via light ones up to a certain order in the inverse (heavy) mass parameters, thus resulting in an explicit form of the effective action (up to a certain order in the inverse mass expansion) for light superfields, only.

As a representative example we carried out a detailed analysis for the so-called minimal model. This model contains two (gauge neutral) chiral superfields, one heavy and one massless, while the classical kählerian potential contains only the renormalizable (canonical) terms and the classical chiral potential is renormalizable with the self-interaction term of the light fields and an interaction term linear in the heavy field. The results of these calculations demonstrate that the leading order decoupling effects at the loop-level grow logarithmically with the heavy mass scale. In addition, we also analysed the loop corrections to the chiral potential, and found that at the two-loop level corrections to the heavy fields do appear, which are, however, suppressed in the effective action of light fields by inverse powers of the heavy mass scales.

In the subsequent section we carried out the analysis of the non-minimal models with a more general structure of the kählerian and chiral potentials, that involve one light and one massive field. While the analysis of these models cannot be carried out explicitly for a general choice of the classical kählerian and chiral potential, we developed techniques to determine the effective action as a series expansion in the inverse powers of the heavy mass parameter. We have also addressed the decoupling effects for $N = 1$ supersymmetric gauge theories with heavy and light chiral fields charged under the gauge group. In particular, we analysed the $U(1)$ -Abelian (minimal) model with charged heavy and light chiral superfields, a minimal (gauge invariant) käherian potential and a chiral potential with an interaction term linear in a heavy field. The leading corrections to the kählerian potential are independent of the gauge coupling, however higher order terms (both for the one-loop corrected kählerian and chiral potential) are proportional to the gauge coupling. In addition we also demonstrated that the leading correction to the gauge function is proportional to the gauge coupling, which can be reinterpreted as a standard threshold effect due to the heavy charged chiral superfields. (Note however that in the leading order the gauge function corrections do not depend on the chiral coupling of the heavy chiral superfields to the light ones.)

While our specific results are, of course, in agreement with the decoupling theorem, the actual quantitative form of the decoupling effects may have important physics implications. As it was stressed in the introduction, in an effective field theory (without reference to an underlying fundamental theory) the couplings are free parameters, fixed by (low-energy) experiments, and thus the above decoupling effects can indeed be all absorbed into rescalings of the effective light fields. However, within string theory the original couplings are calculable at the string scale and the corrections due to decoupling can now have sizable calculable predictions for the low energy couplings. The results of our calculations explicitly confirm that in $N = 1$ supersymmetric theories the decoupling effects at loop-level are of order log M, and in specific cases they can significantly modify the low energy predictions of the effective couplings.

In particular, within a class of perturbative string models with an anomalous $U(1)$, after the vacuum restabilization, there are in general renormalizable interactions between the light and heavy fields, with mass of order $M \sim M_{string} \sim (10^{17})$ GeV. Thus, the above oneloop decoupling effects can significantly change the low energy predictions for the couplings at low energies, e.g., the electro-weak scale ($\mu \sim 1 \text{ TeV}$). Let us repeat the analysis for an example, in a class of string models discussed in [\[10](#page-55-0)]–[\[18](#page-56-0)], where typical values of the couplings, calculated at M_{string} , are $\lambda = g = g_{gauge}$. Here $g_{gauge} \sim 0.8$ is the value of the gauge coupling at at M_{string} . Renormalization group equations then determine the values of λ and g couplings at low energies μ . However, due to the one-loop decoupling effects there is now also an additional correction in the effective coupling g for the light fields; it is of the order of $\lambda(\mu)^2/(16\pi^2)\log M^2/\mu^2 \sim 0.26$. (We used the typical values $M \sim M_{string} \sim 10^{17}$

GeV, $\mu \sim 1$ TeV and $\lambda(\mu) \sim 0.8$.) This specific example indicates that for a class of fourdimensional string vacua the actual prediction for the tri-linear couplings at low energy could be corrected by $10 - 50\%$.

Let us conclude with a remark about the leading contributions at higher loop levels. These contributions are expected to be of the form (cf.[\[26\]](#page-56-0)): the form

$$
|\phi|^2\log^n\big(\frac{|\phi|^2}{\mu^2}\big).
$$

When choosing a renormalization condition of the form $\mu^2 = const \times M^2$ (analogous to the one used in Section 3) the typical contribution of these terms would be of the type

$$
|\phi|^2\log^n{(\frac{|\phi|^2}{M^2})}
$$

thus implying that the decoupling effects at the higher loop leading decoupling effects could be just as significant, and they would further modify low-energy predictions of the effective theory of light fields.

Acknowledgments. M.C. acknowledges discussions with L. Everett and P. Langacker. The work by I.L.B. and A.Yu.P. was supported in part by INTAS grant, INTAS-96-0308; RFBR grant, project No 99-02-16617; RFBR-DFG grant, project No 99-02-04022; GRACE-NAS grant, project No 97-6.2-34. The work of M.C. was supported in part by U.S. Department of Energy Grant No. DOE-EY-76-02-3071. I.L.B. and M.C. would like to thank the organizers of the 32nd International Symposium Ahrenshoop on the Theory of Elementary Particles, where the work was initiated, for hospitality.

Appendix A

To find an exact form of the solution of([3.4\)](#page-13-0) for the model([3.3\)](#page-12-0) it is more convenient to study a system of equations for components of matrices A and C . Therefore we arrive at

$$
\frac{1}{i}\dot{A}_{11} = -C_{11}(\lambda\Phi + g\phi) - C_{12}\lambda\Phi
$$
\n
$$
\frac{1}{i}\dot{C}_{11} = (\lambda\bar{\Phi} + g\bar{\phi}) - \frac{1}{i}A_{11}(\lambda\bar{\Phi} + g\bar{\phi})\Box - A_{12}\lambda\bar{\phi}\Box
$$
\n
$$
\frac{1}{i}\dot{A}_{12} = -C_{11}\lambda\phi - C_{12}M
$$
\n
$$
\frac{1}{i}\dot{C}_{12} = \lambda\bar{\phi} - A_{11}\bar{\phi}\Box - A_{12}M\Box
$$
\n
$$
\frac{1}{i}\dot{A}_{21} = -C_{21}(\lambda\Phi + g\phi) - C_{22}\lambda\phi
$$
\n
$$
\frac{1}{i}\dot{C}_{21} = \lambda\bar{\phi} - A_{21}(\lambda\Phi + g\phi)\Box - A_{22}\lambda\bar{\phi}\Box
$$
\n
$$
\frac{1}{i}\dot{A}_{22} = -C_{21}\lambda\phi - C_{22}M
$$
\n
$$
\frac{1}{i}\dot{C}_{22} = M - A_{21}\lambda\bar{\phi}\Box - A_{22}M\Box
$$
\n(A.1)

with an analogous system of equations for \tilde{A}, \tilde{C} . It is straightforward to solve this system of equation by the method described described in subsection 2.3. It allows one to obtain the following form of the coefficients A_{11} , A_{22} , \tilde{A}_{11} , \tilde{A}_{22} (i.e. only those contribute to the $tr \log \Delta$):

$$
A_{11} = \tilde{A}_{11} = \frac{N_3}{(N_3 - N_1)\square} \cosh(i\omega_1 s) - \frac{N_1}{(N_3 - N_1)\square} \cosh(i\omega_2 s) - \frac{1}{\square}
$$

\n
$$
A_{22} = \tilde{A}_{22} = \frac{N_1}{(N_1 - N_3)\square} \cosh(i\omega_1 s) - \frac{N_3}{(N_1 - N_3)\square} \cosh(i\omega_2 s) - \frac{1}{\square}
$$
\n(A.2)

Here $\omega_1 = \sqrt{P + \sqrt{Q}}\sqrt{\Box}$, $\omega_2 = \sqrt{P - \sqrt{Q}}\sqrt{\Box}$, $N_1 = \frac{\omega_1^2 - (|\lambda \Phi + g\phi|^2 + 2\lambda^2 |\phi|^2)\Box}{\lambda \bar{\phi}(\lambda \Phi + g\phi) + M\lambda \phi}$, $N_3 = \frac{\omega_1^2 - (|\lambda \Phi + g\phi|^2 + 2\lambda^2 |\phi|^2) \Box}{\lambda \bar{\phi}(\lambda \Phi + g\phi) + M\lambda \phi}$, and

$$
P = |\lambda \Phi + g\phi|^2 + 2\lambda^2 |\phi|^2 + M^2
$$

\n
$$
Q = (|\lambda \Phi + g\phi|^2 - M^2)^2 + 4|(\lambda \Phi + g\phi)\lambda \overline{\phi} + \lambda \phi M|^2
$$
\n(A.3)

Let us denote

$$
R_1 = \sqrt{P + \sqrt{Q}}, \ R_2 = \sqrt{P - \sqrt{Q}} \tag{A.4}
$$

The trace of $e^{is\tilde{\Delta}}$ is determined by the matrix trace of A which reads as

$$
tr A = A_{11} + A_{22} = \frac{1}{\Box} (\cosh(isR_1\sqrt{\Box}) + \cosh(isR_2\sqrt{\Box}) - 2)
$$
 (A.5)

The one-loop kählerian effective potential has the form

$$
K^{(1)} = -\frac{i}{2} \int_0^\infty \frac{ds}{s} (A_{11}(s) + A_{22}(s)) U(x, x'; s)|_{x=x'} \tag{A.6}
$$

since $A_{11} = \tilde{A}_{11}$, $A_{22} = \tilde{A}_{22}$. Here $U(x, x'; s) = e^{is\Box} \delta^4(x - x')$ (cf. [\[22](#page-56-0), [23](#page-56-0)]). Expanding $A_{11}(s) + A_{22}(s)$ as a power series in \Box we find the one-loop kählerian effective potential in the form

$$
K^{(1)} = -\frac{i}{2} \int_0^\infty \frac{ds}{s} \sum_{n=0}^\infty \left(\frac{(iR_1)^{2n+2}}{(2n+2)!} + \frac{(iR_2)^{2n+2}}{(2n+2)!} \right) s^{2n+2} \Box^n U(x, x'; s)|_{x=x'}
$$
\n(A.7)

As usual (see [\[22, 23\]](#page-56-0))

$$
\Box^{n}U(x, x'; s)|_{x=x'} = \frac{(-1)^{n}(n+1)!}{16\pi^{2}(is)^{n+2}}
$$

Thus the one-loop kählerian effective potential can be cast in the form

$$
K^{(1)} = -\frac{1}{32\pi^2} \int_0^\infty \frac{ds}{s} \sum_{n=0}^\infty \left(\frac{R_1^{2n+2}}{(2n+2)!} + \frac{R_2^{2n+2}}{(2n+2)!} \right) (n+1)! (-1)^n \frac{(is)^{2n+2}}{(is)^{n+2}} \tag{A.8}
$$

Since this integral is divergent we use the dimensional regularization by introducing the regularization parameter ϵ with the subsequent changes: $\int_0^\infty \frac{ds}{s} \to \int_0^\infty \frac{ds}{s^{1-\epsilon}}$ $rac{ds}{s^{1-\epsilon}}$. As a result we arrive at

$$
K^{(1)} = -\frac{1}{32\pi^{2}} \Big[R_{1}^{2(1+\epsilon)} \int_{0}^{\infty} \frac{d(isR_{1}^{2})}{(isR_{1}^{2})^{1-\epsilon}} \sum_{n=0}^{\infty} \frac{(isR_{1}^{2})^{n}(n+1)!(-1)^{n}}{(2n+2)!} + R_{2}^{2(1+\epsilon)} \int_{0}^{\infty} \frac{d(isR_{2}^{2})}{(isR_{2}^{2})^{1-\epsilon}} \sum_{n=0}^{\infty} \frac{(isR_{2}^{2})^{n}(n+1)!(-1)^{n}}{(2n+2)!} \Big] \tag{A.9}
$$

Using (cf. [\[22](#page-56-0), [23](#page-56-0)])

$$
\sum_{n=1}^{\infty} s^n \frac{(-1)^{n-1} n!}{(2n)!} = s \int_0^1 dt e^{-\frac{1}{4}s(1-t^2)}
$$

one obtains

$$
K^{(1)} = -\frac{1}{16\pi^2} \left\{ (R_1^2)^{1+\epsilon} \int_0^\infty \frac{ds}{s^{1-\epsilon}} \int_0^1 dt e^{-\frac{1}{4}s(1-t^2)} + (R_2^2)^{1+\epsilon} \int_0^\infty \frac{ds}{s^{1-\epsilon}} \int_0^1 dt e^{-\frac{1}{4}s(1-t^2)} \right\}
$$
(A.10)

Here, we have made a redefinition $s \to sR_1^2$ in the first term and $s \to sR_2^2$ in the second one along with a subsequent Wick rotation. Furthermore

$$
(R_1^2)^{1+\epsilon} \int_0^\infty \frac{ds}{s^{1-\epsilon}} \int_0^1 dt e^{-\frac{1}{4}s(1-t^2)} = (R_1^2)^{1+\epsilon} \Gamma(\epsilon) \int_0^1 dt (1 - \epsilon \log(\frac{1-t^2}{4}) + O(\epsilon^2)) =
$$

= $(R_1^2)^{1+\epsilon} [\frac{1}{\epsilon} + \gamma - \int_0^1 dt \log(\frac{1-t^2}{4}) + O(\epsilon)]$ (A.11)

Here $O(\epsilon)$ denotes terms of order one and/or higher in ϵ , and $O(\epsilon^2)$ denotes terms of orders two and/or higher in ϵ . Therefore

$$
K^{(1)} = -\frac{1}{32\pi^2} \left\{ \frac{R_1^2 + R_2^2}{\epsilon} + R_1^2 \log \frac{R_1^2}{\mu^2} + R_2^2 \log \frac{R_2^2}{\mu^2} \right\}
$$
(A.12)

Here the first term contains all one-loop divergences of the theory which can be cancelled by a counterterm of the form

$$
K_{counter}^{(1)} = \frac{1}{32\pi^2 \epsilon} (R_1^2 + R_2^2)
$$
\n(A.13)

The terms containing γ and $\int_0^1 dt \log(\frac{1-t^2}{4})$ $\frac{-t^2}{4}$ are removed by a suitable choice of the renormalizationparameter μ . Using the exact expression for R_1 and R_2 (([A.3\)](#page-47-0) and ([A.4\)](#page-47-0), respectively) we arrive at the result([3.5\)](#page-13-0).

Appendix B

We note that the result (3.5) for the one-loop correction in the kählerian effective potential can also be obtained via a diagram technique.

Let us study possible diagrams contributing to the one-loop kählerian effective potential in this model. The propagators of the superfields have the standard form[[20\]](#page-56-0)

$$
\langle \phi(z_1)\bar{\phi}(z_2) \rangle = -\frac{1}{\Box} \delta^8(z_1 - z_2) \n\langle \Phi(z_1)\bar{\Phi}(z_2) \rangle = -\frac{1}{\Box - M^2} \delta^8(z_1 - z_2) \n\langle \Phi(z_1)\Phi(z_2) \rangle = -\frac{M\bar{D}^2}{4\Box(\Box - M^2)} \delta^8(z_1 - z_2)
$$
\n(B.1)

Thepossible vertices (see ([3.1\)](#page-12-0)) are $\frac{g}{3!}\phi^3$ and $\lambda \Phi \phi^2$. It is easy to see that there are the following types of possible supergraphs. The first type of supergraphs consists of $\langle \phi \phi \rangle$ propagatorswith an alternating background ϕ and $\bar{\phi}$ (cf. [[28](#page-56-0), [29\]](#page-56-0)). Here the external legs correspond to alternating $g\phi$ and $g\phi$.

The result represents itself as a sum of all such supergraphs. The diagram of such a form with $2n$ external legs corresponds to a ring containing n links of the form

A contribution of such a link is of the form

$$
R = \frac{g^2 |\phi|^2}{k^4} \frac{D_{l-1}^2}{4} \delta_{l-1,l} \frac{\bar{D}_l^2}{4} \delta_{l,l+1}
$$
 (B.2)

Here, superfields W'' , \bar{W}'' are treated as constants, and external momenta are put to zero. Let us denote the contribution of a supergraph with $2n$ external lines as I_{2n} . It is easy to see that

$$
I_{2n} = \frac{1}{2n} \int \prod_{k=1}^{2n} d^4 \theta_1 ... d^4 \theta_{2n} \int \frac{d^4 k}{(2\pi)^4} \frac{1}{2n} R^n =
$$

\n
$$
= \frac{1}{2n} \int \prod_{k=1}^{2n} d^4 \theta_1 ... d^4 \theta_{2n} \int \frac{d^4 k}{(2\pi)^4} g^2 |\phi|^{2n} \left(\frac{1}{k^2}\right)^{2n} \times
$$

\n
$$
\times \frac{D_1^2}{4} \delta_{12} \frac{\bar{D}_2^2}{4} \delta_{23} ... \frac{D_{2n-1}^2}{4} \delta_{2n-1,2n} \frac{\bar{D}_{2n}^2}{4} \delta_{2n,1}
$$
(B.3)

Here $2n$ is a symmetric factor (see [[28](#page-56-0)]).

Straightforward D−algebra transformations lead to

$$
I_{2n} = \int d^4\theta \int \frac{d^4k}{\left(2\pi\right)^4 k^2} \frac{1}{2n} \left(-\frac{g^2 |\phi|^2}{k^2}\right)^n \tag{B.4}
$$

Here we used the rule $D^2 \bar{D}^2 D^2 = -16k^2 D^2$.

The total contribution of all these diagrams which is further denoted as $K_0^{(1)}$ $_0^{(1)}$ is a sum of all I_{2n} (cf. [[28](#page-56-0)]), i.e.

$$
K_0^{(1)} = \sum_{n=1}^{\infty} I_{2n} = \int d^4 \theta \frac{d^4 k}{(2\pi)^4 k^2} \sum_{n=1}^{\infty} \frac{1}{2n} \left(-\frac{g^2 |\phi|^2}{K_{\Phi\bar{\Phi}}^2 k^2} \right)^n
$$
(B.5)

Then, since

$$
\sum_{n=1}^{\infty} \frac{1}{2n} (-a)^n = -\log(1+a),
$$

 $K_0^{(1)}$ $\int_0^{(1)}$ is of the form

$$
K_0^{(1)} = \sum_{n=1}^{\infty} I_{2n} = -\int d^4\theta \frac{d^4k}{\left(2\pi\right)^4 k^2} \log\left(1 + \frac{g^2 |\phi|^2}{k^2}\right) \tag{B.6}
$$

We can integrate over angular coordinates which allows us to change d^4k for $\pi^2 r dr$ where $r = k²$. Then, since the integral over r is divergent we carry out dimensional regularization introducing the regularization parameter ϵ . Namely, we change $\pi^2 r dr \to \pi^2 r^{1+\epsilon/2} dr$. As a result we arrive at

$$
K_0^{(1)} = -\mu^{-\epsilon} \int d^4\theta \frac{dr r^{\epsilon/2}}{32\pi^2} \log\left(1 + \frac{g^2 |\phi|^2}{K_{\Phi\bar{\Phi}}^2 r}\right)
$$
(B.7)

Then,

$$
\int_0^\infty dr r^{\epsilon/2} \log(1+\frac{A}{r}) = -A^{1+\epsilon} \Gamma(-1-\epsilon/2) = -\frac{2}{\epsilon}A - A \log \frac{A}{\mu^2} + O(\epsilon)
$$

Inserting $A = g^2 |\phi|^2$ we arrive at the one-loop contribution to the effective action

$$
K_0^{(1)} = -\frac{1}{32\pi^2} g^2 |\phi|^2 (\frac{2}{\epsilon} + \log \frac{g^2 |\phi|^2}{\mu^2})
$$
 (B.8)

 $(see [24]).$ $(see [24]).$ $(see [24]).$

The most effective method to study diagrams that contain vertices proportional to $\Phi\phi^2$ consists of introducing the propagator for the light superfield ϕ_q which depends on the backgroundlight superfield ϕ and corresponds to the classical action ([3.3\)](#page-12-0). This propagator is denoted by a bold line as

$$
\underline{\hspace{1cm}} = \underline{\hspace{1cm}} + \underline{\hspace{1cm}} +
$$

A summation of this chain allows one to show that the total propagator of the light quantum superfield depends on the background light superfield in the following way

$$
\langle \phi_q(z_1)\overline{\phi}_q(z_2)\rangle = -\frac{\overline{D}_1^2 D_2^2}{16(\Box - g^2|\phi|^2)} \delta_{12} \tag{B.9}
$$

Thepropagator for heavy superfield Φ is given by $(B.1)$ $(B.1)$ $(B.1)$. Then we can see that aside from the diagrams discussed above there are two other types of supergraphs. The first one consists of an arbitrary number of repeating chains of the form $\langle \Phi \Phi \rangle \langle \phi \bar{\phi} \rangle$, i.e.

where double lines correspond to an external alternating $\lambda \Phi$ and $\lambda \bar{\Phi}$, and bold one to $\langle \phi \bar{\phi} \rangle$ propagators.A contribution from a supergraph consisting of n such chains, because of $(B.1)$ $(B.1)$ $(B.1)$, is of the form

$$
K_n = \frac{1}{n} \left(\frac{\lambda^2 |\Phi|^2 \bar{D}^2 D^2}{16(\Box - g^2 |\phi|^2)(\Box - M^2)} \right)^n
$$
(B.10)

The second type consists of an arbitrary number of repeating chains of the form $\langle \phi \phi \rangle \langle \phi \bar{\phi} \rangle$ which can be written as

Here a dashed-and-dotted line denotes $\langle \Phi \Phi \rangle$ and a jagged line denotes $\langle \bar{\Phi} \bar{\Phi} \rangle$. The external double line denote background superfield Φ. A contribution of a diagram consisting of n such chains is of the form

$$
L_n = \frac{1}{n} \left(\frac{M^2 \lambda^8 |\Phi|^8 \Box^2 \bar{D}^2 D^2}{16(\Box - g^2 |\phi|^2)^4 (\Box - M^2)^4} \right)^n
$$
(B.11)

The total one-loop correction in the kählerian effective potential is a sum of $K_0^{(1)}$ $_0^{(1)}$ ([B.8](#page-50-0)), and all the K_n and L_n contributions. After a summation with a subsequent Fourier transformation and an integration over momenta with a subsequent subtraction of the one-loop divergences we arrive at the one-loop correction in the kählerian effective potential (3.5) .

Appendix C

Let us consider a calculation of the one-loop kählerian effective potential for the case when light superfields are pure background ones and heavy superfields are pure quantum ones.

The computation of the one-loop effective action reduces to the problem of evaluating the operator

$$
\Omega(s) = e^{is\tilde{\Delta}'} = \exp\left[i(H\{D^2, \bar{D}^2\} + \frac{1}{4}W''\bar{D}^2 + \frac{1}{4}\bar{W}''D^2)\right]
$$
(C.1)

Representing $\Omega(s)$ in the standard form (see [[22](#page-56-0), [23](#page-56-0)])

$$
\Omega(s) = 1 + \frac{1}{16} A(s) D^2 \bar{D}^2 + \frac{1}{16} \tilde{A}(s) \bar{D}^2 D^2 + \frac{1}{8} B^{\alpha}(s) D_{\alpha} \bar{D}^2 +
$$

+
$$
\frac{1}{8} \tilde{B}_{\dot{\alpha}}(s) \bar{D}^{\dot{\alpha}} D^2 + \frac{1}{4} C(s) D^2 + \frac{1}{4} \tilde{C}(s) \bar{D}^2
$$
(C.2)

we have the equation

$$
i\frac{\partial\Omega}{\partial s}=\Omega\tilde{\Delta}
$$

and therefore we arrive at the system of equations:

$$
\frac{1}{16i}\dot{A} = H + HA\Box + \frac{1}{16}W''C
$$
\n
$$
\frac{1}{4i}\dot{C} = \frac{1}{4}\bar{W}'' + 4HC\Box + \frac{1}{4}\bar{W}''A\Box
$$
\n
$$
\frac{1}{8i}\dot{B}^{\alpha} = \frac{1}{8}W''\tilde{B}_{\dot{\alpha}}\partial^{\alpha\dot{\alpha}}
$$
\n(C.3)

The initial conditions for these equations and analogous ones for \tilde{A} , \tilde{B} , \tilde{C} are $A = B = C =$ $\tilde{A} = \tilde{B} = \tilde{C} = 0$ at $s = 0$. Hence one finds that $B^{\alpha}(s) = \tilde{B}_{\dot{\alpha}}(s) = 0$. Solutions of other equations are of the form

$$
A(s) = \tilde{A}(s) = \frac{e^{16iH\Box s}}{2\Box} (\exp(is\sqrt{W''\overline{W''}\Box}) + \exp(-is\sqrt{W''\overline{W''}\Box})) - \frac{1}{\Box}
$$

\n
$$
C(s) = e^{16iH\Box s} \frac{\overline{W''}}{\sqrt{W''\overline{W''}\Box}} \sinh(is\sqrt{W''\overline{W''}\Box)
$$

\n
$$
\tilde{C}(s) = e^{16iH\Box s} \frac{W''}{\sqrt{W''\overline{W''}\Box}} \sinh(is\sqrt{W''\overline{W''}\Box})
$$
\n(C.4)

The one-loop kählerian effective potential has the form

$$
K^{(1)} = -i \int_0^\infty \frac{ds}{s} (A(s) + \tilde{A}(s)) U(x, x'; s)|_{x=x'} \tag{C.5}
$$

where $U(x, x'; s) = e^{is\Box} \delta^4(x - x')$ (cf. [\[22](#page-56-0), [23\]](#page-56-0)). Using (C.4), the expression (C.5) can be rewritten as

$$
K^{(1)} = -i \int_0^\infty \frac{ds}{s} \left(\frac{e^{16iH\Box s}}{2\Box} \left(\exp(is\sqrt{W''\bar{W}''\Box}) + \exp(-is\sqrt{W''\bar{W}''\Box}) \right) - \frac{1}{\Box} \right) \times
$$

$$
\times e^{is\Box} \delta^4 (x - x')|_{x = x'}
$$
 (C.6)

Let us denote $\sqrt{W''\bar{W}''\Box}$ as Q.

Since

$$
\frac{e^{16iH\Box s}}{2\Box}(\exp(is\sqrt{W''\overline{W''}\Box}) + \exp(-is\sqrt{W''\overline{W''}\Box})) - \frac{1}{\Box} =
$$
\n
$$
= \frac{(\cosh(iQs) - 1)}{\Box}e^{16isH\Box} + \frac{e^{16isH\Box} - 1}{\Box}
$$
\n(C.7)

we can rewrite the expression (C.6) as

$$
K^{(1)} = -i \int_0^\infty \frac{ds}{s} \left\{ \frac{\cosh(iQs) - 1}{\Box} e^{16iH\Box s} + \frac{(e^{16iH\Box s} - 1)}{\Box} \right\} \times
$$

$$
\times e^{is\Box} \delta^4 (x - x')|_{x = x'}
$$
 (C.8)

which equals to

$$
K^{(1)} = -i \int_0^\infty \frac{ds}{s} \left\{ \frac{\cosh(iQs) - 1}{\Box} e^{i(16H + 1)\Box s} \delta^4(x - x')|_{x = x'} + \frac{(e^{16iH\Box s} - 1)}{\Box} e^{is\Box} \delta^4(x - x')|_{x = x'} \right\}
$$
(C.9)

Then,

$$
\frac{\cosh(iQs) - 1}{\Box} = \sum_{n=1}^{\infty} \frac{\left(is\sqrt{W''\bar{W}''}\right)^{2n}}{(2n)!} \Box^{n-1}
$$
\n
$$
\frac{\left(e^{16iH\Box s} - 1\right)}{\Box} = \sum_{n=1}^{\infty} \frac{\left(16iHs\right)^n}{n!} \Box^{n-1}
$$
\n(C.10)

In analogy with[[22, 23](#page-56-0)] we conclude that

$$
\Box^{n-1} e^{is\Box} \delta^4(x - x')|_{x=x'} = \frac{(-1)^{n-1} n!}{16\pi^2(is)^{n+1}}
$$

$$
\Box^{n-1} e^{is(1+16H)\Box} \delta^4(x - x')|_{x=x'} = \frac{1}{(1+16H)^{n+1}} \frac{(-1)^{n-1} n!}{16\pi^2(is)^{n+1}}
$$
(C.11)

The last expression can be obtained from the first one by the change $s \to s(1 + 16H)$. Then,

$$
K^{(1)} = -i \int_0^\infty \frac{ds}{s} \left\{ \sum_{n=1}^\infty \frac{(2is\sqrt{W''\bar{W}''})^{2n}}{(2n)!} \frac{1}{(1+16H)^{n+1}} \frac{(-1)^{n-1}n!}{16\pi^2(is)^{n+1}} + \sum_{n=1}^\infty \frac{(16iHs)^n}{n!} \frac{(-1)^{n-1}n!}{16\pi^2(is)^{n+1}} \right\}
$$
(C.12)

It is equal to

$$
K^{(1)} = -i\frac{1}{16\pi^2} \int_0^\infty \frac{ds}{s} \left\{ \frac{1}{(1+16H)is} \sum_{n=1}^\infty \left(\frac{isW''\bar{W}''}{1+16H} \right)^n \frac{(-1)^{n-1}n!}{(2n)!} - \frac{16H}{(1+16H)is} \right\}
$$
(C.13)

The integral $\int_0^\infty \frac{ds}{s}$ $\frac{ds}{s}$ is divergent, therefore we introduce the regularization parameter ϵ and change $\int_0^\infty \frac{ds}{s} \to \int_0^\infty \frac{ds}{s^{1-}}$ $\frac{ds}{s^{1-\epsilon}}$ Then, the last term in the expression (C.13) can be rewritten as

$$
-\frac{1}{16\pi^2} \int_0^\infty \frac{ds}{s^{2-\epsilon}} \frac{16H}{(1+16H)}
$$

and

$$
\int_0^\infty \frac{ds}{s^{2-\epsilon}} = \int_0^\infty ds s^{\epsilon-2} e^{-sp^2} |_{p^2=0} = (p^2)^{1-\epsilon} \Gamma(-1+\epsilon) |_{p^2=0}
$$

This term vanishes at $\epsilon \neq 0$ and can be defined to be equal to zero at $\epsilon = 0$.

Therefore the one-loop kählerian effective potential has the form

$$
K^{(1)} = -i\frac{1}{16\pi^2} \frac{1}{(1+16H)} \int_0^\infty \frac{ds}{s^{2-\epsilon}} \left\{ \sum_{n=1}^\infty \left(\frac{isW''\bar{W}''}{1+16H} \right)^n \frac{(-1)^{n-1}n!}{(2n)!} \right\} \tag{C.14}
$$

These power series can be summed up. Namely, after the change $s \to s \frac{iW''\bar{W}''}{1+16H}$ $\frac{W''W''}{1+16H}$ and a subsequent Wick rotation we arrive at

$$
K^{(1)} = -i\frac{1}{16\pi^2} \frac{1}{(1+16H)} \left\{ \frac{W''\bar{W}''}{1+16H} \right\}^{1+\epsilon} \int_0^\infty \frac{ds}{s^{2-\epsilon}} \left\{ \sum_{n=1}^\infty s^n \frac{(-1)^{n-1}n!}{(2n)!} \right\} \tag{C.15}
$$

It is known (cf.[[22, 23\]](#page-56-0)) that

$$
\sum_{n=1}^{\infty} s^n \frac{(-1)^{n-1} n!}{(2n)!} = s \int_0^1 dt e^{-\frac{1}{4}s(1-t^2)}
$$

Hence

$$
K^{(1)} = -\frac{1}{16\pi^2} \frac{1}{(1+16H)} \left\{ \frac{W''\bar{W}''}{1+16H} \right\}^{1+\epsilon} \int_0^\infty \frac{ds}{s^{1-\epsilon}} \int_0^1 dt e^{-\frac{1}{4}s(1-t^2)} \tag{C.16}
$$

and

$$
\int_0^\infty \frac{ds}{s^{1-\epsilon}} \int_0^1 dt e^{-\frac{1}{4}s(1-t^2)} = \Gamma(\epsilon) \int_0^1 dt (1 - \epsilon \log(\frac{1-t^2}{4}) + O(\epsilon^2)) =
$$

= $\frac{1}{\epsilon} + \gamma - \int_0^1 dt \log(\frac{1-t^2}{4}) + O(\epsilon)$
 $\left\{ \frac{W''\bar{W}''}{1+16H} \right\}^{1+\epsilon} = \frac{W''\bar{W}''}{1+16H}(1 + \epsilon \log \frac{W''\bar{W}''}{1+16H} + O(\epsilon^2))$ (C.17)

Here $O(\epsilon)$ denotes terms of order one and/or higher in ϵ , and $O(\epsilon^2)$ denoted terms of order two and/or higher in ϵ . Therefore

$$
K^{(1)} = -\frac{1}{32\pi^2} \left\{ \frac{W''\bar{W}''}{\left(1+16H\right)^2 \epsilon} + \frac{W''\bar{W}''}{\left(1+16H\right)^2} \left[\log\left\{ \frac{W''\bar{W}''}{\mu^2(1+16H)}\right\} \right] \right\}
$$
(C.18)

And $1 + 16H = F_{\Phi\bar{\Phi}}$. Here the first term contains all the one-loop divergences of the theory and can be cancelled by a suitable counterterm. The terms containing γ and $\int_0^1 dt \log(\frac{1-t^2}{4})$ $\frac{-t^2}{4}$ are removed by a suitable choice of the renormalization parameter μ .

Asa result we arrive at the expression (4.17) (4.17) for the one-loop kählerian effective potential.

References

[1] T. Appelquist, J. Carazzone, Phys.Rev. D11 (1975), 2856.

- [2] K. Symanzik, Comm. Math. Phys. 34 (1973), 7.
- [3] E. Witten, Nucl. Phys. B104, 445 (1976); J.C. Collins, F. Wilczek and A. Zee, Phys. Rev. D18, 242 (1978); B.A. Ovrut and H.T. Schnitzer, Phys. Rev. D21 3369 (1980). J.C. Collins, Renormalization (Gordon and Breach, N.Y.), 1982.
- [4] D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. B271, 93 (1986).
- [5] L. Dixon, E. Martinec, D. Friedan and S. Shenker, Nucl. Phys. B282, 13 (1987); M. Cvetič, Phys. Rev. Lett. 59, 2829 (1987);D. Bailin and A. Love, Phys. Lett. **B260**, 56 (1991); S. Kalara, J. L. Lopez and D.V. Nanopoulos, Nucl. Phys. B353, 650 (1991).
- [6] M.B. Green, J.H. Schwarz and E. Witten, Superstring Theory (Cambridge Univ. Press, 1987), Vol.2.
- [7] M. Dine, N. Seiberg and E. Witten, Nucl. Phys. B289,585 (1986); L. Dixon and V. Kaplunovsky, unpublished.
- [8] M. Cvetič, L. Everett, P. Langacker and J. Wang, Blowing up Type I Z3 Orientifold, Preprint [hep-th/9903051](http://arxiv.org/abs/hep-th/9903051).
- [9] J. Atick, L. Dixon and A. Sen, Nucl. Phys. B292, 109 (1987); M. Dine, I. Ichinose and N. Seiberg, Nucl. Phys. B293, 253 (1987); M. Dine and C. Lee, Nucl. Phys. B336, 317 (1990).
- [10] G. Cleaver, M. Cvetič, J.R. Espinosa, L. Everett and P. Langacker, Nucl. Phys. **B525**, 3 (1998); G. Cleaver, M. Cvetič, J.R. Espinosa, L. Everett and P. Langacker, Flat direction in three generation free fermionic string models, Preprint [hep-th/9805133,](http://arxiv.org/abs/hep-th/9805133) to be published in Nucl. Phys. B.
- [11] M. Cvetič, L. Everett and J. Wang, Nucl. Phys. **B538**, 52 (1998).
- [12] G. Cleaver, M. Cvetič, J.R. Espinosa, L. Everett and P. Langacker, Phys. Rev. D59, 55 (1999); Physics Implications of Flat Directions in Free Fermionic Superstring Models II: Renormalization Group Analysis, Preprint [hep-ph/9811355](http://arxiv.org/abs/hep-ph/9811355), to be published in Phys. Rev. D.
- [13] C.P. Burgess, A. Font and F. Quevedo, Nucl. Phys. B272, 661 (1986); A. Font and F. Quevedo, Phys. Lett. B184, 45 (1987).
- [14] E. Katehou and G.G. Ross, Nucl. Phys. B299, 484 (1988).
- [15] M. Cvetič, L. Everett, and J. Wang, Nucl. Phys. **B538**, 52 (1999).
- [16] N. Arkani-Hamed, M. Dine and S. Martin, Dynamical supersymmetry breaking in models with a Green-Schwarz mechanism, Preprint [hep-ph/9803432](http://arxiv.org/abs/hep-ph/9803432).
- [17] N. Arkani-Hamed, G.F. Giudice, M.A. Luty, and R. Rattazzi, Phys. Rev. D58 (1998), 115005; S. Dimopoulos, G. Dvali, R. Rattazzi, and G. Giudice, Nucl. Phys. B510, 12 (1998); G.F. Giudice and R. Rattazzi, Nucl. Phys. B511, 25 (1998).
- [18] I.L. Buchbinder, M. Cvetič and A.Yu. Petrov, Implication of decoupling effects for one-loop corrected effective actions for superstring theory, Preprint [hep-th/9903243.](http://arxiv.org/abs/hep-th/9903243)
- [19] G.B. Cleaver, A.E. Faraggi and D.V. Nanopoulos, A Minimal Superstring Standard Model. 1. Flat Directions, Preprint [hep-ph/9904301](http://arxiv.org/abs/hep-ph/9904301).
- [20] I.L. Buchbinder and S.M. Kuzenko, Ideas and Methods of Supersymmetry and Supergravity or a Walk Through Superspace. IOP Publishing, Bristol and Philadelphia, 1995.
- [21] I.L. Buchbinder, S.D. Odintsov and I.L. Shapiro, Effective Action in Quantum Gravity. IOP Publishing, Bristol and Philadelphia, 1992.
- [22] I.L. Buchbinder, S.M. Kuzenko and J.V. Yarevskaya, Nucl. Phys. B411, 665 (1994).
- [23] I.L. Buchbinder, S.M. Kuzenko and J.V. Yarevskaya, Yadernaya Fizika (Phys. Atom. Nucl) 56, 202 (1993) (in Russian).
- [24] P. West, Phys. Lett. B258, 375, 1991. P. West, Phys. Lett. B261, 396 (1991); I. Jack, D.R.T. Jones and P.West, Phys. Lett. B258, 382 (1991). J. Dunbar, I. Jack and D.R.T. Jones, Phys. Lett. B261, 62 (1991).
- [25] I.L. Buchbinder, S.M. Kuzenko and A.Yu. Petrov, Phys. Lett. B321, 372 (1994).
- [26] I.L. Buchbinder, S.M. Kuzenko, and A.Yu. Petrov, Yadernaya Fizika (Phys. Atom. Nucl.) 59, 157 (1996) (in Russian).
- [27] I.L. Buchbinder and A.Yu. Petrov, Phys. Lett. B461, 209 (1999).
- [28] A. Pickering and P. West, Phys. Lett. B383, 54 (1996).
- [29] M.T. Grisaru, M. Roček and R. von Unge, Phys.Lett. **B383**, 415 (1996).
- [30] I.N. Mc Arthur and T.D. Gargett, Nucl. Phys. B494, 525 (1997); N.G. Pletnev, A.T. Banin. Covariant technique of derivative expansion of one-loop effective action. *Preprint* [hep-th/9811031.](http://arxiv.org/abs/hep-th/9811031)