

Spectrum generating algebra and coherent states of the C_λ -extended oscillator

C. Quesne *

*Physique Nucléaire Théorique et Physique Mathématique,
Université Libre de Bruxelles, Campus de la Plaine CP229,
Boulevard du Triomphe, B-1050 Brussels, Belgium*

Abstract

C_λ -extended oscillator algebras, generalizing the Calogero-Vasiliev algebra, where C_λ is the cyclic group of order λ , have recently proved very useful in the context of supersymmetric quantum mechanics and some of its variants. Here we determine the spectrum generating algebra of the C_λ -extended oscillator. We then construct its coherent states, study their nonclassical properties, and compare the latter with those of standard λ -photon coherent states, which are obtained as a special case. Finally, we briefly review some other types of coherent states associated with the C_λ -extended oscillator.

1 Introduction

Coherent states (CS) of the harmonic oscillator [1] are known to have properties similar to those of the classical radiation field. They may be defined in various ways, for instance as eigenstates of the oscillator annihilation operator b . With the corresponding creation operator b^\dagger and the number operator $N_b \equiv b^\dagger b$, the latter satisfies the commutation relations

$$[N_b, b^\dagger] = b^\dagger, \quad [N_b, b] = -b, \quad [b, b^\dagger] = I. \quad (1)$$

In contrast, generalized CS associated with various algebras [2] may have some nonclassical properties, such as photon antibunching or sub-Poissonian photon statistics, and squeezing. As examples of such CS, we may quote the eigenstates of b^2 , which were introduced as even and odd CS or cat states [3], and are a special case of generalized CS associated with the Lie algebra $\text{su}(1,1)$ [4]. We may also mention the eigenstates of b^λ ($\lambda > 2$) or kitten states [5], which may be generated in λ -photon processes.

*Directeur de recherches FNRS; E-mail: cquesne@ulb.ac.be

Other examples are provided by nonlinear CS associated with a deformed oscillator (or f -oscillator). The latter is defined in terms of creation, annihilation, and number operators, $a^\dagger = f(N_b)b^\dagger$, $a = bf(N_b)$, $N = N_b$, satisfying the commutation relations [6, 7]

$$[N, a^\dagger] = a^\dagger, \quad [N, a] = -a, \quad [a, a^\dagger] = G(N), \quad (2)$$

where f is some Hermitian operator-valued function of the number operator and $G(N) = (N + 1)f^2(N + 1) - Nf^2(N)$. Nonlinear CS, defined as eigenstates of a [7, 8, 9], of a^2 [10], or of an arbitrary power a^λ ($\lambda > 2$) [11], have been considered in connection with nonclassical properties. It has been shown that for a particular class of nonlinearities the first ones are useful in the description of a trapped ion [8].

In the present communication, we shall consider some multiphoton CS, which may be associated with the recently introduced C_λ -extended oscillator [12]. The latter may be considered as a deformed oscillator with a \mathbb{Z}_λ -graded Fock space and has proved very useful in the context of supersymmetric quantum mechanics and some of its variants [12, 13]. In particular, we shall deal here in detail with CS of the C_λ -extended oscillator spectrum generating algebra [14], which are a special case of the CS of Ref. [11] and exhibit some nonclassical properties.

2 The C_λ -extended oscillator algebra

The C_λ -extended oscillator algebra (where $C_\lambda = \mathbb{Z}_\lambda$ is the cyclic group of order λ) was introduced as a generalization of the Calogero-Vasiliev algebra, defined by

$$[N, a^\dagger] = a^\dagger, \quad [a, a^\dagger] = I + \alpha_0 K, \quad \{K, a^\dagger\} = 0, \quad (3)$$

and their Hermitian conjugates, where α_0 is some real parameter subject to the condition $\alpha_0 > -1$, and K is some Hermitian operator. The latter may be realized as $K = (-1)^N$, so that the second equation in (3) becomes equivalent to $[a, a^\dagger] = I + \alpha_0 P_0 + \alpha_1 P_1$, where $\alpha_0 + \alpha_1 = 0$ and $P_0 = \frac{1}{2}[I + (-1)^N]$, $P_1 = \frac{1}{2}[I - (-1)^N]$ project on the even and odd subspaces of the Fock space \mathcal{F} , respectively.

When partitioning \mathcal{F} into λ subspaces $\mathcal{F}_\mu \equiv \{|k\lambda + \mu\rangle \mid k = 0, 1, \dots\}$, $\mu = 0, 1, \dots, \lambda - 1$, instead of two, the Calogero-Vasiliev algebra is replaced by the C_λ -extended oscillator algebra, defined by [12]

$$[N, a^\dagger] = a^\dagger, \quad [a, a^\dagger] = I + \sum_{\mu=0}^{\lambda-1} \alpha_\mu P_\mu, \quad a^\dagger P_\mu = P_{\mu+1} a^\dagger, \quad (4)$$

and their Hermitian conjugates, where $P_\mu = \lambda^{-1} \sum_{\nu=0}^{\lambda-1} \exp[2\pi i \nu(N - \mu)/\lambda]$ projects on \mathcal{F}_μ , $\sum_{\mu=0}^{\lambda-1} P_\mu = I$, and α_μ are some real parameters subject to the conditions $\sum_{\mu=0}^{\lambda-1} \alpha_\mu = 0$ and $\sum_{\nu=0}^{\mu-1} \alpha_\nu > -\mu$, $\mu = 1, 2, \dots, \lambda - 1$. Taking this form of P_μ into account, it is clear that the C_λ -extended oscillator algebra (4) is a special case of deformed oscillator algebra, as defined in (2).

The operators N , a^\dagger , a are related to each other through the structure function $F(N) = N + \sum_{\mu=0}^{\lambda-1} \beta_\mu P_\mu$, $\beta_\mu \equiv \sum_{\nu=0}^{\mu-1} \alpha_\nu$, which is a fundamental concept of deformed oscillators: $a^\dagger a = F(N)$, $aa^\dagger = F(N+1)$ [6, 7]. Comparing with Eq. (2), we get $f(N) = (F(N)/N)^{1/2}$.

The Fock space basis states $|n\rangle = |k\lambda + \mu\rangle = \mathcal{N}_n^{-1/2} (a^\dagger)^n |0\rangle$, where $a|0\rangle = 0$, $k = 0, 1, \dots$, and $\mu = 0, 1, \dots, \lambda - 1$, satisfy the relations

$$N|n\rangle = n|n\rangle, \quad a^\dagger|n\rangle = \sqrt{F(n+1)}|n+1\rangle, \quad a|n\rangle = \sqrt{F(n)}|n-1\rangle. \quad (5)$$

Due to the restrictions on the range of the parameters α_μ given below Eq. (4), $F(\mu) = \beta_\mu + \mu > 0$ so that all the states $|n\rangle$ are well defined.

The C_λ -extended oscillator Hamiltonian is defined by [12]

$$H_0 = \frac{1}{2} \{a, a^\dagger\}. \quad (6)$$

Its eigenstates are the states $|n\rangle = |k\lambda + \mu\rangle$ and their eigenvalues are given by $E_{k\lambda+\mu} = k\lambda + \mu + \gamma_\mu + \frac{1}{2}$, where $\gamma_\mu \equiv \frac{1}{2}(\beta_\mu + \beta_{\mu+1})$. In each \mathcal{F}_μ subspace of \mathcal{F} , the spectrum of H_0 is harmonic, but the λ infinite sets of equally spaced energy levels, corresponding to $\mu = 0, 1, \dots, \lambda - 1$, are shifted with respect to each other by some amounts depending upon the parameters $\alpha_0, \alpha_1, \dots, \alpha_{\lambda-1}$.

3 Spectrum generating algebra of the C_λ -extended oscillator

One can generate the whole spectrum of the C_λ -extended oscillator Hamiltonian (6) from the eigenstates $|\mu\rangle$, $\mu = 0, 1, \dots, \lambda - 1$, by using the operators [14]

$$J_+ = \frac{1}{\lambda} (a^\dagger)^\lambda, \quad J_- = \frac{1}{\lambda} a^\lambda, \quad J_0 = \frac{1}{\lambda} H_0 = \frac{1}{2\lambda} \{a, a^\dagger\}. \quad (7)$$

They satisfy the commutation relations

$$[J_0, J_\pm] = \pm J_\pm, \quad [J_+, J_-] = f(J_0, P_\mu), \quad [J_0, P_\mu] = [J_\pm, P_\mu] = 0, \quad (8)$$

where $f(J_0, P_\mu)$ (which has nothing to do with the function $f(N)$ of Eq. (2)) is a $(\lambda - 1)$ th-degree polynomial in J_0 with P_μ -dependent coefficients, $f(J_0, P_\mu) = \sum_{i=0}^{\lambda-1} s_i(P_\mu) J_0^i$. The spectrum generating algebra (SGA) of the C_λ -extended oscillator is therefore a C_λ -extended polynomial deformation of $\text{su}(1,1)$: in each \mathcal{F}_μ subspace, it reduces to a standard polynomial deformation of $\text{su}(1,1)$ [15].

Its Casimir operator can be written as

$$C = J_- J_+ + h(J_0, P_\mu) = J_+ J_- + h(J_0, P_\mu) - f(J_0, P_\mu), \quad (9)$$

where $h(J_0, P_\mu)$ is a λ th-degree polynomial in J_0 with P_μ -dependent coefficients, $h(J_0, P_\mu) = \sum_{i=0}^{\lambda} t_i(P_\mu) J_0^i$. Each \mathcal{F}_μ subspace is the carrier space of a unitary irreducible representation (unirrep) of the SGA, characterized by an eigenvalue c_μ of

C , and by the lowest eigenvalue $(\mu + \gamma_\mu + \frac{1}{2})/\lambda$ of J_0 . The explicit expressions of $f(J_0, P_\mu)$, $h(J_0, P_\mu)$, and c_μ are given in Ref. [14].

For $\lambda = 2$, for which the C_λ -extended oscillator algebra reduces to the Calogero-Vasiliev algebra, the SGA (7), (8) reduces to the Lie algebra $\text{su}(1,1)$, for which $f(J_0) = -2J_0$, $h(J_0) = -J_0(J_0 + 1)$, and $c = (1 + \alpha_\mu)(3 - \alpha_\mu)/16$ [16].

Nonlinearities make their appearance for $\lambda = 3$, for which

$$\begin{aligned} f(J_0, P_\mu) &= -9J_0^2 - J_0 \sum_{\mu} (\alpha_\mu + 2\alpha_{\mu+1})P_\mu - \frac{1}{12} \sum_{\mu} (1 + \alpha_\mu)(5 - \alpha_\mu)P_\mu, \\ h(J_0, P_\mu) &= -J_0 \left[3J_0^2 + \frac{1}{2}J_0 \sum_{\mu} (9 + \alpha_\mu + 2\alpha_{\mu+1})P_\mu + \frac{1}{12} \sum_{\mu} (23 + 10\alpha_\mu \right. \\ &\quad \left. + 12\alpha_{\mu+1} - \alpha_\mu^2)P_\mu \right], \\ c_\mu &= \frac{1}{72}(1 + \alpha_\mu)(5 - \alpha_\mu)(3 + \alpha_\mu + 2\alpha_{\mu+1}). \end{aligned} \quad (10)$$

For $\alpha_\mu = 0$ corresponding to $a^\dagger = b^\dagger$, $a = b$, the operators (7) close a polynomial deformation of $\text{su}(1,1)$, with $f(J_0)$ and $h(J_0)$ expressed in terms of some binomial coefficients and Stirling numbers [14].

4 Coherent states associated with the C_λ -extended oscillator spectrum generating algebra

As CS associated with the C_λ -extended oscillator SGA, let us consider generalizations of the Barut-Girardelle CS of $\text{su}(1,1)$ [4], to which they will reduce in the case $\lambda = 2$. These are the eigenstates $|z; \mu\rangle$ of the operator J_- defined in (7),

$$J_-|z; \mu\rangle = z|z; \mu\rangle, \quad z \in \mathbb{C}, \quad \mu = 0, 1, \dots, \lambda - 1. \quad (11)$$

Here μ distinguishes between the λ independent (and orthogonal) solutions of equation (11), belonging to the various subspaces \mathcal{F}_μ . The CS $|z; \mu\rangle$ may be considered as special cases of the nonlinear CS of Ref. [11], since Eq. (11) is equivalent to $a^\lambda|z; \mu\rangle = \lambda z|z; \mu\rangle$, for $a = bf(N_b)$ and $f(N_b)$ as given in Sec. 2.

It can be shown [14] that the states (11) satisfy Klauder's minimal set of conditions for generalized CS [17]: they are normalizable, continuous in the label z , and they allow a resolution of unity. The other discrete label μ is analogous to the vector components of vector (or partially) CS [18].

The states $|z; \mu\rangle$ can be written in either of the alternative forms

$$|z; \mu\rangle = [N_\mu(|z|)]^{-1/2} \sum_{k=0}^{\infty} \frac{(z/\lambda^{(\lambda-2)/2})^k}{[k! (\prod_{\nu=1}^{\mu} (\bar{\beta}_\nu + 1)_k) (\prod_{\nu'=\mu+1}^{\lambda-1} (\bar{\beta}_{\nu'})_k)]^{1/2}} |k\lambda + \mu\rangle, \quad (12)$$

$$|z; \mu\rangle = [N_\mu(|z|)]^{-1/2} {}_0F_{\lambda-1}(\bar{\beta}_1 + 1, \dots, \bar{\beta}_\mu + 1, \bar{\beta}_{\mu+1}, \dots, \bar{\beta}_{\lambda-1}; zJ_+/\lambda^{\lambda-2}) |\mu\rangle, \quad (13)$$

where $\bar{\beta}_\mu \equiv (\beta_\mu + \mu)/\lambda$, $(a)_k$ denotes Pochhammer's symbol, and the normalization factor $N_\mu(|z|)$ can be expressed in terms of a generalized hypergeometric function,

$$N_\mu(|z|) = {}_0F_{\lambda-1}(\bar{\beta}_1 + 1, \dots, \bar{\beta}_\mu + 1, \bar{\beta}_{\mu+1}, \dots, \bar{\beta}_{\lambda-1}; y), \quad y \equiv |z|^2/\lambda^{\lambda-2}. \quad (14)$$

Their unity resolution relation can be written as

$$\sum_\mu \int d\rho_\mu(z, z^*) |z; \mu\rangle \langle z; \mu| = I, \quad (15)$$

where $d\rho_\mu(z, z^*)$ is a positive measure, given in terms of a generalized hypergeometric function and a Meijer G -function by

$$\begin{aligned} d\rho_\mu(z, z^*) &= {}_0F_{\lambda-1}(\bar{\beta}_1 + 1, \dots, \bar{\beta}_\mu + 1, \bar{\beta}_{\mu+1}, \dots, \bar{\beta}_{\lambda-1}; y) h_\mu(y) |z| d|z| d\phi, \\ h_\mu(y) &= \frac{G_{0\lambda}^{\lambda 0}(y | 0, \bar{\beta}_1, \dots, \bar{\beta}_\mu, \bar{\beta}_{\mu+1} - 1, \dots, \bar{\beta}_{\lambda-1} - 1)}{\pi \lambda^{\lambda-2} \left(\prod_{\nu=1}^\mu \Gamma(\bar{\beta}_\nu + 1) \right) \left(\prod_{\nu'=\mu+1}^{\lambda-1} \Gamma(\bar{\beta}_{\nu'}) \right)}, \end{aligned} \quad (16)$$

with y defined in Eq. (14).

In the $\lambda = 2$ case, the functions ${}_0F_1$ and G_{02}^{20} of Eqs. (13), (14), and (16) being proportional to modified Bessel functions $I_{2\nu}(2|z|)$ and $K_{2\nu}(2|z|)$, $\nu = (\alpha_0 - 1 + 2\mu)/2$, respectively, the CS defined in (11) reduce to Barut-Girardello $\text{su}(1,1)$ CS [4] for the appropriate unirreps, as it should be.

For $\alpha_\mu = 0$ corresponding to $a^\dagger = b^\dagger$, $a = b$, the CS defined in (11) reduce to the eigenstates of b^λ or standard λ -photon CS [5],

$$|z; \mu\rangle = [N_\mu(|z|)]^{-1/2} \sum_{k=0}^{\infty} \left(\frac{\mu!}{(k\lambda + \mu)!} \right)^{1/2} (\lambda z)^k |k\lambda + \mu\rangle, \quad (17)$$

satisfying the resolution of unity (15) with $h_\mu(y)$ given by

$$h_\mu(y) = \lambda^{\mu-\lambda+2} (\pi\mu!)^{-1} y^{(\mu-\lambda+1)/\lambda} \exp(-\lambda y^{1/\lambda}). \quad (18)$$

The states (17) can be rewritten in the alternative form

$$|z; \mu\rangle = \left(\frac{\mu!}{E_{\lambda, \mu+1}(\lambda^2 |z|^2)} \right)^{1/2} E_{\lambda, \mu+1}(\lambda^2 z J_+) |\mu\rangle, \quad (19)$$

where $E_{\alpha, \beta}(x) \equiv \sum_{k=0}^{\infty} x^k / \Gamma(\alpha k + \beta)$ is a generalized Mittag-Leffler function. Hence, they provide a simple example of the Mittag-Leffler CS considered in Ref. [19].

5 Nonclassical properties of coherent states

The CS $|z; \mu\rangle$ may be considered as exotic states in quantum optics. Their properties may be analyzed in two different ways, by considering either “real” photons, described by the operators b^\dagger, b satisfying the canonical commutation relation, as given in Eq. (1), or “dressed” photons, described by the operators a^\dagger, a of Eq. (2), which may appear in some phenomenological models explaining some non-intuitive observable phenomena.

5.1 Photon statistics

Since $N = N_b$, the photon number statistics is not affected by the choice made for the type of photons. A measure of its deviation from the Poisson distribution is the Mandel parameter

$$Q = \frac{\langle(\Delta N)^2\rangle - \langle N\rangle}{\langle N\rangle}, \quad \Delta N \equiv N - \langle N\rangle, \quad (20)$$

which vanishes for the Poisson distribution, and is positive or negative according to whether the distribution is super-Poissonian (bunching effect) or sub-Poissonian (antibunching effect).

It is well known that for $\lambda = 2$, the standard even (resp. odd) CS, corresponding to $\alpha_0 = \alpha_1 = 0$ or $a^\dagger = b^\dagger, a = b$ and $\mu = 0$ (resp. $\mu = 1$), are characterized by a super-Poissonian (resp. sub-Poissonian) number distribution. It can be shown [14] that for the even (resp. odd) CS associated with the Calogero-Vasiliev algebra, i.e., for $\lambda = 2, \alpha_0 = -\alpha_1 \neq 0$ and $\mu = 0$ (resp. $\mu = 1$), this trend is enhanced for positive (resp. negative) values of α_0 . However, as shown in Fig. 1, for negative (resp. positive) values of α_0 and sufficiently high values of $|z|$, the opposite trend can be seen.

For higher values of λ , more or less similar results are obtained for $\mu = 0$, on one hand, and $\mu \neq 0$, on the other hand. However the behaviour of Q becomes more complicated for intermediate values of μ [14].

5.2 Squeezing effect

5.2.1 “Dressed” photons

Let us define the deformed quadratures x and p as

$$x = \frac{1}{\sqrt{2}}(a^\dagger + a), \quad p = \frac{i}{\sqrt{2}}(a^\dagger - a). \quad (21)$$

In any state belonging to \mathcal{F}_μ , their dispersions $\langle(\Delta x)^2\rangle$ and $\langle(\Delta p)^2\rangle$ satisfy the uncertainty relation

$$\langle(\Delta x)^2\rangle\langle(\Delta p)^2\rangle \geq \frac{1}{4}|\langle[x, p]\rangle|^2 = \frac{\lambda^2}{4}(\bar{\beta}_{\mu+1} - \bar{\beta}_\mu)^2, \quad (22)$$

where the right-hand side becomes smaller than the conventional value $1/4$ if $\alpha_0 < 0$ for $\mu = 0$ or $-2 < \alpha_\mu < 0$ for $\mu = 1, 2, \dots$, or $\lambda - 1$.

In \mathcal{F}_μ , the role of the vacuum state is played by the number state $|\mu\rangle = |0; \mu\rangle$, which is annihilated by J_- . The corresponding dispersions are given by

$$\langle(\Delta x)^2\rangle_0 = \langle(\Delta p)^2\rangle_0 = \frac{\lambda}{2}(\bar{\beta}_{\mu+1} + \bar{\beta}_\mu). \quad (23)$$

Comparing with the uncertainty relation (22), we conclude that the state $|\mu\rangle$ satisfies the minimum uncertainty property in \mathcal{F}_μ , i.e., gives rise to the equality in (22), only for $\mu = 0$ because $\bar{\beta}_0 = 0$ and $\bar{\beta}_\mu > 0$ for $\mu = 1, 2, \dots, \lambda - 1$. On the other hand, the dispersions in the vacuum may be smaller than the conventional value $1/2$ for $\mu = 0, 1, \dots, \lambda - 2$.

Let us restrict ourselves to the CS $|z; 0\rangle$, which satisfies for $z = 0$ the minimum uncertainty property. The quadrature x (resp. p) is said to be squeezed to the second order in $|z; 0\rangle$ if $X \equiv \langle(\Delta x)^2\rangle/\langle(\Delta x)^2\rangle_0$ (resp. $P \equiv \langle(\Delta p)^2\rangle/\langle(\Delta p)^2\rangle_0$) is less than one. Similarly, it is said to be squeezed to the fourth order if $Y \equiv \langle(\Delta x)^4\rangle/\langle(\Delta x)^4\rangle_0$ (resp. $Q \equiv \langle(\Delta p)^4\rangle/\langle(\Delta p)^4\rangle_0$) is less than one.

For $\lambda = 2$, X and P , or Y and Q , are related with each other by the transformation $\text{Re}z \rightarrow -\text{Re}z$. Moreover X and Y are minimum for real, negative values of z . On Fig. 2, they are displayed for such values. We note a large squeezing effect over the whole range of real, negative values of z for positive values of α_0 (for which the conventional uncertainty relation is respected).

For $\lambda > 2$, there is no second-order squeezing, but for $\lambda = 4$, a small fourth-order squeezing is obtained in accordance with the results for standard λ -photon CS [5].

5.2.2 “Real” photons

Let us now define the quadratures x and p as

$$x = \frac{1}{\sqrt{2}}(b^\dagger + b), \quad p = \frac{i}{\sqrt{2}}(b^\dagger - b). \quad (24)$$

Their dispersions $\langle(\Delta x)^2\rangle$ and $\langle(\Delta p)^2\rangle$ satisfy the usual uncertainty relation. Considering again the CS $|z; 0\rangle$, on Fig. 3 we observe for the ratios X and P more or less similar trends as noted in the case of “dressed” photons.

6 Concluding remarks

In the present contribution, we determined the SGA of the C_λ -extended oscillator and studied some CS associated with it, namely the eigenstates of its lowering generator J_- .

Other types of CS may be considered and will be studied in a forthcoming publication. Let us mention here two of them:

1. The eigenstates of the C_λ -extended oscillator annihilation operator a :

$$a|z; \mu\rangle = z|z; \mu\rangle. \quad (25)$$

These generalize the paraboson CS, which correspond to $\lambda = 2$ [20].

2. The solutions of the equation

$$\left[a^{\lambda-\alpha} - z (a^\dagger)^\alpha \right] |z; \mu\rangle = 0, \quad \alpha = 0, 1, \dots, \left\lfloor \frac{\lambda}{2} \right\rfloor, \quad \mu = 0, 1, \dots, \lambda - \alpha - 1. \quad (26)$$

For $\alpha = 0$, these are the eigenstates of a^λ , which are directly related to those of J_- , considered here. Moreover, for $\lambda = 2$ and $\alpha = 1$, they reduce to the Perelomov $\text{su}(1,1)$ CS [2].

References

- [1] R.J. Glauber, *Phys. Rev.* **131** (1963) 2766.
- [2] A.P. Perelomov, *Generalized Coherent States and Their Applications* (Springer, Berlin, 1986).
- [3] V.V. Dodonov, I.A. Malkin, V.I. Man'ko, *Physica* **72** (1974) 597.
- [4] A.O. Barut, L. Girardello, *Commun. Math. Phys.* **21** (1971) 41.
- [5] V. Bužek, I. Jex, Tran Quang, *J. Mod. Opt.* **37** (1990) 159.
- [6] C. Daskaloyannis, *J. Phys. A* **24** (1991) L789.
- [7] A.I. Solomon, *Phys. Lett. A* **196** (1994) 29.
- [8] R.L. de Matos Filho, W. Vogel, *Phys. Rev. A* **54** (1996) 4560.
- [9] V.I. Man'ko, G. Marmo, F. Zaccaria, E.C.G. Sudarshan, *Phys. Scr.* **55** (1997) 528.
- [10] S. Mancini, *Phys. Lett. A* **233** (1997) 291; S. Sivakumar, *Phys. Lett. A* **250** (1998) 257.
- [11] X.-M. Liu, *J. Phys. A* **32** (1999) 8685.
- [12] C. Quesne, N. Vansteenkiste, *Phys. Lett. A* **240** (1998) 21.
- [13] C. Quesne, N. Vansteenkiste, *Helv. Phys. Acta* **72** (1999) 71; C_λ -extended oscillator algebras and some of their deformations and applications to quantum mechanics, preprint math-ph/0003025, to be published in *Int. J. Theor. Phys.*

- [14] C. Quesne, *Phys. Lett. A* **272** (2000) 313.
- [15] A.P. Polychronakos, *Mod. Phys. Lett. A* **5** (1990) 2325; M. Roček, *Phys. Lett. B* **255** (1991) 554.
- [16] T. Brzeziński, I.L. Egusquiza, A.J. Macfarlane, *Phys. Lett. B* **311** (1993) 202.
- [17] J. Klauder, *J. Math. Phys.* **4** (1963) 1058.
- [18] J. Deenen, C. Quesne, *J. Math. Phys.* **25** (1984) 2354; D.J. Rowe, *J. Math. Phys.* **25** (1984) 2662.
- [19] J.-M. Sixdeniers, K.A. Penson, A.I. Solomon, *J. Phys. A* **32** (1999) 7543.
- [20] J.K. Sharma, C.L. Mehta, E.C.G. Sudarshan, *J. Math. Phys.* **19** (1978) 2089; J.K. Sharma, C.L. Mehta, N. Mukunda, E.C.G. Sudarshan, *J. Math. Phys.* **22** (1981) 78.