

Erratum

Friedrich Wagemann
 Institut Girard Desargues – ESA 5028 du CNRS
 Université Claude Bernard Lyon-I
 43, bd du 11 Novembre 1918
 69622 Villeurbanne Cedex FRANCE
 tel.: +33.4.72.43.11.90
 fax: +33.4.72.43.00.35
 e-mail: wagemann@desargues.univ-lyon1.fr

October 30, 2018

F. Wagemann: Some Remarks on the Cohomology of Krichever-Novikov Algebras, *Lett. Math. Phys.* **47** (1999), 173–177

There is a mistake in the statement of the main lemma. It should read:

Lemma 1 *Let $\Sigma_r = \Sigma \setminus \{p_1, \dots, p_r\}$ be an open Riemann surface of finite type.*

Suppose that Σ_r is trivializable, i.e. that the algebraic tangent bundle of the corresponding affine curve is trivial. Then $\overline{\text{Mer}}(\Sigma_r) = \text{Hol}(\Sigma_r)$,

in full words,

in the topology of uniform convergence on the compact sets K_n , a function with essential singularities in the points p_k may be approximated by functions with poles in the points.

The proof given in the article obviously works only for this case, for an open Riemann surface is always holomorphically trivializable, but the globally non-zero vector field is meromorphic only under the above hypothesis.

The trivializability condition can be expressed as the hypothesis that the divisor associated to the tangent bundle of the affine algebraic curve lies in the linear span of the points p_1, \dots, p_r . This is only valid in very rare cases, the points p_1, \dots, p_r being required to lay at special places, except for genus 0 and 1 where it is always true ($r \geq 1$).

The question whether this lemma remains true even for non-trivializable Riemann surfaces of finite type is - to our knowledge - open.

Acknowledgements:

The author thanks A. Rieman for very useful discussions on the subject.