Quantization of Kinematics on Configuration Manifolds¹

H.-D. Doebner Arnold Sommerfeld Institut für Mathematische Physik Technische Universität Clausthal Leibnizstr. 10 D-38678 Clausthal (Fed. Rep. Germany)

P. Šťovíček and J. Tolar Doppler Institute Faculty of Nuclear Sciences and Physical Engineering Czech Technical University Břehová 7 CZ-115 19 Prague 1 (Czech Republic)

Abstract

The review is devoted to topological global aspects of quantal description. The treatment concentrates on quantizations of kinematical observables — generalized positions and momenta. A broad class of quantum kinematics is rigorously constructed for systems, the configuration space of which is either a homogeneous space of a Lie group or a connected smooth finite-dimensional manifold without boundary. The class also includes systems in an external gauge field for an Abelian or a compact gauge group. Conditions for equivalence and irreducibility of generalized quantum kinematics are investigated with the aim of classification of possible quantizations. Complete classification theorems are given in two special cases. It is attempted to motivate the global approach based on a generalization of imprimitivity systems called quantum Borel kinematics. These are classified by means of global invariants — quantum numbers of topological origin. Selected examples are presented which demonstrate the richness of applications of Borel quantization. The review aims to provide an introductory survey of the subject and to be sufficiently selfcontained as well, so that it can serve as a standard reference concerning Borel quantization for systems admitting localization on differentiable manifolds.

¹Reviews in Mathematical Physics, Vol. 13, No. 4 (2001), pp. 1–47.

Contents

1	Introduction						
2	Mac 2.1 2.2 2.3 2.4 2.5 2.6	Ckey's system of imprimitivity The formalism of quantum mechanics Symmetry and quantum mechanics Localization, systems of imprimitivity Quantization on homogeneous spaces Infinitesimal action on a G-space Examples	6 6 9 9 12				
3	Quantum Borel kinematics: localization 1						
	3.1	Generalized system of imprimitivity	16				
	3.2	Quantum Borel kinematics	17				
	3.3	Quasi-invariant measures, projection–valued measures	18				
	3.4	An example	20				
4	Quantum Borel kinematics: external gauge fields						
	4.1	External magnetic field	21				
	4.2	Construction of a class of generalized systems of imprimitivity	21				
	4.3	Construction in the associated vector bundle	22				
	4.4	The Case $G = U(r)$	25				
	4.5	Unitary equivalence of generalized systems of imprimitivity	26				
	4.6	Irreducibility of generalized systems of imprimitivity	28				
	4.7	Quantum Borel kinematics with vanishing external field	29				
5	Quantum Borel kinematics: classification						
	5.1	Classification of generalized systems of imprimitivity via cocycles	30				
	5.2	Differentiable quantum Borel kinematics	31				
	5.3	Canonical representation of differentiable QBK^r	32				
	5.4	Classification of differentiable QBK^r 's	33				
	5.5	Classification of elementary differentiable quantum Borel kinematics	33				
	5.6	Classification of quantum Borel kinematics of type 0	35				
	5.7	Elementary quantum Borel kinematics with vanishing external field	36				
	5.8	Examples	37				

1 Introduction

The successful development of quantum theory in this century shows convincingly that it provides perhaps the most universal language for the description of physical phenomena. In quantum theory, as in any other physical theory, two fundamental aspects can be distinguished: the mathematical formalism and the physical interpretation.

At the basis of the most common mathematical formalism of quantum mechanics lies the notion of a complex separable Hilbert space \mathcal{H} of, in general, infinite dimension. Normed vectors in \mathcal{H} correspond to pure states of a quantum system, whereas quantal observables are represented by self-adjoint operators in \mathcal{H} .

However, only the rules of a *physical interpretation* enable one to use quantum theory for the description of physical systems. The principal general rule is Born's statistical interpretation of the wave function. For each physical system, or at least for a certain class of them, it is further necessary to specify which operators in \mathcal{H} are associated with physical observables measured by certain measuring devices. This means in particular that at least the operators of kinematical observables (position and momentum), and the dynamical evolution law of the system are to be specified.

An important tool for the derivation of quantum models are quantization methods. The primary aim of quantization of a given classical system is to associate self-adjoint operators with classical observables. As a rule, two main methods are used. The first one is based on Bohr's correspondence principle: the physical meaning of quantum operators is found by looking at their classical counterparts. In this way non-relativistic quantum mechanics was formulated by quantization of classical Hamiltonian mechanics, quantum theory of electromagnetic field by quantization of the Maxwell theory, etc. [42].

The correspondence principle can, of course, be the leading rule for quantization, if the observables already existed in a classical form. What should be done in the case of quantum observables without a classical analogue like the spin? Here the second method is often applicable, which uses *invariance principles* connected with the symmetries of the system. By Noether's theorem the operators corresponding to conserved quantities can be found as generators of some projective representation of the symmetry group in \mathcal{H} . As a far reaching application of this approach let us mention the relativistic quantum theory of elementary particles based on the irreducible unitary representations of the Poincaré group.

Both methods were used from the very first days of quantum theory, always taking into account specific physical properties of the systems considered. The first method usually appears in non-relativistic quantum mechanics as canonical quantization [42], for systems with the Euclidean configuration space \mathbb{R}^n . The position coordinates q_j and the canonically conjugate momenta p_k are quantized into self-adjoint position and momentum operators Q_j , P_k (in a separable Hilbert space \mathcal{H}), satisfying canonical commutation relations. This was originally discovered and mathematically formulated independently by W. Heisenberg and E. Schrödinger in 1925–26. The uniqueness of the mathematical formulation up to unitary equivalence was then guaranteed by the Stone-von Neumann Theorem.

Quantum mechanics on \mathbb{R}^n became very soon a successful theory which has been able to correctly describe experimental findings in vast areas of quantum physics. However, in some cases it was necessary to look for a formulation of quantum mechanics when the configuration space of a system was not Euclidean [52]. For instance, in connection with the studies of rotational spectra of molecules and of deformed nuclei, quantum rotators were introduced as fundamental quantum models with configuration spaces S^1 (the circle), S^2 (the 2-sphere)

and SO(3) (the rotation group). The textbook treatment of spinning top models (quantum mechanics of angular momentum) presents a successful application of the approach via invariance principles.

There were also attempts to enforce canonical quantization in cases where global Cartesian coordinates do not exist on the configuration manifold M. A formal quantization of generalized coordinates q_j and conjugate momenta p_k was suggested [52] on a manifold M with the Riemann structure (metric tensor g_{jk} with determinant g > 0):

$$Q_j = q_j, \quad P_k = -i\hbar \frac{\partial}{\partial q_k} - \frac{i\hbar}{4} \frac{\partial}{\partial q_k} (\ln g).$$
 (1.1)

Note that the additional term in P_k 's makes them symmetric operators in $\mathcal{H} = L^2(M, d\mu)$ with respect to the Riemann measure $d\mu = \sqrt{g} d^n q$ on M. The main difficulty encountered here is that operators (1.1) are not globally defined since, in general, q_j are only local coordinates. It is therefore desirable to invent quantization methods which employ global geometric objects.

On several occasions the formalism of quantum mechanics in connection with non-trivial topology of the configuration manifold lead to new non-classical effects. A deep and in its time not completely understood and recognized accomplishment in this direction was Dirac's famous investigation [6] of a quantum charged particle (charge e) in the external magnetic field of a point-like magnetic monopole (magnetic charge g). If the singular Dirac monopole is placed at the origin of a Cartesian coordinate system in \mathbb{R}^3 , one deals in fact with quantum mechanics on a topologically non-trivial effective configuration manifold $\mathbb{R}^3\setminus\{(0,0,0)\}$ (the three-dimensional Euclidean space with the origin excluded). Here the formalism of quantum mechanics in connection with non-trivial topology of the configuration manifold leads to an unexpected topological quantum effect originating from a peculiar behaviour of the phase of a wave function: Dirac discovered that a quantal description exists only under the condition that the dimensionless quantity $eg/2\pi\hbar$ is an integer.

Another phenomenon of this kind was noticed in 1959 by Y. Aharonov and D. Bohm [1]. The origin of the Aharonov–Bohm effect can be traced to a shift of the phase of wave function due to an external magnetic flux imposed on a charged particle. Here the effective Aharonov–Bohm configuration space is $\mathbb{R}^3 \setminus \mathbb{R}$, the three–dimensional Euclidean space with a straight line excluded. In both mentioned cases the topologies of the configuration spaces differ from the trivial topology of the Euclidean space and play decisive rôle in quantum theory.

These remarks about the early history of quantum mechanics clearly point to the need for a systematic development of *global quantization methods*. For systems with sufficiently *symmetric* configuration or phase spaces, two modern approaches in the theory of group representations can be applied:

- 1. Mackey's quantization on homogeneous configuration manifolds M = G/H [10, 23, 50]. Essentially, it is equivalent to the construction of systems of imprimitivity for a (locally compact, separable) group G, based on M = G/H.
- 2. The method of coadjoint orbits which play the role of homogeneous phase spaces [20, 47].

In the case of configuration or phase manifolds *without* geometric symmetries, two programs of global quantization were suggested:

3. Borel quantization on configuration manifolds [4, 11] which extends the notion of Schrödinger systems [27].

4. Geometric quantization on symplectic phase manifolds [20, 47].

These methods have been elaborated to differing degrees of sophistication and have, in general, different classes of classical systems as their domains of applicability. Borel quantization is built on configuration spaces and reflects the topology of M. For physical applications it is important that it yields both important classification theorems and explicit relations for quantization of kinematical observables. Like canonical quantization, it is a two step procedure. In a first step the kinematics, i.e. position and momentum observables on M, is quantized. The time dependence is introduced in a second step with a quantum analogue of a second order Riemannian dynamics on M [4]. In its most general form it leads to Doebner–Goldin non–linear Schrödinger equations [8].

Concerning other quantization methods respecting global properties of configuration or phase spaces we should especially mention:

- 5. The Feynman path integral method (it was used, e.g., in [28] for M = SO(3) and in [21] for configuration spaces of identical particles).
- 6. Quantization by deformation of classical mechanics [5].
- 7. Dirac quantization of systems with constraints in phase space [41].

This review article is devoted to the mathematical exposition of quantum Borel kinematics. This method yields quantizations of kinematics for systems admitting localization on connected smooth finite-dimensional configuration manifolds without boundary. We restrict our consideration exclusively to paracompact manifolds which (by Whitney's embedding theorem) can be regarded as submanifolds of \mathbb{R}^n .

In Chap. 2, the Hilbert space formalism of quantum mechanics, Wigner's Theorem on symmetry transformations, and the notion of Mackey's system of imprimitivity are briefly surveyed. The notion of quantum Borel kinematics is introduced in Chap. 3. In Chap. 4, a family of quantum Borel kinematics is constructed. This geometrical construction of quantum kinematics (Sect. 4.2) is based on the notion of a generalized system of imprimitivity for the family of one-parameter groups of diffeomorphisms (Sect. 3.1). It represents a generalization of quantum Borel kinematics of Ref. [4], especially in admitting an external gauge field with an arbitrary Abelian or compact structure group G. Thus the construction involves associated \mathbb{C}^r -bundles with finite-dimensional fibres \mathbb{C}^r . Chap. 4 is also devoted to questions of unitary equivalence and irreducibility (Sects. 4.5 and 4.6) of this class of quantum kinematics. Important special case of the vanishing external field is treated in Sect. 4.7. The classification of quantum Borel kinematics cannot be considered to be complete. In Chap. 5 theorems are stated which fully characterize them as well as two cases of complete classifications elementary quantum Borel kinematics (Sect. 5.5) and quantum Borel r-kinematics of type 0 (Sect. 5.6). In these cases it is shown that the first and the second singular homology groups of the configuration manifold M are involved and provide the necessary topological tools for classification of quantizations.

We have payed particular attention to a selection of proper examples which complement each chapter and demonstrate the richness of possible applications. From these examples, we mention a new derivation of the Dirac quantization condition from rotational symmetry [30] (Ex. 2.4), a topological description of the Aharonov–Bohm effect (Ex. 5.2), classification of elementary quantum Borel kinematics on arbitrary two–dimensional compact orientable manifolds (Ex. 5.3) as well as in the real projective space — topologically non–trivial part of the configuration space of the system of two identical particles [9] (Ex. 5.4).

2 Mackey's system of imprimitivity

2.1 The formalism of quantum mechanics

In quantum mechanics, a separable Hilbert space \mathcal{H} is associated with a quantum system we are going to describe. States of the system are represented by von Neumann's statistical (density) operators — bounded self-adjoint positive operators in \mathcal{H} with unit trace. The set of states \mathcal{W} introduced in this way is convex; its extremal points are called *pure states*. The pure states are just the projectors on one-dimensional subspaces of \mathcal{H} .

We assume² with [53] that, to a measurement on the system, taking values in a set X endowed with a σ -algebra $\mathcal{B}(X)$ of measurable subsets, there corresponds a projection-valued measure F on X. To any measurable set $S \in \mathcal{B}(X)$, a projector F(S) is related such that F(X) = 1 and $F(\bigcup_{i=1}^{\infty} S_i) = \sum_{i=1}^{\infty} F(S_i)$, provided $S_i \cap S_j = \emptyset$ for $i \neq j$. If the system is in a state $U \in \mathcal{W}$, then the formula

$$p_U(S) = \text{Tr}(UF(S))$$

gives the probability that the result of the measurement belongs to the set $S \in \mathcal{B}(X)$. The map $p_U: \mathcal{B}(X) \to \mathbb{R}$ is evidently a probability measure on X.

From the above considerations it is clear that in the quantal formalism crucial role is played by an orthocomplemented lattice of projectors onto subspaces of the Hilbert space \mathcal{H} [53]. This lattice will be denoted by \mathcal{L} ; partial ordering of \mathcal{L} is defined as follows: $F_1 \leq F_2$ if and only if $F_1F_2 = F_1$; the complement: $F^{\perp} = \mathbb{1} - F$. Clearly, $F_1 \leq F_2$ if and only if the corresponding subspaces are in inclusion; F^{\perp} projects on the orthogonal complement. Important properties of the lattice of projectors are:

- (i) For any countable set F_1, F_2, \ldots of elements from \mathcal{L} there exist $\bigvee_n F_n$ and $\bigwedge_n F_n$ in \mathcal{L} ;
- (*n*) For $F_1, F_2 \in \mathcal{L}$ and $F_1 \leq F_2$ there exists an element $P \in \mathcal{L}$ such that $P \leq F_1$ and $P \vee F_1 = F_2$. This element is equal to $P = F_1^{\perp} F_2 = (\mathbb{1} F_1) F_2 = F_2 F_1$ and is unique with this property.

An orthocomplemented lattice satisfying (i), (ii) is called a *logic*.

2.2 Symmetry and quantum mechanics

Let the configuration manifold — which we shall always denote by M — be a G—space of a symmetry group G. This means that an action of G on M is given, i.e., to each element g of G there corresponds a transformation g of M onto itself such that:

- (1) e.u = u,
- (2) $g_1(g_2.u) = (g_1g_2).u$, where $g_1, g_2 \in G$, $u \in M$.

²We are going to use such spectral measures for position measurements in configuration space and assume tacitly that they are ideal and the state after the measurement can be described by a projection $E(S)\psi$ of the original state ψ . There are several options for a description of a non-ideal localization — e.g. with the use of positive operator-valued measures [40], Ch. 3.

³ The element $A = \bigvee_n F_n$ is defined by the following properties: $1)A \geq F_n$ for all n; 2) if B is any element of \mathcal{L} such that $F_n \leq B$ for all n, then $A \leq B$. In an analogous fashion, the element $C = \bigwedge_n F_n$ is defined by: $1)C \leq F_n$ for all n; 2) if D is any element of \mathcal{L} such that $F_n \geq D$ for all n, then $C \geq D$.

Some important assertions, in particular Mackey's Imprimitivity Theorem, can be stated provided the group G is locally compact and separable (i.e. with countable basis of the topology). In the following we shall restrict our considerations to the case when G is a finite-dimensional connected Lie group. It is just these groups that very often appear in physical applications. Let the manifold M be also connected and smooth, and the mapping $(g,u) \mapsto g.u$ be infinitely differentiable (C^{∞}) .

Now we would like to associate, to each symmetry transformation g of the configuration space M, a symmetry transformation of the quantum mechanical description. To be specific, we introduce two notions.

Definition 2.1. An automorphism of a logic \mathcal{L} is a one-to-one mapping $\alpha: \mathcal{L} \to \mathcal{L}$ which satisfies

- (i) $\alpha(1) = 1$,
- (ii) $\alpha(F^{\perp}) = \alpha(F)^{\perp}$,
- (iii) $\alpha(\bigvee_{n=1}^{\infty} F_n) = \bigvee_{n=1}^{\infty} \alpha(F_n).$

Definition 2.2. A convex automorphism of the set of states W is a one-to-one mapping $\beta: W \to W$ with the following property: given positive real numbers c_1, c_2, \ldots such that $\sum_n c_n = 1$, then $\beta(\sum_n c_n U_n) = \sum_n c_n \beta(U_n)$.

Now we can state the conditions on symmetry transformations of the quantum mechanical description in the following form:

- (a) There exists a homomorphism $\alpha: g \mapsto \alpha(g)$ from the group G into the group of automorphisms of the logic \mathcal{L} , $\alpha(g): \mathcal{L} \to \mathcal{L}: F \mapsto F^g$.
- (b) There exists a homomorphism $\beta: g \mapsto \beta(g)$ from the group G into the group of convex automorphisms of the set of states \mathcal{W} , $\beta(g): \mathcal{W} \to \mathcal{W}: U \mapsto U^g$.
- (c) The probability does not change under the symmetry transformations, i.e.

$$\operatorname{Tr}(U^g F^g) = \operatorname{Tr}(UF), \qquad U \in \mathcal{W}, \ F \in \mathcal{L}.$$

(d) Given a projection–valued measure F on a set X (endowed with a σ -algebra of measurable subsets) which corresponds to measurements on the system with values in X, there exists a homomorphism $\gamma: g \mapsto \gamma(g)$ from the group G into the group of measurable and one-to-one mappings from X onto itself. Every $\gamma(g)$ induces an automorphism of the σ -algebra, $\mathcal{B}(X) \to \mathcal{B}(X): S \mapsto S^g$. We demand

$$F(S^g) = F(S)^g. (2.1)$$

Automorphisms of the logic and convex automorphisms of the set of states are described by *Wigner's theorem*:

Theorem 2.1 ([53], Chap. VII.3). Let \mathcal{H} be a separable infinite-dimensional Hilbert space. Then:

1) All automorphisms of the logic \mathcal{L} are of the form

$$\alpha(F) = TFT^{-1}, \qquad F \in \mathcal{L},$$

where T is a fixed unitary or antiunitary operator in \mathcal{H} . Two such operators induce the same automorphism of the logic if and only if they differ by a phase factor.

2) All convex automorphisms of the set of states W are of the form

$$\beta(U) = TUT^{-1}, \qquad U \in \mathcal{W},$$

where T is a fixed unitary or antiunitary operator in \mathcal{H} . Two such operators induce the same convex automorphism of the set of states if and only if they differ by a phase factor.

Theorem 2.1 and conditions (a), (b) imply that to each action $g \in G$ a pair of operators T(g), T'(g) is associated, both being unitary or antiunitary. To fulfil condition (c), operators T(g), T'(g) may differ by a phase factor at most. Hence these operators can be identified,

$$T(g) = T'(g), \qquad g \in G.$$

Since we consider only connected Lie groups, all operators T(g) will be unitary⁴. Let us denote by $U(\mathcal{H})$ the group of unitary operators in \mathcal{H} with strong topology. The centre Z of this group consists of operators $z.\mathbb{1}$, $z \in T^1$, where T^1 denotes the compact Lie group of complex numbers of unit modulus. The quotient group

$$P(\mathcal{H}) = U(\mathcal{H})/Z$$

is called the *projective group of the Hilbert space* \mathcal{H} . The conditions (a) — (c) can be summarized in one requirement

(abc) There exists a homomorphism $h: G \to P(\mathcal{H})$.

Moreover, we shall demand h to be measurable. In this case h is even continuous ([54], Chap. VIII.5).

The condition (abc) can be reformulated with the use of the notion of a projective representation. Let $\pi: U(\mathcal{H}) \to P(\mathcal{H})$ be the canonical homomorphism. For each given homomorphism $h: G \to P(\mathcal{H})$ there exists a measurable mapping $V: G \to U(\mathcal{H}), \ V(e) = \mathbb{1}, \ h = \pi \circ V$. The mapping V is called the *projective representation* of G. Two projective representations V, V' are called *equivalent* if there exists a measurable mapping $z: G \to T^1$ such that V'(g) = z(g)V(g). The homomorphism h obviously determines the projective representation uniquely up to this equivalence.

Given a projective representation V, there exists a measurable mapping

$$m: G \times G \to T^1$$

such that

$$V(a)V(b) = m(a,b)V(ab), \qquad a,b \in G.$$

The factor m(a,b) is called a multiplier of G; by definition it fulfils

$$m(ab,c)m(a,b) = m(a,bc)m(b,c), \qquad m(a,e) = m(e,a) = 1.$$

Two multipliers m, m' are equivalent if there exists a measurable mapping $z: G \to T^1$ such that

$$m'(a,b) = z(ab)^{-1}z(a)z(b)m(a,b).$$

⁴This follows, on the one hand, from the fact that the composition of two antiunitary operators is unitary, and, on the other hand, from the fact that in some neighbourhood N of the unit element e there exists $a \in N$ to each $b \in N$ such that $b = a^2$; it is well known that G is generated by the elements of N.

By definition a multiplier is exact (or trivial), if it is equivalent to 1. The set of all multipliers with pointwise multiplication forms an Abelian group; trivial multipliers form its invariant subgroup. The corresponding quotient group is referred to as the multiplier group for G; we shall denote it by $\mathcal{M}(G)$. ⁵

2.3 Localization, systems of imprimitivity

The discussion of condition (d) of Sect. 2.2 was postponed to this section, since its analysis requires the description of a concrete measurement on the system. For localizable systems the position measurements play a distinguished rôle. Results of position measurements are points of the configuration space M, i.e. X = M. So it is natural to consider the σ -algebra $\mathcal{B}(M)$ of Borel subsets of M as the σ -algebra of measurable sets 6 .

The starting point of Mackey's quantization and of quantum Borel kinematics is the notion of localization of a quantum system on a configuration manifold M. It is mathematically modeled by a projection-valued measure $E: S \mapsto E(S)$ mapping Borel subsets S of M ($S \in \mathcal{B}(M)$) into projection operators E(S) on a separable Hilbert space \mathcal{H} subject to the usual axioms of localization. For convenience, these axioms are given below:

$$\begin{split} E(S_1 \cap S_2) &= E(S_1) \cdot E(S_2), \\ E(S_1 \cup S_2) &= E(S_1) + E(S_2) - E(S_1 \cap S_2), \\ E(\bigcup_{i=1}^{\infty} S_i) &= \sum_{i=1}^{\infty} E(S_i) \quad \text{for mutually disjoint} \quad S_i \in \mathcal{B}(M), \\ E(M) &= \mathbb{1}. \end{split}$$

For a given subset $S \in \mathcal{B}(M)$, the projection E(S) corresponds to a measurement which determines whether the system is localized in S; its eigenvalues 1 (0) correspond to situations when the system is found completely inside (outside) S, respectively.

According to (d), each action $g \in G$ induces a Borel transformation of M onto itself. As already mentioned, the other three conditions (abc) imply the existence of a projective representation V of G. Hence (2.1) can be written in the form

$$E(g.S) = V(g)E(S)V(g)^{-1}, (S^g \equiv g.S).$$
 (2.2)

Definition 2.3. A pair (V, E) where V is a (projective) representation of a group G and E is a projection–valued measure on a G-space M, is called a *(projective) system of imprimitivity* for the group G, if (2.2) holds for all $g \in G$, $S \in \mathcal{B}(M)$. Two projective systems of imprimitivity are *equivalent* if the corresponding projective representations are equivalent and if the projection–valued measures are equal.

2.4 Quantization on homogeneous spaces

Stronger results can be obtained if the symmetry group G of M is sufficiently rich. More precisely, we shall turn our attention to homogeneous spaces. By definition, M is a homogeneous G-space if G acts transitively on M, i.e., to each pair of points $u, u' \in M$ there exists a transformation $g \in G$ such that g.u = u'. Let us fix a point $u_0 \in M$. The isotropy subgroup

⁵For details see [54], Chap. X.

 $^{^6\}sigma$ -algebra $\mathcal{B}(M)$ is generated by open subsets of manifold M.

of u_0 in G will be denoted by H. It is well known that H is a closed Lie subgroup of the Lie group G. The space G/H of left cosets gH, $g \in G$, endowed with factor topology, can be given a differentiable (C^{∞}) structure, thus becoming a smooth manifold, and the mapping $\pi: g \mapsto g.u_0$ induces a diffeomorphism of G/H onto M ([45], Chap. II.3, II.4). Having identified G/H with M, the group G acts on M in the natural way, $a: gH \mapsto agH$. The quadruple $(G, \pi, M; H)$ can be viewed as a principal fibre bundle. Let us note here that the requirement on G to be connected is not very restrictive provided M is connected:

Proposition 2.2 ([45], Chap. II.4). Let G be a finite dimensional Lie group acting transitively on a connected smooth manifold M and let G^0 be the connected component of unity in G. Then G^0 acts transitively on M, too.

As shown by Mackey [23, 50], the transitive systems of imprimitivity (i.e. based on homogeneous spaces G/H) can be completely classified. The notions of irreducibility, unitary equivalence, direct sum decomposition, etc., can be taken over for the systems of imprimitivity in exact analogy with these notions for (projective) unitary representations ([50], Chap. 1.2). In order to investigate questions of irreducibility, direct sum decomposition, etc., a commuting ring C(V, E) is considered, which consists of all bounded operators in \mathcal{H} commuting with E(S), V(g) for all $S \in \mathcal{B}(M)$, $g \in G$. We have for instance the property that a system of imprimitivity (V, E) is irreducible if and only if the ring C(V, E) consists of multiples of the unit operator $\mathbb{1}$ only (Schur's Lemma).

Now, following G.W. Mackey, we are going to describe the canonical construction of transitive systems of imprimitivity. Let G be a locally compact group satisfying the second axiom of countability, H its closed subgroup. On the coset space G/H there exists a quasi-invariant measure defined on the σ -algebra of Borel subsets.

A measure μ on G/H is called *quasi-invariant* with respect to the action of G, if for all $g \in G$ the measures μ and $\mu \circ g: S \mapsto \mu(g.S)$ are mutually absolutely continuous. Moreover, all quasi-invariant σ -finite measures on G/H are mutually absolutely continuous ([54], Chap. VIII.4).

We fix a measure μ from this class. Further, let m be a multiplier of G and let L be a projective unitary representation of H with multiplier m restricted to H in a separable Hilbert space \mathcal{H}^L . Then we construct the Hilbert space \mathcal{H} as the space of vector-valued functions $\psi: G \to \mathcal{H}^L$ satisfying

- a) $a \mapsto \langle \psi(a), f \rangle$ is a Borel function on G for all $f \in \mathcal{H}^L$;
- b) $\psi(ah) = m(a,h)L_h^{-1}\psi(a), h \in H;$
- c) $\|\psi\| < \infty$, where $\|\cdot\|$ is the norm induced by the inner product

$$(\psi, \psi') = \int_{G/H} \langle \psi(a), \psi'(a) \rangle d\mu(u);$$

the integral is well-defined since, because of b), the inner product $\langle \psi(a), \psi'(a) \rangle$ in \mathcal{H}^L remains constant on the left cosets u = aH.

Henceforth we shall identify two functions on G/H which are equal μ -almost everywhere. Then the projection–valued measure $S \mapsto E^L(S)$ on G/H is canonically defined by

$$[E^{L}(S)\psi](a) = \tilde{\chi}_{S}(a)\psi(a),$$

where

$$\tilde{\chi}_S(a) = \begin{cases} 1 \text{ for } aH \in S \\ 0 \text{ for } aH \notin S \end{cases}$$

The projective m-representation of $G, g \mapsto V^L(g)$, is given by

$$[V^{L}(g)\psi](a) = \sqrt{\frac{\mathrm{d}\mu}{\mathrm{d}(\mu \circ g)}(g^{-1}u)} \, m(a^{-1}, g) \, \psi(g^{-1}a),$$

where $d\mu/d(\mu \circ g)$ is the Radon–Nikodým derivative. The pair (V^L, E^L) is called a *canonical* system of imprimitivity and its equivalence class does not depend on the choice of a quasi-invariant measure μ .

Theorem 2.3 (The Imprimitivity Theorem [23]). Let G be a locally compact group satisfying the second axiom of countability, H its closed subgroup and m a multiplier of G. Let a pair (V, E) be a projective system of imprimitivity for G based on G/H with multiplier m. Then there exists an m-representation L of H such that (V, E) is equivalent to the canonical system of imprimitivity (V^L, E^L) . For any two m-representations L, L' of the subgroup H the corresponding canonical systems of imprimitivity are equivalent if and only if L, L' are equivalent. The commuting rings $C(V^L, E^L)$ and C(L) are isomorphic.

The Imprimitivity Theorem shows how to obtain all systems of imprimitivity up to unitary equivalence, provided the multiplier group $\mathcal{M}(G)$ is known. More facts about the multiplier group can be given in the case when G is a connected and simply connected Lie group. Then every multiplier is equivalent to a multiplier of class C^{∞} . These multipliers can be expressed in the form $\exp(ip)$ where p is called an *infinitesimal multiplier*. Let us introduce a coboundary operator δ on real skew–symmetric multilinear forms on the Lie algebra \mathcal{G} via

$$\delta p(X_1, \dots, X_n) = \sum_{k < j} (-1)^{k+j-1} p([X_k, X_j], X_1, \dots, \hat{X}_k, \dots, \hat{X}_j, \dots, X_n),$$

where p is any (n-1)-form. Then the Abelian group of infinitesimal multipliers of G is isomorphic to the second cohomology group $H^2(\mathcal{G}, \mathbb{R})$.

Examples

- (i) $H^2(\mathbb{R}^s, \mathbb{R})$ is isomorphic to the additive group of real skew–symmetric 2–forms on \mathbb{R}^s ; given a 2-form p, then $m:(x,y)\mapsto \exp[ip(x,y)]$ is a multiplier; $\mathcal{M}(\mathbb{R}^1)=\{1\}$.
- (ii) $\mathcal{M}(T^s) = \{1\}, T^s = \mathbb{R}^s / \mathbb{Z}^s.$
- (iii) If G is a connected and simply connected semi-simple Lie group, then $\mathcal{M}(G) = \{1\}$.

We note that any connected Lie group G can be replaced by its (connected and simply connected) universal covering Lie group \tilde{G} . The action of $\tilde{g} \in \tilde{G}$ on M is given by $\tilde{g}: u \mapsto \pi(\tilde{g}).u$ where $\pi: \tilde{G} \mapsto G$ is the covering homomorphism. As Example 2.3 (see Sect.2.5) will show, the transition to the covering group can lead to richer results with reasonable physical interpretation.

Remark. Detailed descriptions of general foundations of quantum mechanics can be found in [53], Chap. VI and VII; of systems of imprimitivity in [54], Chap. IX, [23] and [10]; of multipliers in [54], Chap. X.

2.5 Infinitesimal action on a G-space

Let G be a connected Lie group and M a (not necessarily homogeneous) G-space. A one-parameter subgroup of G is a one-dimensional Lie subgroup including its parametrization, $a: \mathbb{R} \to G: t \mapsto a(t)$. There is a one-to-one correspondence between elements A of a Lie algebra G and one-parameter subgroups $\{a(t)\}$ which can be expressed by a'(0) = A ([47], Chap. I.6.4). This correspondence can be used to define the mapping $\exp: G \to G: A \mapsto a(1)$; then one has $a(t) = \exp(tA)$, and \exp is a local diffeomorphism at the unit element of G. To each $A \in G$ there corresponds a one-parameter subgroup $\{a(t) = \exp(tA)\}$ and a flow on M, $(t,u) \mapsto a(t).u$; the corresponding vector field on M will be denoted by D_A . If $F^u: G \to M$ is the mapping $G \mapsto G$ is the mapping $G \mapsto G$ in the terminology of [38], an infinitesimal action is the mapping $G \to \mathcal{X}(M): A \mapsto D_A$, where $\mathcal{X}(M)$ denotes the infinite-dimensional Lie algebra of smooth vector fields on M.

Let N be the subgroup of ineffectively acting elements from G. If $N = \{e\}$, then G is said to act *effectively* on M. N is closed and normal, the factor group G/N is a Lie group acting effectively on M. Manifold M can be considered as a G/N-space if the action is given by $G/N \ni gN: u \mapsto g.u$. In this way the ineffectively acting elements can be eliminated⁷.

Theorem 2.4 ([38], Chap. III.3.7). The infinitesimal action $A \mapsto D_A$ is linear. For all $A, B \in \mathcal{G}$ one has

$$[D_A, D_B] = -D_{[A,B]}.$$

Hence the image of this mapping is a finite-dimensional Lie subalgebra in $\mathcal{X}(M)$. The kernel is \mathcal{N} , the Lie algebra of the group N of ineffective elements from G.

The **proof** of the first part is based on a straightforward calculation ([38], Chap. III.3.7). For the last assertion we observe that, if $D_A = 0$ and ϕ^A is the corresponding flow, then $a(t).u = \phi^A(t, u) = u$ so $\{a(t)\} \subset N$.

Now let us consider the opposite situation. Suppose we are given a finite-dimensional Lie subalgebra \mathcal{G} in $\mathcal{X}(M)$ such that all vector fields from \mathcal{G} are complete. Let $\tilde{\mathcal{G}}$ be the Lie algebra with the same vector space as \mathcal{G} but with a Lie bracket $[.,.]^{\sim} = -[.,.]$. If \tilde{G} is the connected and simply connected Lie group with Lie algebra $\tilde{\mathcal{G}}$ (\tilde{G} is unique up to isomorphism), then according to [38], Chap. III.4.7, Theorem 6, to each $u \in M$ there exists an open neighbourhood B_u and a uniquely defined local action of \tilde{G} on B_u (i.e., an action defined only for elements from some neighbourhood U_e of the unity) such that the associated infinitesimal action is identical with the mapping $\tilde{\mathcal{G}} \ni X \mapsto X \mid_{B_u}$. The neighbourhood U_e can be chosen small enough for exp to be a diffeomorphism on it. For $X \in \tilde{\mathcal{G}}$, ϕ^X —the corresponding flow, $v \in B_u$, $t \in \mathbb{R}$ sufficiently small, we have $\exp(t.X).v = \phi^X(t,v)$. Since all vector fields $X \in \tilde{\mathcal{G}}$ are complete, U_e can be chosen independently of $u \in M$. In this way we obtain a local homomorphism from \tilde{G} into the group of diffeomorphisms of M. Since \tilde{G} is connected and simply connected, the domain of the local homomorphism can be unambiguously extended to the whole \tilde{G} ([39], Chap. II and VII).

2.6 Examples

Example 2.1. $M = \mathbb{R}^1$, $G = \mathbb{R}^1$ – the group of translations. Both the multiplier group $\mathcal{M}(\mathbb{R}^1)$ and the isotropy subgroup are trivial. So in this case exactly one irreducible system of imprimitivity exists (up to unitary equivalence). Let \mathcal{H} be a Hilbert space and (Q, P) be a

⁷However, for N discrete it does not seem reasonable to eliminate N in this way; see Ex. 2.3.

pair of self-adjoint operators in \mathcal{H} satisfying the commutation relation $QP - PQ = i\hbar \mathbb{1}$. Then if $E: S \mapsto E(S)$ is the spectral projection-valued measure of Q, and $V(t) = \exp(-itP/\hbar)$, then the commutation relation is equivalent to the identity

$$V(t)E(S)V(-t) = E(t+S).$$

So the Imprimitivity Theorem implies the Stone-von Neumann theorem (cf. [50], Chap. 2.5).

Example 2.2. $M = \mathbb{R}^2$, $G = \mathbb{R}^2$ – the group of translations. To each element $A = (A_1, A_2) \in \mathcal{G} = \mathbb{R}^2$ there corresponds a vector field $D_A = (A_1 \partial/\partial x_1) + (A_2 \partial/\partial x_2) \in \mathcal{X}(\mathbb{R}^2)$. The isotropy subgroup is trivial. $\mathcal{M}(\mathbb{R}^2)$ is isomorphic to the group \mathbb{R} under addition: if $B \in \mathbb{R}$, then

$$m_B: \mathbb{R}^2 \times \mathbb{R}^2 \to U(1): (x,y) \mapsto \exp\left[-i\frac{eB}{2\hbar}(x_1y_2 - x_2y_1)\right]$$

is a multiplier (e is an arbitrary fixed non-zero constant). The inequivalent irreducible systems of imprimitivity (V^B, E^B) are labelled by $B \in \mathbb{R}$: the Hilbert space is $\mathcal{H}^B = L^2(\mathbb{R}^2, \mathrm{d}x_1\mathrm{d}x_2)$ and we find $E^B(S): \psi \mapsto \chi_S.\psi$ and

$$V^{B}(\exp(tA))\psi = \exp(-\frac{i}{\hbar}tP^{B}(D_{A}))\psi, \qquad t \in \mathbb{R},$$

where $P^{B}(X)$ is a self-adjoint operator,

$$P^{B}(X)\psi = (-i\hbar X - e\alpha(X))\psi,$$

 $\alpha = (B/2)(x_1 dx_2 - x_2 dx_1)$, i.e. $d\alpha = B dx_1 \wedge dx_2$, and $\psi \in C_0^{\infty}(\mathbb{R}^2)$. The real number B can be given physical meaning: a particle with electric charge e moves on the plane \mathbb{R}^2 in an external magnetic field which is perpendicular to the plane and has constant value B (the sign reflects the orientation).

Example 2.3. $M = S^1$, $G = \mathrm{U}(1)$ — the group of rotations of the circle S^1 . Both $\mathcal{M}(\mathrm{U}(1))$ and the isotropy subgroup are trivial; hence there exists exactly one irreducible system of imprimitivity (up to unitary equivalence). Now let us replace $G = \mathrm{U}(1)$ by its universal covering $\tilde{G} = \mathbb{R}$. Then $\mathcal{M}(\mathbb{R}) = \{1\}$ and the isotropy subgroup $H = 2\pi \mathbb{Z}$. The irreducible unitary representations of H are labelled by elements $z \in \mathbb{R}/2\pi \mathbb{Z}$: for $\Phi \in \mathbb{R}$ let

$$L^{\Phi}: 2\pi k \mapsto \exp(\frac{i}{\hbar}ke\Phi), \quad k \in \mathbb{Z};$$

if $(e/\hbar)(\Phi - \Phi') \in 2\pi \mathbb{Z}$, then $L^{\Phi} = L^{\Phi'}$.

We shall describe the system of imprimitivity for given $\Phi \in \mathbb{R}$. The Hilbert space \mathcal{H}^{Φ} consists of (equivalence classes of) functions $\psi \colon \mathbb{R} \to \mathbb{C}$ such that

$$\psi(x+2\pi k) = \exp(-\frac{i}{\hbar}ke\Phi)\psi(x), \quad (k \in \mathbb{Z})$$

almost everywhere; the inner product is defined by

$$(\psi, \psi') = \int_a^{a+2\pi} \psi(x) \overline{\psi'(x)} dx, \qquad a \in \mathbb{R}.$$

We have

$$V^{\Phi}(t) = \exp(-\frac{i}{\hbar}tP^{\Phi}),$$

where $t \in \mathbb{R}$ and

$$P^{\Phi} = -i\hbar \mathrm{d}/\mathrm{d}x$$

is self-adjoint. The mapping

$$W: \mathcal{H}^{\Phi} \to \mathcal{H}^{0}: \psi(x) \mapsto \exp(i\frac{e\Phi x}{2\pi\hbar}).\psi(x)$$

is unitary, \mathcal{H}^0 can be identified with $L^2(S^1, d\varphi)$. We find

$$WP^{\Phi}W^{-1} = -i\hbar \frac{\mathrm{d}}{\mathrm{d}\varphi} - \frac{e\Phi}{2\pi}, \qquad \varphi \in [0, 2\pi).$$

A possible physical interpretation is connected with the Aharonov-Bohm effect [1]: a particle in \mathbb{R}^3 with electric charge e is moving on the circle $x_1^2 + x_2^2 = 1$, $x_3 = 0$, and external magnetic flux Φ is concentrated along the x_3 -axis passing through the centre of the circle. If $(e/2\pi\hbar)(\Phi - \Phi')$ is an integer, then the two quantum kinematics with fluxes $\Phi \neq \Phi'$ lead to the same observable results (e.g., the same interference pattern).

Example 2.4. $M = S^2$, the symmetry group G = SO(3) is replaced by the quantum mechanical symmetry group SU(2) [10] acting on S^2 in the usual way

$$\begin{pmatrix} x_3 & x_1 - ix_2 \\ x_1 + ix_2 & -x_3 \end{pmatrix} \mapsto T \begin{pmatrix} x_3 & x_1 - ix_2 \\ x_1 + ix_2 & -x_3 \end{pmatrix} T^*,$$

where $T \in SU(2)$ and points $(x_1, x_2, x_3) \in S^2 \subset \mathbb{R}^3$ $(\sum_i x_i^2 = 1)$ are identified with the matrices $\sum_i x_i \sigma_i$; $\sigma_1, \sigma_2, \sigma_3$ are the Pauli spin matrices. The isotropy subgroup H of the north pole (0,0,1) consists of the diagonal matrices $\begin{pmatrix} \tau & 0 \\ 0 & \overline{\tau} \end{pmatrix}, \tau \in T^1 \simeq U(1)$, hence $H \equiv U(1)$.

If T is parametrized by $\alpha, \beta \in \mathbb{C}$, $\bar{\alpha}\alpha + \bar{\beta}\beta = 1$,

$$T = T(\alpha, \beta) = \begin{pmatrix} \alpha - \bar{\beta} \\ \beta & \bar{\alpha} \end{pmatrix},$$

then the projection $\pi : \mathrm{SU}(2) \to S^2 \simeq \mathrm{SU}(2)/\mathrm{U}(1)$ is given by

$$\pi(T) = T\sigma_3 T^* = (2 \Re(\bar{\alpha}\beta), 2 \Im(\bar{\alpha}\beta), \bar{\alpha}\alpha - \bar{\beta}\beta).$$

The quadruple $(SU(2), \pi, S^2; U(1))$ constitutes a non-trivial principal bundle known as the Hopf fibration. We shall explicitly write local trivializations of this bundle on sets $U_n = S^2 \setminus \{s\}$, $U_s = S^2 \setminus \{n\}$, where n = (0, 0, 1) and s = (0, 0, -1) are the north and the south pole, respectively. A local trivialization is determined by a selected smooth local section. We choose (in spherical coordinates ϑ , φ)

$$\rho_n: (\vartheta, \varphi) \mapsto T(\cos(\vartheta/2), e^{i\varphi} \sin(\vartheta/2)), \quad O \le \vartheta < \pi,$$

$$\rho_s: (\vartheta, \varphi) \mapsto T(e^{-i\varphi}\cos(\vartheta/2), \sin(\vartheta/2)), \quad O < \vartheta \le \pi.$$

Since SU(2) is simple, connected and simply connected, its multiplier group is trivial. The irreducible representations L^n of the isotropy subgroup $H = \mathrm{U}(1)$ are labeled by integers $n \in \mathbb{Z}$, $L^n : \tau \mapsto \tau^n$. In order to write down explicit expressions for the operators P(X) of generalized momenta, it is convenient to work in the complex line bundle associated (via L^n) with the principal bundle. Then the Hilbert space \mathcal{H} of the canonical system of imprimitivity corresponding to L^n consists of measurable sections ψ in the complex line bundle; each section ψ can be identified with a pair of functions (ψ_n, ψ_s) , where

$$\psi_{n,s} \in L^2(U_{n,s}, \sin \vartheta d\vartheta d\varphi); \quad \psi_s(u) = e^{-in\varphi}\psi_n(u)$$

for almost all $u \in U_n \cap U_s$. We choose $i\sigma_1$, $i\sigma_2$, $i\sigma_3$ as basis of the Lie algebra su(2). The element $-(i/2)\sigma_3$ induces the vector field J_3 on S^2 , $J_3 = x_1(\partial/\partial x_2) - x_2(\partial/\partial x_1)$. Further, the relation

$$\exp(-\frac{i}{\hbar}tP(J_3)) = V(\exp(-\frac{i}{2}t\sigma_3)), \quad t \in \mathbb{R},$$

defines a self-adjoint operator $P(J_3)$ in \mathcal{H} . Operators $P(J_1)$, $P(J_2)$ corresponding to vector fields J_1 , J_2 can be obtained by cyclic permutations. If for $\lambda \in \mathbb{R}^3$, $J = \sum_i \lambda_i J_i$ is a vector field and if $\psi = (\psi_n, \psi_s)$ is a smooth local section, then the self-adjoint operator P(J) is determined by a pair of operators $P_n(J)$, $P_s(J)$; a straightforward calculation yields

$$P_{n,s}(J)\psi(u) = (-i\hbar J - e\alpha_{n,s}(J) - \frac{n\hbar}{2}(\lambda \cdot u))\psi(u),$$

where $u \in U_{n,s}$, $\lambda . u = \sum_{i} \lambda_{i} u_{i}$; 1-forms $i\alpha_{n,s}$ are the localizations on sets $U_{n,s}$ of the connection 1-form $i\alpha$ in the associated bundle. In spherical coordinates

$$\alpha_{n,s} = \frac{n\hbar}{2e} (\pm 1 - \cos \theta) d\varphi,$$

$$i\alpha_s = i\alpha_n + \frac{\hbar}{e} (\mathrm{d}e^{-in\varphi})e^{in\varphi}.$$

(The constant e is again arbitrary, non-zero, but fixed.) On the intersection $U_n \cap U_s$ we find

$$\beta = d\alpha_n = d\alpha_s = \frac{n\hbar}{2e} \sin \theta d\theta \wedge d\varphi.$$

We put

$$g = \int_{S^2} \beta$$
, i.e. $eg = 2\pi n\hbar$.

The situation may have the following physical interpretation: a particle with charge e is moving in the magnetic field of the *Dirac monopole* with magnetic charge g placed at the origin O in \mathbb{R}^3 , so on the sphere $S^2 \subset \mathbb{R}^3$ there is the external magnetic field $B = B(u) = (g/4\pi)u$, $u \in S^2 \subset \mathbb{R}^3$. The relation $eg = 2\pi n\hbar$, $n \in \mathbb{Z}$, coincides with the *Dirac quantization condition* [6, 15]. Operators $P(J_k)$ are the well-known conserved total angular momentum operators for a charged particle moving in the Dirac monopole field.⁸

Example 2.5. $M = \mathbb{R}^1$, G is the group of orientation preserving affine transformations of \mathbb{R}^1 . Having identified M with $\mathbb{R} \times \{1\} \subset \mathbb{R}^2$, G acts on M according to

$$\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} : \begin{pmatrix} x \\ 1 \end{pmatrix} \mapsto \begin{pmatrix} ax + b \\ 1 \end{pmatrix}, \quad a > 0, b \in \mathbb{R}.$$

⁸A more detailed discussion of this example was given in [30]. The first treatments of the magnetic monopole using a connection in a fibre bundle appeared in [19, 36, 37].

The isotropy subgroup H of the origin O consists of all matrices with b=0. The irreducible unitary representations of H are of the form $L^c: a \mapsto a^{ic}, c \in \mathbb{R}$. The Lie algebra \mathcal{G} consists of all matrices

$$A = \begin{pmatrix} A_1 & A_2 \\ 0 & 0 \end{pmatrix}, \qquad A_1, A_2 \in \mathbb{R}.$$

An element $A \in \mathcal{G}$ induces a vector field $D_A = (A_1x + A_2)\mathrm{d}/\mathrm{d}x \in \mathcal{X}(\mathbb{R}^1)$. Let us investigate $\mathcal{M}(G)$: G is connected and simply connected, and a general real skew–symmetric 2-form p on \mathcal{G} is of the form $p:(A,B) \mapsto K(A_1B_2 - A_2B_1)$ for some constant $K \in \mathbb{R}$; but it is easily verified that p(A,B) = q([A,B]) where $q:A \mapsto KA_2$ is a 1-form on \mathcal{G} , so $\mathcal{M}(G)$ is trivial (see Sect. 2.3). Thus the irreducible systems of imprimitivity are labelled by $c \in \mathbb{R}$. Explicitly, $\mathcal{H}^c = L^2(\mathbb{R}, \mathrm{d}x), E^c(S)$ acts via multiplication by indicator function $\chi_S(x)$, and

$$[V^{c}(a,b)\psi](x) = a^{-1/2}e^{ic\ln a}\psi(\frac{x-b}{a}).$$

The self-adjoint generalized momentum operators $P^{c}(D_{A}), A \in \mathcal{G}$, defined by

$$V^{c}(\exp(tA)) = \exp(-itP^{c}(D_{A})/\hbar),$$

are of the form

$$P^{c}(X) = -i\hbar(X + \frac{1}{2}\operatorname{div}X) - \hbar c \operatorname{div}X, \qquad X = D_{A}.$$

3 Quantum Borel kinematics: localization

3.1 Generalized system of imprimitivity

Generally, for a given smooth manifold M there is, a priori, no geometric symmetry group. As indicated in [4, 11, 12, 27], the investigation of vector fields on M is a meaningful starting point. We denote by $\mathcal{X}(M)$ the Lie algebra of smooth vector fields on M, by $\mathcal{X}_0(M)$ its subalgebra of compactly supported vector fields, by $\mathcal{X}_c(M)$ the family of all complete vector fields, $\mathcal{X}_0(M) \subset \mathcal{X}_c(M)$. The flow ϕ^X of a complete vector field X represents a one–parameter group of diffeomorphisms $\{\phi_t^X\}_{t\in\mathbb{R}}$ of M, also called a dynamical system on M. And, vice versa, every dynamical system is a flow of some (uniquely determined) complete vector field

$$[X\psi](u) = \left[\frac{\mathrm{d}}{\mathrm{d}t}(\psi \circ \phi_t^X)(u)\right]_{t=0}.$$

The family of dynamical systems on M will be denoted by D(M). The following theorem summarizes some well-known facts from differential geometry [48], [51].

Theorem 3.1. Let $f: M \to M'$ be a diffeomorphism. Then $f': \mathcal{X}(M) \to \mathcal{X}(M')$, where $(f'.X)_{f(u)} = \mathrm{d}f_u(X_u)$, is a Lie algebra isomorphism; the restriction $f': \mathcal{X}_0(M) \to \mathcal{X}_0(M')$ is also a Lie algebra isomorphism; $f': \mathcal{X}_c(M) \to \mathcal{X}_c(M')$ is a bijection. The mapping

$$f^{D}: D(M) \to D(M'): \{\phi_{t}\}_{t \in IR} \mapsto \{f \circ \phi_{t} \circ f^{-1}\}_{t \in IR}$$

is bijective and $f^D(\phi^X) = \phi^{f'.X}$.

For every $\phi^X \in D(M)$ the manifold M becomes a G-space for the group $G = \mathbb{R}$. Attempting to generalize Mackey's quantization (Sects. 2.3, 2.4) we require that there exist: a

Hilbert space \mathcal{H} , a projection–valued measure E on M and unitary representations V^X in \mathcal{H} of the flows $\phi^X \in D(M)$ such that

$$V^{X}(t) E(S) V^{X}(-t) = E(\phi_{t}^{X}.S),$$
 (3.1)

where the objects \mathcal{H} , E do not depend on the choice of $\phi^X \in D(M)$. Equation (3.1) is just a generalization of (2.2). Geometric shifts of Borel sets $S \in \mathcal{B}(M)$ by flows ϕ_t^X along complete vector fields X are represented in \mathcal{H} by unitary operators V(t) such that (3.1) holds.

Generalized momentum operators can then be introduced via Stone's Theorem as (essentially self-adjoint) infinitesimal generators P(X) of the one-parameter groups of unitary operators — shifts in \mathcal{H} of the localized quantum system,

$$V^X(t) = \exp(-\frac{i}{\hbar}P(X)t), \qquad t \in \mathbb{R}.$$

3.2 Quantum Borel kinematics

The quantization of 'classical' Borel kinematics $(\mathcal{B}(M), \mathcal{X}_c(M))$ thus requires [11] the imprimitivity condition (3.1) for the unitary representation of the flow of each complete vector field individually. Then we can state

Definition 3.1. Quantum Borel kinematics is a pair (V, E), where E is a projection-valued measure on M in a separable Hilbert space \mathcal{H} , and V associates with each $\phi^X \in D(M)$ a homomorphism $V^X \colon \mathbb{R} \to U(\mathcal{H})$ such that the following conditions are satisfied:

- 1) Equation (3.1) holds for all $t \in \mathbb{R}$, $X \in \mathcal{X}_c(M)$, $S \in \mathcal{B}(M)$;
- 2) The mapping $P: X \mapsto P(X)$ from the Lie algebra $\mathcal{X}_0(M)$ into the space of essentially self-adjoint operators with common invariant dense domain in \mathcal{H} is a Lie algebra homomorphism:

$$P(X + aY) = P(X) + aP(Y), \tag{3.2}$$

$$[P(X), P(Y)] = -i\hbar P([X, Y];$$
 (3.3)

3) Locality condition. If two flows $\phi^{X_i} \in D(M)$, i = 1, 2, after restriction on the set $(-a, a) \times S$, a > 0, $S \in \mathcal{B}(M)$, coincide, then the mappings $\mathbb{R} \times \mathcal{H} \to \mathcal{H}: (t, \psi) \mapsto V^{X_i}(t) \psi$ coincide on the domain $(-a, a) \times \mathcal{H}_S$, where \mathcal{H}_S is the subspace of \mathcal{H} projected out by E(S).

If in 2) only linearity (3.2) is required, we shall call (V, E) a generalized system of imprimitivity for D(M).⁹ It will describe quantum Borel kinematics with external gauge field.

It follows from condition 3) that, if V^X is known for all $X \in \mathcal{X}_0(M)$, then it is determined for all $X \in \mathcal{X}_c(M)$; 3) further implies that P(X) are differential operators. Condition 2) may sometimes be too restrictive since (3.3) excludes a non-vanishing external gauge field on M; in this connection see [12, 13, 33] and also Ex. 2.2, Ex. 2.4 and Sect. 4.7.

The projection–valued measure E induces in a natural way a quantization Q of classical (smooth) real functions $f: M \to \mathbb{R}$ on configuration space (e.g. coordinate functions, potentials, etc.). Not necessarily bounded, self–adjoint quantum position operators Q(f) are uniquely determined by their spectral decompositions

$$Q(f) = \int_{-\infty}^{\infty} \lambda \, \mathrm{d}E_{\lambda}^{f} \,,$$

The family D(M) acts transitively on M (we suppose M to be connected, without boundary): for any two points $u, v \in M$ there exists $X \in \mathcal{X}_0(M)$, $\phi^X \in D(M)$ such that $\phi^X(1, u) = v$.

where the spectral function E_{λ}^f is given by the spectral measure $E^f(\Delta) = E(f^{-1}(\Delta))$ on subsets $\Delta = (-\infty, \lambda)$ of \mathbb{R} . Equation (3.1) is then replaced by

$$V^{X}(t) Q(f) V^{X}(-t) = Q(f \circ \phi_{-t}^{X}), \tag{3.4}$$

where $f \in C^{\infty}(M, \mathbb{R})$, and implies a generalization of the Heisenberg commutation relations in terms of coordinate-independent objects

$$[Q(f), P(X)] = i\hbar Q(X.f) \quad \text{on} \quad \mathcal{D} \subset \mathcal{H}. \tag{3.5}$$

It is assumed that operators P(X), Q(f) have a common invariant dense domain \mathcal{D} in \mathcal{H} . If an obvious relation

$$[Q(f), Q(g)] = 0 \quad \text{on} \quad \mathcal{D}$$
(3.6)

for all $f, g \in C^{\infty}(M, \mathbb{R})$ is still added, then (3.6), (3.5) and (3.3) define a *Schrödinger system* in the sense of [27].

We can say that Borel quantization on a smooth configuration manifold M associates the generalized position Q(f) and momentum operators P(X) with smooth functions $f \in C^{\infty}(M, \mathbb{R})$ and smooth vector fields $X \in \mathcal{X}(M)$, respectively. These quantum kinematical observables on M are globally defined, hence Borel quantization incorporates the global structure of M.

Remark. The natural infinite–dimensional Lie algebra structure (3.6), (3.5) and (3.3) of quantum Borel kinematics should be compared with the non–relativistic local current algebra for a Schrödinger second quantized field over $M = \mathbb{R}^3$ studied in [18]:

$$[\rho(f_1), \rho(f_2)] = 0,$$

 $[\rho(f), J(X)] = i\hbar \rho(Xf),$
 $[J(X), J(Y)] = -i\hbar J([X, Y]),$

where $f, f_i \in C^{\infty}(\mathbb{R}^3, \mathbb{R}), X, Y \in \mathcal{X}(\mathbb{R}^3)$. There is an apparent algebraic correspondence of the local density operator $\rho(f)$ with Q(f) and the local current operator J(X) with P(X). In both cases the representations of the algebraic structures yield possible quantum kinematics. However, in contrast to quantum Borel kinematics, where Q(f) is a multiplication operator, local current algebra is more general, since also representations where $\rho(f_1f_2)$ is not equal to $\rho(f_1)\rho(f_2)$ are admitted.

3.3 Quasi-invariant measures, projection–valued measures

The question of existence and uniqueness of a measure which is *quasi-invariant* with respect to all diffeomorphisms $\phi_1: u \mapsto \phi(1, u)$ for which $\phi \in D(M)$, is answered by

Theorem 3.2. The family of quasi-invariant measures on $\mathcal{B}(M)$ is non-empty and, moreover, all measures in this family are mutually equivalent and form a unique invariant measure class.¹⁰ After completion, those subsets in M which have measure zero are exactly measure zero sets in the sense of Lebesgue.

Proof: The fact that the family of sets of zero measure in the sense of Lebesgue is invariant under diffeomorphisms is well known [44]. The existence part of the theorem can be seen as follows. Having embedded M in \mathbb{R}^m (Whitney's Theorem), we can consider a tubular

¹⁰The invariant measure class is called the *Lebesque measure class*.

neighborhood M^{ε} of M in the normal bundle ([44], Chap. 2.3). Denoting by $\pi: M^{\varepsilon} \to M$ the associated submersion, we can define $\mu(S) = \lambda^m(\pi^{-1}(S))$ for $S \in \mathcal{B}(M)$, where λ^m denotes the Lebesgue measure in \mathbb{R}^m ; then μ is quasi-invariant. The assertion about uniqueness of the invariant measure class for $M = \mathbb{R}^n$ follows from the fact that the family of diffeomorphisms ϕ_1 includes all translations and the assertion for the group of translations is known ([54], Chap. II.3). In general, M can be covered by a countable family of open sets, each of which is diffeomorphic to \mathbb{R}^n and so the assertion is true again. \square

Measure μ will be called differentiable if the mapping $\mathbb{R} \to \mathbb{R}$: $t \mapsto \mu(\phi_t.S)$ is smooth for all $S \in \mathcal{B}(M)$, $\phi \in D(M)$. For instance, measure μ used in the proof of Theorem 3.2 is differentiable. Every manifold is locally orientable; having fixed an orientation on an open set $U \subset M$ (dim M = n), then to every differentiable measure μ exactly one n-form ω exists on U such that

$$\int_{U} f \cdot \omega = \int_{U} f(u) d\mu(u) \quad \text{for all } f \in C_0^{\infty}(U);$$

 $\omega_u(X_1,\ldots,X_n)>0$ for every positively oriented basis in T_uU . For each $X\in\mathcal{X}(U)$ we define a function div_uX on U by

$$\operatorname{div}_{\mu} X.\omega = k! \operatorname{d}(i_{X}\omega), \tag{3.5}$$

where the (n-1)-form $i_X\omega$ is defined by

$$(i_X\omega)(X_1,\ldots,X_{n-1}) = \omega(X,X_1,\ldots,X_{n-1}).$$

Equivalently, if $\phi = \phi^X \in D(M)$, then

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\mathrm{d}\mu}{\mathrm{d}(\mu \circ \phi_t)} \right) (\phi_{-t} \cdot u) \mid_{t=0} = -\mathrm{div}_{\mu} X. \tag{3.6}$$

In local coordinates, if $\omega = \rho(u)du_1 \wedge \ldots \wedge du_n$, $\rho > 0$, then

$$\operatorname{div}_{\mu} X = X \cdot \ln \rho + \sum_{i} \frac{\partial X_{j}}{\partial u_{j}}.$$

The structure of projection-valued measures on M is well known. According to [54], Chap. IX.4, we have a canonical representation of a localized quantum system: Let E be a projection-valued measure on M in a separable Hilbert space \mathcal{H} . Then there exist two sequences $\{\mathcal{K}_r\}$, $\{\nu_r\}$, $r=\infty,1,2,\ldots$, the first one consisting of Hilbert spaces, the other of measures on M such that dim $\mathcal{K}_r=r$ and ν_r , ν_s are mutually singular¹¹ for $r\neq s$. The projection-valued measure E is unitarily equivalent to the measure E^0 which acts via multiplication by indicator functions of subsets in the Hilbert space $\mathcal{H}^0=\bigoplus_r \mathcal{H}_r$, where \mathcal{H}_r are the Hilbert spaces of vector-valued functions from E to E to E to non-zero, the projection-valued measure E is called homogeneous. Due to the transitivity of actions of the family E to the following theorem holds (for details see [54], Chap. IX.5, IX.6):

Theorem 3.3. If (V, E) is a generalized system of imprimitivity for D(M), then E is homogeneous. The unique non-zero measure ν_r belongs to the Lebesgue measure class on M.

Thus the canonical representation of a localized quantum system on M involves a smooth measure, i.e. a measure induced by the Lebesgue measure of the coordinate charts. An r-homogeneous localized quantum system of degree r > 1 can be interpreted as a quantum

¹¹ i.e., there exist $S_r, S_s \in \mathcal{B}(M), S_r \cup S_s = M, S_r \cap S_s = \emptyset, \nu_r(S_s) = \nu_s(S_r) = 0.$

system with *internal degrees of freedom*; a 1-homogeneous localized quantum system will be called *elementary* as its E's are related to elementary spectral measures. We shall need also

Theorem 3.4. ([54], Chap. IX.2). Let μ be a Borel measure on M, \mathcal{K} a Hilbert space, $\mathcal{H} = L^2(M, \mathcal{K}, \mu)$, E a projection–valued measure on $\mathcal{B}(M)$ acting in \mathcal{H} via multiplication by indicator functions. Then any bounded operator B in \mathcal{H} commuting with E(S) for all $S \in \mathcal{B}(M)$ ($B \in \mathcal{C}(E)$) is of the form

$$B: \psi(u) \mapsto b(u).\psi(u),$$

where b is a Borel mapping from M into the space of bounded operators in \mathcal{K} such that $\sup_{u \in M} |b(u)| < \infty$. Function b is determined by B uniquely on M modulo a set of μ -measure zero.

3.4 An example

Quantum Borel kinematics are rather diverse (even for trivial configuration space \mathbb{R}^n), as the following example [33] shows. Let $M = \mathbb{R}^n$ with a fixed basis. To every vector field $X = \sum_k X_k \partial/\partial x_k$ we relate a matrix-valued function

$$A(X): \mathbb{R}^n \to \mathbb{R}^{n,n}, \ [A(X)_u]_{i,j} = \frac{\partial X_i(u)}{\partial x_i}.$$

It is straightforward to verify

$$[A(X), A(Y)] - X.A(Y) + Y.A(X) = -A([X, Y]).$$
(3.7)

Let L be a skew-Hermitean representation of the Lie algebra $\mathrm{gl}^+(n, \mathbb{R})$ in a Hilbert space \mathcal{H}^L . We define operators Q(f), P(X) in $\mathcal{H} = L^2(\mathbb{R}^n, \mathcal{H}^L, \mathrm{d} x_1 \dots \mathrm{d} x_n)$ by

$$Q(f)\psi = f.\psi, \quad P(X)\psi = -i\hbar[X + \frac{1}{2}\operatorname{div}X - L(A(X))].$$

Then using (3.7) and the identity X.L(A(Y)) = L(X.A(Y)), the pair (P,Q) can be shown to be a quantum Borel kinematic. Choosing the representation L in $\mathcal{H}^L = \mathbb{C}$ to be given by

$$L(A) = -ic\operatorname{tr} A,$$

where c is a real constant, we obtain

$$L(A(X)) = -ic \operatorname{div} X.$$

This is just an example of the divergence term which we shall encounter in Sect. 4.7.

Remark. Consider the surjective mapping

$$p: \operatorname{gl}(n, \mathbb{R}) \to \operatorname{sl}(n, \mathbb{R}): A \mapsto A - \frac{1}{n} (\operatorname{tr} A) \mathbb{1}_n,$$

where \mathbb{I}_n is the unit $n \times n$ -matrix. This mapping permits to associate with every representation L' of $sl(n, \mathbb{R})$ a representation $L = L' \circ p$ of $gl(n, \mathbb{R})$. Then our mapping P is the infinitesimal form of a representation of the group of diffeomorphisms of \mathbb{R}^n induced from $SL(n, \mathbb{R})$ (see [18] for n = 3). For general results concerning the "divergence–like" terms in the framework of systems of imprimitivity for the group of diffeomorphisms, see [31].

4 Quantum Borel kinematics: external gauge fields

4.1 External magnetic field

In order to motivate our construction of quantum Borel kinematics with external field via generalized systems of imprimitivity for D(M), let us consider quantum kinematics on \mathbb{R}^3 , for a charged particle in an external magnetic field \vec{B} . Since $\operatorname{div} \vec{B} = 0$, the Poincaré lemma implies that there exists a vector potential \vec{A} such that $\vec{B} = \operatorname{rot} \vec{A}$. The Hilbert space is $\mathcal{H} = L^2(\mathbb{R}^3, \mathrm{d}^3x)$, and the Hamiltonian

$$H = -\frac{\hbar^2}{2m} \sum_{j} \left(\frac{\partial}{\partial x_j} - i \frac{e}{\hbar} A_j \right)^2.$$

If another potential $\vec{A'}$, $\operatorname{rot} \vec{A'} = \vec{B}$, is chosen, then according to the Poincaré theorem there exists a real function λ such that $\vec{A'} = \vec{A} + \operatorname{grad} \lambda$. The Hamiltonian H is transformed as $H \mapsto H' = WHW^{-1}$, where W is the unitary mapping

$$W: \mathcal{H} \to \mathcal{H}': \psi \mapsto \exp(\frac{ie}{\hbar}\lambda)\psi.$$

This common quantum mechanical scheme can be reformulated in geometric language [36]: \mathcal{H} is the space of measurable sections in the trivial Hermitian complex line bundle $\mathbb{R}^3 \times \mathbb{C}^1$ associated with the principal bundle $\mathbb{R}^3 \times \mathbb{T}^1$;

$$i(e/\hbar)\alpha, \qquad \alpha = \sum A_j dx_j,$$

is a localized connection 1-form;

$$i(e/\hbar)\beta$$
, $\beta = d\alpha = B_1 dx_2 \wedge dx_3 + \text{cvcl.}$

is the curvature 2-form;

$$\vec{x} \mapsto \exp(\frac{ie}{\hbar}\lambda(\vec{x}))$$

is a transition function in the principal bundle relating two different trivializations.

In general, if a magnetic field is given by a closed 2-form β on manifold M, a vector potential 1-form α such that $\beta = d\alpha$ need not exist. However, following [36], one can always define vector potentials α_k locally, i.e. on open sets U_k (diffeomorphic to \mathbb{R}^n) such that $\{U_k\}$ is an open covering of M. We require that vector potentials α_k , α_j be related on the intersection $U_k \cap U_j$ by a gauge transformation

$$i\alpha_j = i\alpha_k + \frac{\hbar}{e}(\operatorname{d}\exp(\frac{ie}{\hbar}\lambda))\exp(-\frac{ie}{\hbar}\lambda) = i(\alpha_k + \operatorname{d}\lambda),$$

where $\exp(\frac{ie}{\hbar}\lambda(\vec{x}))$ is a transition function. Then $d\alpha_j = d\alpha_k$. In this way we shall construct a principal bundle with typical fibre T^1 and connection $\{i\alpha_j\}$.

4.2 Construction of a class of generalized systems of imprimitivity

We choose a measure μ from the Lebesgue measure class on M and four objects (P,G,Γ,L) , where: $P \equiv (P,\pi,M,G)$ (or shortly P(M,G)) is a principal bundle over M; its typical fibre G is an Abelian or compact Lie group; Γ is a connection in P, and L is a unitary representation of G in a finite-dimensional Hilbert space $\mathcal K$ with inner product <.,.>. We construct a separable Hilbert space $\mathcal H$ like in Sect. 2.3, consisting of vector-valued functions $\psi:P\mapsto \mathcal K$, such that

- 1. $x \mapsto \langle \psi(x), f \rangle$ is a Borel function on P for all $f \in \mathcal{K}$,
- 2. $\psi(xg) = L_a^{-1}\psi(x), g \in G$ (ψ is an equivariant function).
- 3. $\|\psi\| < \infty$, where $\|.\|$ is the norm induced by the inner product

$$(\psi, \psi') = \int_{M} \langle \psi(x), \psi'(x) \rangle d\mu(u).$$

The integral is well defined as the integrand remains constant on the fibres. Two functions ψ, ψ' are identified if they coincide almost everywhere. The projection–valued measure E on M is defined via multiplication by indicator functions:

$$E(S)\psi = \tilde{\chi}_S \psi, \quad \tilde{\chi}_S(x) = \begin{cases} 1, \, \pi(x) \in S \\ 0, \, \pi(x) \notin S \end{cases} \tag{4.1}$$

For $\phi \in D(M)$ we define the unitary representation of the additive group \mathbb{R}

$$[V^{\phi}(t)\psi](x) = \sqrt{\frac{\mathrm{d}\mu}{\mathrm{d}(\mu \circ \phi_t)}}(\phi_{-t}u)\psi(\tilde{\phi}_{-t}x), \quad u = \pi(x). \tag{4.2}$$

Here $\tilde{\phi}$ denotes the horizontal lift of the flow ϕ on M.

It is easily verified that the constructed pair (V, E) is a generalized system of imprimitivity for D(M) in the sense of Definition 3.1. The linearity of mapping P required by this definition will be investigated in Sect. 4.3 (see Eq. (4.7)).

The equivalence class of (V, E) does not depend on measure μ ; if μ' is another measure from the Lebesgue class, then $\mathcal{H} \to \mathcal{H}' : \psi \mapsto \sqrt{\mathrm{d}\mu/\mathrm{d}\mu'} \,\psi$ is the desired unitary mapping. For simplicity we shall suppose μ to be differentiable. We assumed G to be Abelian or compact in order to deal only with finite-dimensional unitary representations L of G. For dim $\mathcal{K} = \nabla$ we identify \mathcal{K} with \mathbb{C}^r endowed with the standard inner product. We say that the pair (V, E) is a generalized system of imprimitivity specified by the quadruple (P, G, Γ, L) .

We note that, for two diffeomorphic manifolds, there is a one–to–one correspondence between the equivalence classes of generalized systems of imprimitivity constructed in this way.

4.3 Construction in the associated vector bundle

The associated vector bundle $(F, \bar{\pi}, M; \mathbb{C}^r)$ will be constructed in the standard way (see e.g. [48, 51]). We introduce equivalence relation

$$(x;\xi) \sim (xg; L_g^{-1}\xi), \qquad g \in G,$$

on $P \times \mathbb{C}^r$ and put $F = P \times \mathbb{C}^r / \sim$; the projection is $\bar{\pi} : F \to M : [x; \xi] \mapsto \pi(x)$. Each fibre becomes an r-dimensional Hilbert space:

$$\lambda[x;\xi] + [x;\xi'] = [x;\lambda\xi + \xi'],$$

< $[x;\xi], [x;\xi'] > = < \xi, \xi' >, \qquad \pi(x) = u.$

The inner product is well–defined since L is unitary. For the same reason we can relate a unitary mapping \hat{x} to each $x \in P$:

$$\hat{x}: \mathbb{C}^r \to F_{\pi(x)}: \xi \mapsto [x; \xi]$$

and we have $\widehat{xg} = \hat{x} \circ L_q$.

Let \mathcal{H} be the Hilbert space of measurable sections in the associated vector bundle, having finite norm induced by the inner product

$$(\sigma, \tau) = \int_{M} \langle \sigma(u), \tau(u) \rangle d\mu(u). \tag{4.3}$$

We can define a unitary mapping $T: \mathcal{H} \to \bar{\mathcal{H}}: \psi \mapsto \sigma$ in a natural way:

$$\sigma(u) = [x; \psi(x)], \qquad \pi(x) = u.$$

The definition of T is correct since $(x; \psi(x)) \sim (xg; L_g^{-1}\psi(x))$ and $L_g^{-1}\psi(x) = \psi(xg)$. The inverse mapping $T^{-1}: \sigma \mapsto \psi$ is given by $\psi(x) = \hat{x}^{-1} \circ \sigma(\pi(x))$. Having performed the unitary transform T we replace (V, E) by a generalized system of imprimitivity (\bar{V}, \bar{E}) in $\bar{\mathcal{H}}$. We shall compute explicit expressions.

Clearly, \bar{E} acts via multiplication by indicator functions

$$\bar{E}(S)\sigma = \chi_S.\sigma. \tag{4.4}$$

In order to express \overline{V} we must first describe induced connection in the associated vector bundle F. Having Hermitian structure on the fibres, we consider only Hermitian connections on F, i.e. connections for which all linear isomorphisms $C(u_t): F_{u_0} \to F_{u_t}$ (shortly C_t , see below), which belong to curves u_t in M, are unitary. As is well known, there is a one-to-one correspondence between Hermitian connections and Hermitian covariant derivatives in F. A covariant derivative ∇ acting on smooth sections, $\nabla_X : \operatorname{Sec} F \to \operatorname{Sec} F$, $X \in \mathcal{X}(M)$, is Hermitian, if it satisfies (in addition to four conditions [48] defining the covariant derivative) the identity

$$X < \sigma, \tau > = < \nabla_X \sigma, \tau > + < \sigma, \nabla_X \tau >;$$

Sec F denotes the linear space of smooth sections in F.

A connection Γ in P induces a Hermitian connection in F; given a piecewise smooth curve u_t in M and its lift x_t in P, then C_t : $F_{u_0} \to F_{u_t}$, $C_t = \hat{x_t} \circ \hat{x_0}^{-1}$ is the desired unitary mapping. Its definition does not depend on the starting point x_0 , since $(\widehat{x_t g}) \circ (\widehat{x_0 g})^{-1} = \hat{x_t} \circ \hat{x_0}^{-1}$, and $x_t g$ is another horizontal lift of the curve u_t . Now the Hermitian covariant derivative is defined by the limit

$$(\nabla_X \sigma)(u_0) = \lim_{t \to 0} \frac{1}{t} [C_t^{-1} \circ \sigma(u_t) - \sigma(u_0)], \quad \dot{u}_0 = X. \tag{4.5}$$

We are now in the position to give explicit formula for $\bar{V}^{\phi}(t)$:

$$\bar{V}^{\phi}(t)\sigma(u) = C(\phi_{-t}u)^{-1} \left[\sqrt{\frac{\mathrm{d}\mu}{\mathrm{d}(\mu \circ \phi_t)}}\sigma\right](\phi_{-t}u). \tag{4.6}$$

Let Sec₀ F denote the linear space of smooth and compactly supported sections in F; the subspace Sec₀ F is dense in $\bar{\mathcal{H}}$. As before, a self-adjoint operator $\bar{P}(X)$ is defined by

$$\bar{V}^{\phi^X}(t) = \exp(-\frac{i}{\hbar}\bar{P}(X)t);$$

using (4.6), (4.5) and (3.6) we find

$$\bar{P}(X)\sigma = -i\hbar(\nabla_X + \frac{1}{2}\mathrm{div}_\mu X)\sigma, \quad \sigma \in \mathrm{Sec}_0 F.$$
 (4.7)

Note that expression (4.7) implies that $\bar{P} \sim P$ is linear in $X \in \mathcal{X}_{l}(\mathcal{M})$.

Using unitary representation L we can associate a principal bundle $\tilde{P}(M, \mathrm{U}(r))$ to the principal bundle P(M,G). The construction is similar to that for F. On $P \times \mathrm{U}(r)$ we introduce the equivalence relation $(x;a) \sim (xg; L_g^{-1}a), g \in G$, and put $\tilde{P} = P \times \mathrm{U}(r)/\sim, \tilde{\pi}: [x;a] \mapsto \pi(x)$. Lie group $\mathrm{U}(r)$ acts on \tilde{P} via $[x;a].b = [x;ab], b \in \mathrm{U}(r)$. The mapping $f: P \to \tilde{P}: x \mapsto [x;e]$ is a bundle homomorphism — we have $f(xg) = f(x)L_g$. With the help of this homomorphism we can transform the connection Γ in P into a connection $\tilde{\Gamma}$ in \tilde{P} , as described in the following theorem.

Theorem 4.1. ([51], Chap. II.5; [48]). Let $f: P(M,G) \to \tilde{P}(M,\tilde{G})$ be a principal bundle homomorphism and Γ a connection in P. Then there exists a unique connection $\tilde{\Gamma}$ in \tilde{P} such that the tangent mapping f_* maps every horizontal subspace of connection Γ onto a horizontal subspace of $\tilde{\Gamma}$.

The generalized system of imprimitivity (\tilde{V}, \tilde{E}) specified by $(\tilde{P}, U(r), \tilde{\Gamma}, id)$, with $id: U(r) \to U(r)$ being the identity mapping (fundamental representation), is equivalent to (V, E). We describe the corresponding unitary mapping. If $\psi \in \mathcal{H}$, there is a unique vector $\tilde{\psi} \in \tilde{\mathcal{H}}$ such that $\tilde{\psi}(f(x)) = \psi(x)$. Indeed, if f(x) = f(x'), then x' = xg and $L_g = e$, hence $\psi(x) = \psi(x')$ for each $\psi \in \mathcal{H}$. The mapping defined in this way is unitary and transforms E in \tilde{E} since $\tilde{\pi}(f(x)) = \pi(x)$; V is transformed in \tilde{V} since f preserves the connection.

We can again associate a vector bundle \tilde{F} to the principal bundle \tilde{P} using the fundamental representation id. Both \tilde{F} and F have the same base space and the same typical fibre. In fact, F and \tilde{F} can be identified by the mapping

$$W: F \to \tilde{F}: [x; \xi] \mapsto [f(x); \xi],$$

where the homomorphism $f: P \to \tilde{P}$ was described above. W is well–defined since

$$[f(xg); L_g^{-1}\xi] = [f(x)L_g; L_g^{-1}\xi] = [f(x); \xi].$$


W is surjective because $(\tilde{x};\xi) \sim (\tilde{x}a^{-1};a\xi) = (f(x);a\xi)$ for each $\tilde{x} = [x;a] \in \tilde{P}$. W is injective because, for each $u \in M$, the induced mapping $W_u : F_u \to \tilde{F}_u$ is unitary.

There is again a Hermitian covariant derivative $\tilde{\nabla}$ in \tilde{F} which corresponds to the Hermitian connection $\tilde{\Gamma}$ in \tilde{P} . We can briefly say that the following diagram commutes:

$$P(M,G),\Gamma \xrightarrow{f} \tilde{P}(M,U(r)),\tilde{\Gamma}$$

$$F(M,\mathbb{C}^r),\nabla$$

For the corresponding generalized systems of imprimitivity we find the following commuting diagram:



All these generalized systems of imprimitivity are mutually unitarily equivalent.

4.4 The Case G = U(r)

With the help of the identical representation id of $\mathrm{U}(r)$ in \mathbb{C}^r we can associate a vector bundle $F(M,\mathbb{C}^r)$ to every principal bundle $P(M,\mathrm{U}(r))$. To each $x\in P$ there corresponds the unitary mapping \hat{x} which we henceforth denote by the same letter x, namely $x:\mathbb{C}^r\to F_{\pi(x)}:\xi\mapsto [x;\xi]$. Conversely, let $w:\mathbb{C}^r\to F_{\pi(x)}$ be a unitary mapping. Then w necessarily has the form $\xi\mapsto [x;\xi^w]$, where $\xi\mapsto \xi^w$ is a unitary mapping $\mathbb{C}^r\to \mathbb{C}^r$. Hence there exists $a\in \mathrm{U}(r)$ such that $w\xi=[x;a\xi]=[xa;\xi]=xa\xi$. Every unitary mapping $\mathbb{C}^r\to F_u$ which is represented by unitary mapping $xa:\mathbb{C}^r\to F_u$ is equal to the composition $x\circ a$ since $[xa;\xi]=[x;a\xi]$. Further, two points $x,x'\in P$ coincide as unitary mappings if and only if x=x'. So we can return from the vector bundle $F(M,\mathbb{C}^r)$ back to the principal bundle $P(M,\mathrm{U}(r))$. The fibre P_u over $u\in M$ consists of unitary mappings $\mathbb{C}^r\to F_u$ and the structure group $G=\mathrm{U}(r)$ acts on P by composition $a:x\mapsto x\circ a$.

Definition 4.1. Two principal bundles P(M,G), P'(M,G) over the same base space and with the same structure group are said to be *isomorphic* if there exists a diffeomorphism $f: P \to P'$ fulfilling

- 1. $\pi'(f(x)) = \pi(x)$,
- 2. $f(x \circ a) = f(x)a$ for all $a \in G$.

Two Hermitian vector bundles $F(M, \mathbb{C}^r)$, $F'(M, \mathbb{C}^r)$ are said to be *isomorphic* if there exists a diffeomorphism $W: F \to F'$ such that the restrictions $W_u: F_u \to F'_u$ are unitary mappings for all $u \in M$.

Lemma 4.1. Two principal bundles both with structure group U(r) are isomorphic if and only if the corresponding associated vector bundles (with typical fibres \mathbb{C}^r) are isomorphic.

Proof. If $f: P \to P'$ is an isomorphism, then $W: F \to F': [x; \xi] \mapsto [f(x); \xi]$ is an isomorphism of the associated vector bundles. Conversely, let $x \in P$ be a unitary mapping $\mathbb{C}^r \to F_u$; then $W_u \circ x$ is unitary mapping $\mathbb{C}^r \to F'_u$ and there exists a unique $x' \in P'_u$ such that $x' = W_u \circ x$. The mapping $P \to P': x \mapsto W_{\pi(x)} \circ x$ is the desired isomorphism. \square

As already shown, a connection in a principal bundle can be carried over to the associated vector bundle. If G = U(r), L = id, this procedure can be inverted. Let us take a Hermitian connection in the vector bundle $F(M, \mathbb{C}^r)$ associated with principal bundle P(M, U(r)). The connection relates a family of unitary mappings $C_t : F_{u_0} \to F_{u_t}$ to every piecewise smooth curve u_t on M. If $x_0 \in P$, $\pi(x_0) = u_0$, then $x_t = C_t \circ x_0$ will be the lift in P of the curve u_t with the initial point x_0 . We have $(C_t \circ x_0)a = C_t \circ (x_0a)$. This lifting prescription determines a unique connection Γ in the principal bundle P. This correspondence between the connections in P and the Hermitian connections in F is one—to—one.

Definition 4.2. Connections Γ , Γ' in principal bundles P(M,G), P'(M,G), respectively, are said to be *isomorphic*, if there exists an isomorphism $F: P \to P'$ which maps connection Γ in connection Γ' . Hermitian covariant derivatives ∇ , ∇' in Hermitian vector bundles $F(M, \mathbb{C}^r)$, $F'(M, \mathbb{C}^r)$, respectively, are said to be *isomorphic* if there exists an isomorphism $W: F \to F'$ such that $\nabla' = W \nabla W^{-1}$; more precisely,

$$\nabla_X' \sigma(u) = W_u(\nabla_X W_u^{-1} \sigma(u)).$$

Lemma 4.2. Connections Γ, Γ' in principal bundles P(M, U(r)), P'(M, U(r)), respectively, are isomorphic if and only if the corresponding Hermitian covariant derivatives in the associated vector bundles (with typical fibres \mathbb{C}^r) are isomorphic.

Proof. According to Lemma 4.1, to every isomorphism $f: P \to P'$ there exists an isomorphism $W: F \to F'$ fulfilling $f(x) = W_{\pi(x)} \circ x$, and conversely. If f transforms Γ in Γ' and if x_t is the lift in P of a curve u_t with starting point x_0 , then $x'_t = f(x_t)$ is the lift in P' of the same curve with starting point $f(x_0)$. We have: if $C_t = x_t \circ x_0^{-1}: F_{u_0} \to F_{u_t}$, then $C'_t = x'_t \circ x_0^{-1} = W_{u_t}C_tW_{u_0}^{-1}$. Using the last relation we obtain $\nabla' = W\nabla W^{-1}$. Conversely, the relation $\nabla' = W\nabla W^{-1}$ implies $C'_t = W_{u_t}C_tW_{u_0}^{-1}$ and so $x'_t = W_{\pi(x_t)} \circ x_t$. this means that the isomorphism $f: x \mapsto W_{\pi(x)} \circ x$ preserves the connection. \square

4.5 Unitary equivalence of generalized systems of imprimitivity

Theorem 4.2. Two generalized systems of imprimitivity (V_j, E_j) specified by quadruples $(P_j, G_j, \Gamma_J, L_j)$, j = 1, 2, are unitarily equivalent, if and only if the corresponding covariant derivatives in the associated vector bundles are isomorphic.

Proof. First we have the necessary condition that the Hilbert spaces of representations L_1 , L_2 should have the same dimension, say r. According to the results of Sect. 4.3 we can equivalently take the generalized systems of imprimitivity (\bar{V}_j, \bar{E}_j) , j = 1, 2, constructed in the associated vector bundles and investigate their unitary equivalence.

Thus consider a unitary mapping $W: \bar{\mathcal{H}}_1 \to \bar{\mathcal{H}}_2$ relating the two generalized systems of imprimitivity. Let $U(F_1, F_2)$ denote the fibre bundle over M with each fibre over $u \in M$ consisting of unitary mappings $F_{1u} \to F_{2u}$. From the equality $\bar{E}_2(S) = W\bar{E}_1(S)W^{-1}$ it follows that W is induced by a measurable section $u \mapsto W(u)$ in the bundle $U(F_1, F_2)$. From the equality $\bar{V}_2^{\phi}(t) = W\bar{V}_1^{\phi}(t)W^{-1}$ and from relation (4.6) one deduces that

$$C_2(\phi_t u) = W(\phi_t u)C_1(\phi_t u)W^{-1}(u)$$
(4.8)

holds for all $\phi \in D(M)$, all $t \in \mathbb{R}$ and for almost all (depending on t) $u \in M$. Here $C_j(\phi_t u)$, j = 1, 2, are unitary mappings corresponding to the curve $u_t = \phi_t u$.

We shall show that the measurable section $u \mapsto W(u)$, after proper redefinition on a set of measure zero, is smooth. It is sufficient to verify this assertion locally, i.e. to investigate the case $M = \mathbb{R}^n$.

Let us consider all constant vector fields on \mathbb{R}^n , the corresponding flows and t = 1. Thus we have for each $v \in \mathbb{R}^n$ and almost all $u \in \mathbb{R}^n$ the equality

$$W(u+v) = C_2(u+v)W(u)C_1^{-1}(u+v), (4.9)$$

where $C_j(w): F_{ju} \to F_{jw}$, j = 1, 2. It follows from Fubini's Theorem that, for almost all u, the equality (4.9) holds true for almost all v. We need one such u. Then we can say that for almost all $w \in \mathbb{R}^b$ we have

$$W(w) = C_2(w)W(u)C_1^{-1}(w).$$

Here the right-hand side depends differentiably on w and this proves our assertion that the section W can be defined in smooth manner. From the identity (4.8)

$$\nabla_X^2 = W \nabla_X^1 W^{-1} \quad \text{for all} \quad X \in \mathcal{X}(M)$$
 (4.10)

now follows immediately.

The converse part of the proof is easier. If covariant derivatives ∇^1 , ∇^2 are isomorphic, there exists a smooth section W in $U(F_1, F_2)$ which defines a unitary mapping $\bar{\mathcal{H}}_1 \to \bar{\mathcal{H}}_2$: $\sigma(u) \mapsto W(u)\sigma(u)$. This unitary mapping carries the generalized system of imprimitivity (\bar{V}_1, \bar{E}_1) over to (\bar{V}_2, \bar{E}_2) . Then it suffices to notice that (4.10) implies (4.8). \square

Corollary 4.1. Two generalized systems of imprimitivity specified by quadruples $(P_j, U(r), \Gamma_j, id), j = 1, 2$, are equivalent if and only if the connections Γ_1 , Γ_2 are isomorphic.

Notation. Let us consider a principal bundle P(M,G) with connection Γ , and fix a point $x_0 \in P, \pi(x_0) = u_0$. Then, for every piecewise smooth closed curve τ on M with base point u_0 there exists a unique element $g \in G$ such that x_0 is a starting point and x_0g the end point of the lift of curve τ . We denote this element by $c(\tau)$.

Theorem 4.3. Let (V_j, E_j) be generalized systems of imprimitivity specified by quadruples $(P_j, G_j, \Gamma_j, L_j)$, j = 1, 2, and let points $x_1 \in P_1$, $x_2 \in P_2$, $u_0 \in M$ satisfy $\pi_1(x_1) = \pi_2(x_2) = u_0$. The generalized systems of imprimitivity (V_1, E_1) , (V_2, E_2) are equivalent if and only if there exists a unitary mapping $W : \mathcal{H}^{L_1} \to \mathcal{H}^{L_2}$ such that for all piecewise smooth closed curves τ on M with the base point u_0 ,

$$L_2(c_2(\tau)) = WL_1(c_1(\tau))W^{-1}$$
(4.11)

is valid.

Proof. We start with two remarks.

- 1. Taking x' = xa instead of x, $\pi(x) = \pi(x') = u_0$, then $c'(\tau) = a^{-1}c(\tau)a$.
- 2. According to Sect. 4.3, to a principal bundle P(M,G) with connection Γ we can associate a principal bundle P(M, U(r)) with connection $\tilde{\Gamma}$; there also exists a homomorphism $f_0: P \to \tilde{P}$ preserving the connection and fulfilling $f_0(xg) = f_0(x)L_g$. If $\tilde{x} = f_0(x)$, $\tilde{\pi}(\tilde{x}) = \pi(x) = u_0$, then $\tilde{c}(\tau) = L(c(\tau))$. The systems of imprimitivity for these two principal bundles are equivalent.

It follows from these two remarks that the choice of a point $x \in P$, $\pi(x) = u_0$, plays no role and, moreover, we can restrict our considerations to the case $G_1 = G_2 = \mathrm{U}(r)$, $L_1 = L_2 = \mathrm{id}$. We are thus considering principal bundles $P_j(M, \mathrm{U}(r))$ with connections Γ_j , j = 1, 2. It suffices to show that condition (4.11) is valid if and only if connections Γ_1 , Γ_2 are isomorphic.

If $f: P_1 \to P_2$ is an isomorphism preserving the connection and if we choose $x_2 = f(x)$, then $c_1(\tau) = c_2(\tau)$ holds for every closed curve τ with base point u_0 .

Conversely, let $c_2(\tau) = bc_1(\tau)b^{-1}$ for some $b \in U(r)$. After having substituted x_1b for x_1 we can suppose $c_1(\tau) = c_2(\tau)$. We define a partial mapping f on some subset of P_1 into P_2 : if u(t), $0 \le t \le 1$, is a piecewise smooth curve in M with starting point u_0 and if $x_1(t)$, $x_2(t)$ are the lifts of this curve in P_1 , P_2 with starting points x_1 , x_2 , respectively, we put $f(x_1(1)) = x_2(1)$. Let us investigate the case when two curves u(t), v(t), $0 \le t \le 1$, $u(0) = v(0) = u_0$, have the same end point u(1) = v(1). Let $x_j(t)$, $y_j(t)$, j = 1, 2 be the lifts of curves u(t), v(t) with starting points x_1 , x_2 , respectively. There exists a unique $a \in G$ such that $y_1(1) = x_1(1)a$. Let $\tau(t)$, $0 \le t \le 1$, be a closed curve with base point u_0 , coinciding with u(t) for $0 \le t \le 1$ and with v(2-t) for $1 \le t \le 2$. Then we find $c_1(\tau) = a^{-1} = c_2(\tau)$ and hence $y_2(1) = x_2(1)a$. Thus function f is well-defined. The domain of f consists of all points in P_1 which can be connected with x_1 by a horizontal curve. But this domain can be extended to the whole P_1 by the relation f(xa) = f(x)a. In this way f becomes an isomorphism of principal bundles $P_1 \to P_2$ and, by construction, preserves the connection. \square

4.6 Irreducibility of generalized systems of imprimitivity

Let us consider principal bundle P(M,G) with connection Γ . We associate a subgroup Φ_x of G to each point $x \in P$; Φ_x consists of all $g \in G$ such that the points x, xg lie on a common horizontal curve in P. The group Φ_x is called the *holonomy group* of the connection Γ with the base point x. It has following properties:

- 1. If $x' \in P$ can be connected with x by a horizontal curve, then $\Phi_{x'} = \Phi_x$;
- 2. $\Phi_{xb} = b^{-1}\Phi_x b$.

So all holonomy groups are conjugate subgroups in G and we need not specify the base point. The restricted holonomy group Φ^0 is the subgroup of Φ corresponding to horizontal lifts of those closed curves which are homotopic to 0. The groups Φ , Φ^0 – being subgroups of G – are topological groups.

Theorem 4.4. ([51], Chap. II.3, [48, 43]). The restricted holonomy group Φ^0 is a connected Lie group, and it coincides with the arcwise connected component of unity in Φ . Moreover, the quotient Φ/Φ^0 is finite or countable.

The holonomy group Φ itself need not be a Lie group. But it can be equipped with a new topology which induces the original topology on Φ^0 and the quotient group Φ/Φ^0 is discrete. In this topology one can verify that the inclusion $i:\Phi\hookrightarrow G:g\mapsto g$ is a homomorphism of Lie groups.

Definition 4.3. We say that a structure group G of a principal bundle P(M, G) is reducible to a Lie group G', if there exists a principal bundle P'(M, G') and homomorphism $f: P' \to P$ such that $f(x'g') = f(x')f_0(g')$, $\pi(f(x')) = \pi'(x')$ with $f_0: G' \to G$ being an injective homomorphism of Lie groups. Moreover, if bundles P, P' are endowed with connections Γ , Γ' , respectively, and the homomorphism f preserves connection, we say that connection Γ is reducible to connection Γ' .

Theorem 4.5. Let a generalized system of imprimitivity (V, E) be specified by a quadruple (P, G, Γ, L) . If connection Γ is reducible to a connection Γ' on a principal bundle P'(M, G') and $L^0 = L \circ f_0$ denotes the representation of the Lie group G', then the generalized system of imprimitivity specified by the quadruple (P', G', Γ', L^0) is equivalent to (V, E).

Proof. The assertion can be proved by a method completely analogous to that used at the end of Sect. 4.3. The desired unitary mapping can be constructed in terms of the injective homomorphism $f: P' \to P$; $\psi'(x') = \psi(f(x'))$.

Theorem 4.6 ([51], Chap. II.6, [48]). Let P(M, G) be a principal bundle, Φ the holonomy group of connection Γ in P. Then the structure group G is reducible to a connection in the reduced principal bundle $P'(M, \Phi)$, the holonomy group of which is identical with Φ .

Theorem 4.7. Let (V, E) be a generalized system of imprimitivity specified by a quadruple (P, G, Γ, L) , Φ be the holonomy group of connection Γ , L^0 be the restriction of representation L to the subgroup Φ . Then the commuting algebras C(V, E), $C(L^0)$ are isomorphic.

Proof. In view of Theorems 4.5 and 4.6 it suffices to consider the case $G = \Phi$. Let \tilde{A} be a bounded operator $\mathcal{H} \to \mathcal{H}$. If \tilde{A} commutes with all E(S), $S \in \mathcal{B}(M)$, then, according to Theorem 3.4, it is of the form $\psi(x) \mapsto A(x)\psi(x)$, where $x \mapsto A(x)$ is some measurable mapping from P into the space of operators in \mathbb{C}^r such that $A(xg) = L_g^{-1}A(x)L_g$ is true on almost all fibres. Moreover, if \tilde{A} commutes with all $V^{\phi}(t)$, then $A(x) = V^{\phi}(t)A(x)V^{\phi}(t)^{-1} = A(\tilde{\phi}_t \cdot x)$ holds almost everywhere. We shall show that function $x \mapsto A(x)$ can be considered smooth after a redefinition on a set of measure zero. To show this local property, we can consider

 $M = \mathbb{R}^n$, $P(M,G) = \mathbb{R}^n \times G$. Then for vectors $v \in S^{n-1} \subset \mathbb{R}^n$ the mapping $y \mapsto A(y)$ remains constant (almost everywhere) on horizontal lifts of straight lines in \mathbb{R}^n with directions v and passing through u for almost all u. More precisely, if $\phi_t^v(u) = u + tv$, then for all $(t,v) \in \mathbb{R} \times S^{n-1}$ and for almost all $u \in \mathbb{R}^n$ (depending on t and v) the equality

$$A(u;g) = A(\tilde{\phi}_t^v(u;g)) \tag{4.12}$$

is valid for each $g \in G$. Using the Fubini Theorem we find that for almost all $u \in \mathbb{R}^n$, (4.12) holds for all $g \in G$ and almost all $(t,v) \in \mathbb{R} \times S^{n-1}$. Let us fix u with this property; we can assume u = 0. Then we construct an auxiliary section σ in P. For $w \in \mathbb{R}^n$ we lift the curve u(t) = tw, $0 \le t \le 1$, in the given connection, choosing (0;e) for the starting point, and we put $\sigma(w)$ to be equal to the final point of the lifted curve. Let A = A(0;e). Then for almost all $w \in \mathbb{R}^n$ and all $g \in G$, $A(\sigma(w)g) = L_g^{-1}AL_g$ holds. This proves our assertion that $x \mapsto A(x)$ can be considered smooth. Namely, the mapping $(w,g) \mapsto \sigma(w)g$ is an automorphism of the principal bundle $\mathbb{R}^n \times G$.

So let us suppose that function $x \mapsto A(x)$ is smooth and again use the equation $A(x) = A(\tilde{\phi}_t \cdot x)$, now being valid for all t, x. We find that the linear mapping $dA_x : T_x P \to \mathbb{C}^{r,r}$, if restricted to the horizontal subspace, is zero. Hence A(x) is constant on horizontal curves in P. Since the holonomy group coincides with the structure group, arbitrary two points in P can be connected by a horizontal curve. Thus A(x) = A for all $x \in P$ and we have $A = A(x) = A(xg) = L_g^{-1}AL_g$. In this way we have associated a unique operator $A \in \mathcal{C}(L^0)$ to each $\tilde{A} \in \mathcal{C}(V, E)$.

Conversely, one can relate a unique $\tilde{A} \in \mathcal{C}(V, E)$ to each $A \in \mathcal{C}(L^0)$ by means of the relation $\tilde{A}\psi(x) = A \cdot \psi(x)$. The one-to-one correspondence $\tilde{A} \leftrightarrow A$ is the desired isomorphism. \square Corollary 4.2. (V, E) is irreducible if and only if L^0 is irreducible.

4.7 Quantum Borel kinematics with vanishing external field

Let \mathcal{G} denote the Lie algebra of the group G, $\mathrm{u}(r)$ the Lie algebra of the group $\mathrm{U}(r)$ (consisting of skew–Hermitian $(r \times r)$ –matrices), and Ω the curvature 2–form of connection Γ . Ω is a 2–form on P taking values in \mathcal{G} such that $R_a^*\Omega = \mathrm{ad}(a^{-1})\Omega$ holds for all $a \in G$, where $R_a : P \to P : x \mapsto xa$. By composition with the representation L' of \mathcal{G} we obtain a 2–form $L' \circ \Omega$ taking values in $\mathrm{u}(r)$. Under the homomorphism of principal bundles $f : P \to \tilde{P}$, $\tilde{P} = \tilde{P}(M, \mathrm{U}(r))$ (see Sect. 4.3) $L' \circ \Omega$ is mapped into the curvature form $\tilde{\Omega}$ of the connection $\tilde{\Gamma}$.

For a vector bundle F associated to P let End F denote a vector bundle over M, with fibres (over $u \in M$) consisting of linear endomorphisms of fibres (from F_u into F_u). 2-forms w on P taking values in u(r) and satisfying $R_a^*w = \operatorname{ad}(a^{-1})w$, $a \in \operatorname{U}(r)$, are in one-to-one correspondence with 2-forms K on M taking skew-adjoint values in the space of sections $\operatorname{Sec}(\operatorname{End} F)$; the correspondence is expressed by the relation

$$K_u(X,Y) = x \circ w_x(X^*,Y^*) \circ x^{-1},$$

where $\tilde{\pi}(x) = u$ and X^* , Y^* are horizontal lifts of X, Y with respect to the connection $\tilde{\Gamma}$. In this way to $\tilde{\Omega}$ a 2-form R is related and

$$R(X,Y) = [\nabla_X, \nabla_Y] - \nabla_{[X,Y]} \tag{4.13}$$

holds ([51], Chap. III.5). R is the *curvature* of the covariant derivative ∇ . A simple calculation using (4.7) then leads to

$$[P(X),P(Y)]=-i\hbar P([X,Y])+(-i\hbar)^2R(X,Y),$$

since $P \approx \tilde{P}$.

If R = 0 (this is true if and only if $L' \circ \Omega = 0$), we say that the *external field vanishes* on the manifold. We see that the field vanishes if and only if

$$[P(X), P(Y)] = -i\hbar P([X, Y]),$$

see Sect. 3.2, eq. (3.3).

We recall the mapping $c: \tau \mapsto c(\tau)$ introduced in Sect. 4.5. If the field vanishes on the manifold, then the value $L(c(\tau)) \in U(r)$ depends only on the homotopy class of the curve τ . In this manner we obtain a representation L^c of the fundamental group $\pi_1(M)$ of manifold M. Let $(M^c, \pi^c, M; \pi_1(M))$ be the universal covering of M. Since dim $M^c = \dim M$, there exists exactly one flat connection Γ^c on the principal bundle $(M^c, \pi^c, M; \pi_1(M))$. Moreover, according to Theorem 4.3 (Eq. (4.11)) the generalized system of imprimitivity (V^c, E^c) specified by the quadruple $(M^c, \pi^c, \Gamma^c, L^c)$ is equivalent to (V, E). Conversely, since the connection Γ^c is flat, the field on M will vanish for every generalized system of imprimitivity (V^c, E^c) , no matter which representation L^c of $\pi_1(M)$ is chosen. So we arrive at a canonical form for generalized systems of imprimitivity (or quantum Borel kinematics) with vanishing field (with flat connection). This form was already studied in detail (see [2, 3] and references therein). The results of Sections 4.5 and 4.6 imply in this case:

Theorem 4.8. Two generalized systems of imprimitivity (or two quantum Borel kinematics) with vanishing field (V_j^c, E_j^c) , j = 1, 2, are equivalent if and only if the representations L_1^c , L_2^c are equivalent. The commuting algebras $\mathcal{C}(V^c, E^c)$, $\mathcal{C}(L^c)$ are isomorphic.

5 Quantum Borel kinematics: classification

5.1 Classification of generalized systems of imprimitivity via cocycles

The generalized systems of imprimitivity described by (4.1), (4.2) are not of the most general form. Starting from a characterization of Mackey's systems of imprimitivity in terms of cocycles ([54], Theorem. 9.11), the following theorem was proved in [26]:

Theorem 5.1. Any r-homogeneous generalized system of imprimitivity on M is unitarily equivalent to a canonical one (V, E), with $\mathcal{H} = L^2(M, \mathbb{C}^r, \mu)$ for some smooth measure μ on M,

$$(E(S)\psi)(u) = \chi_S(u)\psi(u)$$

for all $\psi \in \mathcal{H}$ and $S \in \mathcal{B}(M)$, and

$$[V^{\phi^X}(t)\psi](u) = \xi^X(t, \phi_{-t}^X(u)) \sqrt{\frac{\mathrm{d}\mu}{\mathrm{d}(\mu \circ \phi_t^X)} (\phi_{-t}^X(u))} \psi(\phi_{-t}^X(u))$$
 (5.1)

for all $\psi \in \mathcal{H}$ and all $X \in \mathcal{X}_c(M)$. Equivalence classes of r-homogeneous generalized systems of imprimitivity are in one-to-one correspondence with equivalence classes of cocycles $[\xi^X]$.

Here ξ^X is a *cocycle* of \mathbb{R} relative to the Lebesgue measure class on M with values in U(r), i.e. a Borel measurable map $\xi^X : \mathbb{R} \times M \mapsto U(r)$ with

$$\xi^X(0,u) = 1,$$

$$\xi^X(s+t,u) = \xi^X(s,\phi_t^X(u))\xi^X(t,u)$$

for almost all $u \in M$ and almost all $s, t \in \mathbb{R}$. Two cocycles ξ_j^X , j = 1, 2, are called *equivalent* (cohomologous), if there is a Borel function $\zeta : \mathbb{R} \mapsto U(r)$, such that for all $X \in \mathcal{X}_c(M)$ and $t \in \mathbb{R}$, $u \in M$

$$\xi_2^X(t, u) = \zeta(\phi_t^X(u))\xi_1^X(t, u)\zeta(u)^{-1}.$$

Unfortunately, the classification given in Theorem 5.1 is not easy to handle, since the calculation of cocycles is rather tedious. To be more specific, one has to impose further conditions on the operators under consideration.

5.2 Differentiable quantum Borel kinematics

To gain further insight into the structure of the shift operators (5.1), one can perform a formal calculation. In particular, assume for the moment that the cocycles of the representation are *smooth* maps from $\mathbb{R} \times M$ into U(r). Then, by formal differentiation of (5.1) with respect to t at t = 0, an expression for the generalized momentum operator P(X) is obtained,

$$P(X)\psi = -i\hbar[X\psi + \frac{1}{2}\operatorname{div}_{\mu}(X)\cdot\psi] - i\hbar\rho(X)\cdot\psi, \tag{5.2}$$

where $\psi \in C_0^{\infty}(M, \mathbb{C}^r)$ and

$$\rho(X)(u) := \frac{\mathrm{d}}{\mathrm{d}t} \xi^X(t, \phi_{-t}^X(u))|_{t=0}.$$

The first two terms on the right hand side of (5.2) are linear in the vector field $X \in \mathcal{X}(M)$. Though the set of complete vector fields $\mathcal{X}_c(M)$ is not a linear space — the sum of two complete vector fields may not be complete — it contains the "large" linear subset $\mathcal{X}_0(M)$ of vector fields with compact support for which one can demand linearity (or demand 'partial' linearity at least for all *complete* linear combinations of complete vector fields), cf. Theorem 3.1). Thus as a first additional assumption on P(X) we require (3.2), i.e. $\rho(X)$ to be linear in that case.

Using the formal expression (5.2), the commutator of P(X) and Q(f) is obviously obtained again in the form (3.5). Finally, for the commutator [P(X), P(Y)] we obtain

$$[P(X), P(Y)] = -i\hbar P([X, Y]) - \hbar^2 R(X, Y), \tag{5.3}$$

where

$$R(X,Y) = [\rho(X), \rho(Y)] + X\rho(Y) - Y\rho(X) - \rho([X,Y]). \tag{5.4}$$

If ρ were a localized connection 1–form, (5.4) would represent the local definition of a curvature 2–form R of a \mathbb{C}^r –bundle over M. The Jacobi identity for generalized momenta would then give us precisely the Bianchi identity DR = 0, where D is the covariant differential defined by the connection.

In order to arrive at this point we had to assume differentiability of the shift operators (5.1) and of the functions in the domain of momentum operators. Now there are different ways of defining differentiable structures and thus differentiability on the set $M \times \mathbb{C}^r$; for a discussion of this point we refer to [3, 14]. On the other hand, we have already interpreted R as a curvature 2-form that is in general related to a connection on a \mathbb{C}^r -bundle over M. This line of reasoning leads us to the following definition [14, 26]:

Definition 5.1. Let M be a differentiable manifold, (V, E) an r-homogeneous generalized system of imprimitivity on M for r = 1, 2, ..., and R a differential 2-form on M with values in Hermitian operators on \mathbb{C}^r . Let Q(f) and P(X) denote the corresponding generalized position and momentum operators, respectively. Then:

- 1) The quadruple (\mathcal{H}, Q, P, R) is called an (R-compatible) quantum Borel r-kinematics (QBK^r) , if P is (partially) linear, satisfies (5.3), (5.4), and the common invariant domain \mathcal{D} of Q(f)'s and P(X)'s for $f \in C^{\infty}(M)$ and $X \in \mathcal{X}_c(M)$ is dense in \mathcal{H} . It is local, if P is local, and elementary, if r = 1.
- 2) A quantum Borel kinematics is called *differentiable*, if it is equivalent to (\mathcal{H}, Q, P, R) of standard form, constructed from the following ingredients:
 - 1. Lebesgue measure μ on M;
 - 2. Hermitian vector bundle $F(M; \mathbb{C}^r)$ over M with fibres diffeomorphic to \mathbb{C}^r equipped with Hermitian inner product $\langle ., . \rangle$;
 - 3. A 2–form R with skew–adjoint values in the endomorphism bundle $\operatorname{End} F = F \otimes F^*$;
 - 4. The Hilbert space \mathcal{H} is realized as the Hilbert space $L^2(F, <.,.>, \mu)$ of sections of F, i.e. (measurable) mappings $\sigma: M \to F$ such that $\bar{\pi} \circ \sigma = \mathrm{id}_M$ and with finite norm induced by the inner product

$$(\sigma, \tau) = \int_{M} \langle \sigma(u), \tau(u) \rangle d\mu(u);$$

- 5. The common invariant domain \mathcal{D} for Q, P contains the set $\operatorname{Sec}_0 F$ of smooth sections of F with compact support and $P(X)\operatorname{Sec}_0 F \subset \operatorname{Sec}_0 F$;
- 6. The position operators Q(f) have the usual form of the Schrödinger representation

$$Q(f)\sigma = f \cdot \sigma, \qquad \forall f \in C^{\infty}(M, \mathbb{R}), \quad \sigma \in \operatorname{Sec}_0 F.$$

5.3 Canonical representation of differentiable QBK^r

For local differentiable quantum Borel kinematics the formal calculations can be made precise. According to Sect. 5.2 it only remains to derive the representation of generalized momenta P(X) in a standard form. This is the content of

Theorem 5.2 ([3, 14]). Let (\mathcal{H}, Q, P, R) be a local differentiable quantum Borel kinematics on M in a standard form. Then there is

1. a Hermitian connection ∇ with curvature R on F, i.e. a connection compatible with the inner product

$$X < \sigma, \tau > = < \nabla_X \sigma, \tau > + < \sigma, \nabla_X \tau > ,$$

2. a covariantly constant self-adjoint section Φ of End $F = F \otimes F^*$, the bundle of endomorphisms of F,

such that for all $X \in \mathcal{X}_c(M)$ and all $\sigma \in \operatorname{Sec}_0 F$

$$P(X)\sigma = -i\hbar\nabla_X\sigma + \left(-\frac{i\hbar}{2}I + \Phi\right)Q(\operatorname{div}_{\nu}X)\sigma, \quad \sigma \in \mathcal{D}.$$
 (5.5)

Moreover, R is a curvature 2-form on F satisfying the Bianchi identity

$$DR = 0,$$

where D denotes the covariant differential defined by the connection ∇ .

The canonical form of generalized momenta (5.2) shows that by imposing R-compatibility the quantum system on M is influenced by an external classical field on M. In general, this field is defined by a curvature 2-form on F, and thus a curvature $\tilde{\Omega}$ on the associated $\mathrm{U}(r)$ -principal bundle $\tilde{P}(M,\mathrm{U}(r))$. We could think of this $\tilde{\Omega}$ as a classical gauge (Yang-Mills) field. The simplest example was provided in Sect. 4.1 for $M=\mathbb{R}^3$, r=1, where the 2-form $e\beta$ was interpreted as a coupling constant times a magnetostatic field on \mathbb{R}^3 , with the Bianchi identity corresponding to the Maxwell equation $\mathrm{div}\vec{B}=0$. Up to a coupling constant the connection ∇ generalizes the notion of a vector potential. For $M=\mathbb{R}^3$ the global connection form α corresponds to the vector potential \vec{A} and the 2-form $\beta=\mathrm{d}\alpha$ to the magnetic field $\vec{B}=\mathrm{rot}\vec{A}$ of Maxwell's theory.

5.4 Classification of differentiable QBK^r 's

The canonical form given in Theorem 5.2 indicates that a classification of differentiable quantum Borel kinematics amounts to a classification of Hermitian \mathbb{C}^r -bundles with connection over M and covariantly constant self-adjoint sections of the corresponding endomorphism bundle.¹² This is the content of

Theorem 5.3. Two local differentiable quantum Borel kinematics $(\mathcal{H}_j, Q_j, P_j, R_j)$, j = 1, 2, in canonical form of Theorem 5.2 are equivalent, if and only if there is a strong, unitary, and connection (and thus curvature) preserving bundle isomorphism $I: F_1 \to F_2$ mapping Φ_j into each other, i.e.

$$\nabla_2 = I \circ \nabla_1 \circ I^{-1}, \quad R_2 = I \circ R_1 \circ I^{-1}, \quad \Phi_2 = I \circ \Phi_1 \circ I^{-1}.$$

Unfortunately, there are no general existence and classification theorems of Hermitian \mathbb{C}^r -bundles with connection. Looking back to Sect. 4, there a rather big class of local differentiable QBK^r's is constructively defined, however with $\Phi = 0$. Hence even in these cases, the additional classification of covariantly constant self-adjoint sections has to be found as well. This last problem was solved only in certain special cases — elementary quantum Borel kinematics and type 0 or type U(1) QBK^r's — described in the following sections.

5.5 Classification of elementary differentiable quantum Borel kinematics

The problem of existence and classification of *elementary*, i.e. r=1 local differentiable quantum Borel kinematics in terms of global geometrical properties (cohomology groups) of the underlying manifold M was completely solved [3, 14]. It is based on a theorem [20, 19] concerning existence and classification of complex line bundles with hermitian connection.

Theorem 5.4 Let M be a connected differentiable manifold and $B \in \Lambda^2(M)$ be a closed 2-form on M with dB = 0. Then there exists a complex line bundle F with hermitian connection ∇ of curvature $R = \frac{i}{\hbar}B$ if and only if R satisfies the integrality condition

$$\frac{1}{2\pi i} \int_{\sigma} R = \frac{1}{2\pi \hbar} \int_{\sigma} B \in \mathbb{Z}$$

for all closed 2–surfaces σ in M. In terms of cohomology theory, the de Rham class of $R/(2\pi i) = B/(2\pi \hbar)$ has to be integral,

$$[\frac{1}{2\pi\hbar}B]\in H^2(M,\mathbb{Z}).$$

 $^{^{12}}$ The zero section of End F always exists and is covariantly constant and self–adjoint.

Hence non–isomorphic equivalence classes of principal bundles over a manifold M with the structure group U(1) are labeled by elements of the second cohomology group $H^2(M, \mathbb{Z})$. The Lie algebra of U(1) coincides with the imaginary axis $i \mathbb{R}$. Since U(1) is Abelian, the vector bundle End $F = M \times \mathbb{C}$ is trivial. So the curvature R is a purely imaginary 2–form on $M, R = \Omega$, where Ω is the curvature 2–form of connection Γ . If we put

$$\beta = -\frac{i\hbar}{e}R,$$

 β can be interpreted as the 2-form of external magnetic field on M. For an arbitrary 2-cycle σ of the singular homology on M, $\partial \sigma = 0$, we have

$$\exp(\int_{\sigma} \Omega) = 1.$$

This leads to the *Dirac quantization condition* on the magnetic field

$$\int_{\sigma} \beta = g, \quad \text{with} \quad \frac{eg}{2\pi} = n\hbar,$$

where $n \in \mathbb{Z}$. We may interpret this result that the 2-cycle σ — besides the usual magnetic field satisfying $\int_{\sigma} \beta = 0$ — encloses a *Dirac magnetic monopole* with quantized magnetic charge g.

Furthermore, the various inequivalent choices of $(F, <.,.>, \nabla)$ for fixed curvature R are parametrized by

$$H^1(M, U(1)) = \pi_1(M)^*,$$

where $\pi_1(M)^*$ denotes the group of characters of the fundamental group of M.

We should emphasize that $H^1(M, \mathrm{U}(1)) = \pi_1(\mathrm{M})^*$ classifies pairs of hermitian line bundles (F, < ., . >) and compatible connections ∇ . This implies that the curvature 2-forms of two equivalent complex line bundles with hermitian connection are identical. The classification of complex line bundles F themselves — disregarding their connection — is given by elements of the Čech cohomology $H^1(M, \mathrm{U}(1)) = H^2(\mathrm{M}, \mathbb{Z})$: two complex line bundles are equivalent if and only if their Chern classes in $H^2(M, \mathbb{Z})$ coincide, i.e. the curvature 2-forms admissible on these two bundles are in the same integral de Rham cohomology class of $H^2(M, \mathbb{Z})$.

Theorem 5.4 is the basis of a classification theorem that goes back to [3] for the flat (B=0) case and was extended to the case of external fields B by [14]:

Theorem 5.5. The equivalence classes of elementary local differentiable quantum Borel kinematics are in one–to–one correspondence to elements of the set

$$H^2(M, \mathbb{Z}) \times H^1(M, \mathrm{U}(1)) \times \mathbb{R}.$$

For the proof it remains to classify the inequivalent choices of covariantly constant self-adjoint sections Φ of End F. For a line bundle the endomorphism bundle is actually trivial: As the transition functions $\varphi_{jk}: U_j \cap U_k \to \mathrm{U}(1)$ of F commute with complex numbers, the induced transition functions of End $F = F \otimes F^*$ become trivial, $z \mapsto \varphi_{jk}z\varphi_{jk}^* = z$, hence End $F = M \times \mathbb{C}$. Thus the sections of this bundle correspond to complex functions on M. Furthermore, the induced connection on M is the trivial connection on $M \times \mathbb{C}$ given by the Lie derivative. Thus covariantly constant self-adjoint sections Φ of End F are real multiples of the identity,

$$\Phi = \hbar c \cdot \mathrm{id}_{\mathbf{Sec}}, \quad \mathbf{c} \in \mathbb{R}.$$

Obviously, c is not changed under strong bundle isomorphisms I, so each value of $c \in \mathbb{R}$ for a given Hermitian line bundle determines an inequivalent local differentiable quantum Borel kinematics. \Box

Finally let us note that elementary quantum Borel kinematics with c=0 are, in terms of constructions of Sect. 4, described by generalized systems of imprimitivity (V, E) specified by the quadruple $(P, U(1), \Gamma, id)$.

5.6 Classification of quantum Borel kinematics of type 0

The whole variety of quantizations could be read off the formula (5.5). In order to get a more transparent result we define a QBK^r of type 0 [25] by

$$\Phi = \hbar c.id_{\mathbf{Sec.}F}, \qquad c \in \mathbb{R}.$$

Then we obtain an identical formula for P(X) as in QBK¹ [4]:

$$P(X)\sigma = -i\hbar \nabla_X \sigma + \left(-\frac{i\hbar}{2} + \hbar c\right)(\operatorname{div}_{\mu} X) \cdot \sigma, \quad \sigma \in \mathcal{D}.$$
 (5.6)

As proved in [25], on every smooth manifold M there exists a differentiable QBK^r of type 0. Let us note that for r = 1, the type 0 QBK¹'s classify all possible Borel quantizations [4]; this is not the case, however, for r > 1.

Finally, a complete classification of QBK^r's of type 0 was possible in the case of *flat* connection (R = 0) [4, 33, 25]. It turns out that it is essentially the question of the topology of M. The corresponding investigations can be summarized in

Theorem 5.6 The set of classes of unitarily equivalent QBK^r's of type 0 with flat connection on M can be bijectively mapped onto the set of pairs (\mathcal{L}, c) , where $c \in \mathbb{R}$ and \mathcal{L} denotes the isomorphism class of flat \mathbb{C}^r -bundles over M.

Since there is a one-to-one correspondence between the isomorphism classes of flat \mathbb{C}^r -bundles over M and flat U(r)-principal bundles over M, we can use Milnor's

Lemma [24]. A U(r)-principal bundle over M admits a flat connection if and only if it is induced from the universal covering bundle of M by a homomorphism of the fundamental group $\pi_1(M)$ into U(r).

Thus, disregarding the real constant c, the set of inequivalent quantizations of type 0 with \mathbb{C}^r -valued wave functions is isomorphic to the set

$$\operatorname{Hom}(\pi_1(M), \operatorname{U}(r))$$

of (the conjugacy classes of) r-dimensional unitary representations of the fundamental group of M.

In the case r=1, i.e. of quantizations with complex-valued wave functions, the topological part of the classification reduces to $\operatorname{Hom}(\pi_1(M), \operatorname{U}(1))$, i.e. to the set of one-dimensional unitary representations of $\pi_1(M)$ [3, 4, 12, 25].¹³ Since the commutator subgroup $\Gamma(\pi_1(M))$ (generated by elements $aba^{-1}b^{-1}$) belongs to the kernels of all such one-dimensional representations and since the singular homology group $H_1(M, \mathbb{Z})$ is isomorphic to $\pi_1(M)/\Gamma(\pi_1(M))$ (the Hurewicz isomorphism), inequivalent QBK¹'s are labeled by elements of the character group of $H_1(M, \mathbb{Z})$.

¹³This result was obtained independently also in the Feynman path integral approach [21] and in geometric quantization [20].

Finally, let us describe the general structure of the Abelian group $H_1(M, \mathbb{Z})$ for compact M. It has a decomposition $H_1(M, \mathbb{Z}) = F \oplus T$, where the free Abelian group F is $F = \mathbb{Z} \oplus \cdots \oplus \mathbb{Z}$ (b_1 terms), with b_1 being the first Betti number of M, and the torsion Abelian group is $T = \mathbb{Z}_{\tau_1} \oplus \cdots \oplus \mathbb{Z}_{\tau_k}$ with \mathbb{Z}_{τ_i} being cyclic groups of orders τ_i (torsion coefficients) such that $\tau_{i+1}/\tau_i = \text{positive integer}$. Thus the characters of $H_1(M, \mathbb{Z})$ can be parametrized by $(b_1 + k)$ -tuples

 $[e^{2\pi i\theta_1}, \cdots, e^{2\pi i\theta_{b_1}}; e^{2\pi im_1/\tau_1}, \cdots, e^{2\pi im_k/\tau_k}]$

with the numbers $\theta_l \in [0,1)$, $l = 1, \ldots, b_1$, and $m_i = 0, 1, \ldots, \tau_i - 1$, $i = 1, \ldots, k$, classifying inequivalent quantum Borel 1-kinematics on M.

5.7 Elementary quantum Borel kinematics with vanishing external field

It is remarkable that elementary quantum Borel kinematics with vanishing external magnetic field find application in quantum mechanics (Aharonov-Bohm effect [1]). According to Sect. 4.3 they can be labeled by one–dimensional unitary representations of the fundamental group $\pi_1(M)$, and consequently, as already explained in Sect. 4.9, inequivalent systems of imprimitivity with vanishing magnetic field are labeled by elements of the character group of $H_1(M, \mathbb{Z})$.

Quantum Borel kinematics in the case of trivial fibration $P = M \times U(1)$ was studied in detail in [11]. In this case a localized connection 1–form $i(e/\hbar)\alpha$ can be defined globally on the whole manifold M; the closed 1–form α represents the vector potential of the vanishing magnetic field. A covariant derivative on $M \times \mathbb{C}^1$ has the form

$$\nabla_X = X - i \frac{e}{\hbar} \alpha(X), \quad X \in \mathcal{X}(M).$$

Two such covariant derivatives ∇^1 , ∇^2 are isomorphic if and only if there exists a function $f: M \to T^1$ such that

$$\alpha_2 = \alpha_1 - i \frac{\hbar}{e} \frac{df}{f}.$$

Following the terminology of [11, 27], we say that the 1-forms $(e/\hbar)\alpha_j$, j=1,2 are logarithmically cohomologous; $\lambda=-i(df)/f$ is said to be logarithmically exact. A 1-form λ is logarithmically exact if and only if

$$\exp(i\int_{\gamma}\lambda)=1$$
, i.e. $\int_{\gamma}\lambda=2\pi n,\ n\in\mathbb{Z}$,

holds for all 1–cycles γ of the singular homology. If γ is a periodic cycle, i.e. $p\gamma = \partial \sigma$ for some $p \in \mathbb{Z}$ and 2-cycle σ , then

$$0 = \int_{\sigma} d\lambda = \int_{p\gamma} \lambda = p \int_{\gamma} \lambda,$$

because a logarithmically exact form is closed.

So it suffices to check independent, non-periodic 1-cycles. Their number is $b_1(M)$, the first Betti number of manifold M. Now, quantum Borel kinematics on M in the case of trivial fibration and of vanishing magnetic field is determined by a closed 1-form α . Its cohomology class is in turn — according to the de Rham Theorem — determined by $b_1(M)$ periods

$$\Phi_j = \int_{\gamma_j} \alpha.$$

For each $j = 1, ..., b_1(M)$, Φ_j represents an external magnetic flux outside the manifold M and passing through the j-th independent cycle γ_j . Two potentials α_1 , α_2 determine the same quantum Borel kinematics if

$$\frac{e}{\hbar}(\Phi_j^{(1)} - \Phi_j^{(2)}) \in 2\pi \mathbb{Z}$$
 for all j .

We conclude that in the considered special case the family of all inequivalent quantum Borel kinematics can be parametrized by elements

$$z = (z_1, \dots, z_{b_1(M)}) \in \mathrm{U}(1) \times \dots \times \mathrm{U}(1), \quad z_j = \exp(\mathrm{i}\mathrm{e}\Phi_j/\hbar).$$

5.8 Examples

In the examples we concentrate on elementary quantum Borel kinematics.

Example 5.1. $M = \mathbb{R}^3$. Since $H^2(\mathbb{R}^3, \mathbb{Z}) = 0$, only the trivial principal bundle exists over M. Every smooth function $f : \mathbb{R}^3 \to \mathrm{U}(1)$ can be written in the form $f = \exp(ie\lambda/\hbar)$ with λ being a smooth real function. Two 1-forms are logarithmically cohomologous if and only if they are cohomologous. Thus (an equivalence class of) quantum kinematics is obtained by taking (a cohomology class of) a closed 2-form β of magnetic field. There exists exactly one equivalence class of quantum Borel kinematics (i.e. with vanishing magnetic field) — the standard one with $\alpha = 0$.

Example 5.2. $M = \mathbb{R}^3 \backslash \mathbb{R}$ is the 3-dimensional real space \mathbb{R}^3 with the x_3 -axis excluded (Aharonov-Bohm configuration). Then $H^2(M, \mathbb{Z}) = 0$, and there exists only the trivial principal bundle over M. There exists exactly one (up to homology) independent cycle in M and it is non-periodic, i.e. $H_1(M, \mathbb{Z}) = \mathbb{Z}, H^1(M, \mathbb{R}) = \mathbb{R}$. Inequivalent quantum Borel kinematics are labeled by $z \in \mathrm{U}(1), z = \exp(ie\Phi/\hbar)$; Φ denotes the magnetic flux supported by the excluded line \mathbb{R} . The corresponding vector potential 1-form α can be chosen in the form

$$\alpha = \frac{\Phi}{2\pi} \left(-\frac{x_2}{x_1^2 + x_2^2} dx_1 + \frac{x_1}{x_1^2 + x_2^2} dx_2 \right).$$

From the point of view of quantum mechanics a charged particle cannot distinguish between the flux $\Phi = 2\pi n\hbar/e$, $n \in \mathbb{Z}$, and the zero one. In fact, this is the effect discovered by Y.Aharonov and D.Bohm [1]. Let us note that the Aharonov–Bohm effect in the presence of two solenoids or n solenoids placed along a straight line was studied in [32].

Example 5.3. M be a compact orientable surface. It is known that, up to diffeomorphism, such M are classified by the Euler characteristic $\chi(M)=2-2p$, where $p=0,1,2,\ldots$ is the genus of M. For given p, we denote $M=K_p$; it is modeled by a 2-sphere with p handles ([46], Chap. 9.3). Since $H^2(K_p, \mathbb{Z}) = \mathbb{Z}$, there exist countably many principal bundles over K_p , labeled by $n \in \mathbb{Z}$. According to our physical interpretation, n can by related to the magnetic charge of a Dirac monopole enclosed by the closed surface, $g=(2\pi\hbar/e)n$. Further, $H_1(K_p, \mathbb{Z}) = \mathbb{Z}^{2p}$; thus inequivalent quantum Borel kinematics (with vanishing magnetic field) are labeled by elements of the dual group $U(1)^{2p}$. Each quantum Borel kinematics depends on 2p external magnetic fluxes, each handle carrying two of them. On a 2-torus, generalized quantum kinematics was studied in [13], quantum Borel kinematics in [29].

Example 5.4. $M = \mathbb{R}P^n = S^n/\{\pm\}$ is a real projective space, n > 2. We can identify:

$$S \in \mathcal{B}(\mathbb{R}P^n) \leftrightarrow S \in \mathcal{B}(S^n), \qquad S = -S; X \in \mathcal{X}(\mathbb{R}P^n) \leftrightarrow X \in \mathcal{B}(S^n), \qquad X_{-u} = -X_u;$$

$$H^2(\mathbb{R}P^n, \mathbb{Z}) = \mathbb{Z}_2, \qquad H_1(\mathbb{R}P^n), \mathbb{Z}) = \mathbb{Z}_2.$$

Table 1: Examples of elementary quantum Borel kinematics.

Quantum system	M	$\pi_1(M)$	$H_1(M,Z)$	$H^2(M,Z)$	Topological quantum numbers
Spinless particle in \mathbb{R}^3	R^3	$\{e\}$	0	0	_
Aharonov–Bohm configuration	$R^3 \backslash R$	Z	Z	0	$\vartheta \in [0,1)$
Dirac's monopole	$R^3 \backslash O = R_+ \times S^2$	$\{e\}$	0	Z	$n \in Z$
2 distinguishable particles in \mathbb{R}^3	$R^3 \times R_+ \times S^2$	$\{e\}$	0	Z	$n \in Z$
2 indistinguishable particles in \mathbb{R}^3	$R^3 \times R_+ \times RP^2$	S_2	Z_2	Z_2	$m \in Z_2$
Rigid body	$R^3 \times SO(3)$	Z_2	Z_2	Z_2	$m \in Z_2$
Symmetric top	S_2	$\{e\}$	0	Z	$n \in Z$
Rotator with fixed axis	S^1	Z	Z	0	$\vartheta \in [0,1)$
Particle on orientable surface of genus p	K_p	$\pi_1(K_p)$	Z^{2p}	Z	$n \in Z,$ $\vartheta_1 \dots \vartheta_{2p} \in [0,1)$

Hence there exist two inequivalent principal bundles over M and two inequivalent QBK's in mutually inequivalent fibrations. The QBK's can be explicitly described in the following way: the Hilbert spaces \mathcal{H}_+ , \mathcal{H}_- are chosen as subspaces in $L^2(S^n, d\mu)$ (with measure μ invariant under the transformation $u \to -u$), $\psi \in \mathcal{H}_{\pm}$ if and only if $\psi(-u) = \pm \psi(u)$; the two inequivalent systems of imprimitivity (c=0) are defined for both signs + and - by the operators

$$E(S) = \chi_S, P(X) = -i\hbar(X + \frac{1}{2}\operatorname{div}_{\mu}X), \quad S \in \mathcal{B}(\mathbb{R}P^n), X \in \mathcal{X}(\mathbb{R}P^n))$$

acting in \mathcal{H}_+ and \mathcal{H}_- , respectively. The real projective space $\mathbb{R}P^n$ appears in quantum mechanics e.g. as a (topologically non-trivial) part of the effective configuration space of two indistinguishable point-like particles localized in the (n+1)-dimensional Euclidean space \mathbb{R}^{n+1} . The two cases with signs + and - correspond in quantum mechanics to the cases of bosonic and fermionic statistics, respectively. More details can be found in [9]; see also [21, 22]. It should be stressed that the case n=2 presents unexpected features: in [22] it was found for a system of two particles in two dimensions that there is a continuous family of quantizations describing new statistics which interpolate between fermions and bosons. These anomalous or fractional statistics were later discovered independently by [16, 17] and by [34, 35], who actually coined the term 'anyons' for the corresponding particles.

ACKNOWLEDGEMENTS

J. T. and P. Š. are grateful to Prof. Dr. H.D. Doebner for the kind hospitality extended to them on various occasions at the Arnold Sommerfeld Institute in Clausthal. The authors thank A. Bóna for typing the manuscript. Partial support of the Grant Agency of Czech Republic (contract No. 202/96/0218) is acknowledged. The list of references is by no means complete and we apologize to the authors of papers which have not been included.

References

- [1] Y. Aharonov, D. Bohm: "Significance of electromagnetic potentials in the quantum theory", Phys. Rev. 115, 485–491 (1959).
- [2] B. Angermann: "Global geometry and Schrödinger quantization on manifolds", Latin American Summer School of Physics at UNAM, Mexico (1980).
- [3] B. Angermann, H.-D. Doebner: "Homotopy groups and quantization of localizable systems", Proc. Xth Coll. Group Theoretical Methods in Physics, Physica **114**A, 433–439 (1982).
- [4] B. Angermann, H.-D. Doebner, J. Tolar: "Quantum kinematics on smooth manifolds". In: *Non-linear Partial Differential Operators and Quantization Procedures*, Lecture Notes in Mathematics, Vol. 1037, Springer-Verlag, Berlin (1983), pp. 171–208.
- [5] F. Bayen, M. Flato, C. Fronsdal, A. Lichnérowicz, D. Sternheimer: "Deformation theory and quantization", Ann. Phys. (New York) **111**, 61–110, 111–152 (1978).
- [6] P.A.M. Dirac: "Quantized singularities in the electromagnetic field", Proc. Roy. Soc. London A 133, 60–72 (1931).
- [7] H.-D. Doebner, H.J. Elmers, W.F. Heidenreich: "On topological effects in quantum mechanics; The harmonic oscillator in the pointed plane", J. Math. Phys. **30**, 1053–1059 (1989).
- [8] H.-D. Doebner, G.A. Goldin: "On a general nonlinear Schrödinger equation admitting diffusion currents", Phys. Lett. A162, 397–401 (1992).
- [9] H.-D. Doebner, P. Štovíček, J. Tolar: "Quantizations of the system of two indistinguishable particles", Czech. J. Phys. B **32**, 1240–1248 (1982).
- [10] H.-D. Doebner, J. Tolar: "Quantum mechanics on homogeneous spaces", J. Math. Phys. 16, 975–984 (1975).
- [11] H.-D. Doebner, J. Tolar: "On global properties of quantum systems". In: Symmetries in Science (B. Gruber and R. S. Millman, eds.) Plenum Press, New York (1980), pp. 475–486.
- [12] H.-D. Doebner, J. Tolar: "Symmetry and topology of the configuration space and quantization". In: Symmetries in Science II (B. Gruber and R. Lenczewski, eds.) Plenum, New York (1986), pp. 115–126.
- [13] H.-D. Doebner, J. Tolar: "Quantum particle on a torus with an external magnetic field", in: *Quantization and Coherent States Methods* (S.T. Ali, I.M. Mladenov and A. Odzijewicz, eds.) World Scientific, Singapore (1993), pp. 3–10.
- [14] M. Drees: "Zur Kinematik lokalisierter quantenmechanischer Systeme unter Berücksichtigung innerer Freiheitsgrade und äußerer Felder", Ph.D. Thesis, Technical University, Clausthal (1992).
- [15] P. Goddard, D.I. Olive: "Magnetic monopoles in gauge field theories", Rep. Prog. Phys. 41, 1357–1437 (1978).

- [16] G.A. Goldin, R. Menikoff, D.H. Sharp: "Particle statistics from induced representations of a local current group", J. Math. Phys. 21, 650–664 (1980).
- [17] G.A. Goldin, R. Menikoff, D.H. Sharp: "Representations of a local current algebra in non-simply connected space and the Aharonov-Bohm effect", J. Math. Phys. 22, 1664– 1668 (1981).
- [18] G.A. Goldin, R. Menikoff, D.H. Sharp: "Induced representations of the group of diffeomorphisms of \mathbb{R}^3 ", J. Phys. A: Math. Gen. 16, 1827–1833 (1983).
- [19] W. Greub, H.-R. Petry: "Minimal coupling and complex line bundles", J. Math. Phys. 16, 1347–1351 (1975).
- [20] B. Kostant: "Quantization and unitary representations: Part I. Prequantization". Lecture Notes in Mathematics, Vol. 170, Springer-Verlag, Berlin (1970), pp. 87–208.
- [21] M.G.G. Laidlaw, C. Morette-DeWitt: "Feynman functional integrals for systems of indistinguishable particles", Phys. Rev. D 3, 1375–1378 (1970).
- [22] J.M. Leinaas, J. Myrheim: "On the theory of identical particles", Nuovo Cimento B 37, 1–23 (1977).
- [23] G.W. Mackey: "Unitary representations of group extensions I", Acta Math. 99, 265–311 (1958).
- [24] J. Milnor: "On the existence of a connection with curvature zero", Comment. Math. Helv. **32**, 215–223 (1958).
- [25] U.A. Müller, H.D. Doebner: "Borel quantum kinematics of rank k on smooth manifolds", J. Phys. A: Math. Gen. 26, 719–730 (1993).
- [26] P. Nattermann: "Dynamics in Borel quantization: Nonlinear Schrödinger equations vs. master equations", Ph.D. Thesis, Technical University, Clausthal (1997).
- [27] I.E. Segal: "Quantization of nonlinear systems", J. Math. Phys. 1, 468–488 (1960).
- [28] L.S. Schulman: "A path integral for spin", Phys. Rev. 176, 1558–1569 (1968).
- [29] C. Schulte: "Quantum mechanics on the torus, Klein bottle and projective sphere", in: Symmetries in Science IX (B. Gruber ed.) Plenum Press, New York (1997), pp. 313–323.
- [30] P. Štovíček: "Dirac monopole derived from representation theory", Suppl. Rend. Circ. Mat. Palermo, Ser. II, No. 3, 301–306 (1984).
- [31] P. Štovíček: "Systems of imprimitivity for the group of diffeomorphisms I, II", Ann. Global Anal. Geom. 5, 89–95 (1987), 6, 31–37 (1988).
- [32] P. Štovíček: "Scattering on a finite chain of vortices", Duke Math. J. 76, 303–332 (1994).
- [33] P. Šťovíček, J. Tolar: "Topology of the configuration manifold and quantum mechanics", Acta Polytechnica (Prague) Ser. IV, No. 1 (1984), 37–75 (in Czech).
- [34] F. Wilczek: "Magnetic flux, angular momentum and statistics", Phys. Rev. Lett. 48, 1144–1146 (1982).

- [35] F. Wilczek: "Quantum mechanics of fractional spin particles", Phys. Rev. Lett. 49, 957–959 (1982).
- [36] T.T. Wu, C.N. Yang: "Concept of nonintegrable phase factors and global formulation of gauge fields", Phys. Rev. D12, 3845–3857 (1975).
- [37] C.N. Yang: "Magnetic monopoles, fiber bundles, and gauge fields", Annals of the New York Academy of Science **294**, 86–97 (1977).
- [38] N. Bourbaki: Groupes et algèbres de Lie, Hermann, Paris (1968).
- [39] C. Chevalley: *Theory of Lie Groups I*,Princeton University Press, Princeton, N. J. (1946).
- [40] E.B. Davies: Theory of Open Quantum Systems, Academic Press, London (1976).
- [41] P.A.M. Dirac: Lectures on Quantum Mechanics, Belfer Graduate School of Science, Yeshiva University, New York (1964).
- [42] P.A.M. Dirac: *The Principles of Quantum Mechanics*, 4th ed., Oxford University Press, Oxford (1958).
- [43] W. Greub, S. Halperin, R. Vanstone: Connections, Curvature and Cohomology, I III, Academic Press, New York (1973).
- [44] V. Guillemin, A. Pollack: Differential Topology, Prentice-Hall, Inc., Englewood Cliffs, N. J. (1974).
- [45] S. Helgason: Differential Geometry, Lie Groups, and Symmetric Spaces, Academic Press, New York (1978).
- [46] M.W. Hirsch: Differential Topology, Springer-Verlag, Berlin (1970).
- [47] A.A. Kirillov: Elements of Representation Theory, Springer-Verlag, Berlin (1984).
- [48] S. Kobayashi, K. Nomizu: Foundations of Differential Geometry I, Interscience—Wiley, New York (1963).
- [49] S. Lang: Introduction to Differentiable Manifolds, Interscience Publishers, New York (1962).
- [50] G.W. Mackey: Induced Representations and Quantum Mechanics, W.A. Benjamin, Inc., New York (1968).
- [51] K. Nomizu: Lie Groups and Differential Geometry, The Mathematical Society of Japan (1956).
- [52] W. Pauli: Wellenmechanik, Handbuch der Physik Bd. 24, Teil 1, 120 (1933).
- [53] V.S. Varadarajan: Geometry of Quantum Theory I, Van Nostrand, Princeton, N. J. (1968).
- [54] V.S. Varadarajan: Geometry of Quantum Theory II. Quantum Theory of Covariant Systems, Van Nostrand Reinhold Co., New York (1970).